



Cochrane
Library

Cochrane Database of Systematic Reviews

Ab interno supraciliary microstent surgery for open angle glaucoma (Protocol)

Sandhu A, Jayaram H, Hu K, Bunce C, Gazzard G

Sandhu A, Jayaram H, Hu K, Bunce C, Gazzard G.
Ab interno supraciliary microstent surgery for open angle glaucoma.
Cochrane Database of Systematic Reviews 2017, Issue 9. Art. No.: CD012802.
DOI: 10.1002/14651858.CD012802.

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
BACKGROUND	1
OBJECTIVES	2
METHODS	2
ACKNOWLEDGEMENTS	5
REFERENCES	5
APPENDICES	7
CONTRIBUTIONS OF AUTHORS	12
DECLARATIONS OF INTEREST	12
SOURCES OF SUPPORT	12

[Intervention Protocol]

Ab interno supraciliary microstent surgery for open angle glaucoma

Amanjeet Sandhu¹, Hari Jayaram¹, Kuang Hu¹, Catey Bunce², Gus Gazzard¹

¹Moorfields Eye Hospital NHS Foundation Trust, London, UK. ²Department of Primary Care & Public Health Sciences, Kings College London, London, UK

Contact address: Amanjeet Sandhu, Moorfields Eye Hospital NHS Foundation Trust, 162 City Road, London, EC1V 2PD, UK. amanjeet.sandhu@moorfields.nhs.uk.

Editorial group: Cochrane Eyes and Vision Group.

Publication status and date: New, published in Issue 9, 2017.

Citation: Sandhu A, Jayaram H, Hu K, Bunce C, Gazzard G. Ab interno supraciliary microstent surgery for open angle glaucoma. *Cochrane Database of Systematic Reviews* 2017, Issue 9. Art. No.: CD012802. DOI: 10.1002/14651858.CD012802.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

This is a protocol for a Cochrane Review (Intervention). The objectives are as follows:

The main objective is to assess the results at two years of supraciliary drainage devices for OAG in comparison to conventional medical, laser, surgical treatment in terms of efficacy and safety. A secondary objective will be to examine the effects of supraciliary drainage devices with concomitant phacoemulsification in comparison to supraciliary drainage devices alone.

BACKGROUND

Description of the condition

Glaucoma is a chronic progressive optic neuropathy, affecting up to 4% of people by the age of 80 years (Burr 2007). It is the leading cause of irreversible blindness, affecting 60 million people globally (Quigley 2006). This figure is expected to increase to 80 million people by 2020. Open angle glaucoma (OAG) is the commonest type, accounting for three-quarters of cases (Quigley 2006). In one large population cohort, one in six patients with OAG became bilaterally blind (Peters 2013). The only proven way to prevent vision loss is to reduce the pressure inside the eye (intraocular pressure) over the long term (AGIS 2000; CNTG Study Group 1998; Heijl 2002; Kass 2002). Approaches to reducing intraocular pressure (IOP) include medical therapy, laser treatments, and surgery. Commercially available eye-drop preparations have a short-lasting

effect, medical therapy requires eye-drops to be instilled one or more times daily for life. Adherence is very poor, even if use is monitored (Friedman 2009; Okeke 2009). Conventional surgical techniques such as trabeculectomy are associated with significant risks, with more than 40% of patients developing perioperative complications (Kirwan 2013; Lichter 2001) and reoperation being needed in 7% to 18% (Gedde 2012; Kirwan 2013). Therefore, they are often reserved for disease that is progressing despite other treatments (King 2013).

Description of the intervention

Recently, a number of minimally-invasive surgical techniques have been developed with the aim of achieving long-term reduction of IOP with a better safety profile than conventional surgery (Francis 2011). Among them is ab interno supraciliary microstent surgery, the Cypass (Alcon Laboratories, a division of Novartis,

Basel, Switzerland) and the iStent Supra (Glaukos Corporation, Laguna Hills, CA, USA) are current devices. The former is FDA approved and also CE marked in Europe. The latter is undergoing a phase 3 clinical trial with a view to obtaining FDA approval, but is CE marked in Europe.

How the intervention might work

In cases of open angle glaucoma, an increased resistance to outflow is thought to exist not only at the level of the trabecular meshwork but also within the ciliary body part of the uveoscleral pathway. With the uveoscleral pathway thought to contribute up to half of physiological aqueous outflow (Toris 1999), supraciliary microstents such as the Cypass and iStent Supra have been developed to exploit this, leading to an increase in aqueous outflow and a reduction in intraocular pressure.

Why it is important to do this review

Consultation with patients and healthcare professionals has identified a need for better treatments for glaucoma (James Lind Alliance 2013). Minimally-invasive glaucoma procedures carry the possibility of safe and effective long-term reduction of IOP, removing concerns about permanent vision loss due to nonadherence to eye-drops. A single treatment may also be more acceptable to patients than daily and indefinite self-administration of eye-drops.

The evidence base intended to support the use of supraciliary microstents in practice continues to grow. Randomised controlled clinical studies to assess the safety and efficacy of the Cypass and iStent Supra alone have recruited in excess of 1000 patients. However, what is less clear is where this evidence lies in the current landscape of existing interventional options to manage open angle glaucoma, presently including medical, laser, trabeculectomy and other minimally-invasive glaucoma procedures. Since phacoemulsification itself reduces IOP (Mansberger 2012), we will specifically examine the evidence for the efficacy of supraciliary drainage devices when combined with phacoemulsification in comparison to phacoemulsification alone.

With both the Cypass and iStent Supra devices holding a CE mark for use in Europe and the Cypass already FDA approved, the user availability of such supraciliary microstents is expected to grow, increasing the importance of a review that will critically evaluate the current evidence relating to this group of devices.

This Cochrane review will be conducted in parallel with other reviews currently undertaken by the Cochrane Eyes and Vision MIGS Consortium, which includes minimally-invasive glaucoma surgery (MIGS) techniques and devices such as the Trabectome (NeoMedix, Tustin, California) (Hu 2016), Hydrus Schlemm's canal Microstent (Ivantis Inc., Irvine, California) (Otarola 2017), endoscopic cytophotocoagulation (ECP) (Endo Optiks, Waltham, Massachusetts) (Tóth 2017), XEN Glaucoma Implant (AqueSys

Implant, Aliso Viejo, California) (King 2017) and iStent or iStent inject (Glaukos Corporation, Laguna Hills, California) (Le 2017).

OBJECTIVES

The main objective is to assess the results at two years of supraciliary drainage devices for OAG in comparison to conventional medical, laser, surgical treatment in terms of efficacy and safety. A secondary objective will be to examine the effects of supraciliary drainage devices with concomitant phacoemulsification in comparison to supraciliary drainage devices alone.

METHODS

Criteria for considering studies for this review

Types of studies

We will include randomised controlled trials (RCTs) only. We will include reports of RCTs prepared in any language irrespective of their publication status.

Types of participants

Participants will have OAG of any type, including primary and secondary OAG. Closed angle glaucoma will be excluded. As there are no universally-accepted criteria by which glaucoma may be defined, we will permit studies to use their own definitions of glaucoma (provided these are clearly stated). In addition, participants with ocular hypertension, normal tension glaucoma, or possible glaucoma (suspects for glaucoma) will be included. We will not apply any restrictions regarding location, setting, or demographic factors.

Types of interventions

The intervention will be ab interno supraciliary microstent surgery with the Cypass (Alcon Laboratories, a division of Novartis, Basel, Switzerland), iStent Supra (Glaukos Corporation, Laguna Hills, CA, USA) or other supraciliary microstents that are identified during this review.

We will compare supraciliary microstent surgery to:

1. laser treatment (selective laser trabeculoplasty or argon laser trabeculoplasty);
2. other MIGS techniques;
3. conventional glaucoma surgery (trabeculectomy)
4. medical therapy; or

5. in combination with phacoemulsification compared with phacoemulsification alone (since phacoemulsification cataract surgery is known to reduce IOP (Mansberger 2012)).

Types of outcome measures

We will not use the reporting of particular outcomes as a criterion for eligibility for review. We will not exclude studies from review solely on the grounds of an outcome of interest not being reported.

Primary outcomes

The primary outcome will be the proportion of participants who are drop-free (not using eye-drops) at two years after randomisation.

Several different glaucoma outcome measures have been specified as primary outcomes in other Cochrane Reviews and protocols (Ismail 2015). A recent study classified IOP, visual field, safety, and anatomic outcomes as being highly important to glaucoma experts (Ismail 2016). A panel of patients from the Patient and Public Involvement Group of the National Institute for Health Research (NIHR) Biomedical Research Centre for Ophthalmology identified drop-free disease control as a highly valued outcome (unpublished). We chose a participant-centred primary outcome.

Secondary outcomes

Secondary outcomes will be:

1. Mean change in IOP, measured using Goldmann applanation tonometry, from randomisation to two years.
2. The proportions of participants experiencing intra- and postoperative complications from randomisation to two-year follow-up, including but not restricted to the following:
 - Loss of visual acuity (more than two Snellen lines or more than 0.3 logMAR, according to the method of recording visual acuity; or loss of light perception).
 - Bleeding, as recorded by the investigators.
 - Endophthalmitis, as recorded by the investigators.
 - IOP spikes (postoperative rise in IOP, measured using Goldmann applanation tonometry, of more than 10 mmHg compared to the previous assessment, including during the first postoperative month).
 - Secondary surgery, as recorded by the investigators.
3. Change in health-related quality of life measure, from randomisation to two-year follow-up.

Search methods for identification of studies

Electronic searches

The Cochrane Eyes and Vision Information Specialist will search the following electronic databases for randomised controlled trials and controlled clinical trials. There will be no language or publication year restrictions.

- Cochrane Central Register of Controlled Trials (CENTRAL) (which contains the Cochrane Eyes and Vision Trials Register) in the Cochrane Library (latest issue) (Appendix 1);
- MEDLINE Ovid (1946 to present) (Appendix 2);
- Embase Ovid (1980 to present) (Appendix 3);
- ISRCTN registry (www.isrctn.com/editAdvancedSearch) (Appendix 4);
- US National Institutes of Health Ongoing Trials Register ClinicalTrials.gov (www.clinicaltrials.gov) (Appendix 5);
- World Health Organization (WHO) International Clinical Trials Registry Platform (ICTRP) (www.who.int/ictrp) (Appendix 6).

Searching other resources

We will search the reference lists of included studies for other possible studies and will contact any individuals or organisations who, we believe, may have conducted or be conducting relevant RCTs. We will also search the website of the manufacturer (Ivantis Inc., Irvine, California; www.ivantisinc.com) for any information on forthcoming trials.

Data collection and analysis

Selection of studies

Two review authors working independently will screen titles and abstracts of all articles identified by the search using web-based online review management software (Covidence 2015). If abstracts are not available, we will screen full-text articles. Two review authors will independently assess full-text reports of all potentially eligible studies. If there is disagreement regarding eligibility, a third review author will arbitrate. If any full-text reports are rejected, we will record the reasons for this.

Data extraction and management

We will extract data from reports of included studies using a data collection form, which will be developed and piloted on the first five studies included. Two review authors will work independently to extract study characteristics from reports of each study and enter the data into Review Manager 5 (RevMan 5) (Review Manager 2014). If there is disagreement, a third independent review author will arbitrate.

We will collect the following information on the characteristics of included studies (Appendix 7):

- Year of publication.
- Year of study.
- Country of study.
- Sample size.
- Participation rate.
- Method of recruitment.
- Eligibility criteria.
- Diagnostic criteria.
- Method of randomisation.
- Method of masking.
- Number of study arms.
- Types of participants.
- Types of interventions.
- Types of comparators.
- Use of phacoemulsification at the same time as the intervention.

We will collect the following data regarding outcomes (Appendix 7):

- IOP at baseline.
- IOP at follow-up.
- Number of glaucoma medications at baseline.
- Number of glaucoma medications at follow-up.
- Intraoperative complications.
- Postoperative complications or secondary surgery.
- Duration of follow-up.
- Loss to follow-up.
- Intervals at which outcomes were assessed.

Where data on included studies are missing or unclear, we will contact the individuals or organisations involved to obtain clarification. We will collect and use the most detailed numerical data available to facilitate analyses of included studies. We will attempt to obtain these data from individuals or organisations in preference to less precise methods such as extracting numeric data from graphs. If this is necessary, two independent review authors will extract the data and a third review author will arbitrate, in case of disagreement.

Assessment of risk of bias in included studies

We will use the latest version of the Cochrane 'Risk of bias' tool as described in Chapter 8 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011) to assess the risk of bias and assign judgements of this for included studies.

Measures of treatment effect

The primary outcome is the proportion of participants who are drop-free two years after randomisation. We will use a risk ratio as the treatment effect measure. In assessing this effect measure, we will report how prescribing of IOP-lowering eye-drops was determined during follow-up. We will examine whether the people

measuring IOP and those deciding upon the prescribing of IOP-lowering eye-drops were masked to treatment group. We will report mean change in IOP from randomisation to two years after randomisation. Secondary safety outcomes will be reported as risk ratios. Health-related quality of life outcomes will be reported as differences in means or risk ratios for continuous and binary data, respectively.

Unit of analysis issues

We will assess whether included studies have included one or two eyes from each subject and whether or not randomisation has been conducted at the level of the participant or the eye. There is a potential for medical treatments, such as topical beta blockers used for one eye, to influence the outcome in the other eye (Piltz 2000). Surgery to lower IOP in one eye may also affect the IOP of the fellow eye (Radcliffe 2010). Therefore, we will exclude studies that have adopted a paired design.

Dealing with missing data

We will endeavour to minimize missing outcome data by contacting individuals and organisations to obtain them. If the data are unavailable but the level of missing data in each group and reasons for missing data in each group are similar we may simply analyse available-case data if an intention-to-treat (ITT) analysis has not been performed. We will report if authors have conducted their own ITT analysis despite missing data, but we will document whether they provide any justification for the method they have used to deal with missing data and whether they have compared their ITT result with an available-case result.

Assessment of heterogeneity

We will assess the heterogeneity between trials by careful examination of the study reports, assessing forest plots and an examination of the I^2 value. We will consider I^2 values greater than 50% as indicative of substantial heterogeneity, suggestive that meta analysis might not be wise - however, consideration will be given to the consistency of the effect estimates. If all estimates are in the same direction, we might meta-analyse even where heterogeneity is evident; we will comment on the heterogeneity.

Assessment of reporting biases

We will use a funnel plot to assess the risk of publication bias if there are more than 10 trials within our review.

Data synthesis

We will undertake a meta-analysis where data appear clinically, methodologically, and statistically homogeneous. We will check that participants, interventions, comparators, and outcomes are

sufficiently similar to give a clinically meaningful result and that our I^2 result does not indicate considerable inconsistency (i.e. I^2 less than 50%). If all estimates are in the same direction, we might meta-analyse even where heterogeneity is evident but will comment on this. We will use a random-effects model unless there are fewer than three eligible studies, in which case, we will use a fixed-effect model.

Subgroup analysis and investigation of heterogeneity

We will undertake a subgroup analysis. The effect modifier to be examined will be use of phacoemulsification as a cointervention. Phacoemulsification has been shown to reduce IOP (Mansberger 2012). We will therefore analyse whether the effect of Hydrus surgery differs depending on whether phacoemulsification is used as a cointervention.

Sensitivity analysis

We will assess the impact of including studies at high risk of bias for an outcome in one or more key domains.

Summary of findings

We will prepare tables to summarise the findings of the review, including the assessment of the certainty of evidence for all outcomes using the GRADE approach (GRADEpro 2014).

We will report the following outcomes in the 'Summary of findings' table and the comparison groups described under [Types of](#)

[interventions](#): ab interno supraciliary microstent surgery compared with laser treatment, other MIGS techniques, conventional glaucoma surgery (trabeculectomy), medical therapy or in combination with phacoemulsification compared with phacoemulsification alone.

1. Proportion of participants who are drop-free (not using eye-drops) at two years follow-up.
2. Mean change in number of IOP-lowering drops taken per day from baseline to two years follow-up.
3. Mean change in IOP, measured using Goldmann applanation tonometry, from baseline to two years follow-up.
4. Health-related quality of life at two years follow-up.
5. Intraoperative complications.
6. Postoperative complications, up to two years follow-up.
7. Secondary glaucoma surgery, including laser, as recorded by the investigators of the included trials between baseline and two years follow-up.

ACKNOWLEDGEMENTS

Cochrane Eyes and Vision (CEV) will create and execute the electronic search strategies. We thank Nitin Anand and Jennifer Evans for their comments on the published protocol that forms the template for this one (Hu 2016) and Anupa Shah for assisting with the review process.

We thank the members of the MIGS Consortium for their input in this protocol.

REFERENCES

Additional references

AGIS 2000

AGIS. The Advanced Glaucoma Intervention Study (AGIS): 7. The relationship between control of intraocular pressure and visual field deterioration. *American Journal of Ophthalmology* 2000;**130**(4):429–40.

Burr 2007

Burr JM, Mowatt G, Hernandez R, Siddiqui MA, Cook J, Lourenco T, et al. The clinical effectiveness and cost-effectiveness of screening for open angle glaucoma: a systematic review and economic evaluation. *Health Technology Assessment* 2007;**11**(41):1–190.

CNTG Study Group 1998

CNTG Study Group. Comparison of glaucomatous progression between untreated patients with normal-tension glaucoma and patients with therapeutically reduced intraocular pressures. *American Journal of Ophthalmology* 1998;**126**(4):487–97.

Covidence 2015 [Computer program]

Veritas Health Innovation. Covidence. Melbourne: Veritas Health Innovation, 2015.

Francis 2011

Francis BA, Singh K, Lin SC, Hodapp E, Jampel HD, Samples JR, et al. Novel glaucoma procedures: a report by the American Academy of Ophthalmology. *Ophthalmology* 2011;**118**(7):1466–80.

Friedman 2009

Friedman DS, Okeke CO, Jampel HD, Ying GS, Plyler RJ, Jiang Y, et al. Risk factors for poor adherence to eyedrops in electronically monitored patients with glaucoma. *Ophthalmology* 2009;**116**(6):1097–105.

Gedde 2012

Gedde SJ, Herndon LW, Brandt JD, Budenz DL, Feuer WJ, Schiffman JC, et al. Postoperative complications in the Tube Versus Trabeculectomy (TVT) study during five years of follow-up. *American Journal of Ophthalmology* 2012;**153**(5):804–14.

Glanville 2006

Glanville JM, Lefebvre C, Miles JN, Camosso-Stefinovic J. How to identify randomized controlled trials in MEDLINE: ten years on. *Journal of the Medical Library Association* 2006; **94**(2):130–6.

GRADEpro 2014 [Computer program]

GRADE Working Group, McMaster University. GRADEpro GDT. Version accessed prior to 13 September 2017. Hamilton (ON): GRADE Working Group, McMaster University, 2014.

Heijl 2002

Heijl A, Leske MC, Bengtsson B, Hyman L, Bengtsson B, Hussein M, et al. Reduction of intraocular pressure and glaucoma progression: results from the Early Manifest Glaucoma Trial. *Archives of Ophthalmology* 2002; **120**(10): 1268–79.

Higgins 2011

Higgins JP, Altman DG, Sterne JAC, editor(s). Chapter 8: Assessing risk of bias in included studies. In: Higgins JP, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from handbook.cochrane.org.

Hu 2016

Hu K, Gazzard G, Bunce C, Wormald R. Ab interno trabecular bypass surgery with Trabectome for open angle glaucoma. *Cochrane Database of Systematic Reviews* 2016, Issue 8. [DOI: 10.1002/14651858.CD011693.pub2]

Ismail 2015

Ismail R, Azuara-Blanco A, Ramsay CR. Outcome measures in glaucoma: a systematic review of Cochrane reviews and protocols. *Journal of Glaucoma* 2015; **24**(7):533–8.

Ismail 2016

Ismail R, Azuara-Blanco A, Ramsay CR. Consensus on outcome measures for glaucoma effectiveness trials: results from a delphi and nominal group technique approaches. *Journal of Glaucoma* 2016; **25**(6):539–46.

James Lind Alliance 2013

James Lind Alliance Sight Loss, Vision Priority Setting Partnership. Setting priorities for eye research. www.sightlosspsp.org.uk (accessed prior to 13 September 2017).

Kass 2002

Kass MA, Heuer DK, Higginbotham EJ, Johnson CA, Keltner JL, Miller JP, et al. The Ocular Hypertension Treatment Study: a randomized trial determines that topical ocular hypotensive medication delays or prevents the onset of primary open-angle glaucoma. *Archives of Ophthalmology* 2002; **120**(6):701–13.

King 2013

King A, Azuara-Blanco A, Tuulonen A. Glaucoma. *BMJ* 2013; **346**:f3518.

King 2017

King AJ, Hu K, Nikita E, Mulvaney CA, Azuara-Blanco A, Stead R. Subconjunctival draining minimally-invasive glaucoma devices for medically uncontrolled glaucoma.

Cochrane Database of Systematic Reviews 2017, Issue 8. [DOI: 10.1002/14651858.CD012742]

Kirwan 2013

Kirwan JF, Lockwood AJ, Shah P, Macleod A, Broadway DC, King AJ, et al. Trabeculectomy in the 21st century: a multicenter analysis. *Ophthalmology* 2013; **120**(12):2532–9.

Le 2017

Le JT, Bicket AK, Li T. Ab interno trabecular bypass surgery with iStent for open angle glaucoma. *Cochrane Database of Systematic Reviews* 2017, Issue 8. [DOI: 10.1002/14651858.CD012743]

Lichter 2001

Lichter PR, Musch DC, Gillespie BW, Guire KE, Janz NK, Wren PA, et al. Interim clinical outcomes in the Collaborative Initial Glaucoma Treatment Study comparing initial treatment randomized to medications or surgery. *Ophthalmology* 2001; **108**(11):1943–53.

Mansberger 2012

Mansberger SL, Gordon MO, Jampel H, Bhorade A, Brandt JD, Wilson B, et al. Reduction in intraocular pressure after cataract extraction: the Ocular Hypertension Treatment Study. *Ophthalmology* 2012; **119**(9):1826–31.

Okeke 2009

Okeke CO, Quigley HA, Jampel HD, Ying GS, Plyler RJ, Jiang Y, et al. Adherence with topical glaucoma medication monitored electronically. The Travatan Dosing Aid study. *Ophthalmology* 2009; **116**(2):191–9.

Otarola 2017

Otarola F, Hu K, Gazzard G, Bunce C. Ab interno trabecular bypass surgery with Schlemm's Canal Microstent (Hydrus) for open angle glaucoma. *Cochrane Database of Systematic Reviews* 2017, Issue 8. [DOI: 10.1002/14651858.CD012740]

Peters 2013

Peters D, Bengtsson B, Heijl A. Lifetime risk of blindness in open-angle glaucoma. *American Journal of Ophthalmology* 2013; **156**(4):724–30.

Piltz 2000

Piltz J, Gross R, Shin DH, Beiser JA, Dorr DA, Kass MA, et al. Contralateral effect of topical beta-adrenergic antagonists in initial one-eyed trials in the ocular hypertension treatment study. *American Journal of Ophthalmology* 2000; **130**(4):441–53.

Quigley 2006

Quigley HA, Broman AT. The number of people with glaucoma worldwide in 2010 and 2020. *British Journal of Ophthalmology* 2006; **90**(3):262–7.

Radcliffe 2010

Radcliffe NM, Musch DC, Niziol LM, Liebmann JM, Ritch R, Collaborative Initial Glaucoma Treatment Study Group. The effect of trabeculectomy on intraocular pressure of the untreated fellow eye in the collaborative initial glaucoma treatment study. *Ophthalmology* 2010; **117**(11):2055–60.

Review Manager 2014 [Computer program]

Nordic Cochrane Centre, The Cochrane Collaboration.
Review Manager 5 (RevMan 5). Version 5.3. Copenhagen:
Nordic Cochrane Centre, The Cochrane Collaboration,
2014.

Toris 1999

Toris CB, Yablonski ME, Wang YL, Camras CB. Aqueous
humor dynamics in the aging human eye. *American Journal
of Ophthalmology* 1999;**127**(4):407–12.

Tóth 2017

Tóth M, Hu K, Bunce C, Gazzard G. Endoscopic
cyclophotocoagulation (ECP) for open angle glaucoma
and primary angle closure. *Cochrane Database of
Systematic Reviews* 2017, Issue 8. [DOI: 10.1002/
14651858.CD012741

* Indicates the major publication for the study

APPENDICES

Appendix I. CENTRAL search strategy

- #1 MeSH descriptor: [Glaucoma, Open-Angle] explode all trees
- #2 MeSH descriptor: [Intraocular Pressure] explode all trees
- #3 MeSH descriptor: [Ocular Hypertension] explode all trees
- #4 OAG or POAG or IOP or OHT
- #5 simple near/3 glaucoma*
- #6 open near/2 angle near/2 glaucoma*
- #7 chronic near/2 glaucoma*
- #8 secondary near/2 glaucoma*
- #9 low near/2 tension near/2 glaucoma*
- #10 ow near/2 pressure near/2 glaucoma*
- #11 normal near/2 tension near/2 glaucoma*
- #12 normal near/2 pressure near/2 glaucoma*
- #13 pigment near/2 glaucoma*
- #14 MeSH descriptor: [Exfoliation Syndrome] this term only
- #15 exfoliat* near/2 syndrome*
- #16 exfoliat* near/2 glaucoma*
- #17 pseudoexfoliat* near/2 syndrome*
- #18 pseudoexfoliat* near/2 glaucoma*
- #19 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14 or #15 or #16 or #17 or #18
- #20 MeSH descriptor: [Stents] explode all trees
- #21 (micro-bypass* or microbypass* or micro* or bypass*) near/2 stent*
- #22 bypass near/3 (trabecul* or interno)
- #23 (supraciliary or suprachoroidal) near/3 (microstent* or micro stent* or implant* or drainage or device*)
- #24 (Gold Micro Shunt or SOLX Gold Shunt or iStent Supra or Cypass or Aquashunt or STARflo or Esnooper)
- #25 #20 or #21 or #22 or #23 or #24
- #26 #19 and #25

Appendix 2. MEDLINE Ovid search strategy

1. randomized controlled trial.pt.
2. (randomized or randomised).ab,ti.
3. placebo.ab,ti.
4. dt.fs.
5. randomly.ab,ti.
6. trial.ab,ti.
7. groups.ab,ti.
8. or/1-7
9. exp animals/
10. exp humans/
11. 9 not (9 and 10)
12. 8 not 11
13. exp glaucoma open angle/
14. exp intraocular pressure/
15. ocular hypertension/
16. (OAG or POAG or IOP or OHT).tw.
17. (simple\$ adj3 glaucoma\$).tw.
18. (open adj2 angle adj2 glaucoma\$).tw.
19. (primary adj2 glaucoma\$).tw.
20. (chronic adj2 glaucoma\$).tw.
21. (secondary adj2 glaucoma\$).tw.
22. (low adj2 tension adj2 glaucoma\$).tw.
23. (low adj2 pressure adj2 glaucoma\$).tw.
24. (normal adj2 tension adj2 glaucoma\$).tw.
25. (normal adj2 pressure adj2 glaucoma\$).tw.
26. (pigment\$ adj2 glaucoma\$).tw.
27. exfoliation syndrome/
28. (exfoliat\$ adj2 syndrome\$).tw.
29. (exfoliat\$ adj2 glaucoma\$).tw.
30. (pseudoexfoliat\$ adj2 syndrome\$).tw.
31. (pseudoexfoliat\$ adj2 glaucoma\$).tw.
32. or/13-31
33. exp Stents/
34. ((micro-bypass\$ or microbypass\$ or micro\$ or bypass\$) adj2 stent\$).tw.
35. (bypass adj3 (trabecul\$ or interno)).tw.
36. ((supraciliary or suprachoroidal) adj3 (microstent\$ or micro stent\$ or implant\$ or drainage or device\$)).tw.
37. (Gold Micro Shunt or SOLX Gold Shunt or iStent Supra or Cypass or Aquashunt or STARflo or Esnoper).tw.
38. or/33-37
39. 32 and 38
40. 12 and 39

The search filter for trials at the beginning of the MEDLINE strategy is from the published paper by [Glanville 2006](#).

Appendix 3. Embase Ovid search strategy

1. exp randomized controlled trial/
2. exp randomization/
3. exp double blind procedure/
4. exp single blind procedure/
5. random\$.tw.
6. or/1-5
7. (animal or animal experiment).sh.
8. human.sh.
9. 7 and 8
10. 7 not 9
11. 6 not 10
12. exp clinical trial/
13. (clin\$ adj3 trial\$).tw.
14. ((singl\$ or doubl\$ or trebl\$ or tripl\$) adj3 (blind\$ or mask\$)).tw.
15. exp placebo/
16. placebo\$.tw.
17. random\$.tw.
18. exp experimental design/
19. exp crossover procedure/
20. exp control group/
21. exp latin square design/
22. or/12-21
23. 22 not 10
24. 23 not 11
25. exp comparative study/
26. exp evaluation/
27. exp prospective study/
28. (control\$ or prospectiv\$ or volunteer\$).tw.
29. or/25-28
30. 29 not 10
31. 30 not (11 or 23)
32. 11 or 24 or 31
33. open angle glaucoma/
34. intraocular pressure/
35. intraocular hypertension/
36. (OAG or POAG or IOP or OHT).tw.
37. (open adj2 angle adj2 glaucoma\$).tw.
38. (primary adj2 glaucoma\$).tw.
39. (chronic adj2 glaucoma\$).tw.
40. (secondary adj2 glaucoma\$).tw.
41. (low adj2 tension adj2 glaucoma\$).tw.
42. (low adj2 pressure adj2 glaucoma\$).tw.
43. (normal adj2 tension adj2 glaucoma\$).tw.
44. (normal adj2 pressure adj2 glaucoma\$).tw.
45. (pigment\$ adj2 glaucoma\$).tw.
46. exfoliation syndrome/
47. (exfoliat\$ adj2 syndrome\$).tw.
48. (exfoliat\$ adj2 glaucoma\$).tw.
49. (pseudoexfoliat\$ adj2 syndrome\$).tw.
50. (pseudoexfoliat\$ adj2 glaucoma\$).tw.
51. or/33-50

52. Stent/
 53. ((micro-bypass\$ or micro\$ or bypass\$) adj2 stent\$).tw.
 54. (bypass adj3 (trabecul\$ or interno)).tw.
 55. ((supraciliary or suprachoroidal) adj3 (microstent\$ or micro stent\$ or implant\$ or drainage or device\$)).tw.
 56. (Gold Micro Shunt or SOLX Gold Shunt or iStent Supra or Cypass or Aquashunt or STARflo or Esnoper).tw.
 57. or/52-56
 58. 51 and 57
 59. 32 and 58

Appendix 4. ISRCTN search strategy

(Supraciliary microstent OR Suprachoroidal microstent OR Cypass OR iStent Supra OR Gold Micro Shunt OR SOLX Gold Shunt OR Aquashunt OR STARflo OR Esnoper)

Appendix 5. ClinicalTrials.gov search strategy

(Supraciliary microstent OR Suprachoroidal microstent OR Cypass OR iStent Supra OR Gold Micro Shunt OR SOLX Gold Shunt OR Aquashunt OR STARflo OR Esnoper)

Appendix 6. WHO ICTRP search strategy

(Supraciliary microstent OR Suprachoroidal microstent OR Cypass OR iStent Supra OR Gold Micro Shunt OR SOLX Gold Shunt OR Aquashunt OR STARflo OR Esnoper)

Appendix 7. Data on study characteristics

Mandatory items		Optional items
Methods		
Study design	<ul style="list-style-type: none"> · Parallel group RCT <i>i.e. people randomised to treatment</i> · Within-person RCT <i>i.e. eyes randomised to treatment</i> · Cluster RCT <i>i.e. communities randomised to treatment</i> · Cross-over RCT · Other, specify 	Number of study arms Method of randomisation Exclusions after randomisation Losses to follow-up Number randomised/analysed Method of masking How were missing data handled? <i>e.g. available case analysis, imputation methods</i> Reported power calculation (Y/N), <i>if yes, sample size and power</i> Unusual study design/issues
Eyes Unit of randomisation/ unit of analysis	<ul style="list-style-type: none"> · One eye included in study, <i>specify how eye selected</i> · Two eyes included in study, both eyes received same treatment, <i>briefly specify how analysed (best/worst/average/both and</i> 	

(Continued)

	<p><i>adjusted for within person correlation/both and not adjusted for within person correlation) and specify if mixture of one eye and two eyes</i></p> <ul style="list-style-type: none"> · Two eyes included in study, eyes received different treatments, specify if correct pair-matched analysis done 	
Participants		
Country	-	Setting
Total number of participants	<p><i>This information should be collected for total study population recruited into the study. If these data are reported for the people who were followed up only, please indicate.</i></p>	Ethnic group
Number (%) of men and women		Method of recruitment
Average age and age range		Participation rate
Inclusion criteria		Equivalence of baseline characteristics (Y/N)
Exclusion criteria	-	Diagnostic criteria
Interventions		
Intervention (n =) Comparator (n =)	<ul style="list-style-type: none"> · Number of people randomised to this group · Intervention name · Comparator name · Specify whether phacoemulsification, or other intervention, performed at same time as intervention 	Comparator parameters, e.g. dosage of drugs
Outcomes		
Primary and secondary outcomes <i>as defined in study reports</i>	<ul style="list-style-type: none"> · IOP at baseline · IOP at follow-up · Number of glaucoma medications at baseline · Number of glaucoma medications at follow-up · Intraoperative complications · Postoperative complications or secondary surgery · Duration of follow-up · Loss to follow-up · Intervals at which outcomes assessed Adverse events reported (Y/N) 	Planned/actual length of follow-up
Notes		

(Continued)

Date conducted	Specify dates of recruitment of participants mm/yr to mm/yr	Full study name: <i>(if applicable)</i> Date of publication
Sources of funding	-	Reported subgroup analyses (Y/N) Were trial investigators contacted?
Declaration of interest	-	

CONTRIBUTIONS OF AUTHORS

Amanjeet Sandhu and Kuang Hu wrote the protocol. All authors reviewed and approved the protocol.

DECLARATIONS OF INTEREST

The authors are seeking funding to address the subject of this review.

Kuang Hu performs minimally-invasive glaucoma surgery. He has lectured on 'Constructing clinical trials for MIGS - the lack of evidence and what to do about it' at the Moorfields International Glaucoma Symposium 2016, sponsored by Laboratoires Thea, which is contributing an educational grant to Moorfields Eye Hospital.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- National Institute for Health Research (NIHR), UK.
- Richard Wormald, Co-ordinating Editor for Cochrane Eyes and Vision (CEV) acknowledges financial support for his CEV research sessions from the Department of Health through the award made by the National Institute for Health Research to Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology for a Specialist Biomedical Research Centre for Ophthalmology.
- This protocol was supported by the National Institute for Health Research, via Cochrane Infrastructure funding to the CEV UK editorial base.

The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the Systematic Reviews Programme, NIHR, NHS, or the Department of Health.