



1 **A comparison of two astronomical tuning approaches**  
2 **for the Oligocene-Miocene Transition from Pacific Ocean**  
3 **Site U1334 and implications for the carbon cycle**

4

5 Helen M. Beddow<sup>1</sup>, Diederik Liebrand<sup>2,\*</sup>, Douglas S. Wilson<sup>3</sup>, Frits J. Hilgen<sup>1</sup>, Appy  
6 Sluijs<sup>1</sup>, Bridget S. Wade<sup>4</sup>, Lucas J. Lourens<sup>1</sup>

7

8 <sup>1</sup>*Department of Earth Sciences, Faculty of Geosciences, Utrecht University,*  
9 *Heidelberglaan 2, 3584 CS Utrecht, The Netherlands*

10 <sup>2</sup>*Palaeoclimate.Science, 3501 AB, Utrecht (province), The Netherlands*

11 <sup>3</sup>*Department of Earth Science University of California, Santa Barbara, CA 93106-*  
12 *9630, United States*

13 <sup>4</sup>*Department of Earth Sciences, Faculty of Mathematical and Physical Sciences,*  
14 *University College London, Gower Street, London, WC1E 6BT, United Kingdom*

15 \*Corresponding author: [diederik@palaeoclimate.science](mailto:diederik@palaeoclimate.science)

16

17 **Abstract**

18 **Astronomical tuning of sediment sequences requires both unambiguous**  
19 **cycle-pattern recognition in climate proxy records and astronomical**  
20 **solutions, and independent information about the phase relationship**  
21 **between these two. Here we present two astronomically tuned age models**  
22 **for the Oligocene-Miocene Transition (OMT) from Integrated Ocean**  
23 **Drilling Program Site U1334 (equatorial Pacific Ocean) to assess the effect**  
24 **tuning approaches have on astronomically calibrated ages and the geologic**  
25 **time scale. These age models are based on different phase-assumptions**  
26 **between climate proxy records and eccentricity: the first age model is**  
27 **based on an inverse and in-phase assumption of CaCO<sub>3</sub> weight (wt%) to**  
28 **Earth's orbital eccentricity, the second age model is based on an inverse**  
29 **and in-phase assumption of benthic foraminifer stable carbon isotope**  
30 **ratios ( $\delta^{13}C$ ) to eccentricity. The phase-assumptions that underpin these**  
31 **age models represent two end-members on the range of possible tuning**  
32 **options. To independently test which tuned age model and tuning**



33 **assumptions are correct, we assign their ages to magnetostratigraphic**  
34 **reversals identified in anomaly profiles. Subsequently we compute tectonic**  
35 **plate-pair spreading rates based on the tuned ages. These alternative**  
36 **spreading rate histories indicate that the CaCO<sub>3</sub> tuned age model is most**  
37 **consistent with a conservative assumption of constant spreading rates. The**  
38 **CaCO<sub>3</sub> tuned age model thus provides robust ages and durations for**  
39 **polarity chrons C6Bn.1n–C6Cn.1r, which are not based on astronomical**  
40 **tuning in the latest iteration of the Geologic Time Scale. Furthermore, it**  
41 **provides independent evidence that the relatively large (several 10,000**  
42 **years) time lags documented in the benthic foraminiferal isotope records**  
43 **relative to orbital eccentricity, constitute a real feature of the Oligocene-**  
44 **Miocene climate system and carbon cycle. The age constraints from Site**  
45 **U1334 thus provide independent evidence that the delayed responses of**  
46 **the Oligocene-Miocene climate-cryosphere system and carbon cycle**  
47 **resulted from increased nonlinear feedbacks to astronomical forcing.**

#### 48 49 **Keywords**

50 Astronomical tuning, marine carbon cycle, Oligocene Miocene Transition, IODP  
51 Site U1334, equatorial Pacific Ocean, geologic time scale

#### 52 53 **1. Introduction**

54 Astronomically tuned age models are important in studies of Cenozoic climate  
55 change, because they shed light on cause and effect relationships between  
56 insolation forcing and the linear and nonlinear responses of Earth's climate  
57 system (e.g., [Hilgen *et al.*, 2012, Vandenberghe *et al.*, 2012; Westerhold *et al.*,  
58 submitted]). As more Cenozoic paleoclimate records are generated that use  
59 astronomical tuning as the main high-precision dating tool, it is important to  
60 understand the assumptions and limitations inherent in this age-calibration  
61 method, in particular with respect to assumptions related to phase-relationships  
62 between tuning signal and target curves. These phase assumptions have  
63 implications for (i) determining the absolute timing of events, (ii) the  
64 understanding of leads and lags in the climate system, and (iii) the exact  
65 astronomical frequencies that are present in climate proxy records after tuning.



66

67 Previously published astronomically tuned age-models for high-resolution climate  
68 records that span the Oligocene-Miocene Transition (OMT, ~23 Ma), have used  
69 different tuning signal curves for sites from different paleoceanographic settings. In  
70 addition, different tuning target curves have been applied. For example, records from  
71 Sites 926 and 929 from the Ceara Rise (equatorial Atlantic) were tuned using  
72 magnetic susceptibility and/or color reflectance records (i.e., proxies for bulk  
73 sediment carbonate content) as tuning signal curve, and used obliquity as the main  
74 tuning target curve, sometimes with weaker precession and eccentricity components  
75 added (e.g. [Pälike *et al.*, 2006a; Shackleton *et al.*, 1999, 2000; Zachos *et al.*, 2001]).  
76 In contrast, sediments from Site 1090 from the Agulhas Ridge (Atlantic sector of the  
77 Southern Ocean) and Site 1218 from the equatorial Pacific Ocean were tuned using  
78 benthic foraminiferal stable oxygen ( $\delta^{18}\text{O}$ ) and/or carbon ( $\delta^{13}\text{C}$ ) isotope records as  
79 tuning signal (e.g. [Billups *et al.*, 2004; Pälike *et al.*, 2006b]). These records used  
80 different combinations of eccentricity, obliquity and/or precession as tuning targets  
81 (ETP curves).

82

83 More recently, Oligocene-Miocene records from Ocean Drilling Program (ODP) Site  
84 1264 and Middle Miocene records from Integrated Ocean Drilling Program (IODP)  
85 Site U1335 used the Earth's eccentricity solution as the sole tuning target [Laskar *et al.*,  
86 2004], and lithological data, such as elemental estimates based on X-ray  
87 fluorescence (XRF) core scanning records, was used as the sole tuning signal  
88 [Liebrand *et al.*, 2016, Kochhann *et al.*, 2016]. The records from both these sites are  
89 characterized by a very clear expression of eccentricity, either resulting from  
90 productivity dominated cycles (at Site 1264) or dissolution dominated cycles (at Site  
91 U1335). The phase relationships between the ~110-ky cycles and 405-ky cycles (in  
92 case of Site U1335), in lithologic records and eccentricity, were straightforward to  
93 derive [Liebrand *et al.*, 2016, Kochhann *et al.*, 2016] and were in agreement with  
94 those previously derived using benthic foraminiferal  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  records (e.g.,  
95 Zachos *et al.*, 2001, Pälike *et al.*, 2006b). An additional advantage of tuning solely to  
96 eccentricity is that no phase-assumption to either northern or southern hemisphere  
97 precession forcing is needed, and variations in the long-term stability of precession



98 and obliquity due to tidal dissipation and dynamical ellipticity do not affect the  
99 astronomically tuned ages.

100

101 The different approaches to astronomical age calibration of the Oligocene-Miocene  
102 time interval has resulted in large variations in the resulting phase-estimates after  
103 tuning between ~110-ky and 405-ky cycles present in both the eccentricity solution  
104 and in lithologic and climatologic proxy records. To obtain better constraints for the  
105 true phase-relationships of the ~110-ky and 405-ky cycles between benthic  
106 foraminiferal stable isotope records and orbital eccentricity, and to better understand  
107 the implications that initial phase-assumptions for astronomical age calibration have  
108 on absolute ages across the OMT, we need independent dates that are free from tuning  
109 phase-assumptions. Previous studies have successfully used plate-pair spreading rates  
110 to independently date magnetochron reversals and used these ages as independent age  
111 control (e.g., *Hilgen et al.*, 1991, *Lourens et al.*, 2004).

112

113 Here, we present two astronomically tuned age models for previously published high-  
114 resolution benthic foraminiferal  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  records across the OMT from IODP  
115 Site U1334 (eastern equatorial Pacific Ocean) [*Beddow et al.*, 2016]. We select  
116 (estimates of) sediment  $\text{CaCO}_3$  content and benthic  $\delta^{13}\text{C}$  as tuning signals, because  
117 these data represent two end-members in terms of tuning phase assumptions [*Pälike et*  
118 *al.*, 2006, *Liebrand et al.*, 2016]. We evaluate the ramifications of these different  
119 tuning methods for (i) absolute ages of magnetochron reversals, and (ii) the lead and  
120 lags between eccentricity and lithologic/paleoclimate records, by evaluating the  
121 spreading rate histories of a suite of tectonic plate-pairs after assigning the tuned ages  
122 to the magnetostratigraphic reversals in their anomaly profiles. The constraints given  
123 by the long-term evolutions of these potential spreading-rate histories are sufficiently  
124 precise to discriminate between tuning options and phase assumptions.

125

## 126 **2. Materials and Methods**

### 127 **2.1 Site description**

128 Site U1334, located in the eastern equatorial Pacific (4794 meters below sea level  
129 (mbsl), 7°59.998'N, 131°58.408'W), was recovered during IODP Expedition 320  
130 (Fig.1). Upper Oligocene and lower Miocene sediments from Site U1334 were



131 deposited at a paleodepth of ~4200 mbsl and consist of foraminifer- and radiolaria-  
132 bearing nannofossil ooze and chalk [Pälike *et al.*, 2010, 2012]. An expanded  
133 Oligocene-Miocene section with a well-defined magnetostratigraphy was recovered  
134 [Pälike *et al.*, 2010; Channell *et al.*, 2013] (Fig. 2), and a continuous spliced record of  
135 Holes A, B and C was placed on a core composite depth scale below seafloor (CCSF-  
136 A, equivalent to meters composite depth; Fig. 2) [Westerhold *et al.*, 2012a]. Samples  
137 were taken along the splice and all results presented here follow this depth model  
138 [Beddow *et al.*, 2016].

139

## 140 **2.2 Coulometric CaCO<sub>3</sub> and magnetic susceptibility**

141 To obtain a continuous lithological proxy record, we estimate CaCO<sub>3</sub> wt% (hereafter:  
142 CaCO<sub>3</sub> content), by calibrating high-resolution shipboard magnetic susceptibility data  
143 (MS) to lower resolution discrete shipboard coulometric CaCO<sub>3</sub> measurements for  
144 Site U1334 [Pälike *et al.*, 2010]. Minimum MS (SI unit) values correspond to  
145 maximum CaCO<sub>3</sub> values. The correlation between coulometric CaCO<sub>3</sub> measurements  
146 and MS (SI unit) was calculated using a third order polynomial fit, with an  $r^2$  value of  
147 0.79 (Fig. 2), indicating that approximately 80% of the variability in the MS record is  
148 caused by changes in the bulk sediment CaCO<sub>3</sub> content. Middle Miocene CaCO<sub>3</sub>  
149 records from nearby Site U1335 show negatively skewed cycle shapes and have been  
150 interpreted as a dissolution-dominated signal [Herbert, 1994, Kochhann *et al.*, 2016].  
151 In contrast, cycle shapes in the CaCO<sub>3</sub> content record for the Oligocene-Miocene of  
152 Site U1334 are less skewed, suggesting that here CaCO<sub>3</sub> content was predominantly  
153 controlled by a combination of productivity and dissolution.

154

## 155 **2.3 Benthic stable isotope records and magnetostratigraphic age model**

156 We use the benthic foraminifer  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  records of Site U1334, which were  
157 measured on the *Oridorsalis umbonatus* and *Cibicidoides mundulus* benthic  
158 foraminifer species [Beddow *et al.*, 2016]. To construct this mixed-species record, *O.*  
159 *umbonatus* values were corrected to *C. mundulus* values based on ordinary least  
160 squares linear regression that was based on the analysis of 180 pairs of for inter-  
161 species isotope value comparison was applied and  $n$  [Beddow *et al.*, 2016]. The  
162 benthic stable isotope datasets at Site U1334 were placed on a magnetostratigraphic  
163 age model calculated by fitting a third-order polynomial through 14



164 magnetostratigraphic age-depth tie-points (Table 1 and Fig. 4). This  
165 magnetostratigraphic age model yields an initial duration of ~21.9 to 24.1 Ma for the  
166 study interval (Fig. 3) [Channell *et al.*, 2013; Beddow *et al.*, 2016].

167

#### 168 **2.4 Spectral analysis**

169 We use AnalySeries [Paillard *et al.*, 1996] to conduct spectral analyses on the benthic  
170 foraminiferal  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  and the  $\text{CaCO}_3$  datasets in the depth domain, on the  
171 magnetostratigraphic age model [Beddow *et al.*, 2016], and on both astronomically  
172 tuned age model options presented here. Prior to analysis, the data were re-sampled  
173 and trends longer than 6 m, or 600 ky, were removed using a notch-filter (frequency =  
174 0, bandwidth = 0.015). Blackman Tukey spectral analysis was used to identify  
175 dominant periodicities present within the data, which subsequently were filtered using  
176 a Gaussian filter. We applied cross-spectral analysis to identify coherency and phase  
177 relationships between the eccentricity and the  $\text{CaCO}_3$ ,  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  chronologies.  
178 These calculations were performed at 95% significance. Evolutive spectral analyses  
179 were computed using MATLAB.

180

#### 181 **2.5. Reversal ages based on plate-pair spreading rates**

182 Anomaly profiles for tectonic plate pair spreading rates were recorded [Wilson, 1993],  
183 and applied subsequently for testing astronomical age models (e.g., [Hilgen *et al.*,  
184 1991; Krijgsman *et al.*, 1999; Hüsing *et al.*, 2007]). The plate pairs that we have  
185 selected to compute reversal ages for are in order of decreasing spreading rate:  
186 Pacific-Nazca, Pacific-Juan de Fuca, Australia-Antarctic, and Pacific-Antarctic. When  
187 multiple plate pairs show simultaneous changes in spreading rate with the same ratio,  
188 e.g., all are faster by say 15% in a short time interval, this indicates errors in the  
189 astronomical timescale. Data for the Pacific-Nazca pair is limited to the northern part  
190 of the system, which is well surveyed from studies of the separation of the Cocos  
191 plate from the northern Nazca plate during chron C6Bn [Lonsdale, 2005;  
192 Barckhausen *et al.*, 2008]. Pacific-Juan de Fuca data are from immediately north of  
193 the Mendocino fracture zone. Reversal ages based on these spreading rates are also  
194 used in previous timescale calibrations [e.g. Cande and Kent, 1992] despite the fact  
195 that for the Oligocene-Miocene only the Pacific-plate record survives. Wilson [1988]  
196 interpreted a sudden change of spreading-rate gradient for this pair from south faster



197 prior to C6Cn.2n(o) to north faster after that reversal. The dataset for the Australia-  
198 Antarctic pair is similar to that presented by *Cande and Stock* [2004]. It is expanded  
199 from that used by *Lourens et al.* [2004] who assigned reversal ages for 18.52–23.03  
200 Ma based on a spreading rate of 69.9 mm/yr for this plate pair. Data for Pacific-  
201 Antarctic come primarily from recent surveys near the Menard and Vacquier fracture  
202 zones [*Croon et al.*, 2008].

203

### 204 3. Results

#### 205 3.1. Lithologic and paleoclimatic records

206 The synthetic wt% calcium carbonate record ( $\text{CaCO}_3$  est. wt%) ranges between 54%  
207 and 88%, consistent with the  $\text{CaCO}_3$  wt% measurements on discrete samples (Figs. 2,  
208 3). Values decrease to below 70% in the upper Oligocene, between 114.9 and 116.2 m  
209 CCSF-A (Fig. 3). From 116.2 m to 121.9 m CCSF-A, the  $\text{CaCO}_3$  est. wt% varies  
210 between 61 and 83%. Variability is generally twice as large in the lower Miocene  
211 section of the record, between 88.95 and ~102 m CCSF-A, varying by ~40% with  
212 several minima in the record dipping below 70%. There is little variability across the  
213 OMT between ~102 and ~106 m CCSF-A. The benthic oxygen isotope record  
214 captures the large shift towards positive  $\delta^{18}\text{O}$  values at the Oligocene-Miocene  
215 boundary, with peak positive values (2.43‰) occurring at 104.5 CCSF-A (23.03 Ma).  
216 After the boundary, both  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  values show higher amplitude variability, and  
217 a shift towards more positive values [*Beddow et al.*, 2016].

218

#### 219 3.2. Spectral Analysis in the depth domain

220 The power spectra of the  $\text{CaCO}_3$  content record in the depth domain reveal strong  
221 spectral peaks at frequencies of 0.2 cycles/m and 0.65 cycles/m (Fig. 3). These  
222 frequencies broadly correspond to those found in the benthic  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  depth  
223 series at 0.15 cycles/m and 0.65 cycles/m [*Beddow et al.*, 2016]. Smaller spectral  
224 peaks are present in the  $\text{CaCO}_3$  content record at 1.83 cycles/m and 2.8 cycles/m (Fig.  
225 3). High-amplitude cycles with low frequencies are present in all datasets with a 1:4  
226 ratio, suggesting a strong influence from eccentricity forcing (i.e. ~110:405 ky  
227 cycles). This interpretation of strong eccentricity is supported by the application of the  
228 initial magnetostratigraphic age model [*Beddow et al.*, 2016].

229

#### 230 4. Astronomical tunings of Site U1334



#### 231 **4.1 Initial age model**

232 As a starting point for astronomical tuning we use an initial magnetostratigraphic age  
233 model [Beddow *et al.*, 2016; Channel *et al.*, 2013], which is based on the chron  
234 reversal ages of the 2012 Geologic Time Scale (GTS2012) [Vandenbergh *et al.*,  
235 2012; Hilgen *et al.*, 2012]. On this initial age model, evolutive and power spectra  
236 demonstrate that the CaCO<sub>3</sub> content and benthic foraminiferal  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  records  
237 are dominated by ~110 ky and 405 ky eccentricity paced cycles, with short intervals  
238 of significant responses at higher frequencies (Fig. 5). To further assess the influence  
239 of eccentricity on the records from Site U1334, we filter the ~110-ky and 405-ky  
240 cycles of the CaCO<sub>3</sub> est. (%) and  $\delta^{13}\text{C}$  records (Figs. 6a and 7a). In total, we observe  
241 just over five 405-ky cycles in both the filtered CaCO<sub>3</sub> content and  $\delta^{13}\text{C}$  records.  
242 There is a notable difference in the number of filtered ~110-ky cycles present between  
243 these two datasets. We observe twenty-three ~110-ky cycles in the CaCO<sub>3</sub> content  
244 record, and twenty-one in the  $\delta^{13}\text{C}$  record. This is not surprising as the exact number  
245 is often very sensitive to the width of the band-pass filter. Visual assessment of the  
246 number of cycles is not always straightforward, because not every ~110-ky cycle is  
247 expressed equally strong in all data records. In the eccentricity solution for the  
248 interval approximately between 21.9 and 24.1 Ma, we count five and a half 405-ky  
249 cycles and twenty-two ~110-ky cycles. These numbers are largely in agreement with  
250 those obtained from visual assessment and Gaussian filtering.

251

#### 252 **4.2 Astronomical target curve**

253 For our astronomical target curve, we select Earth's orbital eccentricity. Time-series  
254 analyses on the CaCO<sub>3</sub> content, and the benthic  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  records in the depth  
255 domain, and on the initial age model, indicate that eccentricity is the dominant cycle  
256 and that higher-frequency cycles are intermittently expressed (Fig. 7). Additional  
257 reasons to select eccentricity as the sole tuning target for the OMT of Site U1334 are  
258 the uncertain phase relationships of the data records to precession, and the unknown  
259 evolution of tidal dissipation and dynamical ellipticity before 10 Ma [Zeeden *et al.*,  
260 2014]. These parameters affect the long-term stability of both the precession and  
261 obliquity solutions [Lourens *et al.*, 2004; Husing *et al.*, 2007]. We use the most recent  
262 nominal eccentricity solution (i.e., La2011\_ecc3L) [Laskar *et al.*, 2011a, 2001b;  
263 Westerhold *et al.*, 2012b] as tuning target, and for the OMT interval this solution is  
264 not significantly different from the La2004 eccentricity solution [Laskar *et al.*, 2004],





265 which was used to generate previous astronomically tuned high-resolution age models  
266 for this time interval [Pälike *et al.*, 2006a,b].

267

### 268 **4.3. Astronomical age calibration of the OMT from Site U1334**

269 To test different phase-assumptions between the data from Site U1334 and  
270 eccentricity, we first consider the CaCO<sub>3</sub> content record and then the benthic  $\delta^{13}\text{C}$   
271 record as tuning signals. Both tuning options are underpinned by assumptions of a  
272 consistent and linear in-phase relationship between the tuning signal and the target,  
273 eccentricity. Previously tuned climate records for the OMT have shown that these two  
274 datasets represent end-members with respect to phase assumptions, with CaCO<sub>3</sub>  
275 content showing no lag or the smallest lag, and  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  showing increasingly  
276 larger lags to the ~110-ky and 405-ky eccentricity cycles [Liebrand *et al.*, 2016,  
277 Pälike *et al.*, 2006a, Pälike *et al.*, 2006b]. By selecting the CaCO<sub>3</sub> content record and  
278 the benthic  $\delta^{13}\text{C}$  chronology, we span the full range of tuned ages that different phase-  
279 assumptions between eccentricity and proxy data could imply.

280

#### 281 ***4.3.1. Astronomical tuning using the CaCO<sub>3</sub> content record***

282 We use the initial magnetostratigraphic age model as a starting point for a more  
283 detailed calibration of maxima in CaCO<sub>3</sub> content to ~110-ky eccentricity minima.  
284 CaCO<sub>3</sub> maxima generally correspond to positive  $\delta^{18}\text{O}$  values (i.e. cooler, glacial  
285 periods), which are usually linked to eccentricity minima and are therefore  
286 anticorrelated with eccentricity [Zachos *et al.*, 2001; Pälike *et al.*, 2006a; Pälike *et al.*,  
287 2006b]. The CaCO<sub>3</sub> content record has 23 clearly delineated ~110 ky maxima, which  
288 we match directly to minima in the La2011 eccentricity time series (Fig. 6c). In  
289 addition to these well expressed ~110-ky cycles, we take the expression of the 405-ky  
290 cycle into account to establish the tuned age model. The data records from Site U1334  
291 span the interval between 21.96 and 24.15 Ma (2.19 My duration) on the CaCO<sub>3</sub>  
292 tuned age model. Linear sedimentation rates (LRS) vary between 0.9 and 2.2 cm/ky,  
293 with relatively higher sedimentation rates across the OMT (Fig. 6). On average this  
294 yields a sample resolution of 3.6 ky for the benthic isotope records.

295

296 Evolutive analyses of the benthic  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  records on the CaCO<sub>3</sub> tuned age  
297 model indicate that the 405-ky cycle is best expressed. In contrast, the CaCO<sub>3</sub> content  
298 record on this age model reveals that the ~110-ky cycle has the highest amplitudes.



299 Despite the overall clear expression of the 405-ky cycle in the CaCO<sub>3</sub> evolutive  
300 spectrum, this signal is more subdued across the OMT (Fig. 5). Spectral power at the  
301 ~110-ky periodicity increases in all three records in the interval following peak  
302 glacial conditions associated with the OMT. This cycle is particularly pronounced in  
303 the δ<sup>18</sup>O record, and we can identify power at both the 125 ky and the 95 ky  
304 eccentricity cycles in both the CaCO<sub>3</sub> and δ<sup>18</sup>O datasets. We note that this could be a  
305 direct result from using eccentricity as a tuning target. For δ<sup>13</sup>C, the evolutive analysis  
306 and power spectra indicate that ~110 ky cycle is more strongly expressed at the 125-  
307 ky periodicity, compared to the 95-ky component. We find intermittent power present  
308 at a periodicity of ~50 ky/cycle, which is either related to the obliquity cycle that is  
309 offset towards a slightly longer periodicity, or to the first harmonic of the ~110-ky  
310 eccentricity cycle [King, 1996]. The ~50-ky cycle is best expressed in the benthic  
311 δ<sup>18</sup>O record on the CaCO<sub>3</sub> tuned age model, where we identify two main intervals  
312 with significant power at this periodicity, one between ~23.5 and ~23.8 Ma, and the  
313 other between ~22.4 and ~22.6 Ma (Fig. 5).

314

315 Cross-spectral analyses between the data records on the CaCO<sub>3</sub> tuned age model and  
316 eccentricity, indicate that CaCO<sub>3</sub> content, δ<sup>18</sup>O and δ<sup>13</sup>C are significantly coherent  
317 (99%) with eccentricity at the 405-ky, 125-ky and 95-ky eccentricity cycles (Fig. 5).  
318 Phase estimates of benthic δ<sup>18</sup>O with respect to eccentricity indicates a lag of 20–35  
319 ky at the 405 ky period, and 2–18 ky at the ~110 ky periodicity. The δ<sup>13</sup>C record lags  
320 eccentricity by 19–38 ky at the 405-ky cycle, by 5–8 ky at the 125-ky cycle and by 8–  
321 10 ky at the 95-ky cycle (Fig. 5). CaCO<sub>3</sub> is roughly in-phase with eccentricity by 0-7  
322 ky at the 405 ky cycle, 125-ky cycle and 95-ky cycle, which is not surprising, because  
323 it was used to obtain astronomically tuned ages. These phase relationships between  
324 CaCO<sub>3</sub> and eccentricity thus confirm that the in-phase tuning assumption was applied  
325 successfully.

326

#### 327 ***4.3.2. Astronomical tuning using the benthic δ<sup>13</sup>C record***

328 An important consequence of the CaCO<sub>3</sub> tuned age model is that eccentricity-related  
329 variability within the benthic foraminiferal δ<sup>13</sup>C record is not in-phase with  
330 eccentricity (Fig. 7b). On both the initial magnetostratigraphic age model and on the  
331 CaCO<sub>3</sub> tuned age model, the phase-lag, as identified in the filtered records, between  
332 the 405-ky-eccentricity cycle and the 405-ky cycle in δ<sup>13</sup>C increases during the Early



333 Miocene (Figs. 6 and 7). The 405-ky eccentricity pacing of  $\delta^{13}\text{C}$  is a consistent  
334 feature that characterizes the Cenozoic carbon cycle [Holbourn *et al.*, 2004, 2013;  
335 Littler *et al.*, 2014; Pälike *et al.*, 2006a,b; Liebrand *et al.*, 2016]. To date, no large  
336 changes in the phase-relationship of this cycle to eccentricity have been documented.  
337 An increased phase lag in the response of the 405-ky cycle to eccentricity, as is  
338 suggested by the  $\text{CaCO}_3$  tuned age model, could provide further support for a large-  
339 scale reorganization of the carbon cycle across the OMT as has previously been  
340 suggested based on proxy studies [Diester-Haas *et al.*, 2011, Mawbey and Lear,  
341 2013]. Alternatively, the phase-lag of the 405-ky cycle in benthic  $\delta^{13}\text{C}$  to eccentricity  
342 remains relatively small, which would indicate that the tuning assumptions  
343 underpinning the  $\text{CaCO}_3$  tuned age model are flawed.

344  
345 To distinguish between these two contrasting hypotheses, we generate another  
346 astronomically tuned age model. This time, we select the benthic  $\delta^{13}\text{C}$  record as the  
347 tuning signal and assume that the 405-kyr and  $\sim 110$ -ky cycles in benthic  $\delta^{13}\text{C}$  are  
348 continuously in phase with eccentricity across the OMT (Fig. 7d). Approximately five  
349 405-ky cycles are identified in the benthic  $\delta^{13}\text{C}$  record, which facilitate initial visual  
350 alignment to the same cycle in the eccentricity solution. Subsequently, we correlated  
351 the maxima and minima in the of the benthic  $\delta^{13}\text{C}$  record, as identified in Gaussian  
352 filters of this data on the initial magnetostratigraphic age model (Fig. 7a), to those  
353 identified in the filtered component of the eccentricity solution (Fig. 7d).

354  
355 The data records, on the benthic  $\delta^{13}\text{C}$  tuned age model, span the interval between 22.1  
356 and 24.2 Ma (i.e., 2.1 My duration), resulting in an average time step of 3.4 ky for the  
357 benthic stable isotope records. LRS range from 0.7–3.3 cm/ky, with an abrupt and  
358 short-lived increase across the OMT to  $\sim 1.7$  cm/ky. On the  $\delta^{13}\text{C}$  tuned age model, the  
359  $\text{CaCO}_3$  record remains in anti-phase with respect to  $\sim 110$ -ky eccentricity, but the  
360 benthic  $\delta^{13}\text{C}$  tuning results in an alternative alignment  $\text{CaCO}_3$  maxima to eccentricity  
361 minima that result in a  $\sim 110$ -ky shorter duration of the data records (Fig. 6 and 7).  
362 The evolutive analyses and power spectra are broadly consistent with the evolutive  
363 analyses from the  $\text{CaCO}_3$  tuned age model, with dominant 405-ky cyclicity in all three  
364 datasets, an increase in spectral power at  $\sim 110$ -ky eccentricity cycles after the OMT  
365 and intermittent expression of higher frequency astronomical cycles. On the  $\delta^{13}\text{C}$



366 tuned age model, all datasets exhibit a more significant response at the 95-ky short  
367 eccentricity cycle than the 125-ky short eccentricity cycle, in contrast to the CaCO<sub>3</sub>  
368 tuned age model. Significant power at the 41-ky obliquity periodicity is present in the  
369 late Oligocene, between ~ 23.3 and 23.8 Ma.

370

371 Cross-spectral analyses between data records on the  $\delta^{13}\text{C}$  tuned age model and  
372 eccentricity (Fig. 5) indicate that CaCO<sub>3</sub>,  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  are significantly coherent  
373 (99%) with eccentricity at the 405-, 125- and 95-ky eccentricity cycles. Phase  
374 estimates of  $\delta^{18}\text{O}$  with respect to eccentricity (Fig. 5) shows lags of 1–9 ky at the 405-  
375 ky period and of 1–10 ky at the ~125 ky cycle. Benthic  $\delta^{13}\text{C}$  lags eccentricity by 1–8  
376 ky at the 405-ky periodicity and by 2–10 ky at the ~125-ky eccentricity cycle. At the  
377 95-ky eccentricity cycle,  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  lead eccentricity by 1–9 ky. CaCO<sub>3</sub> leads  
378 eccentricity by 15–40 ky at the 405-ky cycle, by 0–14 ky at the ~125-ky cycle, and by  
379 1–13 ky at the ~95-ky cycle.

380

### 381 **5. Spreading rates**

382 To independently test whether the CaCO<sub>3</sub> tuned ages or the benthic  $\delta^{13}\text{C}$  tuned ages  
383 and their underlying phase-assumption, are most appropriate for tuning the deep  
384 marine Oligocene-Miocene records from Site U1334, we use independent ages based  
385 on plate pair spreading rates as a control age. When multiple plate pairs show  
386 simultaneous changes in spreading rate with the same ratio, e.g., all are faster by say  
387 15% in a short time interval, this indicates errors in the timescale. We propose to use  
388 the age model that passes this test most successfully to provide ages for C6Bn.1n (o)  
389 to C6Cn.1r (o) and potentially revise those currently presented in the GTS2012.

390

391 Of the two astronomically tuned age models and GTS2012, the CaCO<sub>3</sub> tuned age  
392 model is most consistent with the least amount of changes in plate-pair spreading  
393 rates (Fig. 8). This suggests that a lithologic proxy record is the most suitable signal  
394 curve for Oligocene-Miocene records from the equatorial Pacific. It may also provide  
395 support for similar astronomical age calibration approaches that have been used for  
396 Middle Miocene [Kochhann *et al.*, 2016] and Eocene-Oligocene [Westerhold *et al.*,  
397 2015] records from the equatorial Pacific Ocean, and for Oligocene-Miocene records  
398 from the South Atlantic Ocean [Liebrand *et al.*, 2016]. Although these studies also



399 used CaCO<sub>3</sub>-controlled lithological proxy records for tuning to eccentricity, we note  
400 that these records show variable amounts of productivity versus dissolution as the  
401 main source of variance in the data.

402

403 On the CaCO<sub>3</sub> tuned age model, the Australia-Antarctica, Pacific-Nazca, and Pacific-  
404 Antarctic plate pairs are all very close to a constant spreading rate, at least prior to  
405 Chron C6Bn. The Juan de Fuca-Pacific plate-pair indicates a sudden decrease in  
406 spreading rate (145 to 105 mm/yr) at ~23 Ma, consistent with expectations [Wilson,  
407 1988]. The implied synchronous changes for the Australia-Antarctica, Pacific-Nazca,  
408 and Pacific-Antarctic plate pairs in the δ<sup>13</sup>C tuned age model, especially the faster  
409 spreading rates ~22.5-23.0 Ma implied by older ages for C6Bn, make this option less  
410 plausible. Differences between the CaCO<sub>3</sub> tuned age model for Site U1334 and  
411 GTS2012 are subtler. The longer duration of C6Cn.3n in the CaCO<sub>3</sub> tuned age model  
412 (106 vs. 62 kyr) eliminates a brief pulse of fast spreading implied by GTS2012,  
413 visible in Figure 8a as positive slopes in age-distance during that chron. Over longer  
414 intervals, CaCO<sub>3</sub> tuned ages remove a slight but synchronous rate slowdown that is  
415 also implied by GTS2012 and which starts at ~23.2 Ma.

416

417 The spreading rates computed using the CaCO<sub>3</sub> tuned age model suggest a duration  
418 for C6Cn.2n of 67 ky. This duration may be up to ~40 ky too short, as is suggested by  
419 the implied fast spreading during this chron (see the positive slopes in Figure 8b).  
420 Although our distance error bars indicate that this discrepancy is only marginally  
421 significant, it provides further support for an age of ~23.06 Ma for the Oligocene-  
422 Miocene boundary, broadly in agreement with independently tuned ages from Site  
423 1264 [Liebrand *et al.*, 2016]. This could indicate an uncertainty in the  
424 magnetostratigraphy at Site U1334, although this is unlikely as the C6Cn.2n reversal  
425 is clearly delineated in the Virtual Geomagnetic Pole (VGP) latitude signal [Channell  
426 *et al.*, 2013]. In both the CaCO<sub>3</sub> content and δ<sup>13</sup>C record, this short interval is difficult  
427 to align to the tuning target (Figs. 5 and 6), because CaCO<sub>3</sub> content values are high,  
428 with little variability and benthic δ<sup>13</sup>C values corresponds to the marked shift towards  
429 higher values at the Oligocene-Miocene carbon maximum [Hodell and Woodruff,  
430 1994]. The 83 kyr duration of C6Cn.2n from the δ<sup>13</sup>C tuned age model is somewhat  
431 more consistent with spreading rates than the 67 kyr duration from the CaCO<sub>3</sub> tuned  
432 age model, and the 118 kyr duration in GTS2012 is even more consistent. If there is a



433 problem with the tuning in both records for this chron, using constant spreading rates  
434 to interpolate from the adjacent CaCO<sub>3</sub> ages would imply reversal ages for the top and  
435 bottom of C6Cn.2n of ~22.95 and ~23.06 Ma. On significant difference the CaCO<sub>3</sub>  
436 tuned ages suggest is that the increase in spreading rates of the Juan de Fuca-Pacific  
437 plate-pair occurred approximately 200 ky later than those ages presented in the  
438 GTS2012 (i.e. during Chron C6Cn.2n. instead of C6Cn.3n, respectively; see Fig 8).  
439 Overall the spreading rates suggest that the CaCO<sub>3</sub> tuned age model is the preferred  
440 age model option.

441

## 442 **6. Discussion**

### 443 **6.1. Age model evaluation**

444 The final eccentricity tuned age models for the OMT time interval differ for two  
445 reasons. Firstly, there are 21 complete 110 ky cycles in the  $\delta^{13}\text{C}$  tuned age model, and  
446 22 in the CaCO<sub>3</sub> content record, making the  $\delta^{13}\text{C}$  tuned age model ~1 eccentricity  
447 cycle shorter in duration. This is a direct result of the patterns observed in the 405 ky  
448 and ~110 ky cycles in the CaCO<sub>3</sub> and  $\delta^{13}\text{C}$  datasets on the initial magnetostratigraphic  
449 age model. The tuned age models are consistent with each other across the positive  
450  $\delta^{18}\text{O}$  isotope excursion during the OMT, with the peak positive value in the  $\delta^{18}\text{O}$   
451 record, and the base of Chron C6Cn.2n (marking the Oligocene-Miocene boundary),  
452 occurring within 10 ky on both age models. They diverge at ~22.7 Ma, where the  
453 CaCO<sub>3</sub> content has an additional ~110 ky cycle on the initial magnetostratigraphic age  
454 model. Here, either the ~110 ky response at 22.7 Ma has not been recorded in the  
455  $\delta^{13}\text{C}$  record or there is a double peak in the CaCO<sub>3</sub> content. If we assign these  
456 contrasting ages to the selection of plate pair anomaly profiles, their spreading rates  
457 histories support the CaCO<sub>3</sub> tuned ages. In the depth domain, the existence of two  
458 distinct ~110-ky minima in the  $\delta^{18}\text{O}$  record between 97.5 and 99 CCSF-A lends  
459 additional support to the CaCO<sub>3</sub> content age model.

460

### 461 **6.2. Phase relationships**

462 The second factor contributing to the difference between the age models is the  
463 different phase relations between  $\delta^{13}\text{C}$  and eccentricity and CaCO<sub>3</sub> and eccentricity,  
464 which account for up to ~30 ky difference between the ages of maxima and minima in  
465 ~110 kyr cycles in the two records. One of the assumptions of our CaCO<sub>3</sub> content  
466 tuning is that it is more likely to be in-phase with eccentricity modulation of



467 precession than the benthic  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$  stable isotope records [Pälike et al.,  
468 2006a,b; Liebrand et al., 2011]. Variations in the  $\delta^{13}\text{C}$  signal are generally considered  
469 to best reflect global ocean signals, but are thought to lag global climate by ~10% on  
470 all periodicities (Table 2) [Billups et al., 2004; Pälike et al., 2006a,b; Liebrand et al.,  
471 2016]. The  $\text{CaCO}_3$  signal, in contrast, most likely represents a more regional, ocean-  
472 basin wide response to insolation because it depends on regional carbonate  
473 productivity, dissolution and/or dilution. These processes affecting the  $\text{CaCO}_3$  content  
474 of the sediment were probably more directly responsive to insolation forcing [Hodell  
475 et al., 2001]. The longer lag time of  $\delta^{13}\text{C}$  with respect to eccentricity in comparison  
476 with  $\text{CaCO}_3$  leads to older ages assigned to ~110 kyr maxima and minima in the  $\delta^{13}\text{C}$   
477 age model. This is particularly notable between 22.7 Ma and 24.2 Ma, when the age  
478 difference between the age models is accounted for only by the difference in phase.  
479 As the spreading rates support the  $\text{CaCO}_3$  tuned ages, this implies that the long phase  
480 lag in the response of  $\delta^{13}\text{C}$  to eccentricity results in less accurate tuned ages for Site  
481 U1334. This suggests that local/regional tuning signals produce more accurate age  
482 models in comparison with globally integrated isotope records, which are known to  
483 produce significant lags relative to eccentricity as a result of non-linear feedbacks  
484 [Pälike et al., 2006b, Zeebe et al., 2017].

485

### 486 6.3. Implications for the carbon cycle

487 Benthic foraminiferal  $\delta^{13}\text{C}$  variations in the open ocean are typically interpreted to  
488 reflect the ratio between global organic and inorganic carbon burial [Shackleton,  
489 1977; Broecker, 1982; Diester-Haas et al., 2013, Mawbey and Lear, 2013].  
490 Astronomical forcing of organic carbon burial is typically expected in the  
491 precessional band because organic carbon burial, notably in the marine realm,  
492 depends on clay fluxes and thus hydrology (Berner et al., 1983). However, the  
493 residence time of carbon (~100 kyr) is so long (Broecker and Peng, 1982) that this  
494 energy is transferred into eccentricity bands (e.g., Pälike et al., 2006; Ma et al., 2011).  
495 Importantly, while the total marine carbon inventory is driven by ocean chemistry, the  
496 phase lag between eccentricity forcing and  $\delta^{13}\text{C}$  should primarily be a function of the  
497 residence time of carbon (Zeebe et al., 2017). Hypothetically, a change in total  
498 organic matter burial will only result in whole-ocean steady state when the  $\delta^{13}\text{C}$  of  
499 buried carbon equals that of the input (through rivers). Because the burial fluxes are



500 small compared to the total carbon inventory, a pronounced time lag between  
501 eccentricity forcing and  $\delta^{13}\text{C}$  is expected (e.g., *Zeebe et al.*, 2017).

502

503 Interestingly, the  $\text{CaCO}_3$  age model for Site U1334 implies that the phase lag between  
504 the 405 ky cycle in the  $\delta^{13}\text{C}$  record and the eccentricity forcing increases across the  
505 OMT. A similar shift in phase is also present in the benthic foraminiferal  $\delta^{13}\text{C}$  record  
506 from 1264 [*Liebrand et al.*, 2011; *Liebrand et al.*, 2016]. In theory (*Zeebe et al.*,  
507 2017), an increase in the phase lag suggests an increase in the residence time oceanic  
508 carbon, either through a rise in the total carbon inventory or a drop in the supply and  
509 burial of carbon. The lengthening of the phase lag of the 405 ky cycle coincides with  
510 a large shift in the benthic foraminiferal  $\delta^{13}\text{C}$  record across the OMT to more positive  
511 values, evidencing a structural relative increase in the supply of  $^{13}\text{C}$ -depleted or drop  
512 in the burial of  $^{13}\text{C}$ -enriched carbon. Reliable reconstructions of  $\text{CO}_2$  are rare across  
513 the OMT ([www.p-co2.org](http://www.p-co2.org)) and the OMT does not seem associated with a large  
514 change in the depth of the Pacific calcite compensation depth (*Pälike et al.*, 2012).  
515 Therefore, additional constraints on atmospheric  $\text{CO}_2$  concentrations and burial fluxes  
516 are required to speculate on the mechanisms associated with the increased phase lag.

517

## 518 7. Conclusions

519 We explore the application of  $\text{CaCO}_3$  content and benthic foraminiferal  $\delta^{13}\text{C}$  records  
520 as tuning signals for the OMT record at Site U1334 in the eastern equatorial Pacific.  
521 These two tunings highlight the importance of carefully considering the implications  
522 of tuning choices and assumptions when creating astronomical age models. Spreading  
523 rate histories provide independent evidence for the astronomically tuned age models,  
524 and are generally in best agreement with the  $\text{CaCO}_3$  tuned age model. This suggests  
525 that lithological signals respond more directly to insolation forcing than stable isotope  
526 signals, for which we find support for a delayed respond to astronomical climate  
527 forcing. The  $\text{CaCO}_3$  based age model thus provides a valuable method to better  
528 understand the (lagged) response in benthic foraminiferal  $\delta^{18}\text{O}$  and  $\delta^{13}\text{C}$ , which are  
529 widely used and reproducible proxies for the global climate/cryosphere system and  
530 (marine) carbon cycle. One important implication of the  $\text{CaCO}_3$  age model is that 405  
531 ky cycle in benthic  $\delta^{13}\text{C}$  shows a distinct phase lag with respect to orbital  
532 eccentricity. Lastly, the  $\text{CaCO}_3$  age model for Site U1334 provides astronomically





533 calibrated ages for C6AAr.3r to C6Cn.1r, which in GTS2012 are not presently  
534 astronomically calibrated. The polarity chron ages from the CaCO<sub>3</sub> tuned ages are  
535 generally older by approximately 60 ky on average than those presented in the  
536 GTS2012. We suggest that these updated early Miocene ages are incorporated in the  
537 next version of the GTS.

538

### 539 **Acknowledgements**

540 This research used samples provided by the Integrated Ocean Drilling Program  
541 (IODP), collected by the staff, crew and scientists of IODP Expedition 320/321. We  
542 thank Dominika Kasjanuk, Arnold van Dijk, Maxim Krasnoperov and Jan Drenth for  
543 laboratory assistance. Linda Hinnov kindly provided her evolutive analysis MATLAB  
544 script. This research was supported by PalaeoClimate.Science (DL), NWO grant  
545 865.10.001 (L.J.L), ERC grant 259627 (A.S.), NERC grant NE/G014817 (B.S.W.),  
546 and a Marie Curie Career Integration Grant “ERAS”. All data are available online  
547 ([www.pangaea.de](http://www.pangaea.de)). (NB. Data will be uploaded after acceptance for publication and  
548 DOI link will be provided).

549

### 550 **Figure Captions**

551 **Figure 1. Locations of ODP and IODP drill sites discussed in this study.** Location  
552 of IODP Site U1334 with reference to ODP Sites 1264, 1218, 926, 929 and 1090.

553

554 **Figure 2. Calibration between the shipboard Magnetic Susceptibility record and**  
555 **shipboard coulometric CaCO<sub>3</sub> measurements to obtain a record of CaCO<sub>3</sub>**  
556 **estimates (wt%). (a)** The Magnetic susceptibility and CaCO<sub>3</sub> content records. **(b)**  
557 The relationship between coulometric CaCO<sub>3</sub> measurements and discrete sample  
558 magnetic susceptibility was calculated using ordinary least squares linear regression,  
559 and yielded an  $r^2$  value of 0.79.

560

561 **Figure 3. Site U1334 datasets, evolutive spectra and power spectra against depth.**  
562 **(a)** Magnetostratigraphy for Site U1334 (*Channell et al., 2013*). **(b)** The CaCO<sub>3</sub>  
563 content record. **(c)** The benthic foraminiferal  $\delta^{18}\text{O}$  record. **(d)** The benthic  
564 foraminiferal  $\delta^{13}\text{C}$  record. **(e)** Evolutive and power spectra of the CaCO<sub>3</sub> content  
565 record. **(f)** Evolutive and power spectra of the benthic foraminiferal  $\delta^{18}\text{O}$  record. **(g)**



566 Evolutive and power spectra of the benthic foraminiferal  $\delta^{13}\text{C}$  record. All data plotted  
567 against the latest available splice (*Westerhold et al.*, 2012)

568

569 **Figure 4. Depth versus age relationships for the different age models for Site**  
570 **U1334.** Magnetochron ages are based on GTS2012 [*Vandenbergh et al.*, 2012;  
571 *Hilgen et al.*, 2012], the initial age model, the  $\text{CaCO}_3$  content age model and the  $\delta^{13}\text{C}$   
572 age model. Magnetochrons are plotted as colored circles, and the lines represent a  
573 third order polynomial fit.

574

575 **Figure 5. Implication of age models on time series analysis. (a-c)**  $\text{CaCO}_3$  on the  
576 initial,  $\text{CaCO}_3$  tuned, and the  $\delta^{13}\text{C}$  tuned age model, respectively. **(d-f)** As in (a-c) but  
577 for benthic foraminiferal  $\delta^{18}\text{O}$ . **(g-i)** As in (a-c) but for benthic foraminiferal  $\delta^{13}\text{C}$ .  
578 Prior to analysis, the  $\text{CaCO}_3$  data are resampled at a time step of 2 ky, the benthic  
579 foraminiferal data are resampled at a time step of 4 ky. For all records, periodicities  
580 larger than 600 ky are notch-filtered out. Coherence and phase estimates are between  
581 eccentricity La2011 solution and benthic isotope datasets. The significance level  
582 represented by the red line for the coherence plots is 99%. For the phase estimates  
583 between the benthic foraminiferal series and eccentricity, eccentricity was multiplied  
584 by  $-1$ .

585

586 **Figure 6. Site U1334  $\text{CaCO}_3$  versus age. (a)** The  $\text{CaCO}_3$  dataset and Gaussian filters  
587 plotted on **(a)** the magnetostratigraphic age model, **(b)** the  $\delta^{13}\text{C}$  tuned age model, and  
588 **(c)** the  $\text{CaCO}_3$  tuned age model. **(d)** Earth's orbital eccentricity solution is plotted in  
589 grey [*Laskar et al.*, 2010, *Laskar et al.*, 2011]. Tie points are represented by red dots  
590 and dashed lines. Gaussian filters were calculated in AnalySeries [*Palliard et al.*,  
591 1996] with the following settings: 405 ky  $-f: 2.5$   $bw 0.8$ ,  $\sim 110$  ky  $-f: 10$ ,  $bw: 3$ . **(e)**  
592 Sedimentation rates are calculated using the  $\text{CaCO}_3$  tuned age model.

593

594 **Figure 7. Site U1334  $\delta^{13}\text{C}$  versus.** The  $\delta^{13}\text{C}$  dataset and Gaussian filters plotted on  
595 **(a)** the magnetostratigraphic age model, **(b)** the  $\text{CaCO}_3$  tuned age model, and **(c)** the  
596  $\delta^{13}\text{C}$  tuned age model. **(d)** Earth's orbital eccentricity solution is plotted in grey  
597 [*Laskar et al.*, 2010, *Laskar et al.*, 2011]. Tie points are represented by red dots and  
598 dashed lines. Gaussian filters were calculated in AnalySeries [*Palliard et al.*, 1996]



599 with the following settings: 405 ky –  $f$ : 2.5  $bw$  0.8, ~110 ky –  $f$ : 10,  $bw$ : 3. (e)  
600 Sedimentation rates are calculated using the  $\delta^{13}\text{C}$  age model.

601

602 **Figure 8. Plate-pair spreading rates based on different age models.** Reduced-  
603 distance plots for the labeled plate pairs implied by (a) the GTS2012, (b) the  $\text{CaCO}_3$   
604 tuned age model and (c) the  $\delta^{13}\text{C}$  tuned age model. Reduced distance is the full  
605 spreading distance ( $D$ ) minus the age ( $A$ ) times the labeled spreading rate ( $R$ , see  $y$ -  
606 axes). Distance scale is plotted inversely with spreading rate so that for true constant  
607 spreading rate, age errors will cause uniform vertical departures from a straight line.  
608 Error bars are 95% confidence. The  $\text{CaCO}_3$  based age model (b) gives the simplest  
609 spreading rate history.

610

611 **Table 1. Comparison of magnetostratigraphic reversal ages.** Chron boundary ages  
612 across the Oligocene Miocene Transition from the published literature and this study.  
613 Age differences are presented on the right hand side.

614

615 **Table 2. Comparison of tuning methods and phase relationships.** List of  
616 astronomically dated Oligocene-Miocene spanning record. Tuning signal and target  
617 curves, and phase relationships to the target curves are compared.

618

## 619 References

620 Barckhausen, U., C. R. Ranero, S. C. Cande, M. Engels and W. Weinrebe (2008),  
621 Birth of an intraoceanic spreading center. *Geology*, 36(10), 767-770.

622

623 Beddow, H. M., D. Liebrand, A. Sluijs, B. S. Wade, and L. J. Lourens (2016), Global  
624 change across the Oligocene-Miocene transition: High-resolution stable isotope  
625 records from IODP Site U1334 (equatorial Pacific Ocean), *Paleoceanography*,  
626 31, doi:10.1002/2015PA002820.

627

628 Berner, R. A., A. C. Lasaga, and R. M. Garrels (1983), The carbonate-silicate  
629 geochemical cycle and its effect on atmospheric carbon dioxide over the past  
630 100 million years. *American Journal of Science*, 283, 641–683, doi:  
631 10.2475/ajs.283.7.641.



- 632
- 633 Billups, K., H. Pälike, J. E. T. Channell, J. C. Zachos, and N. J. Shackleton (2004),  
634 Astronomic calibration of the late Oligocene through early Miocene  
635 geomagnetic polarity time scale, *Earth Planet. Sci. Lett.*, 224, 33–44,  
636 doi:10.1016/j.epsl.2004.05.004.
- 637
- 638 Broecker, W. S. (1982), Glacial to interglacial changes in ocean chemistry, *Progress*  
639 *in Oceanography*, 11, 2, 151-197.
- 640
- 641 Broecker, W. S., T-H Peng (1982), Tracers in the Sea, Lamont-Doherty Geological  
642 Observatory, Columbia University.
- 643
- 644 Cande, S. C., and D. V. Kent (1992), A new geomagnetic polarity time scale for the  
645 Late Cretaceous and Cenozoic, *J. Geophys. Res.*, 97(B10), 13917-13951.
- 646
- 647 Cande, S. C., and J. M. Stock (2004), Pacific-Antarctic-Australia motion and the  
648 formation of the Macquarie plate, *J. Geophys. Int.*, 157, 399-414.
- 649
- 650 Channell, J. E. T., C. Ohneiser, Y. Yamamoto, and M.S. Kesler (2013), Oligocene-  
651 Miocene magnetic stratigraphy carried by biogenic magnetite at sites U1334  
652 and U1335 (equatorial Pacific ocean) *Geochemistry, Geophysics, Geosystems*,  
653 14(2) pp1525-2027doi:10.1029/2012GC004429.
- 654
- 655 Croon, M. B., S. C. Cande, and J. M. Stock (2008), Revised Pacific-Antarctic plate  
656 motions and geophysics of the Menard Fracture Zone: *Geochemistry,*  
657 *Geophysics, Geosystems*, v. 9, Q07001, doi:10.1029/2008GC002019.
- 658
- 659 DeConto, R.M., S. Galeotti, M. Pagani, D. Tracy, K. Schaefer, T. Zhang, D. Pollard,  
660 and J.D. Beerling, (2012). Past extreme warming events linked to massive  
661 carbon release from thawing permafrost. *Nature*, 484(7392), p.87.
- 662
- 663 Diester-Haass, L., K. Billups, and K. Emeis (2011), Enhanced paleoproductivity  
664 across the Oligocene/Miocene boundary as evidenced by benthic foraminiferal



- 665 accumulation rates. *Palaeogeography, Palaeoclimatology, Palaeoecology* 302,  
666 464 - 473 doi:10.1016/j.palaeo.2011.02.006.  
667
- 668 Diester-Haass, L., K. Billups, I. Jacquemin, K.C. Emeis, V. Lefebvre and L. François,  
669 (2013). Paleoproductivity during the middle Miocene carbon isotope events: A  
670 data-model approach. *Paleoceanography*, 28(2), 334-346.  
671
- 672 Gradstein, F. M., J. G. Ogg, M. D. Schmitz and G. M. Ogg (2012), *The geologic time*  
673 *scale 2012*.  
674
- 675 Herbert, T. D (1994), Reading orbital signals distorted by sedimentation: models and  
676 examples. *Orbital forcing and cyclic sequences*, 483-507.  
677
- 678 Hilgen, F. J., L. J. Lourens, and J. A. Van Dam (2012), The Neogene Period. *The*  
679 *geologic time scale*, 2, 923-978.  
680
- 681 Hodell, D. A., and F. Woodruff (1994), Variations in the strontium isotopic ratio of  
682 seawater during the Miocene: Stratigraphic and geochemical implications.  
683 *Paleoceanography* 9, 405-426.  
684
- 685 Hodell, D.A., C.D. Charles and F.J. Sierro (2001), Late Pleistocene evolution of the  
686 ocean's carbonate system. *Earth and Planetary Science Letters*, 192(2), pp.109-  
687 124.  
688
- 689 Holbourn, A., W. Kuhnt, J. T. Simo and Q. Li (2004), Middle Miocene isotope  
690 stratigraphy and paleoceanographic evolution of the northwest and southwest  
691 Australian margins (Wombat Plateau and Great Australian Bight).  
692 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 208(1), 1-22.  
693
- 694 Holbourn, A., W. Kuhnt, S. Clemens, W. Prell, and N. Andersen (2013), Middle to  
695 late Miocene stepwise climate cooling: Evidence from a high-resolution deep-  
696 water isotope curve spanning 8 million years, *Paleoceanography*, 28,  
697 doi:10.1002/2013PA002538.



- 698  
699 Hüsing, S. K., F. J. Hilgen, H. Abdul Aziz, and W. Krijgsman (2007), Completing the  
700 Neogene geological time scale between 8.5 and 12.5 Ma, *Earth and Planetary*  
701 *Science Letters*, 253, 340-358.  
702  
703 King, T. (1996). Quantifying nonlinearity and geometry in time series of climate.  
704 *Quaternary Science Reviews*, 15(4), 247-266.  
705  
706 Krijgsman, W., F. J. Hilgen, I. Raffi, F. J. Sierro and D. S. Wilson (1999),  
707 Chronology, causes and progression of the Messinian salinity crisis. *Nature*,  
708 400(6745), 652-655.  
709  
710 Kochhann, K. G., Holbourn, A., Kuhnt, W., Channell, J. E., Lyle, M., Shackford, J.  
711 K., Andersen, N. (2016). Eccentricity pacing of eastern equatorial Pacific  
712 carbonate dissolution cycles during the Miocene Climatic Optimum.  
713 *Paleoceanography*, 31(9), 1176-1192.  
714  
715 Laskar, J., P. Robutel, F. Joutel, M. Gastineau, A. C. M. Correia, and B. Levrard  
716 (2004), A long-term numerical solution for the insolation quantities of the  
717 Earth. *Astronomy & Astrophysics*, 428(1), 261-285.  
718  
719 Laskar, J., A. Fienga, M. Gastineau, and H. Manche (2011a), La2010: A new orbital  
720 solution for the long term motion of the Earth, *Astronomy and Astrophysics*,  
721 532(A89).  
722  
723 Laskar, J., M. Gastineau, J.-B. Delisle, A. Farrés, and A. Fienga (2011b), Strong  
724 chaos induced by close encounters with Ceres and Vesta, *Astronomy and*  
725 *Astrophysics*, 532(L4), 1-4.  
726  
727 Liebrand, D., L. Lourens, D. A. Hodell, B. de. Boer, R. S. W. van der Wal, and H.  
728 Pälike (2011), Antarctic ice sheet and oceanographic response to eccentricity  
729 forcing in the early Miocene. *Climates of the past*, 7, pp 869 - 880.  
730



- 731 Liebrand, D., H. M. Beddow, L. J. Lourens, H. Pälike, I. Raffi, S. M. Bohaty, F. J.  
732 Hilgen, Mischa J. M. Saes., P.A. Wilson, A. E. van Dijk, D. A. Hodell, D.  
733 Kroon., C. E. Huck and S. J. Batenburg (2016). Cyclostratigraphy and  
734 eccentricity tuning of the early Oligocene through early Miocene (30.1-17.1  
735 Ma): Cibicides mundulus stable oxygen and carbon isotope records from Walvis  
736 Ridge Site 1264. *Earth and Planetary Science Letters*, 450, 392-405.  
737
- 738 Liebrand, D., et al. (2017), Evolution of the early Antarctic ice ages, *Proc. Natl.*  
739 *Acad. Sci. U. S. A.*, 114(15), 3867–3872.  
740
- 741 Littler, K., U. Röhl, T. Westerhold, and J.C. Zachos (2014). A high-resolution benthic  
742 stable-isotope record for the South Atlantic: Implications for orbital-scale  
743 changes in Late Paleocene–Early Eocene climate and carbon cycling. *Earth and*  
744 *Planetary Science Letters*, 401, 18-30.  
745
- 746 Lonsdale, P., (2005), Creation of the Cocos and Nazca plates by fission of the  
747 Farallon plate: *Tectonophysics*, v. 404, p. 237–264, doi: 10.1016/  
748 j.tecto.2005.05.011.  
749
- 750 Lourens, L. J., F. J. Hilgen, N. J. Shackleton, J. Laskar and D. Wilson (2004), ‘Chapter  
751 21: The Neogene Period’. In: Gradstein, F., Ogg, J. and Smith, A., (eds), *A*  
752 *Geologic Time Scale 2004*, Cambridge University Press, Cambridge, pp. 409-  
753 440.  
754
- 755 Ma, W., J. Tian, Q. Li, and P. Wang (2011), Simulation of long eccentricity (400-kyr)  
756 cycle in ocean carbon reservoir during Miocene Climate Optimum: Weathering  
757 and nutrient response to orbital change, *Geophys. Res. Lett.*, 38, L10701,  
758 doi:10.1029/2011GL047680.  
759
- 760 Paillard, D., L. Labeyrie, and P. Yiou (1996), Macintosh program performs time -  
761 series analysis, *Eos Trans. AGU*, 77, 379, doi:10.1029/96EO00259.  
762



- 763 Pälike, H., J. Frazier, and J.C. Zachos (2006a), Extended orbitally forced  
764 palaeoclimatic records from the equatorial Atlantic Ceara Rise. *Quaternary*  
765 *Science Reviews*, 25, 3138–3149.  
766
- 767 Pälike, H., R. N. Norris, J. Herrle, P. A. Wilson, H. K. Coxall, C. H. Lear, N. J.  
768 Shackleton, A. K. Tripathi, and B. S. Wade (2006b), The heartbeat of the  
769 Oligocene climate system, *Science*, 314, 1894–1898, doi:10.1126/  
770 science.1133822.  
771
- 772 Pälike, H., M. W. Lyle, H. Nishi, I. Raffi, K. Gamage, A. Klaus and the Expedition  
773 320/321 Scientists (2010), *Proceedings of the Integrated Ocean Drilling*  
774 *Program*, Volume 320/321. Tokyo (Integrated Ocean Drilling Program  
775 Management International, Inc.).  
776
- 777 Pälike, H., M. W. Lyle, H. Nishi, I. Raffi, A. Ridgwell, K. Gamage, et al (2012), A  
778 Cenozoic record of the equatorial Pacific carbonate compensation depth,  
779 *Nature*, 488(7413), 609-614.  
780
- 781 Shackleton, N. J. (1977), Carbon-13 in Uvigerina: Tropical rain forest history and the  
782 equatorial Pacific carbonate dissolution cycles, in *The Fate of Fossil Fuel CO<sub>2</sub>*  
783 *in the Oceans*, edited by N.R. Andersen and A. Malahoff, 401-427, Plenum,  
784 New York,  
785
- 786 Shackleton, N. J., S. J. Crowhurst, G. P. Weedon, and J. Laskar (1999), Astronomical  
787 Calibration of Oligocene-Miocene Time, *Philosophical Transactions:*  
788 *Mathematical, Physical and Engineering Sciences*, 357(1757), 1907-1929.  
789
- 790 Shackleton, N.J., M. A. Hall, I. Raffi, L. Tauxe, and J. C. Zachos (2000),  
791 Astronomical calibration age for the Oligocene/Miocene boundary. *Geology* 28  
792 (5), 447–450.  
793
- 794 Vandenberghe, N., F. J. Hilgen, and R. P. Speijer (2012), The paleogene period. *The*  
795 *geologic time scale, 2012*, 855-921.  
796





- 797 Westerhold, T., et al. (2012a), Revised composite depth scales and integration of  
798 IODP Sites U1331–U1334 and ODP Sites 1218–1220, in *Proceedings of the*  
799 *Integrated Ocean Drilling Program*, vol. 320/321, edited by H. Pälike et al.,  
800 Integ. Ocean Drill. Progr. Manage. Int., College Station, Tex.  
801
- 802 Westerhold, T., U. Röhl and J. Laskar (2012b), Time scale controversy: Accurate  
803 orbital calibration of the early Paleogene. *Geochemistry, Geophysics,*  
804 *Geosystems*, 13(6).  
805
- 806 Westerhold, T., Röhl, U., Frederichs, T., Agnini, C., Raffi, I., Zachos, J. C., and  
807 Wilkens, R. H.: Astronomical Calibration of the Ypresian Time Scale:  
808 Implications for Seafloor Spreading Rates and the Chaotic Behaviour of the  
809 Solar System?, *Clim. Past Discuss.*, doi:10.5194/cp-2017-15, in review, 2017.  
810
- 811 Wilson, D. S (1988). Tectonic history of the Juan de Fuca ridge over the last 40  
812 million years. *Journal of Geophysical Research: Solid Earth (1978–2012)*,  
813 93(B10), 11863-11876.  
814
- 815 Wilson, D. S. (1993), Confirmation of the astronomical calibration of the magnetic  
816 polarity time scale from rates of sea-floor spreading, *Nature*, 364, 788-790.  
817
- 818 Zachos, J. C., N. J. Shackleton, J. S. Revenaugh, H. Pälike, and B. P. Flower (2001),  
819 Climate response to orbital forcing across the Oligocene – Miocene boundary,  
820 *Science* 292, 274– 278.  
821
- 822 Zeebe, R. E., T. Westerhold, K. Littler, J. C. Zachos (2017), Orbital forcing of the  
823 Paleocene and Eocene carbon cycle, *Paleoceanography*, 32, 440–465,  
824 doi:10.1002/2016PA003054.  
825
- 826 Zeeden, C., F. J. Hilgen, S. K. Hüsing, and L. J. Lourens (2014), The Miocene  
827 astronomical time scale 9–12 Ma: new constraints on tidal dissipation and their  
828 implications for paleoclimatic investigations. *Paleoceanography*, 29(4), 296-  
829 307.



Figure 1






Figure 2






Figure 3







Figure 4






Figure 5



Figure 6






Figure 7








Figure 8






Table 1

Chron (old end)	CCSF-A (m) [Channell et al., 2013]	GTS2004 (Ma) [Lourens et al., 2004]	GTS2012 (Ma) [Hilgen et al., 2012]	Onset (Ma) [Billups et al., 2004]	Onset (Ma) <i>Patike et al.</i> , 2006	Onset (Ma) CaCO <sub>3</sub> based astronomical tuning	Onset (Ma) $\delta^{13}\text{C}$ based astronomical tuning	Difference Between GTS2012 and <i>Billups et al.</i> , 2004 (Myr)	Difference Between GTS2012 and <i>Patike et al.</i> , 2006 (Myr)	Difference Between GTS2012 and CaCO <sub>3</sub> based tuning (Myr)	Difference Between GTS2012 and $\delta^{13}\text{C}$ based tuning (Myr)
<b>C6Bn.1n</b>	89.17	21.936	21.936	21.991	21.998	21.985	22.115	-0.055	-0.062	-0.049	-0.179
<b>C6Bn.1r</b>	89.79	21.992	21.992	22.034	22.062	22.042	22.165	-0.042	-0.070	-0.050	-0.173
<b>C6Bn.2n</b>	94.72	22.268	22.268	22.291	22.299	22.342	22.473	-0.023	-0.031	-0.074	-0.205
<b>C6Br</b>	98.26	22.564	22.564	22.593	22.588	22.621	22.697	-0.029	-0.024	-0.057	-0.133
<b>C6Cn.1n</b>	100.00	22.754	22.754	22.772	22.685	22.792	22.809	-0.018	0.069	-0.038	-0.055
<b>C6Cn.1r</b>	102.50	22.902	22.902	22.931	22.854	22.973	22.970	-0.029	0.048	-0.071	-0.068
<b>C6Cn.2n</b>	103.96	23.030	23.030	23.033	23.026	23.040	23.053	-0.003	0.004	-0.01	-0.023
<b>C6Cn.2r</b>	107.50	23.249	23.233	23.237	23.278	23.212	23.211	-0.004	-0.045	0.021	0.022
<b>C6Cn.3n</b>	108.68	23.375	23.295	23.299	23.340	23.318	23.286	-0.0026	-0.045	-0.023	0.009
<b>C6Cr</b>	119.10	24.044	23.962	23.988	24.022	24.025	24.026	-0.013	-0.060	-0.063	-0.064
<b>C7n.1n</b>	119.58	24.102	24.000	24.013	24.062	24.061	24.066	-0.029	-0.038	-0.061	-0.066
<b>C7n.1r</b>	120.76	24.163	24.109	24.138	24.147	24.124	24.161	-0.029	-0.038	-0.015	-0.052



Table 2

Site	Tuning signal	Tuning target	Lead/lag	Lead/lag	Lead/lag	Lead/lag	Lead/lag	Lead/lag
<b>Site U1334</b> (This study)	CaCO <sub>3</sub> est. %	Eccentricity	Lag ~30 kyrs	Lag ~10 kyrs	Lag ~25-30 kyrs	Lag ~10 kyrs	Lag ~10 kyrs	In phase
<b>Site U1334</b> (This study)	Carbon isotopes	Eccentricity	In phase	Lag ~10 kyrs at 125 kyr, In phase at 96 kyr	In phase	Lag ~10 kyrs	Lag ~10 kyrs	Leads ~10 kyrs
<b>Site 1090</b> (Billups <i>et al.</i> , 2004)	Oxygen isotopes	ETP	Lag ~20 -30 kyrs	Lag ~20 -30 kyrs	In phase	In phase at 125 kyr, ~10 kyr lag at 96 kyr	-	-
<b>Site 926</b> (Palike <i>et al.</i> , 2006a)	Combination of magnetic susceptibility and colour reflectance (SusRef)	ETP	Lag ~35 kyrs	Lag ~30 kyrs	Lag ~10 kyrs	Lag ~20 kyrs	-	-
<b>Site 1218</b> (Palike <i>et al.</i> , 2006b)	Carbon isotopes	ETP	Lag ~30 kyrs	In phase	Lag ~10 kyrs	In phase	-	-
<b>Site 1264</b> (Liebrand <i>et al.</i> , 2016)	CaCO <sub>3</sub> est. (%)	Eccentricity	Lag ~36 kyrs	Lag ~12 kyrs	Lead ~14 kyrs	Lag ~12 kyrs	Unstable phase	In phase