

Full Title: Genetic Variants in *PPARGC1B* and *CNTN4* are Associated with Thromboxane A₂ Formation and with Cardiovascular Event Free Survival in the Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT).

First author's surname and running title: McCarthy; Genetic associations with Thromboxane A₂

Authors: McCarthy NS^{1,2}, Vangjeli C¹, Surendran P^{1,3}, Treumann A⁴, Rooney C¹, Ho E¹, Sever P⁵, Thom S⁵, Hughes AD⁵, Munroe PB⁶, Howard P⁶, Johnson T^{6,7}, Caulfield M⁶, Shields DC³, O'Brien E³, Fitzgerald DJ³, Stanton AV¹.

¹Molecular and Cellular Therapeutics, Royal College of Surgeons in Ireland, RCSI, Dublin, Ireland

²Centre for the Genetic Origins of Health and Disease, University of Western Australia, Perth, Australia

³School of Medicine, UCD Conway Institute, University College Dublin, Dublin.

⁴Newcastle University Protein and Proteome Analysis (NUPPA), University of Newcastle, Newcastle upon Tyne, United Kingdom

⁵International Centre for Circulatory Health, Imperial College London, London, UK

⁶Clinical Pharmacology, William Harvey Research Institute, Barts and the London Medical School, Queen Mary University of London and NIHR Barts Cardiovascular Biomedical Research Unit, London, UK

⁷GlaxoSmithKline, Gunnels Wood Road, Stevenage SG1 2NY, UK

Corresponding author address:

Name: Nina S. McCarthy

Street Address:

Centre for the Genetic Origins of Health and Disease, Level 5,

1 MRF Building,

2 50 Murray St,

3
4 WA 6000

5
6
7 Australia

8
9 Tel: +61(0)424567369

10
11 Email: nina.mccarthy@uwa.edu

12
13
14
15
16 **Total Word Count:** 3,822 (incl abstract, text, tables and figures legends)

17
18
19 **Number of Figures:** 2

20
21 **Number of Tables:** 3

22
23 **Number of Supplementary Files:** 1

24
25
26
27
28 **Journal Subject Terms:**

- 29
30
31 1. Genetic, Association Studies
- 32
33 2. Thrombosis
- 34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Highlights

- This paper describes the first genome wide association study of thromboxane A₂ formation.
- It shows that 5 SNPs in two genes, *PPARGC1B* and *CNTN4*, are associated with elevated thromboxane A₂ formation in 806 Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT) participants and with an excess of cardiovascular events in an independent ASCOT population (n=6,199).
- Results indicate specific protection of aspirin for *PPARGC1B* and *CNTN4*. If confirmed, *PPARGC1B* and *CNTN4* genotyping could potentially provide guidance in the use of aspirin in primary prevention.

Abstract

Background and aims: Elevated urinary 11-dehydro thromboxane B₂ (TxB₂), a measure of thromboxane A₂ formation in vivo, predicts future atherothrombotic events. To further understand this relationship, the genetic determinants of 11-dehydro TxB₂ and their associations with cardiovascular morbidity were investigated in this study.

Methods and Results: Genome-wide and targeted genetic association studies of urinary 11-dehydro TxB₂ were conducted in 806 Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT) participants. The strongest associations were in *PPARGC1B* (rs4235745, rs32582, rs10515638) and *CNTN4* (rs10510230, rs4684343) – these 5 single nucleotide polymorphisms (SNPs) were independently associated with 11-dehydro TxB₂ formation. Haplotypes of 11-dehydro TxB₂ increasing alleles for both *PPARGC1B* and *CNTN4* were significantly associated with 11-dehydro TxB₂, explaining 5.2% and 4.5% of the variation in the whole cohort, and 8.8% and 7.9% in participants not taking aspirin, respectively. In a second ASCOT population (n=6,199), addition of these 5 SNPs significantly improved the covariate-only cox proportional hazards model for cardiovascular events (chisq=14.7, P=0.01). Two of the risk alleles associated with increased urinary 11-dehydro TxB₂ were individually associated with greater incidences of cardiovascular events - rs10515638 (HR =1.31, P=0.01) and rs10510230 (HR=1.25, P=0.007); effect sizes were larger in those not taking aspirin.

Conclusions: *PPARGC1B* and *CNTN4* genotypes are associated with elevated thromboxane A₂ formation and with an excess of cardiovascular events. Aspirin appears to blunt these associations. If specific protection of *PPARGC1B* and *CNTN4* variant carriers by aspirin is confirmed by additional studies, *PPARGC1B* and *CNTN4* genotyping could potentially assist in clinical decision making regarding the use of aspirin in primary prevention.

Key words: Genome wide association study, thromboxane, thrombosis

Introduction

1
2 Thromboxane A₂ (TxA₂) is a potent platelet agonist formed during platelet activation and contributes
3
4 to the risk of arterial thrombosis [1]. Aspirin exerts its major antithrombotic effect by irreversibly
5
6 acetylating platelet cyclo-oxygenase-1 (COX-1), inhibiting production of TxA₂. In high risk patients,
7
8 low-dose aspirin reduces the risk of major cardiovascular events by about 20% [2]. Direct
9
10 measurement of TxA₂ is not feasible as it is rapidly metabolised in vivo to its stable metabolite,
11
12 thromboxane B₂ (TxB₂). Measurement of plasma or urinary levels of 11-dehydro TxB₂ by mass
13
14 spectrometry gives an accurate reflection of in vivo TxA₂ production [3,4]. Increased urinary 11-
15
16 dehydro TxB₂ concentration is an independent predictor of atherothrombotic events even in aspirin-
17
18 treated patients [5,6].
19
20
21
22
23
24
25

26 Measures of TxA₂, including urinary [7] and plasma [8] 11-dehydro TxB₂ have been shown to be
27
28 heritable in Caucasian populations off and on aspirin ($h^2 = 0.5-0.7$ and $0.2-0.4$ respectively). Genetic
29
30 studies have largely focused on the influence of the functional COX-2 single nucleotide
31
32 polymorphism (SNP) rs20417 (-765G>C) on cardiovascular endpoints; a recent meta-analysis
33
34 (n=49,232) reported the minor allele was associated with reduced risk of cardiovascular events and
35
36 lower urinary 11-dehydro TxB₂ [9].
37
38
39
40
41
42

43 To more comprehensively assess genetic contribution to TxA₂ levels, we conducted genome-wide
44
45 and targeted genetic association studies of urinary 11-dehydro TxB₂ in a subset of participants from
46
47 the Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT). In addition, we investigated whether
48
49 variants associated with elevated 11-dehydro TxB₂ levels increased the risk of atherothrombotic
50
51 events in a second subset of ASCOT participants.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Methods

1
2 The ASCOT trial was a randomized, multicentre trial comparing the long-term effects of two
3
4 antihypertensive regimens on myocardial infarction [10]. The population characteristics, genotyping,
5
6 and imputation of the subsets of ASCOT participants investigated in this study are described in full in
7
8 supplementary material online. All patients gave written informed consent. Written informed
9
10 consent and approval by local research ethics committees and/or institutional review boards were
11
12 obtained for ASCOT, the ASCOT DNA Repository, and the Hypertension Associated Cardiovascular
13
14 Disease (HACVD) sub-study.
15
16
17
18
19
20

Genotypes Associated with Elevated Thromboxane A2:

21
22 Fasting urinary 11-dehydro TxB₂ was measured by mass spectrometry in 1,006 participants of the
23
24 HACVD sub-study [11] of the ASCOT trial, and expressed as pg 11-dehydro TxB₂ /mg creatinine to
25
26 normalize for urinary output. Genotyping was performed using the genome-wide Illumina
27
28 HumanCNV370-Duo array (CNV370) and/or the Illumina HumanCVD BeadChip (CVD50k, targeting
29
30 >2,000 genic regions related to cardiovascular disease (CVD)), and quality control exclusions applied
31
32 as described previously [12, 13]. After further excluding all copy number variant (CNV) markers and
33
34 SNPs with minor allele frequency (MAF) <5%, **272,166 genotyped SNPs** (2,031,499 **including** imputed
35
36 SNPs) on the CNV370 chip and 31,570 SNPs on the CVD50K chip were analysed in n=777 and n=544
37
38 individuals with 11-dehydro TxB₂ measurements, respectively (**Supplementary Figure 1**).
39
40
41
42
43
44
45
46

47 For each SNP, a linear regression was performed of genotype (assuming an additive genetic model)
48
49 on urinary 11-dehydro TxB₂ level (a continuous trait) in PLINK v1.07 [14]. Unless otherwise stated,
50
51 analyses were adjusted for the covariates: age, sex, smoking habit (current smokers vs. never & ex-
52
53 smokers), presence of type 2 diabetes, systolic blood pressure (SBP), body mass index (BMI), high
54
55 density lipoprotein (HDL), low density lipoprotein (LDL), randomized anti-hypertensive regimen,
56
57
58
59
60
61
62
63
64
65

1 reported aspirin use, study location (UK/Ireland or Scandinavia), and the first ten vectors from
2 ancestry principal component analysis to avoid confounding due to population stratification. The
3
4 11-dehydro TxB₂ measurement was log transformed to an approximately normal distribution prior to
5
6 analysis and quantile-quantile plots did not indicate any inflation of the test statistics
7
8
9 (Supplementary methods and Supplementary Figures 2 and 3).

10
11
12
13 Haplotypes were imputed in PLINK using the standard E-M algorithm. Bonferroni threshold for the
14
15 CNV370 chip was $P=1.8E-07$ ($0.05/272,166$ genotyped SNPs) to account for multiple testing with a
16
17 5% false positive rate. As many SNPs on the dense CVD50K chip are correlated, correction for the
18
19 21,180 effective tests on the chip is appropriate [13], resulting in a Bonferroni threshold of $P=2.4E-$
20
21 06 ($0.05/21,180$). SNPs associated at $P<1E-03$ were considered suggestive of association on both
22
23
24
25
26 chips.

30 31 **Effect of PPARGC1B and CNTN4 genotypes on Cardiovascular Event Free Survival:**

32
33 All ASCOT participants with DNA ($n=9,063$) were considered for inclusion in the survival analysis. The
34
35 following exclusions were applied: study participants with prior cardiovascular event at baseline
36
37 ($n=1,382$), other non- ischemic/haemorrhagic event during the study ($n=342$), self-declared non-
38
39 European ancestry ($n=370$), missing phenotype data or DNA ($n=770$) (Supplementary Figure 1). The 5
40
41 SNPs independently associated with 11-dehydro TxB₂ were successfully genotyped in 6,199
42
43 participants ($n=3,369$ from Scandinavian centres and $n=2,851$ from UK/Ireland centres). The primary
44
45 endpoint was a composite endpoint including all ischaemic cardiovascular events and procedures,
46
47 defined as any of: fatal and non-fatal myocardial infarction (MI), fatal and non-fatal heart failure,
48
49 fatal and non-fatal ischaemic stroke and transient ischaemic attack, angina (stable and unstable),
50
51 peripheral arterial disease, revascularization procedures, and retinal vascular thrombosis. Where
52
53 participants had multiple events, the earliest qualifying event was used. The study was stopped after
54
55
56
57
58
59 5.5 patient-years of follow-up because of benefits of the amlodipine-based regimen on all-cause
60
61
62
63
64
65

1 mortality and stroke outcomes. Study participants with no qualifying event were censored at the
2 earlier of the date of withdrawal from the study or the date of study termination (median, range: 5.7
3 [1.1-7.1] years).
4
5
6
7
8

9 Survival analyses were conducted using the R packages 'survival' version 3.2.0 and 'survivalROC'
10 version 1.0.3 [15]. The proportional hazards assumption was met for all variables (Supplementary
11 Table 1). Multivariate Cox proportional hazards models were used to collectively analyse all clinical
12 covariates which were associated with the endpoint of cardiovascular event free survival, plus all
13 genotypes. Chi square comparison of the log likelihoods from the covariates-only and full (covariates
14 plus SNPs) model was used to test the alternative hypothesis that the addition of thromboxane-
15 associated SNPs would significantly improve the risk model for CV events. We hypothesized that
16 genotypes associated with increased urinary 11-dehydro TxB₂ in the association study would be
17 inversely correlated with cardiovascular event-free survival, therefore *P* values and confidence
18 intervals reported for individual SNPs are based on a 1-tailed test.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 **Results**

39 **Genotypes Associated with Elevated Thromboxane A₂.**

40 Baseline characteristics of the cohort are shown in Table 1. Approximately half of the patients
41 reported taking aspirin at the time of urinary 11-dehydro TxB₂ measurement. As expected, 11-
42 dehydro TxB₂ levels were significantly lower in those on aspirin.
43
44
45
46
47
48
49
50
51

52 Manhattan plots (Figure 1) show that none of the SNPs was individually associated with 11-dehydro
53 TxB₂ level beyond Bonferroni threshold. There were no associations between COX-1 (*PTGS1*) and
54 COX-2 (*PTGS2*) SNPs and 11-dehydro TxB₂ in this study, even at the suggestive level (*P*<1E-03,
55 Supplementary Table 2).
56
57
58
59
60
61
62
63
64
65

Analysis of CVD50K chip: a *PPARGC1B* haplotype is significantly associated with 11-dehydro TxB₂

The strongest association signal from the CVD50K chip was in the *PPARGC1B* (peroxisome proliferator-activated receptor gamma, coactivator 1 beta) gene (Figure 1), where 14 SNPs were suggestive of association with 11-dehydro TxB₂ (Supplementary Table 3). The most significant SNP at this locus was rs4235745 ($P=4.3E-06$, Table 2). Conditioning on rs4235745 genotype, two SNPs (rs32582 and rs10515638) remained associated with 11-dehydro TxB₂ and were not correlated with each other ($r^2=0.15$, Supplementary Table 4, Supplementary Figure 4). Although rs4235745 and rs32582 were not genotyped on the CNV370 chip, other variants in the gene did show some evidence of association with 11-dehydro TxB₂ (Supplementary Table 5 and Figure 1).

The haplotype of 11-dehydro TxB₂-increasing alleles of the three independently associated SNPs (TTA, frequency 6%) was significantly associated with an increase of 382pg/mg creatinine 11-dehydro TxB₂, surpassing Bonferroni threshold ($P=2.0E-06$, Table 2). This haplotype explained 5.2% of the variation in 11-dehydro TxB₂ in all subjects, 8.8% in subjects not on aspirin, and 1.8% in those on aspirin (multiple R^2 values calculated in R). The 11-dehydro TxB₂-decreasing alleles haplotype (GCC, frequency 66%) was associated with a decrease of 146pg/mg creatinine (Supplementary Table 6).

Analysis of CNV370 chip: a *CNTN4* haplotype is significantly associated with 11-dehydro TxB₂

The strongest association signal from the CNV370 analysis was in the *CNTN4* (contactin 4) gene (Figure 1), where 13 SNPs were suggestive of association with 11-dehydro TxB₂ (Supplementary Table 7). The most significant genotyped SNP at this locus was rs10510230 ($P=5.1E-07$, Table 2) and conditioning on this genotype revealed one other SNP, rs4684343, independently associated with 11-dehydro TxB₂ and not correlated with rs10510230 ($r^2=0.09$, Supplementary Table 8 and Supplementary Figure 5). Only one SNP in the *CNTN4* gene was genotyped on the CVD50k chip

(rs1171387), and this SNP is not very close to either of the top hit *CNTN4* SNPs (rs10510230 or rs4684343). Hence there was very little signal at the *CNTN4* gene on the CVD50K chip (Supplementary Table 5 and Figure 1).

The haplotype of both 11-dehydro TxB₂-increasing alleles of rs10510230 and rs4684343 (CG, frequency 13%) was significantly associated with increased 11-dehydro TxB₂, surpassing Bonferroni threshold ($P=2E-09$, Table 2). This haplotype explained 4.5% of the variation in 11-dehydro TxB₂ in all subjects, 7.9% for subjects not on aspirin, and 1.5% for those on aspirin. The 11-dehydro TxB₂-decreasing alleles haplotype (AA, frequency 44%) was associated with a decrease of 128pg/mg creatinine (Supplementary Table 9).

Interaction with aspirin

The five independently associated *PPARGC1B* and *CNTN4* SNPs showed evidence for interaction with aspirin use in a model including the main SNP and aspirin effects and all covariates, with a greater association between genotype and 11-dehydro TxB₂ in those not taking aspirin (Figure 1, Table 2, Supplementary Tables 10 and 11). The interaction was significant for three of these five SNPs after correction for multiple testing (P value threshold <0.01 ; interaction P -values for *PPARGC1B* SNPs: rs4235745 ($P=3.3E-04$), rs32582 ($P= 3.0E-04$), rs10515638 ($P=0.02$) and for *CNTN4* SNPs: rs4684343 ($P=0.02$), rs10515638 ($P= 6.6E-05$); Supplementary Tables 10 and 11).

Function of 11-dehydro TxB₂ – associated *CNTN4* and *PPARGC1B* SNPs

None of the independently associated SNPs nor their proxies were in exonic or predicted functional regions using the ensemble variant effect predictor (<http://www.ensembl.org/Tools/VEP>), nor were they associated with expression of any genes in any tissues in the GTEx portal (<http://www.gtexportal.org/home/>), (Supplementary Table 12).

Effect of PPARGC1B and CNTN4 genotypes on Cardiovascular Event Free Survival.

Demographic and clinical characteristics of the survival analyses population are shown in Table 1.

Addition of the 11-dehydro TxB₂-associated genotypes improves survival model.

The three *PPARGC1B* and two *CNTN4* independently associated SNPs which comprised the haplotypes significantly associated with 11-dehydro TxB₂ were included in the survival analysis. The addition of these five SNPs significantly improved the covariates-only Cox proportional hazards model of cardiovascular event free survival (chisq comparison of the log likelihoods $P=0.01$, Table 3).

None of the SNPs were associated with any of the cardiovascular risk factors with the exception of a slightly increased risk of type 2 diabetes in rs32582 and rs10510230 carriers (Supplementary Table 13). Results of the univariate survival analysis (Supplementary Table 14) were consistent with those reported in the multivariate analysis. There was a small sample overlap between the genetic association study and the survival analysis; when these individuals were excluded from the survival sample in a sensitivity analysis, it did not alter the overall results (Supplementary Table 15).

Two SNPs independently associated with risk of event

After Bonferroni correction for the 5 SNPs tested (P value threshold <0.01), rs10515638 (*PPARGC1B*) and rs10510230 (*CNTN4*) were associated with increased risk of event in multivariate ($P=0.01$ and $6.9E-03$ respectively) and univariate ($P=0.03$ and $P=8.5E-03$ respectively) survival models. The magnitudes of effect on survival associated with carriage of each minor allele of rs10515638 (*PPARGC1B*, $HR=1.31$) and rs10510230 (*CNTN4*, $HR=1.25$) were similar to or greater than those associated with increments of 10 years age, 20 mmHg systolic BP, 1 mmol LDL-cholesterol, a diagnosis of diabetes mellitus, or use of amlodipine-based antihypertensive therapy in this cohort (Table 3).

Stratification by aspirin use

Kaplan Meier plots of survival analysis stratified by aspirin use for the two significant SNPs show that effect sizes were larger in subjects not taking aspirin compared to those on aspirin (Figure 2 and Supplementary Table 16) and survival receiver operating characteristic (ROC) curves showed that addition of the SNPs improved the area under the curve (AUC) for the all subjects and subjects not on aspirin samples, but not for those on aspirin (Supplementary Figure 5). However, when Aspirin:SNP interaction terms were added to the multivariate survival analysis, none were significant beyond correction for multiple testing (Supplementary Table 17).

Stratification by type of event

Comparing the separate stroke and coronary event analyses, the PPARGC1B SNP (rs10515638) had a larger effect size for coronary events (HR 1.41, P=0.002, 433 events) than for stroke events. By contrast, despite much lower power in the stroke analysis, the CNTN4 SNP (rs10510230) had a larger effect size for stroke (1.31, P=0.17, 57 events) than for coronary events, especially in those not on aspirin (HR 2.71, P=0.003, 26 events), Supplementary Table 18.

The 11-dehydro TxB₂-associated haplotypes are associated with cardiovascular event free survival.

Survival analysis of the PPARGC1B and CNTN4 haplotypes which were associated with the highest 11-dehydro TxB₂ levels (Supplementary Table 19 and Supplementary Figure 6) showed that carriership of one copy of the PPARGC1B haplotype (HR= 1.21, P=0.05) and of the CNTN4 haplotype (HR=1.33, P=0.005), had similar effect sizes on survival to those of the highest effect size SNPs in the haplotypes (rs10515638 (HR=1.31, P=0.01) and rs10510230 (HR=1.25, P=0.007) respectively, Table 3).

Discussion

1 Association analysis of genome-wide and targeted, CVD-based genotype data revealed that a 3-SNP
2 haplotype of *PPARGC1B* and a 2-SNP haplotype of *CNTN4* are significantly associated with urinary
3 levels of 11-dehydro TxB_2 , a surrogate measure for thromboxane A_2 in the HACVD cohort. The 5
4 constituent *PPARGC1B/CNTN4* SNPs were genotyped in a second ASCOT cohort, and addition of
5 these genotypes to clinical and demographic covariates significantly improved the Cox proportional
6 hazards model of cardiovascular event-free survival, with 11-dehydro TxB_2 -increasing alleles
7 associated with reduced event-free survival. In both the genetic association and the survival
8 analyses, stratification by aspirin use showed that most of the genetic associations were more
9 prominent in those not on aspirin. This may suggest that the genetic variants are primarily
10 associated with COX1-synthesized thromboxane A_2 .

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26 **Unfortunately**, suitable data were not available to **formally** replicate the genetic associations with
27
28 11-dehydro TxB_2 in an independent cohort. **However, our finding that the 11-dehydro TxB_2 -**
29
30 **increasing alleles of the same genetic variants increase the risk of cardiovascular events in an**
31
32 **independent population does support our finding that these variants are associated with increased**
33
34 **11-dehydro TxB_2 , (an established cardiovascular risk factor). In addition, Mendel's observation that**
35
36 inheritance of one trait should be independent of the inheritance of other traits asserts that it is
37
38 unlikely the same SNPs would be associated with both 11-dehydro TxB_2 and CHD unless the
39
40 phenotypes are causally linked. Therefore **our finding that the same SNPs are associated with**
41
42 **increased 11-dehydro TxB_2 and increased risk of cardiovascular event (unconfounded by association**
43
44 **with other measured cardiovascular risk factors) supports a causal relationship between 11-dehydro**
45
46 **TxB_2 and CHD and strengthens** the finding by Eickelboom *et al* that urinary 11-dehydro TxB_2 is an
47
48 independent risk factor for atherothrombotic events. Whether the atherosclerosis causes the
49
50 elevated 11-dehydro TxB_2 or vice versa, however, cannot be clearly ascertained from these data.
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 All five 11-dehydro TxB₂-associated SNPs and their proxies are intronic. They are not in strong LD
2 with neighbouring genes and it appears likely therefore that these tagging SNPs implicate *PPARGC1B*
3 and *CNTN4*, however substantial further work is required to ascertain the true causal variants
4 behind the association signals.
5
6
7
8
9

10
11 The *PPARGC1B* gene is a member of the peroxisome proliferator-activated receptor gamma
12 coactivator-1 (*PGC-1*) family, along with its homologue, *PPARGC1A*, and more distant relative PGC-1–
13 related coactivator (*PRC*). *PPARGC1B* and *PPARGC1A* appear to have overlapping function and they
14 exhibit shared, wide tissue expression, especially in heart, skeletal muscle, brain and brown adipose
15 tissue [16]. A variety of metabolic programs are regulated by PGC-1s, including in the heart where
16 PGC-1s help maintain energy homeostasis and activate a broad program of angiogenic factors [17].
17 PGC-1s can co-activate a large range of transcription factors, including the nuclear receptor PPAR γ
18 ([Supplementary Figure 7](#)) which plays an important role in metabolism, is anti-atherosclerotic and
19 anti-inflammatory, and regulates the expression of many genes involved in atherosclerosis including
20 COX-2 [18, 19]. Regulation of COX-2 expression by PPAR γ signalling may be the mechanism whereby
21 polymorphisms in *PPARGC1B* influence thromboxane A₂ levels. However, the larger effect size in
22 subjects not on aspirin suggest the effect is more likely mediated by the primary aspirin target, COX-
23 1, or alternatively through influencing substrate availability or enzymes downstream of COX-1 in the
24 thromboxane A₂ synthetic pathway. It is not known whether PGC-1/ PPAR γ signalling effects
25 expression of COX-1.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

50 Contactin 4 is a member of the immunoglobulin superfamily and is a known neuronal membrane cell
51 adhesion molecule. *CNTN4* expression has been shown in many tissue types other than neuronal,
52 however its role(s) there are not known [20].
53
54
55
56
57
58
59
60
61
62
63
64
65

1 We conducted a search of the term 'cardiovascular in the NHGRI-EBI GWAS catalogue-
2 (<https://www.ebi.ac.uk/gwas/search?query=cardiovascular>, accessed 25/10/17, Supplementary
3 Table 20) in order to put our findings in the context of the vast GWAS data now available. The
4 *CNTN4* gene has been associated with serum uric acid levels in previous genome-wide association
5 and linkage studies, [21, 22], which is itself correlated with a variety of CVD risk factors. There is
6 some evidence for cross talk between COX and uric acid pathways - uric acid has been shown to
7 stimulate COX-2 synthesis [23], for example. Although *PPARGC1B* does not feature in the GWAS
8 catalogue results, genetic polymorphisms in the *PPARGC1B* genes have previously been putatively
9 associated with type 2 diabetes [24, 25].

10
11
12
13
14
15
16
17
18
19
20
21
22
23 A search of the five 11-dehydro TxB₂ SNPs reported in this study in GWAS central,
24 (<http://www.gwascentral.org/> accessed 25/10/17), showed that none of the five SNPs have been
25 included in GWAS of MI in this catalogue (Supplementary Table 21). Some of the SNPs have shown
26 previous nominal associations (P<0.05) with other cardiovascular phenotypes, including associations
27 between rs32583 and ischemic stroke (Supplementary Table 21). It should be noted that the five 11-
28 dehydro TxB₂ -associated SNPs in this study were not present on most of the commonly used GWAS
29 chips; rs4235745 for example is only on the IBC (CVD50K) chip (Supplementary Table 5). In addition,
30 the thromboxane pathway represents one of many risk factors for a cardiovascular disease;
31 therefore, we would not necessarily expect the SNPs reported here to be among the top GWAS hits
32 for CVD.

33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50 Identification of the genetic determinants of thromboxane A₂ levels will help to improve our
51 understanding of the physiology of this important prothrombotic agent. They may also prove to be
52 useful biomarkers for assessing thromboxane levels without the need for urinary thromboxane
53 measurement. If specific protection of *PPARGC1B* and *CNTN4* variant carriers by aspirin is confirmed
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

by additional studies, *PPARGC1B* and *CNTN4* genotyping could potentially assist in clinical decision making regarding the use of aspirin in primary prevention.

Acknowledgements

We thank all ASCOT trial participants, physicians, nurses, and practices for their important contributions to the study.

Funding Sources

The HACVD sub-study was supported by an investigational grant by Pfizer International, New York, NY, USA. The principal funding source for ASCOT was Pfizer, New York, NY, USA. Additional funding for ASCOT was provided by Servier Research Group, Paris, France, and Leo Laboratories, Copenhagen, Denmark. Genotyping was funded by Barts, the London School of Medicine and Dentistry, and by the Centre Nationale de Genotypage Paris. NMCC was funded by the Health Research Board in Ireland under grant number PHD/2007/11.

Disclosures:

TJ is a full time employee of GlaxoSmithKline and owns company stock. NMCC and AVS are named inventors on an RCSI filed patent application entitled "Identification of thrombosis or bleeding risk in an individual with and without antiplatelet therapy", Patent application number 11166142.7 – 2402.

All other authors: none declared.

Table 1. Baseline characteristics

Baseline characteristics of the genetic association cohort (n=806).	All Subjects (n=806)	Not On Aspirin (n=415)	On Aspirin (n=391)	P
11-dehydro TxB ₂ , pg/mg creatinine (median,IQR)	507(305-818)	716(480-1072)	331(224-552)	2.00E-16
Age, years (mean±sd)	64.7±8.1	64.2±8.1	65.2±8.09	0.09
Male (n,%)	626(78%)	336(81%)	290(74%)	0.02
Current smoker (n,%)	186(23%)	94(23%)	92(23%)	0.77
Current diabetes (n,%)	150(19%)	55(13%)	95(24%)	6.00E-05
SBP, mmHg (mean±sd)	135.2±12.0	133.1±12.5	137.4±12.5	2.00E-07
BMI, kg/m ² (mean±sd)	29.2±4.7	29.1±4.7	29.3±4.8	0.53
HDL, mmol/L (mean±sd)	1.3±0.3	1.3 ± 0.4	1.3±0.3	0.16
LDL, mmol/L (mean±sd)	3.0±0.9	3.0 ± 0.9	3.0±0.8	0.17
Amlodipine treatment group (n,%)	407(50%)	215(52%)	192(49%)	0.44
Baseline characteristics of the survival cohort (n=6,199)	All subjects (n=6,199)	Not on aspirin (n=4,355)	On aspirin (n=1,844)	P
Age, years (mean±sd)	61.8±8.4	61.2±8.4	63.0±8.1	2.40E-15
Male (n,%)	4,996(80.6)	3,470(79.7)	1,526(82.7)	5.70E-03
Current smoker (n,%)	2,007(32.4)	1,525(35.0)	482(26.1)	1.00E-11
Diabetes, type 2 (n,%)	1,404(22.6)	894(20.5)	510(27.6)	1.10E-09
SBP, mmHg (mean±sd)	152.7±14.7	153.3±14.9	151.2±14.2	5.40E-07
BMI, kg/m ² (mean±sd)	28.9±4.6	28.8±4.6	29.2±4.6	9.90E-04
HDL, mmol/L (mean±sd)	1.28±0.34	1.30±0.34	1.25±0.32	1.60E-06
LDL, mmol/L (mean±sd)	3.79±0.91	3.78±0.89	3.81±0.95	0.4
UK/Ireland cohort (n,%)	2,830(45.6)	1,534(35.2)	1,296(70.3)	<2.2E-16
Amlodipine treatment group (n,%)	3,153(50.9)	2,231(51.2)	922(50.0)	0.39

17

Follow-up, years (median, IQR)	5.6(5.1-6.0)	5.7(5.2-6.1)	5.5(5.0-6.0)	6.60E-15
n events (composite endpoint)	477(7.7)	189(4.3)	288(15.6)	<2.2E-16
n events per 100 person-years (95%CI)	1.40(1.28-1.53)	0.77(0.67-0.90)	2.98(2.64-3.34)	<2.2E-16
-Coronary revascularisation procedure (n)	149	55	94	<2.2E-16
-Non-fatal MI (incl. silent) + fatal CHD (n)	271	108	163	<2.2E-16
-Ischemic stroke (n)	44	15	29	9.20E-08
-Other (n)	13	11	2	0.46

Amlodipine treatment group: membership of amlodipine (vs atenolol) regime. SBP: Systolic blood pressure; BMI: Body mass index; HDL: High density lipoprotein; LDL: Low-density lipoprotein.

UK/Ireland: UK/Ireland (vs Scandinavian) centres. Composite endpoint: total ischaemic cardiovascular events/procedures as described in methods. MI: myocardial infarction. CHD: coronary heart disease. IQR: Inter quartile range; SD: Standard deviation; 95% CIs for events were calculated using an exact Poisson test. *P*: comparison of on-aspirin and not-on-aspirin groups (Student's t-test or Mann-Whitney U test for normally and non-normally distributed variables respectively, or Pearson's chi-squared test of proportions).

Table 2. *PPARGC1B* and *CNTN4* SNPs independently associated with 11-dehydro TxB₂.

<i>PPARGC1B</i>							all subjects (n=544)			subjects not on aspirin (n=276)		subjects on aspirin (n=268)	
Chr	SNP	Position	MAF	A1/2	r ²	D'	beta	P	P ^{SNP*Asp}	beta	P	beta	P
5	rs10515638	intron 1	0.08	T/G	0.07	0.58	233.1	7.30E-04	0.02	391.9	1.90E-03	74.7	0.12
5	rs4235745	intron 1	0.3	T/C	NA	NA	166.7	4.30E-06	3.30E-04	284.9	2.20E-05	50.5	0.06
5	rs32582	intron 2	0.18	A/C	0.42	0.93	201.1	9.10E-06	3.00E-04	353.1	2.10E-05	45.4	0.18
3-SNP minor allele			0.06	TTA			382.4	2.10E-06		658.7	9.50E-06	105.5	8.00E-02
haplotype													
<i>CNTN4</i>							all subjects (n=772)			subjects not on aspirin (n=396)		subjects on aspirin (n=376)	
Chr	SNP	Position	MAF	A1/2	r ²	D'	beta	P	P ^{SNP*Asp}	beta	P	beta	P
3	rs10510230	intron 1	0.15	C/A	NA	NA	189.8	5.10E-07	6.60E-05	329.5	1.50E-07	37.6	0.34
3	rs4684343	intron 2	0.45	A/G	0.1	0.77	-133.4	8.80E-07	1.90E-02	-194.6	2.50E-05	-65.2	0.02
2-SNP minor/major allele			0.13	CG			249	2.00E-09		456.2	1.30E-10	48.4	0.24
haplotype													

CHR: chromosome, BP: base pair position, MAF: minor allele frequency, beta: change in 11-dehydro TxB₂ associated with one copy of the minor allele /

haplotype. Position: of the SNP in the *PPARGC1B* gene, r²: squared correlation coefficient between the genotype of the SNP and the top hit SNP in the gene

(rs4235745 for PPARGC1B and rs10510230 for CNTN4). D' : the normalised coefficient of linkage disequilibrium between the genotype of the SNP and the top hit SNP in the gene. A1: major allele, A2: minor allele.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 3. Multivariate Cox proportional hazards analysis all significant clinical covariates plus the five 11-dehydro Tx_B₂-associated SNPs with a composite cardiovascular endpoint of all ischaemic cardiovascular events and procedures.

All subjects (n=6,199, 477 ischaemic cardiovascular events)			
	HR	95% CI	P
Age (yrs)	1.02	1.01-1.04	7.70E-05
Male sex	1.54	1.17-2.02	2.00E-03
Current smoker	1.43	1.17-1.74	0.03
Type 2 diabetes	1.12	0.92-1.37	5.30E-04
SBP (mmHg)	1.01	1.00-1.02	0.06
BMI (kg/m ²)	0.98	0.95-1.00	0.25
HDL (mmol/L)	0.44	0.31-0.62	8.90E-05
LDL (mmol/L)	1.21	1.10-1.34	1.80E-06
Amlodipine treatment	0.82	0.68-0.98	1.40E-04
Aspirin use	4.68	3.84-5.71	< 2e-16
UK/Ireland cohort	0.56	0.45-0.68	2.10E-08
rs4235745 (per minor (T) allele)	0.84	0.71-0.99	0.96
rs32582 (per minor (A) allele)	1.00	0.82-1.21	0.48
rs10515638 (per minor (T) allele)	1.31	1.08-1.58	0.01
rs10510230 (per minor (C) allele)	1.25	1.08-1.46	6.90E-03
rs4684343 (per minor (G) allele)	1.04	0.93-1.17	0.28

LRT comparison with covariates-only model: chisq 14.7 on 5 df, p= 0.01

The primary endpoint was a composite endpoint including all ischaemic cardiovascular events and procedures, defined as any of: fatal and non-fatal MI, fatal and non-fatal heart failure, fatal and non-fatal ischaemic stroke and transient ischaemic attack, angina (stable and unstable), peripheral arterial disease, revascularization procedures, and retinal vascular thrombosis. The Cox proportional

1 hazards model included all SNP genotypes and all covariates. Df: degrees of freedom. SNP statistics
2 are based a one-tailed test; where the observed direction of effect is opposite to that predicted, 1-*P*
3
4 is reported and 95% CI excludes the lower 5% of the distribution. For all SNPs, minor alleles were
5
6 associated with increased 11-dehydro TxB₂. SBP: systolic blood pressure. BMI: body mass index. HDL:
7
8 high density lipoprotein cholesterol. LDL: low density lipoprotein cholesterol. Amlodipine treatment
9
10 group: membership of amlodipine (vs atenolol) regime. UK/Ireland: UK/Ireland: UK/Ireland (vs
11
12 Scandinavian) centres participants at UK/Ireland centres vs Scandinavian participants. LRT: likelihood
13
14 ratio test, comprising the chisq (chi squared) comparison of the log likelihoods of the covariates-only
15
16 and covariates-plus-genotypes models.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figures




Figure 1. Association with 11-dehydro TxB₂ for the 31,570 SNPs (MAF \geq 0.05) on the CVD50K chip (A) and 2,031,499 SNPs (MAF \geq 0.05) on the CNV370 chip (B). X-axis: chromosomal location. All SNP associations were adjusted for age, sex, smoking habit, diabetes, systolic blood pressure, BMI, HDL, LDL, aspirin and anti-hypertensive regimen. The lower dashed lines are the threshold for 'suggestive' association ($P < 1E-03$), and the upper dashed lines are the Bonferroni threshold for significance ($P < 2.4E-06$ for the CNV50K chip and $P = 1.8E-07$ for the CNV370K chip). The most strongly associated locus in all subjects on the CVD50K chip, *PPARGC1B*, is coloured in red and on the CNV370 chip, *CNTN4*, is coloured in blue.




Figure 2. Kaplan Meier plot of survival by genotype for the two SNPs independently associated

with **cardiovascular event-free** survival, rs10515638 (PPARGC1B) and rs10510230 (CNTN4).

‘Carriers’: carriers of one or more minor allele. ‘Non carriers’: major allele homozygotes. HR: Hazard ratio for the SNP in the multivariate Cox proportional hazards analysis, and corresponding P value

(Table 3).

References

- 1
2 1. Davi G and Patrono C. Platelet activation and atherothrombosis. *N Engl J Med*.
3
4 2007;357:2482-94.
5
- 6
7 2. Antithrombotic Trialists' (ATT) Collaboration. Aspirin in the primary and secondary prevention of
8
9 vascular disease: collaborative meta-analysis of individual participant data from randomised trials.
10
11 *Lancet*. 2009;373:1849-60.
12
- 13
14 3. Catella F, Healy D, Lawson JA and FitzGerald GA. 11-Dehydrothromboxane B2: a quantitative
15
16 index of thromboxane A2 formation in the human circulation. *Proc Natl Acad Sci U S A*.
17
18 1986;83:5861-5.
19
- 20
21 4. Ciabattoni G, Pugliese F, Davi G, Pierucci A, Simonetti BM and Patrono C. Fractional
22
23 conversion of thromboxane B2 to urinary 11-dehydrothromboxane B2 in man. *Biochim Biophys Acta*.
24
25 1989;992:66-70.
26
- 27
28 5. Eikelboom JW, Hirsh J, Weitz JI, Johnston M, Yi Q and Yusuf S. Aspirin-resistant thromboxane
29
30 biosynthesis and the risk of myocardial infarction, stroke, or cardiovascular death in patients at high
31
32 risk for cardiovascular events. *Circulation*. 2002;105:1650-5.
33
- 34
35 6. Eikelboom JW, Hankey GJ, Thom J, Bhatt DL, Steg PG, Montalescot G, et al. Incomplete
36
37 inhibition of thromboxane biosynthesis by acetylsalicylic acid: determinants and effect on
38
39 cardiovascular risk. *Circulation*. 2008;118:1705-12.
40
- 41
42 7. Faraday N, Yanek LR, Mathias R, Herrera-Galeano JE, Vaidya D, Moy TF, et al. Heritability of
43
44 platelet responsiveness to aspirin in activation pathways directly and indirectly related to
45
46 cyclooxygenase-1. *Circulation*. 2007;115:2490-6.
47
- 48
49 8. Vila L, Martinez-Perez A, Camacho M, Buil A, Alcolea S, Pujol-Moix N, et al. Heritability of
50
51 thromboxane A2 and prostaglandin E2 biosynthetic machinery in a Spanish population. *Arterioscler*
52
53 *Thromb Vasc Biol*. 2010;30:128-34.
54
- 55
56 9. Ross S, Eikelboom J, Anand SS, Eriksson N, Gerstein HC, Mehta S, et al. Association of
57
58 cyclooxygenase-2 genetic variant with cardiovascular disease. *Eur Heart J*. 2014;35:2242-8a.
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
10. Dahlof B, Sever PS, Poulter NR, Wedel H, Beevers DG, Caulfield M, et al. Prevention of cardiovascular events with an antihypertensive regimen of amlodipine adding perindopril as required versus atenolol adding bendroflumethiazide as required, in the Anglo-Scandinavian Cardiac Outcomes Trial-Blood Pressure Lowering Arm (ASCOT-BPLA): a multicentre randomised controlled trial. *Lancet*. 2005;366:895-906.
 11. Stanton A, Fitzgerald D, Hughes A, Mayet J, O'Brien E, Poulter NR, et al. An intensive phenotyping study to enable the future examination of genetic influences on hypertension-associated cardiovascular disease. *J Hum Hypertens*. 2001;15 Suppl 1:S13-8.
 12. Deshmukh HA, Colhoun HM, Johnson T, McKeigue PM, Betteridge DJ, Durrington PN, et al. Genome-wide association study of genetic determinants of LDL-c response to atorvastatin therapy: importance of Lp(a). *J Lipid Res*. 2012;53:1000-11.
 13. Johnson T, Gaunt TR, Newhouse SJ, Padmanabhan S, Tomaszewski M, Kumari M, et al. Blood pressure loci identified with a gene-centric array. *Am J Hum Genet*. 2011;89:688-700.
 14. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007;81:559-75.
 15. R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria, 2010.
 16. Meirhaeghe A, Crowley V, Lenaghan C, Lelliott C, Green K, Stewart A, et al. Characterization of the human, mouse and rat PGC1 beta (peroxisome-proliferator-activated receptor-gamma co-activator 1 beta) gene in vitro and in vivo. *Biochem J*. 2003;373:155-65.
 17. Rowe GC, Jiang A and Arany Z. PGC-1 coactivators in cardiac development and disease. *Circ Res*. 2010;107:825-38.
 18. Inoue H, Tanabe T and Umesono K. Feedback control of cyclooxygenase-2 expression through PPARgamma. *J Biol Chem*. 2000;275:28028-32.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
19. Ahmadian M, Suh JM, Hah N, Liddle C, Atkins AR, Downes M, et al. PPARgamma signaling and metabolism: the good, the bad and the future. *Nat Med.* 2013;19:557-66.
 20. Hansford LM, Smith SA, Haber M, Norris MD, Cheung B and Marshall GM. Cloning and characterization of the human neural cell adhesion molecule, CNTN4 (alias BIG-2). *Cytogenet Genome Res.* 2003;101:17-23.
 21. Voruganti VS, Nath SD, Cole SA, Thameem F, Jowett JB, Bauer R, et al. Genetics of variation in serum uric acid and cardiovascular risk factors in Mexican Americans. *J Clin Endocrinol Metab.* 2009;94:632-8.
 22. Voruganti VS, Kent JW, Jr., Debnath S, et al. Genome-wide association analysis confirms and extends the association of SLC2A9 with serum uric acid levels to Mexican Americans. *Frontiers in genetics* 2013;4:279.
 23. Kanellis J, Watanabe S, Li JH, Kang DH, Li P, Nakagawa T, et al. Uric acid stimulates monocyte chemoattractant protein-1 production in vascular smooth muscle cells via mitogen-activated protein kinase and cyclooxygenase-2. *Hypertension.* 2003;41:1287-93.
 24. Villegas R, Williams SM, Gao YT, Long J, Shi J, Cai H, et al. Genetic variation in the peroxisome proliferator-activated receptor (PPAR) and peroxisome proliferator-activated receptor gamma co-activator 1 (PGC1) gene families and type 2 diabetes. *Ann Hum Genet.* 2014;78:23-32.
 25. Park KS, Shin HD, Park BL, Cheong HS, Cho YM, Lee HK, et al. Putative association of peroxisome proliferator-activated receptor gamma co-activator 1beta (PPARGC1B) polymorphism with Type 2 diabetes mellitus. *Diabet Med.* 2006;23:635-42.