



Contents lists available at ScienceDirect

Vaccine

journal homepage: [www.elsevier.com/locate/vaccine](http://www.elsevier.com/locate/vaccine)

## Acceptability of intranasal live attenuated influenza vaccine, influenza knowledge and vaccine intent in The Gambia

Edwin P. Armitage<sup>a,\*</sup>, Janko Camara<sup>a</sup>, Sulayman Bah<sup>a</sup>, Alice S. Forster<sup>b</sup>, Ed Clarke<sup>a</sup>, Beate Kampmann<sup>a,c</sup>, Thushan I. de Silva<sup>a,c</sup>

<sup>a</sup> Vaccines and Immunity Theme, Medical Research Council Unit The Gambia, PO Box 273, Banjul, Gambia

<sup>b</sup> Research Department of Behavioural Science and Health, University College London, Gower Street, London WC1E 6BT, UK

<sup>c</sup> Centre of International Child Health, Section of Paediatrics, Department of Medicine, Imperial College London, St Mary's Campus, London W2 1PG, UK

### ARTICLE INFO

#### Article history:

Received 30 October 2017

Received in revised form 4 February 2018

Accepted 5 February 2018

Available online xxxxx

#### Keywords:

Influenza

Vaccine acceptability

Intranasal live attenuated influenza vaccine

Knowledge, attitudes and perceptions

Health Belief Model

The Gambia

### ABSTRACT

**Background:** The burden of influenza is increasingly recognised in Africa. The WHO recommends introducing influenza vaccination to high-risk groups: pregnant women, children <5 years, and the elderly. The Gambia currently has no influenza vaccination policy, but the NASIMMUNE study, a clinical trial of intranasal live attenuated influenza vaccines (LAIV) in young children provided an opportunity to study maternal attitudes towards LAIV for the first time in sub-Saharan Africa. We assess acceptability of LAIV, influenza knowledge and attitudes towards influenza vaccination in Gambian women. Additionally, we investigate predictors of willingness to receive influenza vaccine (intent) in pregnancy or seasonally for children <5.

**Methods:** A cross-sectional survey was conducted in Gambian women at two urban health facilities. To assess LAIV acceptability, the exposure group (women whose children had received LAIV during the NASIMMUNE study) were compared to a control group (women whose children were not enrolled in the NASIMMUNE study). Demographics and health belief constructs were analysed as predictors of influenza knowledge and vaccine intent.

**Findings:** The exposure group (n = 150) expressed a higher preference for a nasal spray vaccine than an injection compared to the control group (n = 304) (93.3% vs. 34.9%, OR = 26.15, p < 0.0001). Those in the exposure group who preferred the nasal spray found it less distressing, safer or equally safe, and easier or equally easy to give (all p < 0.001) than injections. Influenza knowledge increased with education level (p = 0.006 for higher education vs. none), and varied between sites (p = 0.0005). Vaccine intent was >98%, but no association with influenza knowledge or difference between groups was observed. Various health belief constructs were associated with vaccine intent.

**Conclusion:** LAIV acceptability was higher in those with first-hand experience. Influenza vaccine intent was also high. Incorporation of seasonal LAIV into the childhood immunisation schedule in The Gambia would be feasible, particularly if combined with community-based health education.

© 2018 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

### 1. Introduction

Influenza is a major cause of acute respiratory infection globally, leading to a significant burden of morbidity and mortality [1]. Seasonal influenza has been well-studied in high- and middle-income countries, but neglected in Africa [2]. The rate of influenza-associated hospitalisations in children <5 years of age is approximately 3-fold higher in low- and middle-income countries (LMIC) compared to high-income countries [3]. One meta-

analysis found that 99% of deaths attributable to influenza-associated acute lower respiratory infections occurred in LMICs [4].

As a result of the increasing awareness of influenza-related disease in LMIC, in 2012 the WHO recommended that countries should consider influenza vaccination in high-risk groups including pregnant women and children <5 years [5]. The Gambia has no influenza vaccination policy and is yet to include influenza vaccination in the Expanded Programme on Immunisation (EPI) [6,7]. The NASIMMUNE study, a systems immunology research study in children aged 24–59 months of intranasal live attenuated influenza vaccine (LAIV) is ongoing (NCT02972957). To our knowledge, this is the first interventional study in The Gambia using intranasal

\* Corresponding author.

E-mail address: [earmitage@mrc.gm](mailto:earmitage@mrc.gm) (E.P. Armitage).

<https://doi.org/10.1016/j.vaccine.2018.02.037>

0264-410X/© 2018 The Authors. Published by Elsevier Ltd.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Please cite this article in press as: Armitage EP et al. Acceptability of intranasal live attenuated influenza vaccine, influenza knowledge and vaccine intent in The Gambia. Vaccine (2018), <https://doi.org/10.1016/j.vaccine.2018.02.037>

LAIV and offers a unique opportunity to study, for the first time, attitudes towards a new mode of vaccine delivery in this setting.

Factors that shape parental views on vaccination impact childhood vaccination rates and having insight into parental perceptions can be useful when introducing new vaccinations to a country [8]. High vaccine uptake in The Gambia through the EPI [9] has led to reductions in invasive disease secondary to *Haemophilus influenzae* type b [10] and *Streptococcus pneumoniae* [11]. Yet as more vaccines are added to the EPI, vaccine hesitancy might be a concern [12–14]. Knowledge and attitudes surrounding influenza and influenza vaccination in The Gambia are unknown. Gaining a better understanding of these factors can help guide future influenza vaccination strategies, including maternal influenza immunisation, shown to be beneficial to both mothers and infants in recent studies from South Africa and Mali [15,16].

The Health Belief Model (HBM) provides a framework of psychosocial constructs that may be determinants of health behaviour [17]. It posits that preventative health behaviours are more likely to be exhibited when an individual perceives they are highly susceptible to the disease, that the disease is serious, that the behaviour is beneficial, there are few barriers, and are cued into action [18]. These constructs have been shown to predict uptake of influenza vaccination and other health behaviours [19–24]. However, emotions experienced at the point of decision-making may be as important in health behaviour as a cognitive assessment of the risk, such as worry and anticipated regret, which have also been shown to be predictive of influenza vaccination [20,25,26]. Few studies have used the HBM to study influenza vaccine behaviour in LMICs [27–31].

We aimed to compare attitudes towards the safety, ease of use, and tolerability of LAIV between mothers with direct experience of their children receiving LAIV and mothers without such experience, and to establish whether their attitudes towards intranasal vaccines differ. Additionally, we aimed to quantify knowledge about influenza and to determine possible relationships between influenza knowledge, socio-demographic factors and willingness to accept influenza vaccination in pregnancy or for their children <5 (vaccine intent). Finally, we aimed to explore the relationship between health belief constructs and influenza vaccine intent in this cohort.

## 2. Methods

### 2.1. Study design

A cross-sectional survey was conducted in women  $\geq 18$  years at two sites (Sukuta and Faji Kunda health centres) five kilometres apart, in the coastal region of The Gambia in August and September 2017. In Sukuta, where NASIMMUNE was conducted, mothers of children who had been vaccinated with the intranasal LAIV (Nasovac-S<sup>®</sup>, Serum Institute of India Pvt. Ltd.) as part of the study were contacted sequentially and invited to an interview, up to a total of 150 (exposure group). A further 154 women from Sukuta who had not participated in the NASIMMUNE study were recruited at the same health centre. Due to wider community sensitization regarding the NASIMMUNE trial carried out in Sukuta, this group may have had some exposure to information about influenza and LAIV, therefore an additional control group of 150 women were recruited at Faji Kunda health centre (where there could have been no exposure to the study or community sensitization). These women were recruited through opportunistic sampling when attending for routine healthcare. The sample size was determined by the availability of mothers of children in the NASIMMUNE study ( $n = 168$ ), with a 1:2 exposure to control ratio. Inclusion criteria

were having at least one child <5, maternal age  $\geq 18$ , resident in the area, and fluency in Mandinka (the most commonly spoken local language in the areas).

### 2.2. Data collection

Face-to-face interviews were conducted by trained field-workers who entered data in real-time into a questionnaire designed in REDCap<sup>™</sup> [32]. The questionnaire was developed using simple terminology and was refined through cognitive pre-testing with field-workers and test participants. The questionnaire included seven sections: inclusion criteria, socio-demographics, vaccine intent, influenza knowledge, health-seeking behaviour, health beliefs, and LAIV acceptability. The influenza knowledge questions were designed *de novo* for this setting, selected and simplified from influenza knowledge questions used previously [31,33–35] and refined following the cognitive pre-testing phase. Questions asked about symptoms, transmission, prevention, treatment, risk, and vaccination (see [supplementary material: Appendix A](#)). The responses were collated to form a score out of 15 points, which was converted to a percentage for analysis (score divided by 15, multiplied by 100).

Influenza vaccine intent was assessed for pregnancy and for children <5 by asking participants to respond to two statements: “If I was pregnant, I would get a flu vaccine if it was free” and “I would get a flu vaccine for my child under 5, every year, if it was free”. Answers were recorded on a 4-point scale where 1 = agree strongly, 2 = agree, 3 = disagree and 4 = disagree strongly, with a “don’t know” option. In follow-up questions, unprompted reasons given for answers were coded into predefined categories by the interviewer.

Health belief constructs were assessed using statements answered on the same 4-point scale. HBM constructs assessed were: perceived susceptibility, severity, benefit, barriers and cues to action. Two additional concepts were also included: worry and anticipated regret. The answers given were converted into binary “agree” or “disagree” responses for analysis.

### 2.3. Ethical considerations

Ethical approval for the study was provided by The Gambia Government/MRC Joint Ethics Committee (SCC1555). Written informed consent was obtained from all participants. It was made clear that answers were confidential and anonymised, that they could withdraw at any time or decline to answer any questions.

### 2.4. Statistical analysis

Analysis was conducted using Stata<sup>®</sup> 12.0. Descriptive statistics were used to compare proportions between groups: Pearson’s chi-squared test or two-tailed Fisher’s exact test (when one category had <5 participants) for categorical data; and Student’s *t*-test or Wilcoxon rank-sum test for normally and non-normally distributed continuous data respectively. Univariate linear regression analysis was performed for predictors of influenza knowledge. Significance-testing was used for selection of variables to include in the multivariate model at a level of  $p < 0.2$  to minimize type II error in selection [36]. The Cochran–Mantel–Haenszel test was used to analyse variations between groups for individual influenza knowledge question responses and to evaluate the difference in preference for intranasal or injection vaccinations between groups. Two-tailed Fisher’s exact test was used to analyse perceptions of intranasal LAIV in the exposed group and associations between vaccine intent and health belief constructs.  $p < 0.05$  was considered statistically significant.

### 3. Results

#### 3.1. Participants' characteristics

The 454 participants' characteristics who answered the survey are displayed in Table 1. There were significant differences between the exposure and control groups with respect to age, parity, education, husband's education and monthly household income.

#### 3.2. LAIV acceptability

When asked "If given a choice for your child between a flu vaccine injection and a nasal spray, which would you prefer?" a signif-

icantly higher proportion of the exposure group stated they would prefer a nasal spray compared to the control group (93.3% vs. 34.9%, OR = 26.15,  $p < 0.0001$ , Table 2). The most common unprompted reasons for preferring the nasal spray were that it is easier to give, less painful and a perception of greater effectiveness (Fig. 1A). The most commonly stated reasons for preferring injections were a greater familiarity with injections and a belief that injections are more effective (Fig. 1B). In the exposure group, stating a preference for the nasal spray was associated with finding the LAIV less distressing ( $p < 0.001$ ), safer or equally safe ( $p < 0.001$ ) and easier or equally easy to give ( $p < 0.001$ ) when compared to injections (Table 2).

**Table 1**  
Socio-demographic and other characteristics of participants by group.

| Participant characteristics              | All n = 454 n (%) | NASIMMUNE (exposure) n = 150 n (%) | Non-NASIMMUNE (control) n = 304 n (%) | P value  |
|--|-------------------|------------------------------------|---------------------------------------|----------|
| <b>Interview site:</b>                   |                   |                                    |                                       |          |
| Sukuta                                   | 304 (67.0)        | 150 (100.0)                        | 154 (50.7)                            | –        |
| Faji Kunda                               | 150 (33.0)        | 0 (0.0)                            | 150 (49.3)                            | –        |
| Mean% influenza knowledge score (SD)     | 68.0 (10.3)       | 69.2 (10.0)                        | 67.4 (10.4)                           | 0.0816*  |
| Vaccine intent in pregnancy              | 447 (98.5)        | 150 (100.0)                        | 297 (97.7)                            | 0.185†   |
| Vaccine intent for children < 5 years    | 448 (98.7)        | 146 (97.3)                         | 302 (99.3)                            | 0.600‡   |
| <b>Socio-demographics:</b>               |                   |                                    |                                       |          |
| Mean age (SD)                            | 28.4 (5.6)        | 29.5 (5.1)                         | 27.9 (5.7)                            | 0.0026*  |
| Median parity (IQR)                      | 3 (2–5)           | 3 (2–5)                            | 3 (2–5)                               | 0.0340** |
| Median household size (IQR)              | 4 (3–6)           | 4 (3–5)                            | 4 (3–6)                               | 0.0852** |
| Currently pregnant                       | 36 (7.9)          | 15 (10.0)                          | 21 (6.9)                              | 0.251‡   |
| Another pregnancy in household           | 111 (24.5)        | 33 (22.0)                          | 78 (25.7)                             | 0.394‡   |
| <b>Marital status:</b>                   |                   |                                    |                                       |          |
| Never married                            | 5 (1.1)           | 1 (0.7)                            | 4 (1.3)                               | –        |
| First (and only) wife                    | 344 (75.8)        | 118 (78.7)                         | 226 (74.3)                            | –        |
| First (not only) wife                    | 37 (8.2)          | 9 (6.0)                            | 28 (9.2)                              | –        |
| Second wife                              | 58 (12.8)         | 18 (12.0)                          | 40 (13.2)                             | –        |
| Third or fourth wife                     | 10 (2.2)          | 4 (2.7)                            | 6 (2.0)                               | 0.734†   |
| <b>Education (English school):</b>       |                   |                                    |                                       |          |
| None                                     | 98 (21.6)         | 27 (18.0)                          | 71 (23.4)                             | –        |
| Arabic school only                       | 78 (17.2)         | 17 (11.3)                          | 61 (20.1)                             | –        |
| Attended primary school                  | 50 (11.0)         | 17 (11.3)                          | 33 (10.9)                             | –        |
| Attended upper school                    | 215 (47.4)        | 86 (57.3)                          | 129 (42.4)                            | –        |
| Higher education                         | 13 (2.9)          | 3 (2.0)                            | 10 (3.3)                              | 0.025†   |
| <b>Occupation:</b>                       |                   |                                    |                                       |          |
| None/house wife                          | 266 (58.6)        | 79 (52.7)                          | 187 (61.5)                            | –        |
| Student                                  | 3 (0.7)           | 1 (0.7)                            | 2 (0.7)                               | –        |
| Self-employed (unskilled)                | 69 (15.2)         | 32 (21.3)                          | 37 (12.2)                             | –        |
| Self-employed (skilled)                  | 65 (14.3)         | 23 (15.3)                          | 42 (13.8)                             | –        |
| Employed (salaried)                      | 51 (11.2)         | 15 (10.0)                          | 36 (11.8)                             | 0.105†   |
| <b>Husband's education:</b>              |                   |                                    |                                       |          |
| None                                     | 42 (9.3)          | 6 (4.0)                            | 36 (11.8)                             | –        |
| Arabic school only                       | 61 (13.4)         | 13 (8.7)                           | 48 (15.8)                             | –        |
| Attended primary school                  | 16 (3.5)          | 7 (4.7)                            | 9 (3.0)                               | –        |
| Attended upper school                    | 259 (57.1)        | 89 (59.3)                          | 170 (55.9)                            | –        |
| Higher education                         | 51 (11.2)         | 23 (15.3)                          | 28 (9.2)                              | –        |
| Don't know                               | 25 (5.5)          | 12 (8.0)                           | 13 (4.3)                              | 0.004‡   |
| <b>Husband's occupation:</b>             |                   |                                    |                                       |          |
| None                                     | 15 (3.3)          | 5 (3.3)                            | 10 (3.3)                              | –        |
| Student                                  | 3 (0.7)           | 1 (0.7)                            | 2 (0.7)                               | –        |
| Self-employed (unskilled)                | 6 (1.3)           | 2 (1.3)                            | 4 (1.3)                               | –        |
| Self-employed (skilled)                  | 161 (35.5)        | 48 (32.0)                          | 113 (37.2)                            | –        |
| Employed (salaried)                      | 253 (55.7)        | 93 (62.0)                          | 160 (52.6)                            | –        |
| Don't know                               | 16 (3.5)          | 1 (0.7)                            | 15 (4.9)                              | 0.120†   |
| <b>Household income (GMD per month):</b> |                   |                                    |                                       |          |
| GMD 500–GMD 4,999                        | 71 (15.6)         | 20 (13.3)                          | 51 (16.8)                             | –        |
| GMD 5,000–GMD 9,999                      | 186 (41.0)        | 70 (46.7)                          | 116 (38.2)                            | –        |
| GMD >10,000                              | 100 (22.0)        | 44 (29.3)                          | 56 (18.4)                             | –        |
| Don't know/unwilling to say              | 97 (21.4)         | 16 (10.7)                          | 81 (26.6)                             | <0.001‡  |

Monthly household income stated in Gambian Dalasis (GMD). 1 USD = 48 GMD at time of writing.

\* Two-group mean-comparison *t*-test (normally distributed).

\*\* Wilcoxon rank-sum test (non-normally distributed).


† Two-tailed Fisher's exact test (when  $n < 5$  in some cases).

‡ Pearson's chi-squared test.

**Table 2**  
Intranasal LAIV preference and acceptability.

| Question                       | Levels      | NASIMMUNE (exposure) n = 150 n (%) |                               | Non-NASIMMUNE (control) n = 304 n (%) | Cochran–Mantel–Haenszel odds ratio (95% CI, p value) |
|--------------------------------|-------------|------------------------------------|-------------------------------|---------------------------------------|--|
| Vaccine delivery preference    | Injection   | 10 (6.7)                           |                               | 198 (65.1)                            | ref  |
|                                | Nasal spray | 140 (93.3)                         |                               | 106 (34.9)                            | 26.15 (11.51, 59.41, p < 0.0001)*                    |
|                                | Levels      | Preferred injections n = 10        | Preferred nasal spray n = 140 |                                       | Two-tailed Fisher's exact test p value               |
| Less distressing               | No          | 7 (70.0)                           | 3 (2.1)                       | –                                     | ref  |
|                                | Yes         | 3 (30.0)                           | 137 (97.9)                    | –                                     | <0.001*  |
| Safer or equally safe          | No          | 8 (80.0)                           | 2 (1.4)                       | –                                     | ref  |
|                                | Yes         | 2 (20.0)                           | 138 (98.6)                    | –                                     | <0.001*  |
| Easier or equally easy to give | No          | 8 (80.0)                           | 1 (0.7)                       | –                                     | ref  |
|                                | Yes         | 2 (20.0)                           | 139 (99.3)                    | –                                     | <0.001*  |

\* Significant p < 0.05. ref = reference category used for baseline comparison.



**Fig. 1.** Respondents' unprompted reasons for answers given. (A) Respondents' unprompted reasons for preferring nasal spray by group (exposure n = 140, control n = 106). (B) Respondents' unprompted reasons for preferring injections by group (exposure n = 10, control n = 198). (C) Respondents' unprompted reasons for accepting an influenza vaccine during pregnancy or for their children < 5 years (pregnant n = 447, children < 5 n = 448).

### 3.3. Influenza knowledge

The mean percentage influenza knowledge score was 68.0% and was not significantly different between exposure and control groups (69.2% vs. 67.4%,  $p = 0.0816$ ), although differences were seen in individual questions (Fig. 2). Participants recruited at Faji Kunda had significantly higher scores than at Sukuta (70.4% vs. 66.8%,  $p = 0.0005$ ). When comparing exposure and control participants from Sukuta, the knowledge was significantly higher in the exposure group (69.2% vs. 64.4%,  $p = 0.0001$ ). Of the control participants from Sukuta, only two (1.3%) had been asked to participate in NASIMMUNE but did not.

The responses to individual knowledge questions are shown in Fig. 2. A significantly higher proportion of exposure participants could name all three core symptoms of influenza (11.3% vs. 3.0%,  $OR = 4.19$ ,  $p = 0.001$ ), while a significantly lower proportion of exposure group participants correctly identified that there is no cure for influenza (11.3% vs. 23.0%,  $OR = 0.43$ ,  $p = 0.004$ ) but thought "medication" could cure influenza (86.7% vs. 71.1%,  $OR = 2.65$ ,  $p < 0.0001$ ). There were no significant differences between groups in any other questions.


Results of the univariate and multivariate linear regression analysis of factors that predict influenza knowledge are presented

in Table 3. In the multivariate linear regression model, there was a significantly higher level of influenza knowledge in participants who had attended Arabic school, upper school or had higher education compared to no education, exhibiting an increased score with each stepwise increase in educational level. Additionally, participants whose husbands were students showed a significantly higher knowledge score than those whose husbands had no occupation ( $\beta = 13.31$ ,  $p = 0.032$ ). The multivariate model explained a significant proportion of the variance in influenza knowledge (adjusted  $R^2 = 0.1165$ ,  $p < 0.0001$ ).

### 3.4. Influenza vaccination intent

Almost all respondents stated that they agreed or agreed strongly with the vaccine intent statements (98.5% and 98.7% for pregnancy and children < 5 respectively) and there was no difference in vaccine intent between groups (Table 1). Unprompted reasons given for intent are shown in Fig. 1C. There was no association between influenza knowledge and vaccine intent for pregnancy or children < 5 as shown in Table 3 ( $p = 0.782$  and  $p = 0.817$  respectively).

The associations between health belief concepts and vaccine intent are presented in Table 4. For children < 5, significant



**Fig. 2.** Individual influenza knowledge question responses by group. Cochran–Mantel–Haenszel odds ratios and p values shown only for questions where groups differed significantly. \*Correct statements.

associations were seen for perceived susceptibility, severity, benefit, and barriers, while for pregnancy, only perceived susceptibility, severity and worry were significantly associated with vaccine intent.

#### 4. Discussion

There are few countries with seasonal influenza vaccination in sub-Saharan Africa, with none currently using intranasal LAIV [37]. The NASIMMUNE study, ongoing in The Gambia, provided an opportunity to study perceptions of intranasal LAIV between mothers whose children had recently received LAIV and women with no experience of LAIV. In the latter group, preference for an intranasal vaccine was moderate at 34.9%, with the most commonly stated reasons for preferring an injection relating to familiarity with injections and beliefs of greater effectiveness. Our results show that in those with direct experience of LAIV, the preference for nasal spray was significantly higher, in keeping with studies in high income countries [38,39]. Women in the exposure group stated, unprompted, that nasal sprays were easier to give than injections and that they were less painful, suggesting that the LAIV given in NASIMMUNE was viewed favourably. These results indicate that a future influenza vaccination programme in The Gambia using intranasal LAIV in children <5 would be received positively, particularly if the introduction were coupled with demonstrations or educational sessions to promote their safety, ease of use and effectiveness.

The influenza knowledge questions covered a range of influenza-related topics including symptoms, transmission, severity, high-risk groups, treatment and vaccination. The educational level in The Gambia is generally low, with many women illiterate, and never attending school (21.6%). The mean knowledge scores were no different between the exposure and control groups, but a good understanding of disease and health-risk concepts was seen overall despite the low formal education level.

Due to community sensitization and possible knowledge transfer between NASIMMUNE study participants and non-study participants in Sukuta, a control group from Faji Kunda who had had no exposure to any trial information about influenza were included. Surprisingly, Faji Kunda had significantly higher knowledge than Sukuta overall. The effect was independent of education, which was broadly similar between sites, although knowledge did

increase significantly with more education at both sites. The reason for the higher scores in Faji Kunda is unclear, but health beliefs within in the two communities may differ significantly and previous studies conducted at Faji Kunda may have influenced the results. Several large vaccine trials have recently recruited children through the Faji Kunda EPI clinic, which have included health, hygiene and nutritional education as part of community sensitization. One study from The Gambia has shown that such activities effectively disseminate information throughout the community [40], while others have shown that community social networks are important to knowledge, particularly when access to external information is limited [41]. Furthermore, educational women's groups can significantly improve health outcomes in low-income settings [42,43]. Community health beliefs may, therefore, be as important as formal education in regard to influenza-related knowledge.

When comparing groups from Sukuta alone, influenza knowledge was significantly higher in the exposure group, suggesting that participation in the NASIMMUNE study may have increased knowledge about influenza and vaccination. However, when asked what can cure influenza, a significantly higher proportion of the exposure group incorrectly answered “medication”, rather than “nothing/no treatment”. This could indicate that involvement in the study led people to erroneously believe that specific anti-influenza medications are available. Children presenting with cough, fever and rhinorrhoea in the follow-up period of the NASIMMUNE study were commonly given paracetamol, which may have been mistakenly regarded as a “cure”, possibly explaining the finding. This is a reminder that clinical trials can inadvertently spread misinformation in these settings and that steps should be taken to mitigate this risk through careful provision of information during informed consent processes and throughout studies [44–47].

Influenza vaccine intent was assessed by asking participants if they would accept an influenza vaccine if it was freely available for themselves during pregnancy or for their children <5. Over 98% of participants responded that they would accept the vaccination. The high intent could be explained by the clinical trials that have been conducted at both health centres for many years, which may have led to a high level of trust in healthcare in these communities. To avoid response bias, participants were reassured during the interview that they should not feel pressured to answer in the affirmative. Nonetheless, the high influenza vaccine intent observed may be, in part, due to acquiescence bias or social

**Table 3**  
Predictors of influenza knowledge.

| Predictors   |                               | Influenza knowledge score mean % | Linear regression   |               |                     |                                    |                     |                      |
|--|-------------------------------|----------------------------------|---------------------|---------------|---------------------|------------------------------------|---------------------|----------------------|
|  |                               |                                  | Univariate analysis |               |                     | Multivariate analysis <sup>†</sup> |                     |                      |
| Continuous/discrete  |                               |                                  | $\beta$             | 95% CI        | p value             | $\beta^{\ddagger}$                 | 95% CI <sup>‡</sup> | p value <sup>‡</sup> |
| Total n = 453 (n = 1 excluded due to missing data in knowledge questions)    |                               |                                  |                     |               |                     |                                    |                     |                      |
| Age (in years)   |                               | -                                | 0.02                | -0.15, 0.19   | 0.788               | -                                  | -                   | -                    |
| Parity   |                               | -                                | -0.67               | -1.13, -0.19  | 0.006 <sup>*</sup>  | -0.16                              | -0.66, 0.35         | 0.546                |
| Number of people living in household   |                               | -                                | -0.25               | -0.66, 0.16   | 0.236               | -                                  | -                   | -                    |
| Categorical  | Levels                        |                                  |                     |               |                     |                                    |                     |                      |
| Group and site   | Sukuta (exposure)             | 69.2                             | ref                 | -             | -                   | ref                                | -                   | -                    |
|  | Sukuta (control)              | 64.4                             | -4.71               | -6.96, -2.46  | <0.001 <sup>*</sup> | -4.36                              | -6.65, -2.07        | <0.001 <sup>**</sup> |
|  | Faji Kunda (control)          | 70.4                             | 1.20                | -1.06, 3.46   | 0.298               | 1.49                               | -0.90, 3.88         | 0.220                |
| Marital status   | Never married                 | 68.0                             | ref                 | -             | -                   | -                                  | -                   | -                    |
|  | First and only wife           | 68.0                             | -0.01               | -9.13, 9.11   | 0.998               | -                                  | -                   | -                    |
|  | First wife (not only)         | 67.9                             | -0.07               | -9.72, 9.57   | 0.988               | -                                  | -                   | -                    |
|  | Second wife                   | 67.2                             | -0.76               | -10.19, 8.68  | 0.875               | -                                  | -                   | -                    |
| Educational level  | Third or fourth wife          | 71.3                             | 3.33                | -7.76, 14.42  | 0.555               | -                                  | -                   | -                    |
|  | None                          | 63.5                             | ref                 | -             | -                   | ref                                | -                   | -                    |
|  | Arabic school only            | 67.4                             | 3.88                | 0.93, 6.83    | 0.010 <sup>*</sup>  | 4.32                               | 1.23, 7.42          | 0.006 <sup>**</sup>  |
|  | Attended primary school       | 67.3                             | 3.88                | 0.48, 7.28    | 0.026 <sup>*</sup>  | 3.40                               | -0.03, 6.83         | 0.052                |
| Occupation   | Attended upper school         | 69.8                             | 6.36                | 3.99, 8.73    | <0.001 <sup>*</sup> | 6.08                               | 3.49, 8.68          | <0.001 <sup>**</sup> |
|  | Higher education              | 76.9                             | 13.45               | 7.71, 19.19   | <0.001 <sup>*</sup> | 9.71                               | 2.84, 16.59         | 0.006 <sup>**</sup>  |
|  | None                          | 67.9                             | ref                 | -             | -                   | ref                                | -                   | -                    |
|  | Student                       | 80.0                             | 12.13               | 0.46, 23.79   | 0.042 <sup>*</sup>  | 5.40                               | -6.53, 17.32        | 0.374                |
| Husband's educational level  | Self-employed (unskilled)     | 66.8                             | -1.11               | -3.83, 1.60   | 0.422               | -1.06                              | -3.76, 1.64         | 0.443                |
|  | Self-employed (skilled trade) | 67.2                             | -0.69               | -3.48, 2.09   | 0.624               | 0.04                               | -2.69, 2.77         | 0.977                |
|  | Employed (salaried)           | 70.3                             | 2.45                | -0.62, 5.52   | 0.117 <sup>*</sup>  | 0.05                               | -3.25, 3.35         | 0.977                |
|  | None                          | 67.1                             | ref                 | -             | -                   | ref                                | -                   | -                    |
| Husband's occupation   | Arabic school only            | 65.9                             | -1.24               | -5.27, 2.78   | 0.545               | -2.62                              | -6.59, 1.35         | 0.195                |
|  | Attended primary school       | 66.2                             | -0.89               | -6.79, 5.00   | 0.766               | -0.37                              | -6.02, 5.28         | 0.898                |
|  | Attended upper school         | 68.2                             | 1.05                | -2.29, 4.39   | 0.538               | -0.65                              | -3.98, 2.69         | 0.704                |
|  | Higher education              | 71.4                             | 4.23                | 0.05, 8.41    | 0.048 <sup>*</sup>  | 0.61                               | -3.73, 4.95         | 0.783                |
| Household income ( GMD per month)  | Don't know                    | 66.1                             | -1.01               | -6.08, 4.06   | 0.696               | -3.23                              | -8.44, 1.98         | 0.223                |
|  | None                          | 65.8                             | ref                 | -             | -                   | ref                                | -                   | -                    |
|  | Student                       | 77.8                             | 12.00               | -0.78, 24.78  | 0.066 <sup>*</sup>  | 13.31                              | 1.17, 25.46         | 0.032 <sup>**</sup>  |
|  | Self-employed (unskilled)     | 70.0                             | 4.22                | -5.54, 13.99  | 0.396               | 5.19                               | -4.32, 14.70        | 0.284                |
|  | Self-employed (skilled trade) | 67.7                             | 1.88                | -3.57, 7.34   | 0.498               | 3.83                               | -1.39, 9.05         | 0.150                |
|  | Employed (salaried)           | 68.1                             | 2.29                | -3.08, 7.66   | 0.403               | 3.23                               | -1.92, 8.37         | 0.218                |
| Vaccination intent   | Don't know <sup>†</sup>       | 68.9                             | 3.11                | -4.27, 10.49  | 0.408               | 5.31                               | -2.08, 12.70        | 0.159                |
|  | GMD 500-GMD 4,999             | 66.5                             | ref                 | -             | -                   | ref                                | -                   | -                    |
|  | GMD 5,000-GMD 9,999           | 68.2                             | 1.59                | -1.23, 4.40   | 0.269               | 1.09                               | -1.64, 3.82         | 0.432                |
|  | GMD >10,000                   | 69.2                             | 2.68                | -0.46, 5.82   | 0.094 <sup>*</sup>  | 1.72                               | -1.36, 4.79         | 0.272                |
| "I would get a flu vaccine in pregnancy if available for free"               | Don't know/unwilling to say   | 67.6                             | 1.15                | -2.00, 4.30   | 0.474               | 1.10                               | -1.98, 4.18         | 0.481                |
|  | Disagree                      | 66.7                             | ref                 | -             | -                   | -                                  | -                   | -                    |
| "I would get a flu vaccine for my child under 5, every year, if it was free" | Agree                         | 67.9                             | 1.28                | -7.80, 10.36  | 0.782               | -                                  | -                   | -                    |
|  | Disagree                      | 66.7                             | ref                 | -             | -                   | -                                  | -                   | -                    |
|  | Agree                         | 68.0                             | 1.37                | -10.27, 13.00 | 0.817               | -                                  | -                   | -                    |

<sup>†</sup> Predictors included in the multivariate linear regression model (p values < 0.2): parity, group/site, educational level, occupation, husband's educational level, husband's occupation and monthly household income.

<sup>‡</sup> Adjusted values.

\* p < 0.2.

\*\* Significant p < 0.05. ref = reference category used for baseline comparison.

**Table 4**  
Health belief constructs as predictors of vaccine intent.

| Health belief construct  | Question wording   | Levels   | Vaccine intent <sup>†</sup> |             | Two-tailed Fisher's exact p value |
|--------------------------|--|----------|-----------------------------|-------------|-----------------------------------|
|                          |  |          | Disagree n (%)              | Agree n (%) |                                   |
| <i>Children &lt; 5</i>   |  |          |                             |             |                                   |
| Perceived susceptibility | “If my child under 5 had not been given the flu vaccine, I would expect them to get flu”   | Disagree | 4 (100.0)                   | 30 (6.7)    | <0.001 <sup>*</sup>               |
|                          |  | Agree    | 0 (0.0)                     | 415 (93.3)  |                                   |
| Perceived severity       | “If my child under 5 caught flu, it would be more mild than in the general public”   | Disagree | 2 (50.0)                    | 188 (42.4)  | 1.000                             |
|                          |  | Agree    | 2 (50.0)                    | 257 (57.8)  |                                   |
|                          | “If my child under 5 caught flu, they might need to be admitted to hospital”   | Disagree | 1 (25.0)                    | 4 (0.9)     | 0.044 <sup>*</sup>                |
|                          |  | Agree    | 3 (75.0)                    | 441 (99.1)  |                                   |
| Perceived benefit        | “If my child under 5 was given the flu vaccine, it would prevent them catching flu”  | Disagree | 3 (75.0)                    | 3 (0.7)     | <0.001 <sup>*</sup>               |
|                          |  | Agree    | 1 (25.0)                    | 441 (99.3)  |                                   |
| Perceived barriers       | “If my child under 5 had been given the flu vaccine, the vaccine could give them flu”  | Disagree | 2 (50.0)                    | 404 (91.0)  | 0.046 <sup>*</sup>                |
|                          |  | Agree    | 2 (50.0)                    | 40 (9.0)    |                                   |
|                          | “The flu vaccine is unsafe for children”   | Disagree | 2 (50.0)                    | 406 (92.1)  | 0.036 <sup>*</sup>                |
|                          |  | Agree    | 2 (50.0)                    | 35 (7.9)    |                                   |
| Cues to action           | “If a nurse or doctor recommended the flu vaccine during pregnancy or for my child under 5, I would agree to get it because of what they said” | Disagree | 1 (25.0)                    | 202 (45.1)  | 0.631                             |
|                          |  | Agree    | 3 (75.0)                    | 246 (54.9)  |                                   |
|                          | “If my friends or relatives recommended the flu vaccine during pregnancy or for my child under 5, I would get it because of what they said”    | Disagree | 3 (100.0)                   | 336 (75.3)  | 1.000                             |
|                          |  | Agree    | 0 (0.0)                     | 110 (24.7)  |                                   |
| Worry                    | “If my child under 5 hadn't been given the flu vaccine, I would worry about them getting flu”  | Disagree | 1 (25.0)                    | 13 (2.9)    | 0.119                             |
|                          |  | Agree    | 3 (75.0)                    | 434 (97.1)  |                                   |
| Anticipated regret       | “If I refused to get the flu vaccine for my child under 5, but then they got sick with flu, I would be angry with myself”                      | Disagree | 1 (25.0)                    | 14 (3.1)    | 0.127                             |
|                          |  | Agree    | 3 (75.0)                    | 433 (96.9)  |                                   |
| <i>Pregnancy</i>         |  |          |                             |             |                                   |
| Perceived susceptibility | “If I was pregnant but hadn't been given the flu vaccine, I would expect to get flu”   | Disagree | 3 (50.0)                    | 30 (6.9)    | 0.006 <sup>*</sup>                |
|                          |  | Agree    | 3 (50.0)                    | 407 (93.1)  |                                   |
| Perceived severity       | “If I was pregnant and caught flu, it would be more mild than in the general public”   | Disagree | 0 (0.0)                     | 200 (45.1)  | 0.036 <sup>*</sup>                |
|                          |  | Agree    | 6 (100.0)                   | 244 (55.0)  |                                   |
| Perceived benefit        | “If I had been given a flu vaccine during pregnancy, it would prevent me catching flu”   | Disagree | 0 (0.0)                     | 5 (1.1)     | 1.000                             |
|                          |  | Agree    | 6 (100.0)                   | 440 (98.9)  |                                   |
|                          | “If I got a flu vaccine during pregnancy, it would protect my baby from getting flu in the first few months of life”                           | Disagree | 0 (0.0)                     | 5 (1.13)    | 1.000                             |
|                          |  | Agree    | 6 (100.0)                   | 439 (98.9)  |                                   |
| Perceived barriers       | “The flu vaccine is unsafe during pregnancy”   | Disagree | 4 (66.7)                    | 402 (91.6)  | 0.090                             |
|                          |  | Agree    | 2 (33.3)                    | 37 (8.4)    |                                   |
|                          | “If I got a flu vaccine during pregnancy, the vaccine could give me flu”   | Disagree | 5 (83.3)                    | 414 (93.7)  | 0.332                             |
|                          |  | Agree    | 1 (16.7)                    | 28 (6.3)    |                                   |
| Cues to action           | “If a nurse or doctor recommended the flu vaccine during pregnancy or for my child under 5, I would agree to get it because of what they said” | Disagree | 1 (16.7)                    | 203 (45.4)  | 0.230                             |
|                          |  | Agree    | 5 (83.3)                    | 244 (54.6)  |                                   |
|                          | “If my friends or relatives recommended the flu vaccine during pregnancy or for my child under 5, I would get it because of what they said”    | Disagree | 4 (66.7)                    | 337 (75.9)  | 0.635                             |
|                          |  | Agree    | 2 (33.3)                    | 107 (24.1)  |                                   |
| Worry                    | “If I was pregnant and hadn't been given the flu vaccine, I would worry about getting flu”   | Disagree | 3 (50.0)                    | 18 (4.0)    | 0.002 <sup>*</sup>                |
|                          |  | Agree    | 3 (50.0)                    | 429 (96.0)  |                                   |
| Anticipated regret       | “If I was pregnant and refused to get the flu vaccine, but then got sick with flu, I would be angry with myself”                               | Disagree | 1 (16.7)                    | 24 (5.4)    | 0.290                             |
|                          |  | Agree    | 5 (83.3)                    | 423 (94.6)  |                                   |

<sup>†</sup> Participants who answered “don't know” to health belief construct or vaccine intent questions omitted.

<sup>\*</sup> Significant p < 0.05.

desirability bias. However, the unprompted stated reasons for their answers confirmed that there is a strong belief that vaccines are safe and effective, and related to vaccines generally rather than influenza vaccines specifically. The high intent is also consistent with the high EPI coverage in The Gambia (95% for BCG and 81–99% for three Diphtheria-Pertussis-Tetanus doses) which is among the highest in Africa [9]. Neither higher influenza knowledge nor higher educational level were associated with vaccine intent. This may be due to the small numbers disagreeing with the intent statements, which also meant that the associations between health belief constructs and vaccine intent could not be quantified, but our results confirm previous findings elsewhere [19,20]. A larger

sample size would be required to determine which health belief constructs are most predictive of vaccine intent in this setting.

A key limitation to this study is that opportunistic sampling was used. The sample may therefore not be representative of the wider community, which could explain the small observed differences in socio-demographic makeup between groups. The higher educational level and income observed in the exposure group may be due to more educated and affluent people being more willing to participate in clinical trials. Future studies should use probability sampling at more sites to better represent the knowledge, attitudes and perceptions towards influenza and vaccination in the country.

Due to the high vaccine intent seen, the study was also underpowered to explore associations between intent and knowledge, so these should be interpreted cautiously. The intent questions specified that the vaccine was offered for free, so vaccine intent if the vaccine was only available for a fee remains unknown. Additionally, prior to being asked about LAIV preference over injections, they were not informed about the effectiveness of each vaccine, which may have altered their answers. Future studies could address these limitations in similar surveys, or use qualitative research methods to explore these attitudes in more depth, to gain an understanding of the underlying beliefs and motivations behind vaccine behaviour in The Gambia.

## 5. Conclusion

Willingness to undertake influenza vaccination during pregnancy, or to get annual seasonal vaccination for children <5 years if freely available was high, as was acceptability of LAIV in those with first-hand experience. Incorporation of the intranasal influenza vaccine into the childhood immunisation schedule in The Gambia in the future would likely be feasible from an acceptability perspective. Knowledge and understanding of health-related concepts surrounding influenza was generally good, though varied between communities, and was significantly related to higher educational levels. Despite reasonable health knowledge in this low-income, low-education setting, more formal education would have a positive impact on influenza and health knowledge, and potentially have wider community benefits as well. Community-based educational interventions may also be beneficial in The Gambia.

## Acknowledgements

We acknowledge the study communities and participants who were interviewed. We also acknowledge the dedicated team of field staff, the work of Isatou Ndow, and the advice of Dr. Nuredin Mohammed during statistical analysis of the study.

## Conflicts of interest

None.

## Funding

The NASIMMUNE study (NCT02972957) is funded by a Wellcome Trust Intermediate Clinical Fellowship award to TdS (110058/Z/15/Z). BK is funded by a number of MRC grants (MR/K007602/1, MC\_UP\_A900/1122). Research at the MRC Unit The Gambia is jointly funded by the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement and is also part of the EDCTP2 programme supported by the European Union. AF is funded by Cancer Research UK (C49896/A17429).

## Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.vaccine.2018.02.037>.

## References

- [1] WHO. Fact Sheet no 211. Influenza (seasonal) Geneva: World Health Organization; 2016. Available online: <<http://www.who.int/mediacentre/factsheets/fs211/en/>> [accessed on 16th Oct 2017].
- [2] Katz MA, Schoub BD, Heraud JM, Breiman RF, Njenga MK, Widdowson MA. Influenza in Africa: uncovering the epidemiology of a long-overlooked disease. *J Infect Dis* 2012;206(Suppl. 1):S1–4.
- [3] Lafond KE, Nair H, Rasooly MH, Valente F, Booy R, Rahman M, et al. Global role and burden of influenza in pediatric respiratory hospitalizations, 1982–2012: a systematic analysis. *PLoS Med* 2016;13:e1001977.
- [4] Nair H, Brooks WA, Katz M, Roca A, Berkley JA, Madhi SA, et al. Global burden of respiratory infections due to seasonal influenza in young children: a systematic review and meta-analysis. *Lancet* 2011;378:1917–30.
- [5] Meeting of the Strategic Advisory Group of Experts on immunization, April 2012 – conclusions and recommendations. *Wkly Epidemiol Rec* 2012;87:201–16.
- [6] Schoub BD, Gessner BD, Ampofo W, Cohen AL, Steffen CA. Afriflu2 – second international workshop on influenza vaccination in the African continent – 8 November 2012, Cape Town (South Africa). *Vaccine* 2013;31:3461–6.
- [7] Duque J, McMorrow ML, Cohen AL. Influenza vaccines and influenza antiviral drugs in Africa: are they available and do guidelines for their use exist? *BMC Public Health* 2014;14:41.
- [8] Lasher H. Advocacy for immunization. Geneva, GAVI: Global Alliance for Vaccines and Immunization; 2001.
- [9] Scott S, Odutola A, Mackenzie G, Fulford T, Afolabi MO, Lowe Jallow Y, et al. Coverage and timing of children's vaccination: an evaluation of the expanded programme on immunisation in The Gambia. *PLoS One* 2014;9:e107280.
- [10] Adegbola RA, Secka O, Lahai G, Lloyd-Evans N, Njie A, Usen S, et al. Elimination of Haemophilus influenzae type b (Hib) disease from The Gambia after the introduction of routine immunisation with a Hib conjugate vaccine: a prospective study. *Lancet* 2005;366:144–50.
- [11] Mackenzie GA, Hill PC, Sahito SM, Jeffries DJ, Hossain I, Bottomley C, et al. Impact of the introduction of pneumococcal conjugate vaccination on pneumonia in The Gambia: population-based surveillance and case-control studies. *Lancet Infect Dis* 2017;17:965–73.
- [12] Meeting of the Strategic Advisory Group of Experts on immunization, October 2014 – conclusions and recommendations. *Wkly Epidemiol Rec* 2014;89:561–76.
- [13] Larson HJ, Jarrett C, Schulz WS, Chaudhuri M, Zhou Y, Dube E, et al. Measuring vaccine hesitancy: the development of a survey tool. *Vaccine* 2015;33:4165–75.
- [14] Schmid P, Rauber D, Betsch C, Lidolt G, Denker ML. Barriers of influenza vaccination intention and behavior – a systematic review of influenza vaccine hesitancy, 2005–2016. *PLoS One* 2017;12:e0170550.
- [15] Madhi SA, Cutland CL, Kuwanda L, Weinberg A, Hugo A, Jones S, et al. Influenza vaccination of pregnant women and protection of their infants. *N Engl J Med* 2014;371:918–31.
- [16] Tapia MD, Sow SO, Tamboura B, Teguete I, Pasetti MF, Kodio M, et al. Maternal immunisation with trivalent inactivated influenza vaccine for prevention of influenza in infants in Mali: a prospective, active-controlled, observer-blind, randomised phase 4 trial. *Lancet Infect Dis* 2016;16:1026–35.
- [17] Becker MH. The health belief model and personal health behavior. San Francisco: Society for Public Health Education; 1974.
- [18] Rosenstock IM, Strecher VJ, Becker MH. Social learning theory and the Health Belief Model. *Health Educ Quart* 1988;15:175–83.
- [19] Chen MF, Wang RH, Schneider JK, Tsai CT, Jiang DD, Hung MN, et al. Using the Health Belief Model to understand caregiver factors influencing childhood influenza vaccinations. *J Community Health Nurs* 2011;28:29–40.
- [20] Weinstein ND, Kwitel A, McCaul KD, Magnan RE, Gerrard M, Gibbons FX. Risk perceptions: assessment and relationship to influenza vaccination. *Health Psychol* 2007;26:146–51.
- [21] Christy SM, Winger JG, Raffanello EW, Halpern LF, Danoff-Burg S, Mosher CE. The role of anticipated regret and health beliefs in HPV vaccination intentions among young adults. *J Behav Med* 2016;39:429–40.
- [22] Gerend MA, Shepherd JE. Predicting human papillomavirus vaccine uptake in young adult women: comparing the health belief model and theory of planned behavior. *Ann Behav Med* 2012;44:171–80.
- [23] Bowyer HL, Forster AS, Marlow LA, Waller J. Predicting human papillomavirus vaccination behaviour among adolescent girls in England: results from a prospective survey. *J Fam Plann Reprod Health Care* 2014;40:14–22.
- [24] Livi S, Zeri F, Baroni R. Health beliefs affect the correct replacement of daily disposable contact lenses: predicting compliance with the Health Belief Model and the Theory of Planned Behaviour. *Cont Lens Anterior Eye* 2017;40:25–32.
- [25] Tucker Edmonds BM, Coleman J, Armstrong K, Shea JA. Risk perceptions, worry, or distrust: what drives pregnant women's decisions to accept the H1N1 vaccine? *Matern Child Health J* 2011;15:1203–9.
- [26] Chapman GB, Coups EJ. Emotions and preventive health behavior: worry, regret, and influenza vaccination. *Health Psychol* 2006;25:82–90.
- [27] Kwong EW, Pang SM, Choi PP, Wong TK. Influenza vaccine preference and uptake among older people in nine countries. *J Adv Nurs* 2010;66:2297–308.
- [28] Tuohetamu S, Pang M, Nuer X, Mahemuti Mohemaiti P, Qin Y, et al. The knowledge, attitudes and practices on influenza among medical college students in Northwest China. *Hum Vaccin Immunother* 2017;13:1688–92.
- [29] Ditsungnoen D, Greenbaum A, Praphasiri P, Dawood FS, Thompson MG, Yoocharoen P, et al. Knowledge, attitudes and beliefs related to seasonal influenza vaccine among pregnant women in Thailand. *Vaccine* 2016;34:2141–6.
- [30] Wong KK, Cohen AL, Norris SA, Martinson NA, von Mollendorf C, Tempia S, et al. Knowledge, attitudes, and practices about influenza illness and



- vaccination: a cross-sectional survey in two South African communities. *Influenza Other Respir Viruses* 2016;10:421–8.
- [31] James PB, Rehman IU, Bah AJ, Lahai M, Cole CP, Khan TM. An assessment of healthcare professionals' knowledge about and attitude towards influenza vaccination in Freetown Sierra Leone: a cross-sectional study. *BMC Public Health* 2017;17:692.
- [32] Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap) – a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009;42:377–81.
- [33] Khun M, Heng C, Md HO, Kasuya H, Sakamoto J. Knowledge, attitudes and practices towards avian influenza A (H5N1) among Cambodian women: a cross-sectional study. *Asian Pac J Trop Med* 2012;5:727–34.
- [34] Oria PA, Arunga G, Lebo E, Wong JM, Emukule G, Muthoka P, et al. Assessing parents' knowledge and attitudes towards seasonal influenza vaccination of children before and after a seasonal influenza vaccination effectiveness study in low-income urban and rural Kenya, 2010–2011. *BMC Public Health* 2013;13:391.
- [35] Yudin MH, Salaripour M, Sgro MD. Pregnant women's knowledge of influenza and the use and safety of the influenza vaccine during pregnancy. *J Obstet Gynaecol Can* 2009;31:120–5.
- [36] Maldonado G, Greenland S. Simulation study of confounder-selection strategies. *Am J Epidemiol* 1993;138:923–36.
- [37] Ortiz JR, Perut M, Dumolard L, Wijesinghe PR, Jorgensen P, Ropero AM, et al. A global review of national influenza immunization policies: analysis of the 2014 WHO/UNICEF Joint Reporting Form on immunization. *Vaccine* 2016;34:5400–5.
- [38] Flood EM, Ryan KJ, Rousculp MD, Beusterien KM, Block SL, Hall MC, et al. A survey of children's preferences for influenza vaccine attributes. *Vaccine* 2011;29:4334–40.
- [39] Dube E, Gagnon D, Kiely M, Boulianne N, Landry M. Acceptability of live attenuated influenza vaccine by vaccine providers in Quebec, Canada. *Hum Vaccin Immunother* 2015;11:956–60.
- [40] Dierickx S, O'Neill S, Gryseels C, Immaculate Anyango E, Bannister-Tyrrell M, Okebe J, et al. Community sensitization and decision-making for trial participation: a mixed-methods study from The Gambia. *Dev World Bioeth* 2017.
- [41] Ackerson K, Zielinski R. Factors influencing use of family planning in women living in crisis affected areas of Sub-Saharan Africa: a review of the literature. *Midwifery* 2017;54:35–60.
- [42] Nair N, Tripathy P, Sachdev HS, Pradhan H, Bhattacharyya S, Gope R, et al. Effect of participatory women's groups and counselling through home visits on children's linear growth in rural eastern India (CARING trial): a cluster-randomised controlled trial. *Lancet Glob Health* 2017;5:e1004–16.
- [43] Tripathy P, Nair N, Sinha R, Rath S, Gope RK, Rath S, et al. Effect of participatory women's groups facilitated by Accredited Social Health Activists on birth outcomes in rural eastern India: a cluster-randomised controlled trial. *Lancet Glob Health* 2016;4:e119–28.
- [44] Carvalho AA, Costa LR. Mothers' perceptions of their child's enrollment in a randomized clinical trial: poor understanding, vulnerability and contradictory feelings. *BMC Med Ethics* 2013;14:52.
- [45] Idoko OT, Diallo A, Sow SO, Hodgson A, Akinsola A, Diarra B, et al. Community perspectives associated with the African PsA-TT (MenAfriVac) vaccine trials. *Clin Infect Dis* 2015;61(Suppl. 5):S416–21.
- [46] Mboizi RB, Afolabi MO, Okoye M, Kampmann B, Roca A, Idoko OT. Recall and decay of consent information among parents of infants participating in a randomized controlled clinical trial using an audio-visual tool in The Gambia. *Hum Vaccin Immunother* 2017;13:2185–91.
- [47] Moodley K, Pather M, Myer L. Informed consent and participant perceptions of influenza vaccine trials in South Africa. *J Med Ethics* 2005;31:727–32.