


CORRECTION

Correction: Impact of a Participatory Intervention with Women's Groups on Psychological Distress among Mothers in Rural Bangladesh: Secondary Analysis of a Cluster-Randomised Controlled Trial

Kelly Clarke, Kishwar Azad, Abdul Kuddus, Sanjit Shaha, Tasmin Nahar, Bedowra Haq Aumon, Mohammed Munir Hossen, James Beard, Anthony Costello, Tanja A. J. Houweling, Audrey Prost, Edward Fottrell

There is an error in [Fig 1](#). Please see the complete, corrected [Fig 1](#) here.



OPEN ACCESS

Citation: Clarke K, Azad K, Kuddus A, Shaha S, Nahar T, Aumon BH, et al. (2017) Correction: Impact of a Participatory Intervention with Women's Groups on Psychological Distress among Mothers in Rural Bangladesh: Secondary Analysis of a Cluster-Randomised Controlled Trial. PLoS ONE 12(8): e0183203. <https://doi.org/10.1371/journal.pone.0183203>

Published: August 10, 2017

Copyright: © 2017 Clarke et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 1. Sample selection procedure for evaluation of the effect of women's groups on postpartum psychological distress. From 25,595 births and deaths ('events') over the 24 study months in 2010 and 2011 we excluded data from: 17,181 events that did not occur within the SRQ-20 data collection periods; 62 events where it was not possible to conduct an interview due to migration or refusal; 1945 events associated with mothers who were temporary residents in the study area; nine events associated with a maternal death; 100 events associated with mothers who had previously delivered during the SRQ-20 data collection periods, either because of multiple births or through repeated births; 23 mothers due to missing SRQ-20 data. In total, 6275 mothers were included in the final sample.

<https://doi.org/10.1371/journal.pone.0183203.g001>

There are errors in [Table 1](#). Please see the complete, correct [Table 1](#) here.

Table 1. Respondent characteristics for intervention and control areas (with and without tea garden residents) at baseline and during the SRQ-20 data collection period.

	BASELINE (JANUARY—DECEMBER 2008)			SRQ-20 DATA COLLECTION (JAN—APRIL 2010 & 2011)		
	Intervention (N = 5027)	Control Excluding tea-garden residents (N = 5013)	Control Including tea-garden residents (N = 5571)	Intervention (N = 2967)	Control Excluding tea garden residents (N = 2823)	Control Including tea garden residents (N = 3308)
Age (years)						
Mean (SD)	24.7 (5.6)	24.6 (5.4)	24.6 (5.4)	24.5 (5.3)	24.8 (5.4)	24.8 (5.4)
Age at first pregnancy (years)						
Mean (SD)	18.4 (2.8)	18.5 (2.7)	18.6 (2.6)	18.4 (2.6)	18.6 (2.6)	18.7 (2.6)
Gravidity						
Mean (SD)	2.7 (1.7)	2.5 (1.6)	2.5 (1.6)	2.5 (1.6)	2.5 (1.6)	2.5 (1.6)
Religion						
Islam (%)	4539 (90.3)	4523 (90.2)	4605 (82.7)	2626 (88.5)	2551 (90.4)	2594 (78.5)
Hindu (%)	479 (9.5)	487 (9.7)	959 (17.2)	341 (11.5)	270 (9.6)	708 (21.4)
Other (%)	9 (0.2)	3 (0.1)	7 (0.1)	0 (0)	1 (0.0)	1 (0.0)
Education						
Never went to school (%)	1261 (25.1)	1091 (21.8)	1437 (25.8)	617 (20.8)	529 (18.7)	810 (24.5)
Primary education (%)	1795 (35.7)	1609 (32.1)	1727 (31.0)	1129 (38.1)	942 (33.4)	1059 (32.0)
Secondary or above (%)	1971 (39.2)	2313 (46.1)	2407 (43.2)	1221 (41.2)	1352 (47.9)	1439 (43.5)
Household assets**						
None (%)	1840 (36.6)	1635 (32.6)	1939 (34.8)	630 (21.2)	534 (18.9)	737 (22.3)
One (%)	1056 (21.0)	1017 (20.3)	1087 (19.5)	802 (27.0)	708 (25.1)	801 (24.2)
Two (%)	722 (14.4)	636 (12.7)	697 (12.5)	438 (14.8)	449 (15.9)	511 (15.5)
Three or more (%)	1409 (28.0)	1723 (34.4)	1846 (33.2)	1097 (37.0)	1132 (40.1)	1259 (38.1)
Perinatal health care utilization						
Facility deliveries (%)	943 (18.8)	1065 (21.3)	1114 (20.0)	820 (27.6)	796 (28.2)	862 (26.1)
4 or more ANC check-ups by formal provider (%)	546 (10.9)	648 (12.9)	676 (12.1)	541 (18.2)	386 (13.7)	398 (12.0)
Counts of births and deaths						
Number of births	4965	4930	5485	2876	2749	3216
Neonatal mortality rate per 1000 livebirths	38.3	35.3	37.2	14.6	28.0	28.3

** Assets included in the variable: radio, electric fan, television, fridge, mobile phone, bicycle, generator and electricity

<https://doi.org/10.1371/journal.pone.0183203.t001>

There are errors in Table 2. Please see the complete, correct Table 2 here.

Table 2. Women’s group participation rates among respondents screened with the SRQ-20.

	Proportion of women in intervention clusters (%) N = 2967	Proportion of women in intervention clusters that attended groups (%) N = 1037
Age group		
< 20 years	505 (17.0)	137 (13.2)
20–29 years	1889 (63.7)	676 (65.2)
30–39 years	549 (18.5)	220 (21.2)
40 years or older	24 (0.8)	4 (0.4)
Religion		
Muslim	2626 (88.5)	933 (90.0)
Hindu	341 (11.5)	104 (10.0)
Other	0 (0)	0 (0)
Maternal education		
Never went to school/less than 1 year	617 (20.8)	231 (22.3)
Primary education/non-formal	1129 (38.1)	441 (42.5)
Secondary or above	1221 (41.2)	365 (35.2)
Household assets		
None (%)	630 (21.2)	249 (24.0)
One (%)	802 (27.0)	265 (25.6)
Two (%)	438 (14.8)	171 (16.5)
Three or more (%)	1097 (37.0)	352 (33.9)
Primigravid	966 (32.6)	253 (24.4)
Experienced a neonatal death or stillbirth	133 (4.5)	53 (5.1)
SRQ-20 score >6	1028 (34.7)	351 (33.9)

<https://doi.org/10.1371/journal.pone.0183203.t002>

There are errors in [Table 3](#). Please see the complete, correct [Table 3](#) here.

Table 3. Summary measures of postpartum psychological distress (SRQ-20 score >6) in intervention and control areas (with and without tea garden residents), by year.

	Mothers (N)	Overall prevalence	District prevalence			Mean cluster score	Mean cluster prevalence	Mean (SD)	Median (IQR)
			Bogra	Faridpur	Moulvibazar				
2010 and 2011									
Intervention	2967	1028 (34.7)	320 (34.7)	296 (27.9)	412 (41.9)	5.2 (1.8)	34.8 (16.8)	5.2 (4.3)	4 (2–8)
Control excluding tea garden residents	2823	1014 (35.9)	286 (30.2)	381 (34.3)	347 (45.4)	5.3 (1.4)	33.7 (14.0)	5.5 (4.5)	4 (2–9)
Control including tea garden residents	3308	1180 (35.7)	286 (30.2)	381 (34.3)	513 (41.1)	5.3 (1.2)	34.3 (12.2)	5.4 (4.6)	4 (2–8)
2010, 2011 and 2012									
Intervention	4260	1354 (31.8)	437 (32.8)	378 (25.4)	539 (37.5)	4.9 (1.7)	32.1 (16.3)	4.9 (4.1)	4 (2–8)
Control excluding tea garden residents	4072	1420 (34.9)	464 (33.5)	484 (30.5)	472 (42.8)	5.2 (1.4)	33.1 (15.0)	5.4 (4.5)	4 (2–8)
Control including tea garden residents	4760	1631 (34.3)	464 (33.5)	484 (30.5)	683 (38.1)	5.2 (1.3)	33.4 (13.1)	5.3 (4.5)	4 (2–8)
2010									
Intervention	1507	624 (41.4)	178 (39.7)	183 (33.7)	263 (51.0)	5.9 (1.8)	41.3 (15.7)	5.9 (4.6)	5 (2–9)
Control excluding tea garden residents	1389	566 (40.8)	143 (32.2)	219 (39.8)	204 (51.7)	5.7 (1.5)	38.5 (15.2)	5.9 (4.6)	5 (2–9)
Control including tea garden residents	1644	647 (39.4)	143 (32.2)	219 (39.8)	285 (43.9)	5.7 (1.3)	38.0 (12.7)	5.8 (4.7)	5 (2–9)
2011									
Intervention	1460	404 (27.7)	142 (30.0)	113 (21.8)	149 (31.8)	4.5 (2.2)	28.1 (21.4)	4.5 (3.9)	4 (1–7)
Control excluding tea garden residents	1434	448 (31.2)	143 (28.4)	162 (28.8)	143 (38.8)	4.8 (1.5)	28.5 (15.1)	5.1 (4.4)	4 (2–8)
Control including tea garden residents	1664	533 (32.0)	143 (28.4)	162 (28.8)	228 (38.1)	4.9 (1.3)	30.3 (14.4)	5.1 (4.4)	4 (1.5–8)
2012									
Intervention	1293	326 (25.2)	117 (28.5)	82 (19.1)	127 (28.1)	4.3 (1.7)	25.6 (16.3)	4.2 (3.6)	3 (1–7)
Control excluding tea garden residents	1249	406 (32.5)	178 (40.7)	103 (21.8)	125 (36.9)	5.1 (2.0)	31.7 (20.8)	5.2 (4.3)	4 (2–8)
Control including tea garden residents	1452	451 (31.1)	178 (40.7)	103 (21.8)	170 (31.4)	5.0 (1.9)	31.2 (19.8)	5.0 (4.3)	4 (2–8)

<https://doi.org/10.1371/journal.pone.0183203.t003>

There is an error in the third sentence of the Statistical Analysis subsection of the Methods. The correct sentence is: We used an SRQ-20 score >6 to report the prevalence of psychological distress because a previous study in urban Bangladesh found that this score discriminated best for psychiatric disorders, though the sensitivity (62%) and specificity (69%) were quite low [28].

There is an error in the second sentence of the Results section. The correct sentence is: In total, data were available for 25,595 births and deaths over the study months in 2010 and 2011.

There is an error in the first sentence of the Impact of the women’s groups on postpartum psychological distress section of the Results. The correct sentence is: Using a cut-off score of >6, the overall prevalence of postpartum psychological distress was 35% (2208/6275) (Table 3).

Reference

1. Clarke K, Azad K, Kuddus A, Shaha S, Nahar T, Aumon BH, et al. (2014) Impact of a Participatory Intervention with Women’s Groups on Psychological Distress among Mothers in Rural Bangladesh: Secondary Analysis of a Cluster-Randomised Controlled Trial. PLoS ONE 9(10): e110697. <https://doi.org/10.1371/journal.pone.0110697> PMID: 25329470