

Apolipoprotein E genotypes and longevity across dementia disorders

Tobias Skillbäck^{a,b} MD, Ronald Lautner^{a,b} MD, Niklas Mattsson^{c,d} MD, PhD, Jonathan M. Schott^e MD, PhD, Katarina Nägga^c MD, PhD, Lena Kilander^f MD, PhD, Anders Wimo^{g,h} MD, PhD, Bengt Winblad^{g,i} MD, Maria Eriksdotter^{g,i} MD, PhD, Kaj Blennow^{a,b} MD, PhD, Henrik Zetterberg^{a,b,j} MD, PhD

^aInstitute of Neuroscience and Physiology, Department of Neurochemistry, the Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

^bClinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

^cClinical Memory Research Unit, Department of Clinical Sciences Malmö, Lund University, Simrisbanvägen 14, Malmö 21224, Sweden.

^dDepartment of Neurology, Skåne University Hospital, Getingevägen 4, Lund 22241, Sweden.

^eDementia Research Centre, UCL Institute of Neurology, London, UK

^fDepartment of Public Health and Caring Sciences/Geriatrics, Uppsala University, Uppsala, Sweden

^gDepartment of Neurobiology, Care Sciences, and Society (NVS), Center for Alzheimer Research, Division of Neurogeriatrics, Karolinska Institutet, Huddinge, Sweden

^hCentre for Research & Development Uppsala University/County Council of Gävleborg, Gävle, Sweden

ⁱDepartment Geriatric medicine, Karolinska university hospital, Huddinge, Sweden

^jDepartment of Molecular Neuroscience, UCL Institute of Neurology, Queen Square, London, UK

Word count: 3049

Corresponding author: Tobias Skillbäck, Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital/Mölndal, S-431 80 Mölndal, Sweden. Telephone: +46 707-909478. E-mail: tobias.skillback@neuro.gu.se

Competing interests: KB and HZ are co-founders of Brain Biomarker Solutions in Gothenburg AB, a GU Venture-based platform company at the University of Gothenburg. KB reports personal fees from Innogenetics and Roche outside of the submitted work.

Keywords: Apolipoprotein E, Alzheimer's disease, dementia with Lewy bodies, vascular dementia, frontotemporal dementia, Parkinson's disease dementia, Creutzfeldt-Jakob disease.

Abstract

INTRODUCTION: The $\epsilon 4$ allele of the apolipoprotein E (*APOE*) gene is a prominent risk factor for Alzheimer's disease (AD), but its implication in other dementias is less well studied.

METHODS: We used a dataset on 2858 subjects (1098 AD, 260 vascular dementia [VaD], 145 mixed AD and VaD [MIX], 90 other dementia diagnoses, 1265 controls) to examine the association of *APOE* polymorphisms with clinical dementia diagnoses, biomarker profiles and longevity.

RESULTS: The $\epsilon 4$ allele was associated with reduced longevity as $\epsilon 4$ vs $\epsilon 3$ homozygotes lived on average 2.6 years shorter ($p=.006$). In AD, $\epsilon 4$ carriers lived 1.0 years shorter than non-carriers ($p=.028$). The $\epsilon 4$ allele was more prevalent in AD, MIX and VaD patients compared to controls, but not in other dementia disorders.

DISCUSSION: The *APOE* $\epsilon 4$ allele is influential in AD, but might also be of importance in MIX and VaD; diseases in which concomitant AD pathology is common.

1. Introduction

Apolipoprotein E (apoE) is a protein involved in lipid metabolism and transport of cholesterol to neurons [1, 2]. There are three common alleles of the *APOE* gene ($\epsilon 2$, $\epsilon 3$ and $\epsilon 4$) that encode three isoforms of the protein (E2, E3 and E4) [3]. Being an $\epsilon 4$ allele carrier is the most prominent genetic risk factor for late onset Alzheimer's disease (AD), and has a reported prevalence of 24% in the Nordic population [4, 5]. Being a homozygous $\epsilon 4$ carrier increases the risk of developing sporadic AD by a factor 10-20 compared to being an $\epsilon 3$ homozygote [6-8]. *APOE* $\epsilon 4$ has also been implicated in other settings including atherosclerosis [9], depression [10], accelerated progression in multiple sclerosis [11] and brain atrophy in HIV patients [12]. The $\epsilon 2$ allele may be a protective trait, lowering the risk of developing AD [13], as compared to $\epsilon 3$, the most common allele. *APOE* has also been targeted for examination in longevity studies, where the $\epsilon 4$ allele has been found associated with early mortality [14].

The association of the *APOE* $\epsilon 2/\epsilon 3/\epsilon 4$ polymorphism with dementias other than AD is less clear. In vascular dementia (VaD), studies on the role of the *APOE* polymorphism has yielded conflicting results, possibly due to lack of consensus on the criteria required for diagnosis and the high overlap of VaD with other dementia disorders, primarily AD [15]. In meta-analysis, an increased VaD risk has been identified in *APOE* $\epsilon 4$ carriers [16], a result that has been confirmed by population-based data [17]. Dementia with Lewy bodies (DLB) and Parkinson's disease dementia (PDD) patients share clinical and biochemical features; however, the impact of *APOE* genotype in these conditions might diverge. Evidence suggests an association of increased risk of DLB with a *APOE* $\epsilon 4$ carrier status [18, 19]. By contrast, the $\epsilon 2$ allele, but not $\epsilon 3$ or $\epsilon 4$, has been associated with increased risk of Parkinson's disease [20], although associations between $\epsilon 4$ and PDD has been reported [21]. Studies of *APOE* in frontotemporal dementia (FTD) have again produced conflicting results, but, in meta-analysis, carriage of the

$\epsilon 4$ allele was found to be associated with a slightly increased risk of disease [22]. In amyotrophic lateral sclerosis, another neurodegenerative disease which commonly overlaps with FTD, evidence suggests that $\epsilon 4$ carriage is associated with earlier onset and faster disease progression [23, 24].

In this study, we explored the prevalence of the *APOE* genotypes across a wide array of dementia disorders in a large dataset drawn from the Swedish dementia registry (Svedem) and the Swedish mortality registry. We hypothesized that (i) *APOE* $\epsilon 4$ would be most prevalent, and $\epsilon 2$ least prevalent in AD, (ii) the $\epsilon 4$ allele would be associated with AD-like CSF biomarker profiles in AD and in other neurodegenerative diseases, and (iii) the $\epsilon 4$ allele would be associated with shorter survival.

2. Methods

2.1 Subject cohort

Four sources of information were joined to prepare the dataset used in this study.

The first source was of a complete set of archived data on all clinical practice *APOE* genotype analyses and CSF A β 42, T-tau and P-tau measurements made at the Sahlgrenska University Hospital, Sweden, from January 2002 to June 2012 extracted from the laboratory database.

The second source of information was the Swedish mortality registry, a national resource maintained by the Swedish National Board of Health and Welfare, keeping complete records on all deaths in Sweden, including causes of death as established by the physician issuing the death certificate. This database was queried for underlying cause (i.e. diagnosis codes) and date of death for the patients in our dataset, and information on 1166 records was retrieved.

The third data source was Svedem (the Swedish dementia registry) an initiative started in 2007, with an aim to improve the quality of diagnostics, treatment and care in dementia by collecting clinical information on dementia patients in Sweden [25, 26]. From this data source information on 557 patients was fetched including clinical diagnosis coded as one of nine preset options: early onset AD (EAD, < 65 years of age), late onset AD (LAD, > 65 years of age), FTD, dementia with Lewy bodies (DLB), Parkinson's with dementia (PDD), VaD, mixed AD and vascular dementia (MIX), dementia not otherwise specified (dementia NOS) and a group for the collected remainders of named dementia diagnoses called "other dementias", including corticobasal syndrome, alcohol-related dementias, and other rare diagnoses. Reporting clinicians are instructed to follow diagnostic guidelines as specified in ICD-10 to secure a unified basis for diagnosis [27]. For the purposes of this study, and since we lacked diagnosis date for the patients drawn from the mortality registry, the EAD and LAD groups were merged into a joint AD group, and the NOS and "other dementias" groups were stripped out of the dataset. The study cohort was further sub-classified into biochemically positive or negative profile according to the international working group (IWG-2) [28]. The following cutoffs for pathological biomarker concentrations were used: $A\beta_{42} \leq 550$ pg/mL, $T\text{-tau} \geq 400$ pg/mL, $P\text{-tau} > 60$ pg/mL for patients < 60 years, and $P\text{-tau} > 80$ for patients ≥ 60 years [29]. Pathological concentrations of $A\beta_{42}$ and at least either T-tau or P-tau were required for a biochemical AD positive classification.

These data sources were cross-referenced using the Swedish personal identity number. Note that there was an overlap between the Svedem and the mortality registry records, where both sources were combined to retrieve data for a subset of the study cohort. Also note that date of death only was available for patients taken from the mortality registry.

The fourth source of information was a set of 1265 healthy control subjects from a previous study by Zetterberg et al. [30]. The controls were all western Swedish residents and the majority (n = 980) were selected from population-based cohort studies, while 285 were recruited by advertisements in newspapers and at senior citizen meetings.

2.2 Biochemical measurements

CSF T-tau and P-tau concentrations were measured using enzyme-linked immunosorbent assays (ELISAs) (INNOTEST hTau Ag and Phospho-tau [181P]; Fujirebio, Ghent, Belgium) as previously described [31, 32]. CSF A β 42 concentration was measured using a sandwich ELISA (INNOTEST β -amyloid[1-42]), specifically constructed to measure A β containing both the 1st and 42nd amino acids, as previously described [33]. The between-assay coefficients of variation (CV) for the T-tau and P-tau tests were 10.35 % and 10.19 % respectively (as determined by internal control samples during the entire study period).

All CSF analyses were performed in clinical routine by board-certified laboratory technicians using procedures accredited by the Swedish Board for Accreditation and Conformity Assessment (SWEDAC). Longitudinal stability in the measurements was ascertained using an elaborate system of internal quality control samples and testing of incoming reagents, and further verified in the Alzheimer's Association CSF Quality Control Program [34].

2.3 APOE genotyping

APOE (gene map locus 19q13.2) genotyping was performed in blood by minisequencing or by TaqMan Single Nucleotide Polymorphism (SNP) Genotyping. Genotypes were obtained for the two SNPs, which are used to unambiguously define ϵ 2, ϵ 3, and ϵ 4 alleles (rs7412 and rs429358).

2.4 Statistics

Age distributions in subject groups were tested with ANOVA, and sex distribution by χ^2 statistics. Allele frequency and distributions in diagnosis groups and biochemical AD subgroups were analyzed by χ^2 -statistics and group differences by Kruskal-Wallis.

Differences in life length between *APOE* genotype groups and between gene carriers and non-carriers were analyzed by ANOVA, as was analysis of differences in age of diagnosis in AD.

2.5 Ethics

The study was approved by the regional ethical committee at the University of Gothenburg.

3. Results

3.1 Dataset description

Demographics of study subjects are detailed in Table 1. The FTD subjects were significantly ($p < .001$) younger than the AD, MIX and VaD subjects. The VaD subjects were older ($p < .05$) than the AD, FTD, CJD and PDD subjects. Gender had no influence on any of the associations reported below.

3.2 *APOE* allele frequencies across diagnoses

Table 2 shows allele frequencies across diagnoses. The AD subjects had a higher $\epsilon 4$ allele frequency than the FTD ($p < .001$), VaD ($p < .001$) and healthy control ($p < .001$) groups. The MIX groups had a higher $\epsilon 4$ allele frequency than the FTD ($p < .002$), VaD ($p < .001$) and the healthy control groups ($p < .001$). The VaD group had a higher $\epsilon 4$ allele frequency than the healthy controls ($p < .001$).

The AD subjects had a lower $\epsilon 3$ frequency than the VaD ($p < .001$) and FTD ($p = .012$) subjects and the healthy controls ($p < .001$). The VaD subjects had a higher $\epsilon 3$ frequency than the MIX ($p = .002$) and the healthy controls ($p = .003$). The MIX subjects had a higher $\epsilon 3$ frequency than the FTD group ($p = .031$).

The $\epsilon 2$ allele frequency was lower in the AD and MIX groups than in the healthy controls ($p < .001$, $p = .03$).

3.3 APOE $\epsilon 2/\epsilon 3/\epsilon 4$ allele carriers across diagnoses

The AD group had a lower $\epsilon 2$ carrier proportion compared to the healthy controls ($p < .05$) (figure 1A). The $\epsilon 3$ carrier proportion was lower in the AD and MIX groups than in the healthy controls and VaD groups ($p < .05$) (figure 1B). The $\epsilon 4$ carrier proportion was higher in the AD and MIX groups than in the FTD, VaD and the healthy controls ($p < .05$), and also in VaD compared to the healthy controls ($p < .05$) (figure 1C). There were no other significant differences in $\epsilon 2$, $\epsilon 3$ and $\epsilon 4$ carrier proportions between subject groups.

3.4 APOE in the biochemically AD-like

Disregarding clinical diagnosis the $\epsilon 4$ allele frequency was higher ($p < .001$), and both the $\epsilon 2$ and $\epsilon 3$ allele frequency was lower ($p < .001$, $p < .001$) in the biochemical AD positive subjects (Figure 2A).

Figure 2 B shows the distribution of *APOE* alleles in the AD group, sub-classified into biochemical AD positive and negative subjects. The $\epsilon 4$ allele frequency was higher in the biochemical AD positive group than in the biochemical AD negative group ($p < .001$), and the $\epsilon 3$ allele frequency was lower ($p < .001$) but the $\epsilon 2$ allele frequency did not significantly differ ($p = .262$).

3.5 APOE and life length

In the full study cohort, excluding the healthy controls, the $\epsilon 4$ homozygous subjects had significantly shorter life length than the $\epsilon 3$ homozygous subjects (mean diff=2.1 years, $p=.026$). There were no other significant differences in life span between *APOE* genotype groups in the full study cohort.

When comparing life length within the individual diagnostic groups, the $\epsilon 4$ homozygous AD subjects had shorter life length those who were $\epsilon 2/\epsilon 4$ (mean diff=4.5 years, $p=.048$), $\epsilon 3/\epsilon 3$ (mean diff=2.6 years, $p=.006$) and $\epsilon 3/\epsilon 4$ (mean diff=2.3 years, $p=.014$) genotypes. There were no statistically significant differences in the other diagnostic groups.

The $\epsilon 4$ carriers in the AD group had significantly (mean diff=1.0 years, $p=.028$) shorter life length than the $\epsilon 4$ non-carriers. There were no other significant differences in mean life length between $\epsilon 4$ carriers and non-carriers in any of the diagnosis groups. In this analysis $\epsilon 2$ carriers were excluded, since $\epsilon 2$ might serve as a protective trait.

3.6 Age at diagnosis in AD

In the subgroup of patients with AD pathology (AD + MIX) where dates of diagnosis were available ($n=434$), patients who were $\epsilon 4$ homozygous were younger at the date of diagnosis than the $\epsilon 3$ homozygous patients (mean diff=3.5 years, $p=.01$), and the patients with an $\epsilon 3\epsilon 4$ configuration (mean diff=3.77 years, $p=.001$).

4. Discussion

We compared the *APOE* $\epsilon 2/\epsilon 3/\epsilon 4$ genotype frequencies across a wide array of dementia

disorders in a large cohort drawn from Swedish health registries. We found that the *APOE* $\epsilon 4$ allele was most common in patients with AD, and that the $\epsilon 4$ allele was associated with a shorter lifespan and a more distinguished AD-like biomarker pattern in AD.

Being an *APOE* $\epsilon 4$ carrier is a well-established and major genetic risk factor for AD [3, 8], and this was further corroborated in this study as the $\epsilon 4$ allele frequency was higher in the AD and MIX subjects than in any other group. The frequency of $\epsilon 4$ homozygous subjects and the proportion of $\epsilon 4$ carriers vs. non-carriers were also highest in AD with the MIX group a close second. Furthermore, the average lifespan of both $\epsilon 4$ carriers and homozygous $\epsilon 4$ subjects in AD was significantly shorter than non-carriers'. Previous studies have found evidence of the $\epsilon 2$ allele being associated with a longer lifespan [35], and that the $\epsilon 4$ allele is linked to a shorter life expectancy [36]; however, although the $\epsilon 4$ allele's link to poor survival in AD might be indirectly implied, it has not been thoroughly investigated [37]. The shorter lifespans of the $\epsilon 4$ carriers might, at least in part, be attributed to an earlier onset of disease, as we found $\epsilon 4$ homozygotes to be younger than $\epsilon 3$ homozygotes and patients with an $\epsilon 3\epsilon 4$ configuration at the date of diagnosis, corroborating previous studies [1, 38]. In fact, some studies have indicated that *APOE* might be a timing gene, rather than merely a risk gene, by showing that although disease onset is brought forward by being an $\epsilon 4$ carrier, the lifetime susceptibility of the disease might be unaffected [39]. We also found that the $\epsilon 4$ allele was associated with a biochemical AD-like profile, corroborating previous studies [40, 41].

In the other diagnostic groups, we found that the VaD subjects had a significantly higher $\epsilon 4$ allele frequency than the healthy controls, as previously shown in the literature [42].

However, these results should be interpreted with caution as AD and VaD are clinically hard to distinguish and concomitant AD pathology is common in VaD [43], as was further corroborated by the correlations between an biochemical AD positive biomarker profile and

the $\epsilon 4$ allele in this study [41]. The FTD subjects, also keeping with previous studies [44], closely resembled the healthy controls in *APOE* allele and genotype frequencies. The DLB and PDD groups were small, which might explain why, although the $\epsilon 4$ allele frequency was found to be higher in these groups compared to the healthy controls, this failed to reach statistical significance. These results were, however, in keeping with previous studies, as the $\epsilon 4$ allele has previously been found to be associated with a greater risk of developing DLB and PDD. The presence of $\epsilon 4$ has also been related to a greater amount of senile plaques and neurofibrillary tangles in these diseases, suggesting that this may be driven by concomitant AD, a promoting effect of α -synuclein on β -amyloid accumulation, or the other way around i.e. a promoting effect of β -amyloid on α -synuclein accumulation [21, 45].

We studied only a small number of patients with CJD; whilst these individuals had had a lower $\epsilon 2$ and higher $\epsilon 4$ allele frequency than the healthy controls, no significant differences between them and the other groups were found, likely reflecting the small sample size. These results are, however, keeping with previous studies. In meta-analysis an association between the $\epsilon 4$ allele and CJD has been shown [46], and synergistic effects between *APOE* $\epsilon 4$ and the *PRNP* gene, the most prominent risk factor of CJD, has been suggested in both AD and CJD [47].

The main strength of this study was the large study population size and the diversity of dementia disorders represented in it, but there are also a number of limitations to this study. The main one was that most *APOE* genotypes analyzed in this study were determined in clinical routine, introducing a risk of circular reasoning as the results of the genotyping might have influenced the physician's diagnostic formulation. Another limitation is that the majority of patients were diagnosed only clinically, which is known to introduce a high misdiagnosis rate, with low specificity figures (44-71%) for AD, and a large proportion (39%) of clinically

diagnosed FTD, LBD and VaD patients showing AD pathology at autopsy [48]. On the other hand, *APOE* genotype is not included in any of the diagnostic criteria for AD or any of the other dementia disorders and there are no guidelines on how to interpret *APOE* genotype results in clinical practice. Some of the diagnostic information in this study was taken from the death certificates of the included subjects, and some from Svedem. The diagnostic information taken from the death certificates might be more unreliable as the issuing physician is typically not an experienced neurologist or expert in dementia, in contrast to the physicians who report into Svedem, all of whom are specially trained in the evaluation of dementias. In the absence of autopsy confirmation, the clinical diagnosis may be less accurate in a proportion of cases. Finally, the lack of detailed genetic information on our study subjects preclude GWAS style analysis of other potential genes and SNPs that might influence the impact of the ApoE genotype in the diseases covered in this study.

5. Conclusions

The *APOE* $\epsilon 4$ allele frequency and $\epsilon 4$ carrier proportion was significantly higher in AD than in healthy controls and the $\epsilon 4$ allele was associated with a shorter lifespan and a more AD-like CSF biomarker pattern in AD. The *APOE* $\epsilon 4$ allele frequency was also higher in MIX and VaD as compared to the healthy controls. The results of this study highlight the significance *APOE* genotyping, and confirm the relationship between *APOE* $\epsilon 4$ and AD. Our finding that *APOE* $\epsilon 4$ genotype influences not only the CSF profile, but also life-span, may have clinical application not only in AD, but also in mixed dementia and VaD.

References

- [1] Riedel BC, Thompson PM, Brinton RD. Age, APOE and sex: Triad of risk of Alzheimer's disease. *The Journal of steroid biochemistry and molecular biology*. 2016;160:134-47.
- [2] Rosenberg RN, Lambracht-Washington D, Yu G, Xia W. Genomics of Alzheimer Disease: A Review. *JAMA neurology*. 2016.
- [3] Verghese PB, Castellano JM, Holtzman DM. Apolipoprotein E in Alzheimer's disease and other neurological disorders. *Lancet neurology*. 2011;10:241-52.
- [4] Skoog I, Horder H, Frandin K, Johansson L, Ostling S, Blennow K, et al. Association between APOE Genotype and Change in Physical Function in a Population-Based Swedish Cohort of Older Individuals Followed Over Four Years. *Frontiers in aging neuroscience*. 2016;8:225.
- [5] Yu JT, Tan L, Hardy J. Apolipoprotein E in Alzheimer's disease: an update. *Annual review of neuroscience*. 2014;37:79-100.
- [6] Hauser PS, Ryan RO. Impact of apolipoprotein E on Alzheimer's disease. *Current Alzheimer research*. 2013;10:809-17.
- [7] Michaelson DM. APOE epsilon4: the most prevalent yet understudied risk factor for Alzheimer's disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2014;10:861-8.
- [8] Strittmatter WJ. Medicine. Old drug, new hope for Alzheimer's disease. *Science*. 2012;335:1447-8.
- [9] Hofman A, Ott A, Breteler MM, Bots ML, Slieter AJ, van Harskamp F, et al. Atherosclerosis, apolipoprotein E, and prevalence of dementia and Alzheimer's disease in the Rotterdam Study. *Lancet*. 1997;349:151-4.
- [10] Feng F, Lu SS, Hu CY, Gong FF, Qian ZZ, Yang HY, et al. Association between apolipoprotein E gene polymorphism and depression. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. 2015;22:1232-8.
- [11] Shi J, Tu JL, Gale SD, Baxter L, Vollmer TL, Campagnolo DI, et al. APOE epsilon4 is associated with exacerbation of cognitive decline in patients with multiple sclerosis. *Cognitive and behavioral neurology : official journal of the Society for Behavioral and Cognitive Neurology*. 2011;24:128-33.
- [12] Wendelken LA, Jahanshad N, Rosen HJ, Busovaca E, Allen I, Coppola G, et al. ApoE epsilon4 is Associated with Cognition, Brain Integrity and Atrophy in HIV Over Age 60. *Journal of acquired immune deficiency syndromes (1999)*. 2016.
- [13] Serrano-Pozo A, Qian J, Monsell SE, Betensky RA, Hyman BT. APOEepsilon2 is associated with milder clinical and pathological Alzheimer disease. *Annals of neurology*. 2015;77:917-29.
- [14] Shadyab AH, LaCroix AZ. Genetic factors associated with longevity: a review of recent findings. *Ageing research reviews*. 2015;19:1-7.
- [15] Giau VV, Bagyinszky E, An SS, Kim SY. Role of apolipoprotein E in neurodegenerative diseases. *Neuropsychiatric disease and treatment*. 2015;11:1723-37.
- [16] Dwyer R, Skrobot OA, Dwyer J, Munafo M, Kehoe PG. Using Alzgene-like approaches to investigate susceptibility genes for vascular cognitive impairment. *Journal of Alzheimer's disease : JAD*. 2013;34:145-54.
- [17] Chuang YF, Hayden KM, Norton MC, Tschanz J, Breitner JC, Welsh-Bohmer KA, et al. Association between APOE epsilon4 allele and vascular dementia: The Cache County study. *Dementia and geriatric cognitive disorders*. 2010;29:248-53.
- [18] Hardy J, Crook R, Prihar G, Roberts G, Raghavan R, Perry R. Senile dementia of the Lewy body type has an apolipoprotein E epsilon 4 allele frequency intermediate between controls and Alzheimer's disease. *Neuroscience letters*. 1994;182:1-2.
- [19] Bras J, Guerreiro R, Darwent L, Parkkinen L, Ansorge O, Escott-Price V, et al. Genetic analysis implicates APOE, SNCA and suggests lysosomal dysfunction in the etiology of dementia with Lewy bodies. *Human molecular genetics*. 2014;23:6139-46.
- [20] Huang X, Chen PC, Poole C. APOE-[epsilon]2 allele associated with higher prevalence of sporadic Parkinson disease. *Neurology*. 2004;62:2198-202.

- [21] Tsuang D, Leverenz JB, Lopez OL, Hamilton RL, Bennett DA, Schneider JA, et al. APOE epsilon4 increases risk for dementia in pure synucleinopathies. *JAMA neurology*. 2013;70:223-8.
- [22] Rubino E, Vacca A, Govone F, De Martino P, Pinessi L, Rainero I. Apolipoprotein E polymorphisms in frontotemporal lobar degeneration: a meta-analysis. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2013;9:706-13.
- [23] Zetterberg H, Jacobsson J, Rosengren L, Blennow K, Andersen PM. Association of APOE with age at onset of sporadic amyotrophic lateral sclerosis. *Journal of the neurological sciences*. 2008;273:67-9.
- [24] Drory VE, Birnbaum M, Korczyn AD, Chapman J. Association of APOE epsilon4 allele with survival in amyotrophic lateral sclerosis. *Journal of the neurological sciences*. 2001;190:17-20.
- [25] Religa D, Spangberg K, Wimo A, Edlund AK, Winblad B, Eriksdotter-Jonhagen M. Dementia diagnosis differs in men and women and depends on age and dementia severity: data from SveDem, the Swedish Dementia Quality Registry. *Dementia and geriatric cognitive disorders*. 2012;33:90-5.
- [26] Wimo A, Religa D, Spangberg K, Edlund AK, Winblad B, Eriksdotter M. Costs of diagnosing dementia: results from SveDem, the Swedish Dementia Registry. *International journal of geriatric psychiatry*. 2013;28:1039-44.
- [27] Sorbi S, Hort J, Erkinjuntti T, Fladby T, Gainotti G, Gurvit H, et al. EFNS-ENS Guidelines on the diagnosis and management of disorders associated with dementia. *European journal of neurology : the official journal of the European Federation of Neurological Societies*. 2012;19:1159-79.
- [28] Dubois B, Feldman HH, Jacova C, Hampel H, Molinuevo JL, Blennow K, et al. Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. *Lancet Neurol*. 2014;13:614-29.
- [29] Rosen C, Farahmand B, Skillback T, Nagga K, Mattsson N, Kilander L, et al. Benchmarking biomarker-based criteria for Alzheimer's disease: Data from the Swedish Dementia Registry, SveDem. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2015;11:1470-9.
- [30] Zetterberg M, Landgren S, Andersson ME, Palmer MS, Gustafson DR, Skoog I, et al. Association of complement factor H Y402H gene polymorphism with Alzheimer's disease. *American journal of medical genetics Part B, Neuropsychiatric genetics : the official publication of the International Society of Psychiatric Genetics*. 2008;147b:720-6.
- [31] Blennow K, Wallin A, Agren H, Spenger C, Siegfried J, Vanmechelen E. Tau protein in cerebrospinal fluid: a biochemical marker for axonal degeneration in Alzheimer disease? *Molecular and chemical neuropathology / sponsored by the International Society for Neurochemistry and the World Federation of Neurology and research groups on neurochemistry and cerebrospinal fluid*. 1995;26:231-45.
- [32] Vanmechelen E, Vanderstichele H, Davidsson P, Van Kerschaver E, Van Der Perre B, Sjogren M, et al. Quantification of tau phosphorylated at threonine 181 in human cerebrospinal fluid: a sandwich ELISA with a synthetic phosphopeptide for standardization. *Neuroscience letters*. 2000;285:49-52.
- [33] Andreasen N, Hesse C, Davidsson P, Minthon L, Wallin A, Winblad B, et al. Cerebrospinal fluid beta-amyloid(1-42) in Alzheimer disease: differences between early- and late-onset Alzheimer disease and stability during the course of disease. *Archives of neurology*. 1999;56:673-80.
- [34] Mattsson N, Andreasson U, Persson S, Carrillo MC, Collins S, Chalbot S, et al. CSF biomarker variability in the Alzheimer's Association quality control program. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2013;9:251-61.
- [35] Garatachea N, Marin PJ, Santos-Lozano A, Sanchis-Gomar F, Emanuele E, Lucia A. The ApoE gene is related with exceptional longevity: a systematic review and meta-analysis. *Rejuvenation research*. 2015;18:3-13.
- [36] Kulminski AM, Arbeev KG, Culminskaya I, Arbeeva L, Ukraintseva SV, Stallard E, et al. Age, gender, and cancer but not neurodegenerative and cardiovascular diseases strongly modulate systemic effect of the Apolipoprotein E4 allele on lifespan. *PLoS genetics*. 2014;10:e1004141.
- [37] Corrada MM, Paganini-Hill A, Berlau DJ, Kawas CH. Apolipoprotein E genotype, dementia, and mortality in the oldest old: the 90+ Study. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2013;9:12-8.

- [38] Sando SB, Melquist S, Cannon A, Hutton ML, Sletvold O, Saltvedt I, et al. APOE ϵ 4 lowers age at onset and is a high risk factor for Alzheimer's disease; A case control study from central Norway. *BMC neurology*. 2008;8:9.
- [39] Khachaturian AS, Corcoran CD, Mayer LS, Zandi PP, Breitner JC. Apolipoprotein E epsilon4 count affects age at onset of Alzheimer disease, but not lifetime susceptibility: The Cache County Study. *Archives of general psychiatry*. 2004;61:518-24.
- [40] Leoni V. The effect of apolipoprotein E (ApoE) genotype on biomarkers of amyloidogenesis, tau pathology and neurodegeneration in Alzheimer's disease. *Clinical chemistry and laboratory medicine : CCLM / FESCC*. 2011;49:375-83.
- [41] Andreasson U, Lautner R, Schott JM, Mattsson N, Hansson O, Herukka SK, et al. CSF biomarkers for Alzheimer's pathology and the effect size of APOE varepsilon4. *Molecular psychiatry*. 2014;19:148-9.
- [42] Sun JH, Tan L, Wang HF, Tan MS, Tan L, Li JQ, et al. Genetics of Vascular Dementia: Systematic Review and Meta-Analysis. *Journal of Alzheimer's disease : JAD*. 2015;46:611-29.
- [43] Jellinger KA, Attems J. Prevalence of dementia disorders in the oldest-old: an autopsy study. *Acta neuropathologica*. 2010;119:421-33.
- [44] Verpillat P, Camuzat A, Hannequin D, Thomas-Anterion C, Puel M, Belliard S, et al. Apolipoprotein E gene in frontotemporal dementia: an association study and meta-analysis. *European journal of human genetics : EJHG*. 2002;10:399-405.
- [45] Carrillo Garcia F, Gil Neciga E, Alberca R, Garcia-Solis D, Millan J, Rodriguez Uranga JJ, et al. [Apolipoprotein E4 in dementia with Lewy bodies]. *Neurologia (Barcelona, Spain)*. 2008;23:152-6.
- [46] Wei Y, Tang Y, He W, Qu Z, Zeng J, Qin C. APOE gene polymorphisms and susceptibility to Creutzfeldt-Jakob disease. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. 2014;21:390-4.
- [47] Calero O, Bullido MJ, Clarimon J, Frank-Garcia A, Martinez-Martin P, Lleo A, et al. Genetic cross-interaction between APOE and PRNP in sporadic Alzheimer's and Creutzfeldt-Jakob diseases. *PLoS one*. 2011;6:e22090.
- [48] Beach TG, Monsell SE, Phillips LE, Kukull W. Accuracy of the clinical diagnosis of Alzheimer disease at National Institute on Aging Alzheimer Disease Centers, 2005-2010. *Journal of neuropathology and experimental neurology*. 2012;71:266-73.

Figure legends

Figure 1. Allele carrier proportions

Proportions of allele carriers vs. non-carriers in the study cohort groups. Diagnosis groups are represented on the x-axis, and percentages on the y-axis.

A. $\epsilon 2$. *Proportions of $\epsilon 2$ carriers.*

B. $\epsilon 3$. *Proportions of $\epsilon 3$ carriers.*

C. $\epsilon 4$. *Proportions of $\epsilon 4$ carriers.*

Figure 2. APOE genotypes

Proportions of APOE genotypes in biochemical AD positives vs. biochemical AD negatives. Genotypes are represented on the x-axis, and percentages on the y-axis.

A. Full study cohort. *Proportions in the full study cohort.*

B. AD group. *Proportions in the collected AD group (LAD+EAD+MIX).*

Table 1. Demographics of study cohort

Diagnosis		HC	AD	MIX	VaD	FTD	DLB	PDD	CJD
Sex	F (n)	792	654	79	107	23	3	6	5
	M (n)	473	444	66	153	26	10	15	2
Age at sampling	n	1265	1098	145	260	49	13	21	7
	Mean (SD)	73 (6)	72 (8)	75 (6)	76 (7)	64 (9)	70 (8)	69 (6)	66 (12)
	Mdn (IQR)		73 (67-78)	75 (71-79)	76 (71-81)	64 (58-72)	72 (64-76)	69 (67-74)	63 (61-78)
A β 42 [ng/L]	n	160	1077	145	254	48	13	21	6
	Mean (SD)	712 (189)	404 (164)	436 (179)	521 (221)	624 (240)	589 (277)	587 (207)	402 (198)
	Mdn (IQR)		380 (290-485)	400 (330-490)	480 (360-620)	577 (500-750)	570 (430-800)	619 (410-687)	355 (260-573)
T-tau [ng/L]	n	160	1017	144	246	48	13	21	7
	Mean (SD)	351 (155)	744 (397)	639 (322)	475 (277)	423 (188)	345 (201)	370 (185)	2844 (1460)
	Mdn (IQR)		697 (485-900)	590 (410-820)	410 (280-596)	389 (290-568)	340 (220-390)	340 (250-394)	3810 (1280-4040)
P-tau [ng/L]	n	130	618	131	163	37	13	18	5
	Mean (SD)	62 (18)	95 (40)	83 (33)	59 (29)	55 (15)	51 (24)	53 (21)	49 (25)
	Mdn (IQR)		89 (67-114)	80 (57-104)	53 (42-66)	54 (43-66)	48 (35-60)	52 (37-60)	36 (34-68)

Table 2. Allele frequencies

Alleles	ε2	ε3	ε4	χ ² p-value	ε2/ε2	ε2/ε3	ε2/ε4	ε3/ε3	ε3/ε4	ε4/ε4	χ ² p-value
HC	188 (7.4%)	1956 (77.3%)	386 (15.3%)	n.a.	10 (0.8%)	140 (11.1%)	28 (2.2%)	760 (60.1%)	296 (23.4%)	31 (2.5%)	n.a.
AD	60 (2.7%)	1195 (54.4%)	941 (42.9%)	***	0 (0.0%)	30 (2.7%)	30 (2.7%)	335 (30.5%)	495 (45.1%)	208 (18.9%)	***
MIX	8 (2.8%)	158 (54.5%)	124 (42.8%)	***	0 (0.0%)	6 (4.1%)	2 (1.4%)	38 (26.2%)	76 (52.4%)	23 (15.9%)	***
VaD	78 (5.6%)	1173 (68.5%)	549 (26.0%)	***	3 (1.2%)	20 (7.7%)	3 (1.2%)	122 (46.9%)	92 (35.4%)	20 (7.7%)	***
FTD	7 (7.1%)	71 (72.4%)	20 (20.4%)		0 (0.0%)	4 (8.2%)	3 (6.1%)	26 (53.1%)	15 (30.6%)	1 (2.0%)	
DLB	2 (7.7%)	17 (65.4%)	7 (26.9%)		0 (0.0%)	1 (7.7%)	1 (7.7%)	6 (46.2%)	4 (30.8%)	1 (7.7%)	
PDD	3 (7.1%)	28 (66.7%)	11 (26.2%)		0 (0.0%)	3 (14.3%)	0 (0.0%)	8 (38.1%)	9 (42.9%)	1 (4.8%)	
CJD	0 (0.0%)	11 (78.6%)	3 (21.4%)		0 (0.0%)	0 (0.0%)	0 (0.0%)	4 (57.1%)	3 (42.9%)	0 (0.0%)	

*** p < .001, ** p < .01, * p < .05

P-values were derived from the comparison of each patient group and controls.









