

TABLE 1. Occurrence of nephrogenic rests by histological type in unilateral Wilms tumors treated with preoperative chemotherapy

Tumor type / Rest status	Completely										Focal		Diffuse		Blastemal	
	necrotic		Epithelial		Stromal		Mixed		Regressive		anaplasia		anaplasia			
	No	%	No	%	No	%	No	%	No	%	No	%	No	%	No	%
None	25	66%	27	71%	51	58%	76	53%	144	62%	6	38%	26	62%	28	74%
Any rest	13	34%	11	29%	37	42%	68	47%	88	38%	10	63%	16	38%	10	26%
PLNR	9	24%	8	21%	7	8%	38	26%	66	28%	9	56%	14	33%	8	21%
ILNR	3	8%	2	5%	24	27%	16	11%	11	5%	0	0%	1	2%	1	3%
Both	1	3%	1	3%	6	7%	11	8%	8	3%	1	6%	1	2%	1	3%
NBL	0	0%	0	0%	0	0%	3	2%	3	1%	0	0%	0	0%	0	0%
Total	38	100%	38	100%	88	100%	144	100%	232	100%	16	100%	42	100%	38	100%

PLNR – perilobar nephrogenic rests; ILNR – intralobar nephrogenic rests; NBL - nephroblastomatosis