

TABLE 2. Occurrence of nephrogenic rests by type, gender and by age at diagnosis of patients with unilateral Wilms tumor

	PLNR		ILNR		NBL		PLNR and ILNR		Any rest		None		Overall	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%
Age at diagnosis in years														
Male														
N	68	43%	29	50%	3	50%	15	50%	115	39%	177	61%	292	46%
Median (range)	3.3	(0.4, 9.8)	1.6	(0.5, 10.0)	3.1	(2.2, 4.4)	2.3	(0.6, 6.6)	2.9	(0.4, 10.0)	3.2	(0.2, 15.9)	3.1	(0.2, 14.8)
Female														
N	91	57%	29	50%	3	50%	15	50%	138	40%	206	60%	344	54%
Median (range)	4.0	(0.2, 13.5)	2.2	(0.02, 12.1)	0.78	(0.4, 2.5)	3.2	(0.7, 5.1)	3.6	(0.02, 13.5)	3.5	(0.3, 16.6)	3.7	(0.2, 16.6)
All patients														
N	159	100%	58	100%	6	100%	30	100%	253	100%	383	100%	636	100%
Median (range)	3.7	(0.2, 13.5)	1.9	(0.02, 12.1)	2.2	(0.4, 4.4)	2.6	(0.6, 6.6)	3.2	(0.02, 13.5)	3.4	(0.2, 16.6)	3.3	(0.2, 16.6)