

Body Mass Index and The Risk of Serious Non-Aids Events and All-Cause Mortality in Treated
Hiv-Positive Individuals: D:A:D Cohort Analysis

Amit C Achhra^{1,2}, Caroline Sabin³, Lene Ryom⁴, Camilla Hatleberg⁴, Antonella d'Aminio
Monforte⁵, Stephane de Wit⁶, Andrew Phillips³, Christian Pradier⁷, Rainer Weber⁸, Peter Reiss⁹,
Wafaa El-Sadr¹⁰, Fabrice Bonnet¹¹, Amanda Mocroft³, Jens Lundgren⁴, Matthew G Law¹ for the
D:A:D study group

Affiliations

¹The Kirby Institute, UNSW Australia, Sydney, Australia

² Internal medicine, New York Presbyterian Hospital/Weill Cornell Medical College, Cornell
University, NY, USA

³ Institute for Global Health, UCL, London, United Kingdom

⁴ CHIP, Department of Infectious Diseases, Section 2100, Rigshospitalet, University of
Copenhagen, Copenhagen, Denmark

⁵ Dipartimento di Scienze della Salute, Clinica di Malattie Infettive e Tropicali, Azienda
Ospedaliera-Polo Universitario San Paolo, Milan, Italy

⁶ Division of Infectious Diseases, Saint Pierre University Hospital, Université Libre de
Bruxelles, Brussels, Belgium

⁷ Department of Public Health, Nice University Hospital, Nice, France

⁸ Division of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich, University of Zurich, Switzerland

⁹ Academic Medical Center, Dept. of Global Health and Div. of Infectious Diseases, University of Amsterdam, and HIV Monitoring Foundation, Amsterdam, The Netherlands

¹⁰ ICAP-Columbia University and Harlem Hospital, New York, USA

¹¹ Université de Bordeaux, ISPED, INSERM U1219 and CHU de Bordeaux, Bordeaux, France

Short title: BMI and risk of non-AIDS events

Conflicts of interest: None declared

Funding

Data on Adverse Events (D:A:D) Study: The D:A:D study was supported by a grant [grant number DNR126] from the Danish National Research Foundation (CHIP & PERSIMUNE); the Highly Active Antiretroviral Therapy Oversight Committee (HAARTOC), a collaborative committee with representation from academic institutions, the European Agency for the Evaluation of Medicinal Products, the United States Food and Drug Administration, the patient community, and pharmaceutical companies with licensed anti-HIV drugs in the European Union: AbbVie, Bristol-Myers Squibb, Gilead Sciences Inc., ViiV Healthcare, Merck & Co Inc. and Janssen Pharmaceuticals. Supported also by a grant from the Dutch Ministry of Health, Welfare and Sport through the Center for Infectious Disease Control of the National Institute for Public Health and the Environment to Stichting HIV Monitoring (ATHENA); by a grant from the Agence nationale de recherches sur le sida et les hépatites virales [ANRS, Action Coordonnée no.7, Cohortes] to the Aquitaine Cohort; The Australian HIV Observational Database (AHOD) is

funded as part of the Asia Pacific HIV Observational Database, a program of The Foundation for AIDS Research, amfAR, and is supported in part by a grant from the U.S. National Institutes of Health's National Institute of Allergy and Infectious Diseases (NIAID) [grant number U01-AI069907] and by unconditional grants from Merck Sharp & Dohme; Gilead Sciences; Bristol-Myers Squibb; Boehringer Ingelheim; Janssen-Cilag; ViiV Healthcare. The Kirby Institute is funded by The Australian Government Department of Health and Ageing, and is affiliated with the Faculty of Medicine, The University of New South Wales; by grants from the Fondo de Investigación Sanitaria [grant number FIS 99/0887] and Fundación para la Investigación y la Prevención del SIDA en Españã [grant number FIPSE 3171/00], to the Barcelona Antiretroviral Surveillance Study (BASS); by the National Institute of Allergy and Infectious Diseases, National Institutes of Health [grants number 5U01AI042170-10, 5U01AI046362-03], to the Terry Bein Community Programs for Clinical Research on AIDS (CPCRA); by primary funding provided by the European Union's Seventh Framework Programme for research, technological development and demonstration under EuroCoord grant agreement n° 260694 and unrestricted grants by Bristol-Myers Squibb, Janssen R&D, Merck and Co. Inc., Pfizer Inc., GlaxoSmithKline LLC, (the participation of centres from Switzerland is supported by The Swiss National Science Foundation (Grant 108787)) to the EuroSIDA study; by unrestricted educational grants of AbbVie, Bristol-Myers Squibb, Gilead Sciences, GlaxoSmithKline, Pfizer, Janssen Pharmaceuticals to the Italian Cohort Naive to Antiretrovirals (The ICONA Foundation); and financed within the framework of the Swiss HIV Cohort Study, supported by the Swiss National Science Foundation (grant #148522) and by the SHCS research foundation.

Abstract

Background

The relationship between body mass index (BMI) (weight (kg)/ height(m²)) and serious non-AIDS events is not well understood.

Methods

We followed D:A:D study participants on antiretroviral therapy from their first BMI measurement to the first occurrence of the endpoint or end of follow-up (N=41,149 followed for 295,147 person-years). The endpoints were cardiovascular disease (CVD); diabetes; non-AIDS-defining cancers (NADCs) and BMI-NADCs (cancers known to be associated with BMI in general population); and all-cause mortality. Using poisson regression models, we analysed BMI as time-updated, lagged by 1 year, and categorized at: 18.5, 23, 25, 27.5 and 30 kg/m².

Results

Participants were largely male (73%) with the mean age of 40 years (SD 9.7) and baseline median BMI of 23.3 (IQR: 21.2-25.7). Overall, BMI showed a statistically significant J-shaped relationship with the risk of all outcomes except diabetes. The relative risk (RR) for the BMI of <18.5 and >30 (95% CI) compared to 23-25, respectively, was as follows: CVD: 1.46(1.15-1.84) and 1.31(1.03-1.67); NADCs: 1.78(1.39-2.28) and 1.17(0.88-1.54); 'BMI-NADCs': 1.29(0.66-2.55) and 1.92(1.10-3.36). For all-cause mortality, there was an interaction by gender (P<0.001): RR in males: 2.47(2.12-2.89) and 1.21(0.97-1.50); and in females: 1.60(1.30-1.98) and 1.02(0.74-1.42). RR remained around 1 for intermediate categories of BMI. The risk of diabetes linearly increased with increasing BMI (P<0.001).

Conclusions

Risk of CVD, a range of cancers, and all-cause mortality increased at low BMI (<18.5) and then tended to increase only at BMI>30 with a relatively low risk at BMI of 23-25 and 25-30. High BMI was also associated with risk of diabetes.

Key words: HIV; Obesity; BMI; AIDS; non-AIDS; all-cause mortality

INTRODUCTION

Excess weight is now increasingly prevalent in HIV-positive individuals receiving antiretroviral therapy (ART).^{1,2} While body mass index, BMI (measured as weight (kg)/height (m²)) is an anthropometric measure and does not directly measure total body fat or biological markers of disease, it remains an easy and low-cost metric to screen for risk of certain conditions.³ However, in HIV-positive individuals, the relationship between BMI and various serious non-AIDS events (SNAEs) is not well studied.

In the general population, increasing BMI is associated with increased risk of various clinical outcomes including cardiovascular disease (CVD)³, diabetes mellitus (DM), several cancers such as those of gastrointestinal tract and endometrium,⁴ and all-cause mortality.⁵ Also, very low BMI has been associated with increased risk of mortality.⁵

HIV-positive individuals are unique in that they are exposed to ongoing inflammation/immune activation, ART toxicities, higher prevalence of life-style risk factors such as smoking and a higher overall risk of various SNAEs than general population.⁶ Further, BMI in this group is itself associated with immunosuppression and certain antiretroviral drugs among other factors.²

In one large cohort study BMI correlated strongly with CD4 count response to ART regardless of baseline CD4 count, with a highest CD4 count response at BMI levels of 25-30 but worse at higher or lower levels.⁷ On the other hand, weight gain following ART initiation has been shown to be harmful to cardio-metabolic health, especially in those with 'normal/overweight' weight at baseline.⁸ It is therefore possible that being HIV-infected modifies the relationship between BMI and various outcomes in this population. However, limited, if any, studies have rigorously evaluated BMI as a risk factor for non-AIDS outcomes.⁹ Better understanding of how BMI relates to SNAEs in HIV-positive individuals is therefore needed. This information will provide the key data to clinicians and the HIV community regarding the health implications of BMI—a well-known, easily and economically measured potential risk factor.

In this paper, we analysed the data from the Data Collection on Adverse Events of Anti-HIV Drugs (D:A:D) study, a large heterogeneous cohort with well-validated outcomes and available data on a wide range of risk factors, to assess the relationship between latest BMI and the subsequent risk of various SNAEs and all-cause mortality.

METHODS

Study population and follow-up

The design of the D:A:D study has been described in detail elsewhere.¹⁰ In brief, it is an observational study of >49,000 HIV-positive people from 11 cohorts from Europe, Australia, and the USA. All participants were under active follow-up in their cohorts at the time of enrolment in the study. The primary study aim was to investigate the associations between use of ART and major non-AIDS events and death. Data are collected prospectively during routine clinic visits;

the standardized data set includes information on demographic factors, AIDS-related events, known risk factors for CVD, laboratory markers for monitoring HIV infection and CVD, and ART. Clinical events are regularly monitored and endpoints are centrally adjudicated.

In this study, we included all participants in the D:A:D cohort who had initiated ART, had at least one BMI measure available and at least one year of further follow-up from study entry. Follow-up commenced from the later of cohort enrolment or ART initiation or first BMI measurement. Non-AIDS cancers were collected systematically from 2004 and were therefore analyzed from 1st Jan 2004 onwards. Follow-up ended on the first occurrence of the respective endpoint or was censored at death or, 1st February 2014 or six months after last follow-up visit. All individuals had no pre-existing diagnoses of CVD, diabetes mellitus (DM) or non-AIDS cancers (NADCs) at study initiation.

Endpoints

The endpoints of interest were: (i) first CVD event, defined as the composite of myocardial infarction (MI), sudden cardiac death or invasive cardiovascular procedure (coronary artery bypass graft, carotid endarterectomy or angioplasty), or confirmed stroke; (ii) DM, defined as fasting glucose > 7.0 mmol/L on at least two occasions or a single value of NGSP haemoglobin A1c > 6.5%, or symptoms with a random glucose > 11.1 mmol/L, or 2-h oral glucose tolerance test > 11.1 mmol/L, or use of antidiabetic drugs (see www.chip.dk for details); (iii) NADCs (other than basal or squamous cell skin cancer, pre-cancers, and relapses); (iv) 'BMI-related NADCs' - a composite outcome of cancers thought to be associated with BMI in the general population (i.e. malignancies of oesophagus, pancreas, colon and rectum, breast, endometrium, kidney, thyroid and gallbladder)^{4,11}; and finally (v) All-cause mortality.

Statistical methods

BMI was the main exposure factor analysed as a time-updated variable lagged by 1-year (i.e. there was at least 1-year time-gap between last BMI measurement and the endpoint, so as to minimize bias from reverse causation) and categorised at the following clinical cut-offs: 18.5, 23, 25, 27.5 and 30 kg/m². The choice of strata was driven by the broad WHO BMI clinical cut-offs (underweight (<18.5), normal (18.5-25), overweight (25-30) and obese (>30))³, with additional cut-offs at 23 and 27.5 to better define the relationship. In sensitivity analyses, we also categorized BMI in deciles.

We used Poisson regression to model the relationship between BMI and each endpoint. Models were adjusted for key confounders not thought to be on the causal pathway for each respective endpoint. Potential confounders were identified using directed acyclic graphs (DAGs).¹² All variables, including age and CD4 count, were analyzed as time-updated (lagged by 1-year) where possible. In case a time-updated variable was not recorded at a given visit, previous known value was carried forward. A missing variable category was created for all variables to ensure all observations were included in models.

For CVD we adjusted for race (White/ Black/ Other/Unknown), sex, mode of transmission (sex between men/ injecting drug use/ sex between men and women/ other), family history of CVD, age, smoking status (current/ past/ never/ unknown), current abacavir use, cumulative protease inhibitor (PI) and nucleoside reverse transcriptase inhibitor (NRTI) use in years, and CD4 count (categorized as ≤ 200 , 201-350, >350 cells/mm³). For CVD outcome, we also present analysis from models additionally adjusted for variables thought to be on a causal pathway as identified in DAGs. These include lipids (total, HDL and LDL cholesterol), systolic blood pressure (SBP) and incident DM.

Similarly models for DM, NADC and all-cause mortality were adjusted for respective confounders (listed at the bottom of Figure-1 in 'Results'). Models for all outcomes were additionally adjusted for calendar year and clinical cohort. Finally, we also checked for interaction between BMI and sex for all outcomes.

The following sensitivity analyses were performed: (i) We categorized BMI according to the deciles of the distribution to allow for the more detailed examination of the relationship between BMI and outcomes; (ii) We lagged the BMI and all other time-updated variables by 2 years instead of one year and also analysed first measured BMI ('baseline' BMI) (to further minimize bias from reverse causation); (ii) and for the CVD and mortality outcomes, models were additionally adjusted for calculated creatinine clearance using the Cockcroft-Gault formula as decreased renal function has been associated with these outcomes.^{13,14} Since data on serum creatinine and weight were not available for a couple of cohorts, we only analysed these variables in sensitivity analysis.

All analyses were performed using STATA version 14 (STATA Corporation, College Station TX, USA).

RESULTS

Participant characteristics

A total of 41,149 individuals with 295,147 person-years of follow-up (PYFU) for the all-cause mortality outcome were included. PYFU varied for each outcome depending on follow-up time. Participants were largely male (73%) with baseline mean age of 40 years and median (IQR) BMI of 23.3 (21.2- 25.7). Table-1 shows key characteristics by baseline BMI category. Prevalence of

smoking and injecting drug use as mode of transmission appeared to be inversely related to the baseline BMI category, while mean total cholesterol and systolic blood pressure tended to positively correlate with BMI. During follow-up, BMI was measured at the median (IQR) of 6 months (4-9 months) interval. For all key confounders (including smoking, lipid levels, systolic blood pressure, hepatitis B and C status) a vast majority ($\geq 85\%$) of individuals had data recorded at least at some point during the follow-up. About 15% of individuals had unknown smoking status throughout the follow-up.

BMI and risk of SNAEs

Table-2 shows the number of people experiencing each outcome, and the incidence rates per 1000 PYFU. Figure 1 (panels a-d) shows incidence rate ratios (IRR or relative risk) from various models for all outcomes by time-updated BMI category. The BMI category of 23-25 was chosen as the reference category based on rates shown in Table-2. Overall BMI was a significant predictor for all outcomes (overall $P < 0.05$ for the BMI variable in all models). The relationship of BMI was *J* or *U*-shaped with the risk of all SNAEs except DM as detailed below. Also, the effect of BMI on all-cause mortality tended to vary by sex (P for interaction between BMI and sex for all-cause mortality: < 0.001) as shown in Table-2 (incidence rates) and Figure 1d (relative risks), but not for all other outcomes.

There were 1398 CVD events (rate: 4.8/1000 PYFU). Compared to those with a BMI of 23-25, IRR (95% CI) for those with a BMI of ≤ 18.5 was 1.46 (1.15 to 1.84), for 18.5-23 was 1.10 (0.96 to 1.27) and for > 30 , it was 1.31 (1.03 to 1.67), in the models adjusted for known confounders. Further adjustment for variables known to be on the causal pathway between BMI and CVD (such as SBP, and lipids) attenuated the relative risk at levels > 30 but not at BMI levels ≤ 18.5 or 18.5-23 (Fig 1a).

For DM, the relationship with BMI was linear, i.e. increasing risk with increasing BMI, with relative risk nearly 3.5 times (IRR 3.39, 95% CI: 2.79 to 4.12) for BMI>30 vs BMI of 23-25 (Fig 1b).

There were 1143 NADCs at a rate of 3.9/1000 PYFU of which 184 (rate: 0.6/1000 PYFU) were BMI-cancers. The IRR (95% CI) for NADC was highest for those with a BMI of ≤ 18.5 : 1.78 (1.39 to 2.28) and for those with a BMI of 18.5-23: 1.30 (1.11-1.53) and tended to be higher among those with a BMI>30 although the 95% CI crossed 1 (1.17, 0.88 to 1.54). However, for the BMI-cancers, BMI >30 was associated with nearly twice the risk compared to BMI 23-25 (IRR 1.90, 1.11 to 1.36) (Fig 1c).

Finally, the rates and IRR (95% CI) of all-cause mortality varied by sex. For males, the J-shaped relationship with BMI was more prominent. For males with a BMI of ≤ 18.5 , the relative risk of mortality was about 2.5 times (IRR 2.47, 2.12 to 2.89) higher and for those with a BMI of 18.5-23 IRR was 1.38 (1.23 to 1.54) compared to those with a BMI of 23-25. The risk then tended to increase only in those with BMI>30: 1.21, 0.98 to 1.50). For females, a BMI <18.5 was also strongly related with the risk of mortality but at a lower relative risk compared to that in males (IRR 1.60, 1.30 to 1.98) and the risk did not increase at higher levels of BMI (fig 1d). Of note, the follow-up data for females at BMI>30 was relatively small (about 8000 PYFU).

Sensitivity analyses

Supplementary Table-S1, <http://links.lww.com/QAI/B162> shows IRRs from all models with BMI categorised using the deciles of the distribution. Results were similar for all outcomes: for all SNAEs (except diabetes), relative risk was highest in the lowest 10% of the BMI distribution (equivalent to BMI<19.5) and tended to increase from the tenth decile (equivalent of BMI>28.7).

Lagging the BMI by 2 years (instead of 1) or analysing 'baseline' BMI instead on time-updated BMI broadly showed similar results (data not shown). Finally, data on creatinine were not uniformly available for all participants. Adjustment for creatinine clearance for CVD and all-cause mortality mildly attenuated IRRs for CVD but did not have any significant effect on the overall results or conclusions (data not shown).

DISCUSSION

In this longitudinal analysis of HIV-positive individuals enrolled in the DAD cohort, we found that the relationship between BMI and the risk of SNAEs including CVD, NADCs, 'BMI-related NADCs' and all-cause mortality was non-linear. In general, low BMI (<18.5, and also <23 in case of NADC and all-cause mortality in males) was associated with the higher risk of these SNAEs. Relative risk of SNAEs then only tended to increase again at BMI levels >30 which was most prominent for CVD, 'BMI-related NADC' and all-cause mortality in men. Finally, for diabetes, the risk increased in a linear fashion with increasing BMI, with relative risk nearly 3.5 times at BMI>30 compared to BMI of 23-25. Categorising BMI according to deciles of the distribution confirmed similar findings and broadly agreed with our choice of clinical cut-offs. Results were also robust to lagging BMI by 2 years or analysing baseline BMI to further minimise reverse causality.

Traditionally studies in HIV/AIDS examined BMI in the context of AIDS in the pre- or early ART era. Only a few recent studies have carefully examined BMI in the context of SNAEs in treated HIV-positive individuals. A study by Koethe et al with about 1200 HIV-positive participants in a single site cohort found that low BMI (<20) was associated with a higher risk of

composite of SNAEs, while intermediate BMI of around 25-30 and high BMI >30 was associated with lower or similar risk to BMI of 20-25.⁹ Our findings are broadly similar to those noted in the latter study; however, we did find elevated risk of several SNAEs at BMI>30. Of note, the relative risk at BMI>30 was small for most outcomes (ranging from 1.31 for CVD, 1.21 for all-cause mortality in men) but was nearly two-fold higher for 'BMI-related NADCs'. However, the study by Koethe et al only used BMI at baseline and did not have consistent data on smoking.

The finding that low BMI is associated with higher risk of death is consistent across studies even in the general population.⁵ In a large individual-data meta-analysis of nearly 4 million adults, BMI <18.5 was associated with about 1.8 fold higher risk of mortality in men compared to BMI of 22.5-25 after adjusting for smoking (corresponding IRR in our study was 2.47).⁵ This association was previously thought to be due to reverse causality (i.e. the disease affecting the BMI). However, in our study, this association remained even after ensuring the BMI precedes the outcome by at least 1 or 2 years. Also smoking tends to be inversely related to BMI and is an important confounder to account for.^{15,16} While our models adjusted for time-updated smoking, residual confounding from smoking cannot be ruled out. We also found higher risk of CVD in individuals with low BMI. Besides possible residual confounding by smoking, it is possible that people with low BMI get less aggressive disease prevention interventions compared to those with higher BMI. Our study did not collect health prevention or health behaviour data to be able to analyse this possibility. For the cancer outcomes, it is possible that weight loss starts several years prior to the diagnosis which may explain their associations with lower BMI.

Findings on intermediate to high BMI and all-cause mortality in the general population have tended to vary, although many of these studies did not adjust for smoking or had a shorter follow-up.^{5,17} However, the large meta-analysis described above, after adjustment for smoking, found relative risk of all-cause mortality in males to be around 1 for BMI 25 to 30, but increasing to 1.47 with BMI of 30-35, and to 2.0 in those with BMI>40.⁵ In our study, risk of all-cause mortality in males was high even at BMI of 18.5-23 (considered near optimal in general population), and then only tended to increase at BMI >30 in males. Relative to those with a low BMI, individuals with intermediate BMI tend to have greater muscle mass, exercise capacity and greater fat stores which may help in surviving stressful disease situations and confer survival advantage in sicker patients.¹⁸ Indeed, a large systematic review of patients with coronary artery disease at baseline found that mortality was highest at BMI <18.5 and lowest for those deemed 'overweight/mildly obese' (BMI ranges 25-30) and the risk increasing only at severely high BMI >35.¹⁹ A cohort study evaluating mortality after coronary bypass graft surgery found similar results.²⁰

We found that the BMI above 30 was associated with elevated risk of CVD which attenuated after adjustment for blood pressure, lipids and incident diabetes. Increasing BMI was also associated with increasing risk of diabetes. This suggests that high BMI could be a useful initial screening tool in individuals where detailed CVD risk factors are not known. Finally, though high BMI >30 was not associated with the higher risk of NADCs, it was associated with higher risk of 'BMI-related NADCs' with overall relative risk of around 2 (compared to BMI 23-25). In the general population, obesity is increasingly being recognised as the risk factor for several cancers, with reasonably consistent evidence for oesophagus, pancreas, colon and rectum, breast, endometrium, kidney, thyroid and gallbladder cancers.^{4,11} While the relative risk for specific

cancers varies, we did not have enough follow-up data to examine effect of BMI on specific cancers.

Our study benefitted from data on HIV-positive females. Findings were similar for females for all outcomes, except for all-cause mortality where the magnitude of risk at extremes of BMI in females was lower compared to males. A US cohort study on HIV-infected women found low BMI as predictor of mortality but not the high BMI (>25 or >30).²¹ Our findings on females in our cohort was consistent with this study. Of note, person-year data on females was limited especially at extremes of BMI which may have limited our power to analyse events in this subgroup.

The strengths of our study include a large heterogeneous cohort, with well-validated (through established protocols) outcomes and longitudinally measured BMI. Also, we were able to account for several key confounders in a time-updated fashion, including smoking and ART agents. Our study does have some limitations however. First, we did not have enough follow-up data and events at very high BMI levels (e.g. >35), which would have helped us further clarify the risks associated with severe obesity. Also, data on other risk factors such as health behaviors (e.g. diet/exercise) or preventive interventions (e.g. cancer screening) were not available. Moreover, other anthropometric measures such as waist:hip ratio which could be more accurate or supplement BMI were not available in this study. Also, even though our study designed ensured BMI measurements preceded diagnoses of SNAEs temporally, role of reverse causality cannot completely be ruled out given chronic nature of SNAEs. Finally, we could not perform a direct comparison between HIV-positive and negative individuals to better understand how BMI interacts with HIV or ART or population demographics and behaviour in determining outcomes.

In summary, we found that low BMI (≤ 18.5 , and in some cases 18.5-23) was associated with high risk of several individual SNAEs as well as all-cause mortality. The relative risk of SNAEs and mortality did not increase at intermediate/moderately high BMI (around 23-30) and only tended to increase at BMI >30 . These findings suggest that BMI of 25-30, thought to be 'overweight/mildly obese' in general population may in fact confer some survival advantage in HIV-positive individuals. Also, these findings seem to suggest that obesity at BMI >30 is likely harmful in the long-term even in HIV-positive individuals. Future studies should assess how short-term and long-term changes in BMI relate to the risk of SNAEs. Ultimately, whether change in BMI or weight gain or loss interventions (depending on current BMI) would actually improve outcomes in this population will need trials on carefully designed behavioral and nutritional interventions.

Acknowledgements D:A:D Study group

D:A:D Participating Cohorts

Aquitaine, France; CPCRA, USA; NICE Cohort, France; ATHENA, The Netherlands; EuroSIDA, Europe; SHCS, Switzerland, AHOD, Australia; HIV-BIVUS, Sweden; St.Pierre Brussels Cohort, Belgium; BASS, Spain, The ICONA Foundation, Italy

D:A:D Steering Committee: Names marked with *, Chair with ϕ

Cohort PIs: W El-Sadr* (CPCRA), G Calvo* (BASS), F Bonnet and F Dabis* (Aquitaine), O Kirk* and A Mocroft* (EuroSIDA), M Law* (AHOD), A d'Arminio Monforte* (ICONA), L Morfeldt* (HivBIVUS), C Pradier* (Nice), P Reiss* (ATHENA), R Weber* (SHCS), S De Wit* (Brussels)

Cohort coordinators and data managers: A Lind-Thomsen (coordinator), R Salbøl Brandt, M Hillebreght, S Zaheri, FWNM Wit (ATHENA), A Scherrer, F Schöni-Affolter, M Rickenbach (SHCS), A Tavelli, I Fanti (ICONA), O Leleux, J Mourali, F Le Marec, E Boerg (Aquitaine), E Thulin, A Sundström (HIVBIVUS), G Bartsch, G Thompsen (CPCRA), C Necsoi, M Delforge (Brussels), E Fontas, C Caissotti, K Dollet (Nice), S Mateu, F Torres (BASS), K Petoumenos, A Blance, R Huang, R Puhr (AHOD), K Grønberg Laut, D Kristensen (EuroSIDA)

Statisticians: CA Sabin*, AN Phillips*, DA Kamara, CJ Smith, A Mcroft*

D:A:D coordinating office: CI Hatleberg, L Ryom, A Lind-Thomsen, RS Brandt, D Raben, C Matthews, A Bojesen, AL Grevsen, JD Lundgren*^ç

Member of the D:A:D Oversight Committee: B Powderly*, N Shortman*, C Moecklinghoff*, G Reilly*, X Franquet*

D:A:D working group experts:

Kidney: L Ryom, A Mcroft*, O Kirk*, P Reiss*, C Smit, M Ross, CA Fux, P Morlat, E Fontas, DA Kamara, CJ Smith, JD Lundgren*^ç

Mortality: CJ Smith, L Ryom, CI Hatleberg, AN Phillips*, R Weber*, P Morlat, C Pradier*, P Reiss*, FWNM Wit, N Friis-Møller, J Kowalska, JD Lundgren*^ç

Cancer: CA Sabin*, L Ryom, CI Hatleberg, M Law*, A d'Arminio Monforte*, F Dabis*, F Bonnet*, P Reiss*, FWNM Wit, CJ Smith, DA Kamara, J Bohlius, M Bower, G Fätkenheuer, A Grulich, JD Lundgren*^ç

External endpoint reviewers: A Sjøl (CVD), P Meidahl (oncology), JS Iversen (nephrology)

Funding: ‘Oversight Committee for The Evaluation of Metabolic Complications of HAART’ with representatives from academia, patient community, FDA, EMA and a consortium of AbbVie, Bristol-Myers Squibb, Gilead Sciences, ViiV Healthcare, Merck and Janssen Pharmaceuticals.

The current members of the 11 Cohorts are as follows:

ATHENA (AIDS Therapy Evaluation Project Netherlands):

Central coordination: P. Reiss*, S. Zaheri, M Hillebregt, F.W.N.M. Wit;

CLINICAL CENTRES (α denotes site coordinating physician) Academic Medical Centre of the

University of Amsterdam: J.M. Prinsα, T.W. Kuijpers, H.J. Scherpbier, J.T.M. van der Meer,

F.W.N.M. Wit, M.H. Godfried, P. Reiss, T. van der Poll, F.J.B. Nellen, S.E. Geerlings, M. van

Vugt, D. Pajkrt, J.C. Bos, W.J. Wiersinga, M. van der Valk, A. Goorhuis, J.W. Hovius, J. van

Eden, A. Henderiks, A.M.H. van Hes, M. Mutschelknauss, H.E. Nobel, F.J.J. Pijnappel, S.

Jurriaans, N.K.T. Back, H.L. Zaaijer, B. Berkhout, M.T.E. Cornelissen, C.J. Schinkel, X.V.

Thomas. Admiraal De Ruyter Ziekenhuis, Goes: M. van den Berge, A. Stegeman, S. Baas, L.

Hage de Looff, D. Versteeg. Catharina Ziekenhuis, Eindhoven: M.J.H. Pronkα, H.S.M.

Ammerlaan, E.S. de Munnik. A.R. Jansz, J. Tjhie, M.C.A. Wegdam, B. Deiman, V. Scharnhorst.

Emma Kinderziekenhuis: A. van der Plas, A.M. Weijnsfeld. Erasmus MC, Rotterdam: M.E. van

der Endeα, T.E.M.S. de Vries-Sluijs, E.C.M. van Gorp, C.A.M. Schurink, J.L. Nouwen, A.

Verbon, B.J.A. Rijnders, H.I. Bax, M. van der Feltz. N. Bassant, J.E.A. van Beek, M. Vriesde,

L.M. van Zonneveld. A. de Oude-Lubbers, H.J. van den Berg-Cameron, F.B. Bruinsma-

Broekman, J. de Groot, M. de Zeeuw- de Man, C.A.B. Boucher, M.P.G Koopmans, J.J.A van

Kampen, S.D. Pas. Erasmus MC–Sophia, Rotterdam: G.J.A. Driessen, A.M.C. van Rossum, L.C. van der Knaap, E. Visser. Flevoziekenhuis, Almere: J. Branger[Ⓐ], A. Rijkeboer-Mes, C.J.H.M. Duijf-van de Ven. HagaZiekenhuis, Den Haag: E.F. Schippers[Ⓐ], C. van Nieuwkoop. J.M. van IJperen, J. Geilings. G. van der Hut. P.F.H. Franck. HIV Focus Centrum (DC Klinieken): A. van Eeden[Ⓐ]. W. Brokking, M. Groot, L.J.M. Elsenburg, M. Damen, I.S. Kwa. Isala, Zwolle: P.H.P. Groeneveld[Ⓐ], J.W. Bouwhuis, J.F. van den Berg, A.G.W. van Hulzen, G.L. van der Blik, P.C.J. Bor, P. Bloembergen, M.J.H.M. Wolfhagen, G.J.H.M. Ruijs. Leids Universitair Medisch Centrum, Leiden: , F.P. Kroon[Ⓐ], M.G.J. de Boer, M.P. Bauer, H. Jolink, A.M. Vollaard, W. Dorama, N. van Holten, E.C.J. Claas, E. Wessels. Maasstad Ziekenhuis, Rotterdam: J.G. den Hollander[Ⓐ], K. Pogany, A. Roukens, M. Kastelijns, J.V. Smit, E. Smit, D. Struik-Kalkman, C. Tearno, M. Bezemer, T. van Niekerk, O. Pontesilli. Maastricht UMC+, Maastricht: S.H. Lowe[Ⓐ], A.M.L. Oude Lashof, D. Posthouwer, R.P. Ackens, J. Schippers, R. Vergoossen, B. Weijenberg-Maes, I.H.M. van Loo, T.R.A. Havenith. MCH-Bronovo, Den Haag: E.M.S. Leyten[Ⓐ], L.B.S. Gelinck, A. van Hartingsveld, C. Meerkerk, G.S. Wildenbeest, J.A.E.M. Mutsaers, C.L. Jansen. MC Slotervaart, Amsterdam: J.W. Mulder, S.M.E. Vrouwenraets, F.N. Lauw, M.C. van Broekhuizen, H. Paap, D.J. Vlasblom, P.H.M. Smits. MC Zuiderzee, Lelystad: S. Weijer[Ⓐ], R. El Moussaoui, A.S. Bosma. Medisch Centrum Leeuwarden, Leeuwarden: M.G.A. van Vonderen[Ⓐ], D.P.F. van Houte, L.M. Kampschreur, K. Dijkstra, S. Faber, J. Weel. Medisch Spectrum Twente, Enschede: G.J. Kootstra[Ⓐ], C.E. Delsing, M. van der Burg-van de Plas, H. Heins, E. Lucas. Noorwest Ziekenhuisgroep, Alkmaar: W. Kortmann[Ⓐ], G. van Twillert[Ⓐ], J.W.T. Cohen Stuart, B.M.W. Diederren, D. Pronk, F.A. van Truijen-Oud, W. A. van der Reijden, R. Jansen. OLVG, Amsterdam: K. Brinkman[Ⓐ], G.E.L. van den Berk, W.L. Blok, P.H.J. Frissen, K.D. Lettinga W.E.M. Schouten, J. Veenstra, C.J. Brouwer, G.F. Geerders, K. Hoeksema, M.J. Kleene, I.B.

van der Meché, M. Spelbrink, H. Sulman, A.J.M. Toonen, S. Wijnands, M. Damen, D. Kwa, E. Witte. Radboudumc, Nijmegen: P.P. Koopmans, M. Keuter, A.J.A.M. van der Ven, H.J.M. ter Hofstede, A.S.M. Dofferhoff, R. van Crevel, M. Albers, M.E.W. Bosch, K.J.T. Grintjes-Huisman, B.J. Zomer, F.F. Stelma, J. Rahamat-Langendoen, D. Burger. Rijnstate, Arnhem: C. Richter, E.H. Gisolf, R.J. Hassing, G. ter Beest, P.H.M. van Bentum, N. Langebeek, R. Tiemessen, C.M.A. Swanink. Spaarne Gasthuis, Haarlem: S.F.L. van Lelyveld, R. Soetekouw, N. Hulshoff, L.M.M. van der Pijlt, J. van der Swaluw, N. Bermon, W.A. van der Reijden, R. Jansen, B.L. Herpers, D. Veenendaal. Medisch Centrum Jan van Goyen, Amsterdam: D.W.M. Verhagen, M. van Wijk. St Elisabeth Ziekenhuis, Tilburg: M.E.E. van Kasteren, A.E. Brouwer, B.A.F.M. de Kruijf-van de Wiel, M. Kuipers, R.M.W.J. Santegoets, B. van der Ven, J.H. Marcelis, A.G.M. Buiting, P.J. Kabel. Universitair Medisch Centrum Groningen, Groningen: W.F.W. Bierman, H. Scholvinck, K.R. Wilting, Y. Stienstra, H. de Groot-de Jonge, P.A. van der Meulen, D.A. de Weerd, J. Ludwig-Roukema, H.G.M. Niesters, A. Riezebos-Brilman, C.C. van Leer-Buter, M. Knoester. Universitair Medisch Centrum Utrecht, Utrecht: A.I.M. Hoepelman, T. Mudrikova, P.M. Ellerbroek, J.J. Oosterheert, J.E. Arends, R.E. Barth, M.W.M. Wassenberg, E.M. Schadd, D.H.M. van Elst-Laurijssen, E.E.B. van Oers-Hazelzet, S. Vervoort, M. van Berkel, R. Schuurman, F. Verduyn-Lunel, A.M.J. Wensing. VUmc, Amsterdam: E.J.G. Peters, M.A. van Agtmael, M. Bomers, J. de Vocht, M. Heitmuller, L.M. Laan, A.M. Pettersson, C.M.J.E. Vandenbroucke-Grauls, C.W. Ang. Wilhelmina Kinderziekenhuis, UMCU, Utrecht: S.P.M. Geelen, T.F.W. Wolfs, L.J. Bont, N. Nauta. COORDINATING CENTRE P. Reiss, D.O. Bezemer, A.I. van Sighem, C. Smit, F.W.N.M. Wit., T.S. Boender, S. Zaheri, M. Hillebregt, A. de Jong, D. Bergsma, P. Hoekstra, A. de Lang, S. Grivell, A. Jansen, M.J. Rademaker, M. Raethke, R. Meijering, S. Schnörr, L. de Groot, M. van den Akker, Y. Bakker, E.

Claessen, A. El Berkaoui, J. Koops, E. Kruijne, C. Lodewijk, L. Munjishvili, B. Peeck, C. Ree, R. Regtop, Y. Ruijs, T. Rutkens, L. van de Sande, M. Schoorl, A. Timmerman, E. Tuijn, L. Veenenberg, S. van der Vliet, A. Wisse, T. Woudstra, B. Tuk.

Aquitaine Cohort (France)

Composition du Conseil scientifique :

Coordination: F. Bonnet, F. Dabis

Scientific committee: M. Dupon, V. Gaborieau, D. Lacoste, D. Malvy, P. Mercié, P. Morlat, D. Neau, JL. Pellegrin, S. Tchamgoué, E. Lazaro, C. Cazanave, M. Vandenhende, M.O. Vareil, Y. Gérard, P. Blanco, S. Bouchet, D. Breilh, H. Fleury, I. Pellegrin, G. Chêne, R. Thiébaud, L. Wittkop, L. Wittkop, O. Leleux, S. Lawson-Ayayi, A. Gimbert, S. Desjardin, L. Lacaze-Buzy, V. Petrov-Sanchez

Epidemiology and Methodology: F. Bonnet, G. Chêne, F. Dabis, R. Thiébaud, L. Wittkop

Infectious Diseases and Internal Medicine: K. André, N. Bernard, F. Bonnet, O. Caubet, L. Caunegre, C. Cazanave, I. Chossat, C. Courtault, FA. Dauchy, S. De Witte, D. Dondia, M. Dupon, P. Duffau, H. Dutronc, S. Farbos, I. Faure, H. Ferrand, V. Gaborieau, Y. Gerard, C. Greib, M. Hessamfar, Y. Imbert, D. Lacoste, P. Lataste, E. Lazaro, D. Malvy, J. Marie, M. Mechain, P. Mercié, E. Monlun, P. Morlat, D. Neau, A. Ochoa, JL. Pellegrin, T. Pistone, I. Raymond, MC. Receveur, P. Rispal, L. Sorin, S. Tchamgoué, C. Valette, MA. Vandenhende, MO. Vareil, JF. Viillard, H. Wille, G. Wirth.

Immunology: I. Pellegrin, P. Blanco

Virology: H. Fleury, Me. Lafon, P. Trimoulet, P. Bellecave, C. Tumiotto

Pharmacology: S. Bouchet, D. Breilh, F. Haramburu, G. Miremeont-Salamé

Data collection, Project Management and Statistical Analyses: MJ. Blaizeau, M. Decoin, C. Hannapier, E. Lenaud et A. Pougetoux; S. Delveaux, C. D'Ivernois, F. Diarra B. Uwamaliya-Nziyumvira, O. Leleux; F. Le Marec, Eloïse Boerg, S. Lawson-Ayayi;

IT department and eCRF development: G. Palmer, V. Conte, V. Sapparrart

AHOD (Australian HIV Observational Database, Australia):

Central coordination: M. Law *, K. Petoumenos, R Puhr, R Huang (Sydney, New South Wales).

Participating physicians (city, state): R. Moore, S. Edwards, J. Hoy, K. Watson, N. Roth, H Lau (Melbourne, Victoria); M Bloch, D. Baker, A. Carr, D. Cooper, (Sydney, New South Wales); M O'Sullivan (Gold Coast, Queensland), D. Nolan, G Guelfi (Perth, Western Australia).

BASS (Spain):

Central coordination: G. Calvo, F. Torres, S. Mateu (Barcelona);

Participating physicians (city): P. Domingo, M.A. Sambeat, J. Gatell, E. Del Cacho, J.

Cadafalch, M. Fuster (Barcelona); C. Codina, G. Sirera, A. Vaqué (Badalona).

The Brussels St Pierre Cohort (Belgium):

Coordination: S. De Wit*, N. Clumeck, M. Delforge, C. Necsoi.

Participating physicians: N. Clumeck, S. De Wit*, AF Gennotte, M. Gerard, K. Kabeya, D.

Konopnicki, A. Libois, C. Martin, M.C. Payen, P. Semaille, Y. Van Laethem.

The Brussels St Pierre Cohort (Belgium):

Coordination: S. De Wit*, N. Clumeck, M. Delforge, C. Necsoi.

Participating physicians: N. Clumeck, S. De Wit*, AF Gennotte, M. Gerard, K. Kabeya, D. Konopnicki, A. Libois, C. Martin, M.C. Payen, P. Semaille, Y. Van Laethem.

CPCRA (USA):

Central coordination: J. Neaton, G. Bartsch, W.M. El-Sadr*, E. Krum, G. Thompson, D. Wentworth;

Participating physicians (city, state): R. Luskin-Hawk (Chicago, Illinois); E. Telzak (Bronx, New York); W.M. El-Sadr (Harlem, New York); D.I. Abrams (San Francisco, California); D. Cohn (Denver, Colorado); N. Markowitz (Detroit, Michigan); R. Arduino (Houston, Texas); D. Mushatt (New Orleans, Louisiana); G. Friedland (New Haven, Connecticut); G. Perez (Newark, New Jersey); E. Tedaldi (Philadelphia, Pennsylvania); E. Fisher (Richmond, Virginia); F. Gordin (Washington, DC); L.R. Crane (Detroit, Michigan); J. Sampson (Portland, Oregon); J. Baxter (Camden, New Jersey).

EuroSIDA (multinational)

Steering Committee: J Gatell, B Gazzard, A Horban, I Karpov, M Losso, A d'Arminio Monforte, C Pedersen, M Ristola, A Phillips, P Reiss, J Lundgren, J Rockstroh

Chair: J Rockstroh

Study Co-leads: A Mocroft, O Kirk

Coordinating Centre Staff: O Kirk, L Peters, C Matthews, AH Fischer, A Bojesen, D Raben, D Kristensen, K Grønberg Laut, JF Larsen, D Podlekareva

Statistical Staff: A Mcroft, A Phillips, A Cozzi-Lepri, L Shepherd, A Schultze, S Amele

The multi-centre study group, EuroSIDA (national coordinators in parenthesis).

Argentina: (M Losso), M Kundro, Hospital JM Ramos Mejia, Buenos Aires.

Austria: (B Schmied), Pulmologisches Zentrum der Stadt Wien, Vienna; R Zangerle, Medical University Innsbruck, Innsbruck.

Belarus: (I Karpov), A Vassilenko, Belarus State Medical University, Minsk, VM Mitsura, Gomel State Medical University, Gomel; D Paduto, Regional AIDS Centre, Svetlogorsk.

Belgium: (N Clumeck), S De Wit, M Delforge, Saint-Pierre Hospital, Brussels; E Florence, Institute of Tropical Medicine, Antwerp; L Vandekerckhove, University Ziekenhuis Gent, Gent.

Bosnia-Herzegovina: (V Hadziosmanovic), Klinicki Centar Univerziteta Sarajevo, Sarajevo.

Croatia: (J Begovac), University Hospital of Infectious Diseases, Zagreb.

Czech Republic: (L Machala), D Jilich, Faculty Hospital Bulovka, Prague; D Sedlacek, Charles University Hospital, Plzen.

Denmark: G Kronborg, T Benfield, Hvidovre Hospital, Copenhagen; J Gerstoft, T Katzenstein, Rigshospitalet, Copenhagen; NF Møller, C Pedersen, Odense University Hospital, Odense; L Ostergaard, Skejby Hospital, Aarhus, L Wiese, Roskilde Hospital, Roskilde; L N Nielsen, Hillerod Hospital, Hillerod.

Estonia: (K Zilmer), West-Tallinn Central Hospital, Tallinn; Jelena Smidt, Nakkusosakond Sisekliinik, Kohtla-Järve.

Finland: (M Ristola), I Aho, Helsinki University Central Hospital, Helsinki.

France: (J-P Viard), Hôtel-Dieu, Paris; P-M Girard, Hospital Saint-Antoine, Paris; C Pradier, E Fontas, Hôpital de l'Archet, Nice; C Duvivier, Hôpital Necker-Enfants Malades, Paris.

Germany: (J Rockstroh), Universitäts Klinik Bonn; R Schmidt, Medizinische Hochschule Hannover; O Degen, University Medical Center Hamburg-Eppendorf, Infectious Diseases Unit, Hamburg; HJ Stellbrink, IPM Study Center, Hamburg; C Stefan, JW Goethe University Hospital, Frankfurt; J Bogner, Medizinische Poliklinik, Munich; G. Fätkenheuer, Universität Köln, Cologne.

Georgia: (N Chkhartishvili) Infectious Diseases, AIDS & Clinical Immunology Research Center, Tbilisi

Greece: (P Gargalianos), G Xylomenos, K Armenis, Athens General Hospital "G Gennimatas"; H Sambatakou, Ippokration General Hospital, Athens.

Hungary: (J Szlávik), Szent László Hospital, Budapest.

Iceland: (M Gottfredsson), Landspítali University Hospital, Reykjavik.

Ireland: (F Mulcahy), St. James's Hospital, Dublin.

Israel: (I Yust), D Turner, M Burke, Ichilov Hospital, Tel Aviv; E Shahr, G Hassoun, Rambam Medical Center, Haifa; H Elinav, M Haouzi, Hadassah University Hospital, Jerusalem; D Elbirt, ZM Stoegeger, AIDS Center (Neve Or), Jerusalem.

Italy: (A D' Arminio Monforte), Istituto Di Clinica Malattie Infettive e Tropicale, Milan; R Esposito, I Mazeu, C Mussini, Università Modena, Modena; F Mazzotta, A Gabbuti, Ospedale S Maria Annunziata, Firenze; V Vullo, M Lichtner, University di Roma la Sapienza, Rome; M Zaccarelli, A Antinori, R Acinapura, M Plazzi, Istituto Nazionale Malattie Infettive Lazzaro

Spallanzani, Rome; A Lazzarin, A Castagna, N Gianotti, Ospedale San Raffaele, Milan; M Galli, A Ridolfo, Osp. L. Sacco, Milan.

Latvia: (B Rozentale), Infectology Centre of Latvia, Riga.

Lithuania: (V Uzdaviniene) Vilnius University Hospital Santariskiu Klinikos, Vilnius; R Matulionyte, Center of Infectious Diseases, Vilnius University Hospital Santariskiu Klinikos, Vilnius.

Luxembourg: (T Staub), R Hemmer, Centre Hospitalier, Luxembourg.

Netherlands: (P Reiss), Academisch Medisch Centrum bij de Universiteit van Amsterdam, Amsterdam.

Norway: (V Ormaasen), A Maeland, J Bruun, Ullevål Hospital, Oslo.

Poland: (B Knysz), J Gasiorowski, M Inglot, Medical University, Wroclaw; A Horban, E Bakowska, Centrum Diagnostyki i Terapii AIDS, Warsaw; R Flisiak, A Grzeszczuk, Medical University, Bialystok; M Parczewski, K Maciejewska, B Aksak-Was, Medical University, Szczecin; M Beniowski, E Mularska, Osrodek Diagnostyki i Terapii AIDS, Chorzow; T Smiatacz, M Gensing, Medical University, Gdansk; E Jablonowska, E Malolepsza, K Wojcik, Wojewodzki Szpital Specjalistyczny, Lodz; I Mozer-Lisewska, Poznan University of Medical Sciences, Poznan.

Portugal: (L Caldeira), Hospital Santa Maria, Lisbon; K Mansinho, Hospital de Egas Moniz, Lisbon; F Maltez, Hospital Curry Cabral, Lisbon.

Romania: (R Radoi), C Oprea, Spitalul de Boli Infectioase si Tropicale: Dr. Victor Babes, Bucarest.

Russia: (A Pantelev), O Pantelev, St Petersburg AIDS Centre, St Peterburg; A Yakovlev, Medical Academy Botkin Hospital, St Petersburg; T Trofimora, Novgorod Centre for AIDS, Novgorod, I Khromova, Centre for HIV/AIDS & and Infectious Diseases, Kaliningrad; E Kuzovatova, Nizhny Novgorod Scientific and Research Institute of Epidemiology and Microbiology named after Academician I.N. Blokhina, Nizhny Novogrod; E Borodulina, E Vdoushkina, Samara State Medical University, Samara.

Serbia: (D Jevtic), The Institute for Infectious and Tropical Diseases, Belgrade.

Slovenia: (J Tomazic), University Clinical Centre Ljubljana, Ljubljana.

Spain: (JM Gatell), JM Miró, Hospital Clinic Universitari de Barcelona, Barcelona; S Moreno, J. M. Rodriguez, Hospital Ramon y Cajal, Madrid; B Clotet, A Jou, R Paredes, C Tural, J Puig, I Bravo, Hospital Germans Trias i Pujol, Badalona; P Domingo, M Gutierrez, G Mateo, MA Sambeat, Hospital Sant Pau, Barcelona; JM Laporte, Hospital Universitario de Alava, Vitoria-Gasteiz.

Sweden: (K Falconer), A Thalme, A Sonnerborg, Karolinska University Hospital, Stockholm; A Blaxhult, Venhälsan-Sodersjukhuset, Stockholm; L Flamholc, Malmö University Hospital, Malmö.

Switzerland: (A Scherrer), R Weber, University Hospital Zurich; M Cavassini, University Hospital Lausanne; A Calmy, University Hospital Geneva; H Furrer, University Hospital Bern; M Battegay, University Hospital Basel; P Schmid, Cantonal Hospital St. Gallen.

Ukraine: A Kuznetsova, Kharkov State Medical University, Kharkov; G Kyselyova, Crimean Republican AIDS centre, Simferopol; M Sluzhynska, Lviv Regional HIV/AIDS Prevention and Control CTR, Lviv.

United Kingdom: (B Gazzard), St. Stephen's Clinic, Chelsea and Westminster Hospital, London; AM Johnson, E Simons, S Edwards, Mortimer Market Centre, London; A Phillips, MA Johnson, A Mocroft, Royal Free and University College Medical School, London (Royal Free Campus); C Orkin, Royal London Hospital, London; J Weber, G Scullard, Imperial College School of Medicine at St. Mary's, London; A Clarke, Royal Sussex County Hospital, Brighton; C Leen, Western General Hospital, Edinburgh.

The following centers have previously contributed data to EuroSIDA:

Infectious Diseases Hospital, Sofia, Bulgaria

Hôpital de la Croix Rousse, Lyon, France

Hôpital de la Pitié-Salpêtrière, Paris, France

Unité INSERM, Bordeaux, France

Hôpital Edouard Herriot, Lyon, France

Bernhard Nocht Institut für Tropenmedizin, Hamburg, Germany

1st I.K.A Hospital of Athens, Athens, Greece

Ospedale Riuniti, Divisione Malattie Infettive, Bergamo, Italy

Ospedale di Bolzano, Divisione Malattie Infettive, Bolzano, Italy

Ospedale Cotugno, III Divisione Malattie Infettive, Napoli, Italy

Dérer Hospital, Bratislava, Slovakia

Hospital Carlos III, Departamento de Enfermedades Infecciosas, Madrid, Spain

Kiev Centre for AIDS, Kiev, Ukraine

Luhansk State Medical University, Luhansk, Ukraine

Odessa Region AIDS Center, Odessa, Ukraine

HivBivus (Sweden):

Central coordination: L. Morfeldt, G. Thulin, A. Sundström.

Participating physicians (city): B. Åkerlund (Huddinge); K. Koppel, A. Karlsson (Stockholm); L. Flamholz, C. Håkangård (Malmö).

The ICONA Foundation (Italy):

BOARD OF DIRECTORS

A d'Arminio Monforte (President), A Antinori, A Castagna, F Castelli, R Cauda, G Di Perri, M Galli, R Iardino, G Ippolito, GC Marchetti, CF Perno, F von Schloesser, P Viale

SCIENTIFIC SECRETARY

A d'Arminio Monforte, A Antinori, A Castagna, F Ceccherini-Silberstein, A Cozzi-Lepri, E Girardi, S Lo Caputo, C Mussini, M Puoti

STEERING COMMITTEE

M Andreoni, A Ammassari, A Antinori, C Balotta, A Bandera, P Bonfanti, S Bonora, M Borderi, A Calcagno, L Calza, MR Capobianchi, A Castagna, F Ceccherini-Silberstein, A Cingolani, P Cinque, A Cozzi-Lepri, A d'Arminio Monforte, A De Luca, A Di Biagio, E Girardi, N Gianotti, A Gori, G Guaraldi, G Lapadula, M Lichtner, S Lo Caputo, G Madeddu, F Maggiolo, G

Marchetti, S Marcotullio, L Monno, C Mussini, S Nozza, M Puoti, E Quiros Roldan, R Rossotti, S Rusconi, MM Santoro, A Saracino, M Zaccarelli.

STATISTICAL AND MONITORING TEAM

A Cozzi-Lepri, I Fanti, L Galli, P Lorenzini, A Rodano, M Shanyinde, A Tavelli

BIOLOGICAL BANK INMI

F Carletti, S Carrara, A Di Caro, S Graziano, F Petrone, G Prota, S Quartu, S Truffa

PARTICIPATING PHYSICIANS AND CENTERS

Italy A Giacometti, A Costantini, V Barocci (Ancona); G Angarano, L Monno, C Santoro (Bari); F Maggiolo, C Suardi (Bergamo); P Viale, V Donati, G Verucchi (Bologna); F Castelli, C Minardi, E Quiros Roldan (Brescia); T Quirino, C Abeli (Busto Arsizio); PE Manconi, P Piano (Cagliari); B Cacopardo, B Celesia (Catania); J Vecchiet, K Falasca (Chieti); A Pan, S Lorenzotti (Cremona); L Sighinolfi, D Segala (Ferrara); F Mazzotta, F Vichi (Firenze); G Cassola, C Viscoli, A Alessandrini, N Bobbio, G Mazzarello (Genova); C Mastroianni, V Belvisi (Latina); P Bonfanti, I Caramma (Lecco); A Chiodera, P Milini (Macerata); A d'Arminio Monforte, M Galli, A Lazzarin, G Rizzardini, M Puoti, A Castagna, G Marchetti, MC Moioli, R Piolini, AL Ridolfo, S Salpietro, C Tincati, (Milano); C Mussini, C Puzzolante (Modena); A Gori, G Lapadula (Monza); N Abrescia, A Chirianni, G Borgia, R Orlando, G Bonadies, F Di Martino, I Gentile, L Maddaloni (Napoli); AM Cattelan, S Marinello (Padova); A Cascio, C Colomba (Palermo); F Baldelli, E Schiaroli (Perugia); G Parruti, F Sozio (Pescara); G Magnani, MA Ursitti (Reggio Emilia); M Andreoni, A Antinori, R Cauda, A Cristaudo, V Vullo, R Acinapura, G Baldin, M Capozzi, S Cicalini, A Cingolani, L Fontanelli Sulekova, G Iaiani, A Latini, I Mastrorosa, MM Plazzi, S Savinelli, A Vergori (Roma); M Cecchetto, F Viviani

(Rovigo); G Madeddu, P Bagella (Sassari); A De Luca, B Rossetti (Siena); A Franco, R Fontana Del Vecchio (Siracusa); D Francisci, C Di Giuli (Terni); P Caramello, G Di Perri, S Bonora, GC Orofino, M Sciandra (Torino); M Bassetti, A Londero (Udine); G Pellizzer, V Manfrin (Vicenza) G Starnini, A Ialungo(Viterbo).

Nice HIV Cohort (France):

Central coordination: C. Pradier*, E. Fontas, K. Dollet, C. Caissotti.

Participating physicians: P. Dellamonica, E. Bernard, J. Courjon, E. Cua, F. De Salvador-Guillouet, J. Durant, C. Etienne, S. Ferrando, V. Mondain-Miton, A. Naqvi, I. Perbost, S. Pillet, B. Prouvost-Keller, P. Pugliese, V. Rio, K. Risso, P.M. Roger.

SHCS (Swiss HIV Cohort Study, Switzerland):

The data are gathered by the Five Swiss University Hospitals, two Cantonal Hospitals, 15 affiliated hospitals and 36 private physicians (listed in <http://www.shcs.ch/180-health-care-providers>).

Members of the Swiss HIV Cohort Study :

Aubert V, Battegay M, Bernasconi E, Böni J, Braun DL, Bucher HC, Calmy A, Cavassini M, Ciuffi A, Dollenmaier G, Egger M, Elzi L, Fehr J, Fellay J, Furrer H (Chairman of the Clinical and Laboratory Committee), Fux CA, Günthard HF (President of the SHCS), Haerry D (deputy of "Positive Council"), Hasse B, Hirsch HH, Hoffmann M, Hösli I, Kahlert C, Kaiser L, Keiser O, Klimkait T, Kouyos RD, Kovari H, Ledergerber B, Martinetti G, Martinez de Tejada B, Marzolini C, Metzner KJ, Müller N, Nicca D, Pantaleo G, Paioni P, Rauch A (Chairman of the Scientific Board), Rudin C (Chairman of the Mother & Child Substudy), Scherrer AU (Head of

Data Centre), Schmid P, Speck R, Stöckle M, Tarr P, Trkola A, Vernazza P, Wandeler G, Weber R*, Yerly S.

Funding

Data on Adverse Events (D:A:D) Study: The D:A:D study was supported by a grant [grant number DNR126] from the Danish National Research Foundation (CHIP & PERSIMUNE); the Highly Active Antiretroviral Therapy Oversight Committee (HAARTOC), a collaborative committee with representation from academic institutions, the European Agency for the Evaluation of Medicinal Products, the United States Food and Drug Administration, the patient community, and pharmaceutical companies with licensed anti-HIV drugs in the European Union: AbbVie, Bristol-Myers Squibb, Gilead Sciences Inc., ViiV Healthcare, Merck & Co Inc. and Janssen Pharmaceuticals. Supported also by a grant from the Dutch Ministry of Health, Welfare and Sport through the Center for Infectious Disease Control of the National Institute for Public Health and the Environment to Sticking HIV Monitoring (ATHENA); by a grant from the Agence nationale de recherches sur le sida et les hépatites virales [ANRS, Action Coordonnée no.7, Cohortes] to the Aquitaine Cohort; The Australian HIV Observational Database (AHOD) is funded as part of the Asia Pacific HIV Observational Database, a program of The Foundation for AIDS Research, amfAR, and is supported in part by a grant from the U.S. National Institutes of Health's National Institute of Allergy and Infectious Diseases (NIAID) [grant number U01-AI069907] and by unconditional grants from Merck Sharp & Dohme; Gilead Sciences; Bristol-Myers Squibb; Boehringer Ingelheim; Janssen-Cilag; ViiV Healthcare. The Kirby Institute is funded by The Australian Government Department of Health and Ageing, and is affiliated with the Faculty of Medicine, The University of New South Wales; by grants from the Fondo de Investigación Sanitaria [grant number FIS 99/0887] and Fundación para la Investigación y la

Prevención del SIDA en Españã [grant number FIPSE 3171/00], to the Barcelona Antiretroviral Surveillance Study (BASS); by the National Institute of Allergy and Infectious Diseases, National Institutes of Health [grants number 5U01AI042170-10, 5U01AI046362-03], to the Terry Bein Community Programs for Clinical Research on AIDS (CPCRA); by primary funding provided by the European Union's Seventh Framework Programme for research, technological development and demonstration under EuroCoord grant agreement n° 260694 and unrestricted grants by Bristol-Myers Squibb, Janssen R&D, Merck and Co. Inc., Pfizer Inc., GlaxoSmithKline LLC, (the participation of centres from Switzerland is supported by The Swiss National Science Foundation (Grant 108787)) to the EuroSIDA study; by unrestricted educational grants of AbbVie, Bristol-Myers Squibb, Gilead Sciences, GlaxoSmithKline, Pfizer, Janssen Pharmaceuticals to the Italian Cohort Naive to Antiretrovirals (The ICONA Foundation); and financed within the framework of the Swiss HIV Cohort Study, supported by the Swiss National Science Foundation (grant #148522) and by the SHCS research foundation.

REFERENCES

1. Koethe JR, Jenkins CA, Lau B, et al. Rising Obesity Prevalence and Weight Gain Among Adults Starting Antiretroviral Therapy in the United States and Canada. *AIDS Res Hum Retroviruses*. **2016**;32(1):50-58.
2. Crum-Cianflone N, Roediger MP, Eberly L, et al. Increasing rates of obesity among HIV-infected persons during the HIV epidemic. *PLoS One*. **2010**;5(4):e10106.
3. World Health Organization (WHO). Obesity and overweight [Accessed 1st Aug 2017]. 2006; <http://www.who.int/mediacentre/factsheets/fs311/en/>.

4. Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *The Lancet*. **2008**;371(9612):569-578.
5. Berrington de Gonzalez A, Hartge P, Cerhan JR, et al. Body-mass index and mortality among 1.46 million white adults. *NEJM*. **2010**;363(23):2211-2219.
6. Sabin CA, Reiss P. Epidemiology of ageing with HIV: what can we learn from cohorts? *AIDS*. **2017**;31 Suppl 2:S121-S128.
7. Koethe et al, An Optimal Body Mass Index Range Associated With Improved Immune Reconstitution Among HIV-Infected Adults Initiating Antiretroviral Therapy *Clinical Infectious Diseases* 2011;53(9):952–960
8. Achhra AC, Mocroft A, Reiss P, et al. Short-term weight gain after antiretroviral therapy initiation and subsequent risk of cardiovascular disease and diabetes: the D:A:D study. *HIV Med*. **2016**;17(4):255-268.
9. Koethe JR, Jenkins CA, Turner M, et al. Body mass index and the risk of incident noncommunicable diseases after starting antiretroviral therapy. *HIV Med*. **2015**;16(1):67-72.
10. Friis-Moller N, Reiss P, et al for the D:A:D Study Group. Class of antiretroviral drugs and the risk of myocardial infarction. *NEJM*. **2007**;356(17):1723-1735.
11. Kyrgiou M, Kalliala I, Markozannes G, et al. Adiposity and cancer at major anatomical sites: umbrella review of the literature. *BMJ*. **2017**;356:j477.
12. Textor J, Hardt J, Knuppel S. DAGitty: a graphical tool for analyzing causal diagrams. *Epidemiology*. **2011**;22(5):745.

13. Ryom L, Lundgren JD, Ross M, et al. Renal Impairment and Cardiovascular Disease in HIV-Positive Individuals: The D:A:D Study. *The Journal of infectious diseases*. 2016;214(8):1212-1220.
14. Cockcroft, D.W. and M.H. Gault. Prediction of creatinine clearance from serum creatinine. *Nephron*. **1976**. 16(1):31-41.
15. Petoumenos K, Law MG. Smoking, alcohol and illicit drug use effects on survival in HIV-positive persons. *Current opinion in HIV and AIDS*. 2016;11(5):514-520.
16. Petoumenos K, Worm S, Reiss P, et al. Rates of cardiovascular disease following smoking cessation in patients with HIV infection: results from the D:A:D study. *HIV Med*. **2011**;12(7):412-421.
16. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA*. 2013;309(1):71-82.
18. Hainer V, Aldhoon-Hainerova I. Obesity paradox does exist. *Diabetes Care*. **2013**;36 Suppl 2:S276-281.
19. Romero-Corral A, Montori VM, Somers VK, et al. Association of bodyweight with total mortality and with cardiovascular events in coronary artery disease: a systematic review of cohort studies. *The Lancet*. **2006**;368(9536):666-678.
20. Filardo G, Hamilton C, Hamman B, Ng HK, Grayburn P. Categorizing BMI may lead to biased results in studies investigating in-hospital mortality after isolated CABG. *J Clin Epidemiol*. **2007**;60(11):1132-1139.

21. Sharma A, Hoover DR, Shi Q, et al. Relationship between Body Mass Index and Mortality in HIV-Infected HAART Users in the Women's Interagency HIV Study. *PLOS ONE*. 2015;10(12):e0143740.

Fig-1: Latest BMI (lagged by 1 year) and the risk of SNAEs.

Footnote: All models adjusted for sex, race, mode of HIV transmission, clinical cohort, calendar year and time-updated age, smoking and CD4 count. Additional adjustments are as follows: Fig-1a: CVD: Model A additionally adjusted for being currently on abacavir, cumulative years on NRTIs and PIs. Model B: As model A plus additionally adjusted for time-updated (lagged by 1 year) diabetes, total, HDL and LDL cholesterol, systolic and diastolic blood pressure. Fig 1b: Models for diabetes adjusted for HCV infection and following time-updated (lagged by 1 year) variables: cumulative years of stavudine use, triglycerides, and HDL cholesterol. Fig 1c and Fig 1d: Models for cancer and all-cause mortality adjusted for hepatitis B or C co-infection. P for interaction between gender and BMI: <0.001 for all-cause mortality. BMI category of 23-25 was the reference category in all models.

Table-1: Baseline characteristics by baseline BMI category


	BMI (kg/m ²) category at baseline						Overall
	<18.5	18.5-23	23-25	25-27.5	27.5-30	>30	
N (%)	1929 (4.7)	17640 (42.9)	9283 (22.6)	6998 (17.0)	2910 (7.1)	2389 (5.8)	41149
Age in years (mean) (SD)	38.8 (9.8)	39.3 (9.3)	40.9 (9.8)	41.8 (9.9)	41.7 (10.2)	41.5 (10.3)	40.4 (9.7)
Male (%)	55.1	72	79.6	80.3	73.1	56.2	73.5
Infected with HIV through sex between men (%)	30.9	46.1	50.5	48.3	38.1	25.9	45
Infected with HIV through Injecting drug use (%)	24.9	17.1	13.4	11.2	9.9	9.9	14.7
Race- white (%)	47.6	52.5	51.2	50.2	45.9	39.4	50.5
Current smoker (%)	52.3	46	38.3	33.4	28.1	24.7	39.9
Family history of CVD (%)	6.6	6.7	7.1	7.2	5.7	7	6.8
CD4 count/mm³ (median) (IQR)	354 (205-535)	406 (269-600)	420 (284-610)	415 (277-601)	427 (290-614)	420 (283-635)	410 (273-635)
Log10 HIV RNA (copies/mL) (median) (IQR)	2.1 (1.7-3.9)	1.9 (1.7-3.8)	1.9 (1.7-3.7)	1.8 (1.7-3.7)	1.8 (1.7-3.6)	2 (1.7-3.8)	1.9 (1.7-3.8)
Hepatitis C coinfection (%)	29	21.7	18.1	14.9	13	12.5	18.9
Hepatitis B coinfection (%)	5.9	5.7	5.4	5.3	5.1	4.3	5.5
Total cholesterol mmol/L (mean) (SD)	4.7 (1.3)	4.9 (1.3)	5 (1.4)	5.1 (1.3)	5.2 (1.5)	5.1 (1.3)	4.9 (1.4)
HDL mmol/L (mean) (SD)	1.3 (0.5)	1.25 (0.5)	1.19 (0.4)	1.17 (0.4)	1.18 (0.5)	1.18 (0.4)	1.22 (0.4)
Systolic BP (mean) (SD)	115 (14.8)	120 (14.1)	124 (14.4)	126 (14.9)	128 (15.8)	129 (6.4)	123 (15.0)

Note: BP= blood pressure, CVD= cardiovascular disease, HDL= high-density lipoprotein. Note: At baseline, mode of transmission was unknown in 7.7%; smoking status was unknown in 20%; Hepatitis B and C status was unknown in 12% and 16% respectively; total cholesterol was missing in 7.8%, HDL cholesterol was missing in 31% and SBP was missing in 20% individuals.


Table-2: Number and incidence rates (95% CI) per 1000 PYFU of events by latest BMI category

Events	Latest BMI (kg/m ²) category						Overall
	<18.5	18.5-23	23-25	25-27.5	27.5-30	>30	
CVD	97 (6.7 [5.5-8.2])	578 (4.8 [4.4-5.2])	298(4.6 [4.0-5.1])	242(4.8 [4.2-5.4])	96(4.3 [3.5-5.3])	87(4.9 [3.9-6.0])	1398 (4.8 [4.6-5.1])
Diabetes	33 (2.3[1.6-3.2])	248 (2.0[1.8-2.3])	253 (4.0[3.4-4.4])	280 (5.6[4.9-6.3])	184 (8.5[7.3-9.8])	209 (12.2[10.6-13.9])	1207 (4.2[3.9-4.4])
Non-AIDS defining cancers (NADCs)	95 (7.9[6.4-9.6])	510 (5.1[4.7-5.6])	223 (4.0[3.6-4.6])	167 (3.8[3.3-4.5])	82 (4.2[3.4-5.3])	66 (4.2[3.2-5.3])	1143 (4.7[4.4-5.0])
BMI-cancers	12 (1.0[0.5-1.7])	75 (0.8[0.6-0.9])	32 (0.6[0.4-0.8])	31 (0.7[0.5-1.0])	13 (0.7[0.3-1.2])	21 (1.3[0.8-2.0])	184 (0.7[0.6-0.9])
All-cause mortality in males	260 (33.2 [29.3-37.5])	1138 (13.0 [12.3-13.8])	443 (8.4 [7.6-9.2])	333 (8.1 [7.3-9.0])	142 (8.7 [7.3-10.2])	102 (10.0 [8.2-12.2])	2418 (11.2 [10.8-11.7])
All-cause mortality in females	116 (16.8 [13.9-20.2])	256 (7.4[6.5-8.3])	94 (7.2[5.8-8.8])	66 (6.2[4.8-7.9])	33 (5.4[3.7-7.5])	42 (5.2[3.8-7.1])	607 (7.6[7.0-8.3])

Note: Table shows N (incidence rates per 1000 PYFU [95% confidence interval]). BMI-cancers are cancers previously shown to be strongly associated with BMI in multiple general population studies (see text for details). All-cause mortality rates shown separately by gender because in regression models there was a significant interaction by gender (P<0.001 for interaction term). PYFU= person years of follow-up.



ACCEPTED



ACCEPTED