

Breastfeeding practices in the UK: is the neighbourhood context important?

Andressa B. Peregrino¹, Richard G. Watt¹, Anja Heilmann¹ and Stephen Jivraj¹

¹Department of Epidemiology and Public Health, UCL, London WC1E 6BT, UK.

Abstract

Breastfeeding is an important public health issue worldwide. Breastfeeding rates in the UK, particularly for exclusive breastfeeding, are low compared to other OECD countries, despite its wide-ranging health benefits for both mother and child. There is evidence that deprivation in the structural and social organisation of neighbourhoods is associated with adverse child outcomes. This study aimed to explore whether breastfeeding initiation, exclusive breastfeeding for at least 3 months and any type of breastfeeding for at least 6 months were associated with neighbourhood context measured by neighbourhood deprivation and maternal neighbourhood perceptions in a nationally representative UK sample. A cross-sectional analysis was conducted using data from the Millennium Cohort Study. Logistic regression was carried out on a sample of 17,308 respondents, adjusting for individual- and familial-level socio-demographic characteristics. Neighbourhood deprivation was independently and inversely associated with breastfeeding initiation. Compared to the least deprived areas, the likelihood of initiating breastfeeding was 40% lower in the most deprived neighbourhoods (OR: 0.60, 95% CI: 0.50-0.72). The relationship between both exclusive and any type of breastfeeding at 3 and 6 months respectively with neighbourhood deprivation after adjustment for potential confounders was not entirely linear. Breastfeeding initiation (OR: 0.78, 95% CI: 0.71-0.85), exclusivity for 3 months (OR: 0.84; 95% CI: 0.75-0.95), and any breastfeeding for 6 months (OR: 0.82, 95% CI: 0.73-0.93) were each reduced by about 20% among mothers who perceived their neighbourhoods lacking safe play areas for children. Policies to improve breastfeeding rates should consider area-based approaches and the broader determinants of social inequalities.

Keywords: breastfeeding, neighbourhood, social environment, deprivation, maternal perception, social inequalities.

Introduction

The promotion of breastfeeding is a fundamentally important public health issue globally. Mothers are advised to breastfeed infants exclusively for the first 6 months of life, thereafter it should be prolonged up to 24 months of age or beyond with complementary nutrition (WHO, 2018). Benefits for the infants encompass protection against sepsis, diarrhoea, respiratory infections (Khan *et al.*, 2015) and gastrointestinal infections (Kramer & Kakuma, 2012). Breastfeeding is also important for the development of nervous and endocrine systems (Ballard & Morrow, 2013). Long-term protection against type 2 diabetes, obesity (Horta *et al.*, 2015) and malocclusions (Peres *et al.*, 2015) have also been reported. Benefits are also extended to the mother protecting against type 2 diabetes, breast and ovarian cancers (Victora *et al.*, 2016).

Societal gains have also been evaluated. If every infant was breastfed until 6 months of age, cognitive deficits could be avoided with consequential global savings of US\$ 300 billion yearly (Rollins *et al.*, 2016). Assuming a moderate increase in breastfeeding rates, Renfrew *et al.* (2012) estimated annual NHS savings of about £48 million through a reduction in breast cancer cases and acute infant diseases.

Exclusive breastfeeding for the first six months of life varies worldwide, however, the global mean remained at around 36% over the period of 2007-2014 (WHO, 2009; WHO, 2015).

In the UK in 2010, the prevalence of breastfeeding fell from 81% at birth to 69% at week one and declined further to 55% at six weeks. At six months, 34% of mothers were still breastfeeding, however, only 1% were breastfeeding exclusively (McAndrew *et al.*, 2012). Additionally, the World Health Statistics 2015 show that exclusive breastfeeding rates for the first six months of life remained unchanged at around 1% and were amongst the lowest worldwide in the 2007-2014 year period (WHO, 2015).

Breastfeeding is a complex behaviour influenced by an array of individual, social and societal factors. One of these is the social environment where the mother resides. The socio-economic and political context stratify individuals socially, shaping intermediary determinants (e.g. living conditions) that may lead to social inequalities in child health (WHO, 2010). Whilst structural determinants of health remain paramount, understanding the mechanisms through which intermediary determinants such as neighbourhoods affect child health could contribute to effective public health interventions aiming to increase breastfeeding rates in the UK.

Neighbourhood effects on child outcomes can be direct or more likely indirect (Leventhal & Brooks-Gunn, 2000). Resources such as access to professional support (e.g. access to a Baby Friendly Initiative facility) and social relationships including social capital may facilitate positive health-related behaviours (Leventhal & Brooks-Gunn, 2000) which include breastfeeding. For instance, Tofani *et al.* (2015) found that low individual- and neighbourhood-level social capital were associated with less healthy diets throughout pregnancy. Additionally, collective efficacy referring to formal and informal control monitoring the behaviour of residents (Leventhal & Brooks-Gunn, 2000) can improve neighbours' perceptions of safety within their neighbourhoods (Uchida *et al.*, 2014), which in turn could contribute to the willingness of mothers to breastfeed in public spaces.

Spatial mobility and virtual networking may have widened their influence on people's everyday lives. However, contemporary neighbourhood's redefinition by Sampson *et al.* (2002) considers spatial dynamics wherein neighbourhoods are influenced by their surrounding areas, thereby revealing that through the lens of social processes neighbourhood boundaries extend further than mere census geography. Furthermore, there is considerable evidence that the neighbourhood remains important across a range of health-related behaviours.

Research examining neighbourhood effects on breastfeeding is limited and has produced some conflicting findings. Cubbin *et al.* (2008) using data collected via mailed questionnaire

and telephone (follow-up) from Florida and Washington found no association between neighbourhood-level deprivation and breastfeeding initiation. Similarly, Lagerberg *et al.* (2011) studied a convenience sample of Swedish mother-child dyads. They revealed no association between any breastfeeding at 4 months and neighbourhood socio-economic status. Conversely, a Swedish study (Almquist-Tangen *et al.*, 2013) and a Canadian study (Brown *et al.*, 2013) found variations in breastfeeding duration and exclusivity respectively by neighbourhood income deprivation. Burdette (2013) using a large sample of American low-income, unmarried, urban mothers revealed that living in a highly educated neighbourhood increased the likelihood of initiating and sustaining breastfeeding, whereas the percentage of immigrants, ethnic diversity and economic deprivation at neighbourhood level did not play a significant role. The use of selective study samples, as well as cultural and economic differences between countries, may explain why there are such contrasts in research findings.

Considering the wide-ranging benefits of breastfeeding along with low rates of exclusive breastfeeding in the UK, this study aims to explore whether breastfeeding initiation, exclusivity and duration were associated with neighbourhood context conceptualised by deprivation and maternal neighbourhood perceptions in a nationally representative sample of UK children.

Methodology

Sample

This study was based on the cross-sectional analysis of data from the first wave of the UK Millennium Cohort Study (MCS). The MCS is a multi-disciplinary project following the lives of approximately 19,000 children born in the UK during the period of 2000-2002. The MCS explores the social ecology in which the family is nested, approaching topics such as parenting, demography and social capital, and includes linked-in data on the neighbourhoods where the

families lived when the cohort baby was born. Therefore, the MCS is well placed to investigate neighbourhood effects on children's health. The sampling frame for the MCS was the electoral ward. A cluster random sample was drawn that was stratified to over-represent economically disadvantaged areas, areas with high proportions of people from ethnic minority backgrounds, and the three smaller countries of the UK. A detailed description of the MCS sampling methodology can be found elsewhere (Hansen, 2012).

At wave one, the children were 9 months old and biological mothers constituted 99.7% of main respondents. Among the 28 fathers who were main respondents, only 6 provided answers to the breastfeeding questions. Therefore, in our analyses cohort babies for whom the main respondent was not the biological mother were excluded. Also excluded were families whose cohort members were twins and triplets.

The MCS wave one gained ethical approval (MREC/01/6/19) from the South West Multi-Centre Research Ethics Committee in 2001 (Hansen, 2012).

Measures

Neighbourhood factors

Neighbourhood-level deprivation was measured via the Indices of Multiple Deprivation (IMD) for England (ODPM, 2004), Scotland (Scottish Executive, 2004), Wales (National Assembly for Wales, 2005) and Northern Ireland (NISRA, 2005). For each UK country, the IMD is constructed in a very similar way, including the following domains: (i) barriers to housing and services, (ii) crime, (iii) income, (iv) employment, (v) health and disability, (vi) living environment deprivation, and (vii) education, skills and training (ODPM, 2004). Linked into the MCS data are IMD rank deciles for each UK country, based on a weighted cumulative model of these domains (Noble et al., 2008). For the purposes of this study, the rank deciles for each UK country were combined into a single variable and categorised into quintiles.

Maternal neighbourhood perceptions were operationalised at the individual level. *Neighbourhood satisfaction* was assessed by asking the mother: ‘How satisfied or dissatisfied are you with the area you live in? By your area, I mean within about a mile or 20 minutes’ walk from here’. The response categories were ‘very satisfied’, ‘fairly satisfied,’ ‘neither satisfied nor dissatisfied’, ‘fairly dissatisfied’ and ‘very dissatisfied’. *Neighbourhood friendliness* measured maternal perceptions about neighbours by asking mothers ‘Please choose the phrase that you feel applies to most of your neighbours’. Response categories were ‘friendly’, ‘neither friendly nor unfriendly’, ‘unfriendly’ and ‘cannot say’. *Neighbourhood safety for the child* was a dichotomous variable (‘yes’/ ‘no’) obtained by asking the mother: ‘Are there any places in your area where children can play safely?’

Breastfeeding outcomes

Thresholds for the breastfeeding outcomes were chosen based on the literature and relevant recommendations by the UK Department of Health and WHO (2018). *Breastfeeding initiation*: Breastfeeding initiation has been defined as the mother putting the baby to the breast or giving her breast milk to the baby within a period of 48 hours after birth (Dyson et al., 2006). In the MCS, breastfeeding initiation was measured by asking the mother ‘Did you ever try to breastfeed [baby]?’ The variable was dichotomised into ‘no’ and ‘yes’. *Exclusive breastfeeding for at least 3 months*: WHO defines exclusive breastfeeding as the infant receiving only breast milk, including milk expressed or from a wet nurse, and no other liquids or solids with the exception of ORS (Oral Rehydration Solutions), drops or syrups consisting of vitamins, mineral supplements, or medicines (WHO, 2017). We derived this variable using information about the age of the child when breastfeeding stopped (‘How old was [baby] when he/she last had breast milk?’) and the age any other type of milk was introduced (‘How old was [baby] when he/she first had formula milk, such as Cow & Gate or SMA?’). Exclusive breastfeeding was dichotomised into ‘none to less than 3 months’ and ‘3 months or more’. It should be noted that at the time of the first wave of the MCS (2000-2002), the UK Department of Health

advised mothers to breastfeed exclusively for at least four months with the introduction of solid foods thereafter (Department of Health, 1994). The 3-month cut-off was chosen because at 4 months only 613 (3.4%) women in the sample were exclusively breastfeeding. *Any breastfeeding for at least 6 months* encompassed babies exclusively, predominantly or partially breastfed (WHO, 2008). Any breastfeeding was dichotomised into ‘none to less than 6 months’ and ‘6 months or more’.

Covariates

As we wanted to know whether neighbourhood factors contributed to breastfeeding rates over and above individual- and familial-level socio-demographic characteristics, it was necessary to control for individual-level socio-demographic background that may confound this relationship. These variables were specified a priori, informed by a review of the literature. Confounding factors comprised household and maternal characteristics. *Household structure* summarised the number of parents within the household dichotomised into ‘two parents’ and ‘single parent’. *Household income* was measured using OECD equivalised weekly family income divided into weighted quintiles. *Household social class* was measured by the National Statistics Socio-economic Classification (Rose *et al.*, 2005). We combined maternal and paternal social class into a single variable wherein the highest social class of either partner was considered and categorised into ‘managerial & professional’, ‘intermediate’, ‘small employers & self-employed’, ‘low supervisory & technical’, ‘semi-routine & routine’, ‘never worked’, and ‘not classifiable’. *Residential mobility* measured the total time the family had been living at the current address as follows: ‘more than 5 years’, ‘more than 1 year up to 5’ and ‘up to 1 year’. *Maternal age* was used as a continuous variable (unit = years) and also as a categorical variable (four age groups). The former was used in the regression analysis and the latter in the descriptive analysis of the sample. *Maternal general health* was derived from the question: ‘How would you describe your health generally?’ The response categories comprised ‘excellent’, ‘good’, ‘fair’ and ‘poor’. *Maternal longstanding illness* was a dichotomous variable

derived by probing the mother: ‘Do you have a longstanding illness, disability or infirmity?’ *Maternal education* was measured using the National Vocational Qualification (NVQ) classification. The derived NVQ variable considers both vocational and academic qualifications. It was recategorised as follows: ‘NVQ levels 4 & 5’ (e.g. degree or higher degree), ‘NVQ level 3’ (e.g. 2+ A levels), ‘NVQ level 2’ (e.g. 5 General Certificate of Secondary Education A–C or 1 A level), ‘NVQ level 1’ (e.g. < 5 General Certificate of Secondary Education D–E) ‘none’ and ‘overseas qualification only’. *Maternal ethnicity* was measured using six categories: ‘White’, ‘Mixed’, ‘Indian’, ‘Pakistani & Bangladeshi’, ‘Black or Black British’, and ‘other ethnic groups’.

Statistical analysis

The data were analysed using STATA version 14. The complex survey design of the MCS was accounted for by using the STATA ‘svy’ command followed by variables for the stratification design, clustering effect and finite population correction, in addition to the MCS overall weights which are the inverse of the predicted probability of participation in a wave combined with the sampling weights.

For descriptive statistics, an initial assessment of neighbourhood exposures and covariates with each breastfeeding outcome was carried out using chi-squared tests accounting for the survey design. The initial assessment of neighbourhood exposures and covariates with each breastfeeding outcome revealed that all differences in the proportions were statistically significant except for breastfeeding initiation and longstanding illness. P-values are not included in Tables 1 and 2 as they were deemed redundant given that trends are clear from the presented data. We conducted a complete case analysis as the rate of missingness for any variable did not exceed 5%, which is considered an appropriate threshold (Schafer, 1999). Characteristics of participants with missing data on exposures and covariates were explored using chi-squared tests accounting for the survey design. For the neighbourhood factors,

multicollinearity was tested and ruled out, and further investigation of possible interactions between maternal neighbourhood satisfaction and IMD revealed no clear associations.

For each of the three breastfeeding outcomes, a series of four multivariable logistic regression models was estimated. We applied a theoretical approach to our model construction using control variables that explain breastfeeding initiation and duration, and no variable was removed based on p-values. The following modelling strategy was developed: (i) Model 1 included the IMD quintiles; (ii) Model 2 additionally adjusted for maternal neighbourhood perceptions (i.e. neighbourhood satisfaction, neighbourhood friendliness and neighbourhood safety for the child); (iii) Model 3 added to Model 2 maternal age and SES variables (i.e. income quintiles, NS-SEC and maternal education), and (iv) the fully adjusted model, additionally including maternal ethnicity, household structure, maternal general health, longstanding illness and residential mobility. The interpretation of the models was carried out with Wald tests. Statistical significance was defined at the 0.05 level. Logistic regression results are presented as odds ratios with their 95% confidence intervals and p-values.

Results

Sample characteristics

Descriptive statistics are displayed in Tables 1 and 2. The exclusion criteria resulted in an original sample comprising 18,239 children. Missingness on outcomes totalled 5 observations (0.01%) that were excluded, resulting in 18,234 participants where missingness on exposures was highest (2.3%) for neighbourhood friendliness. Therefore, we conducted a complete case analysis for which a listwise deletion resulted in a final sample size of 17,308 children. Mothers with missing information for any of the exposures or covariates were more likely to live in deprived neighbourhoods, to belong to families with lower incomes, to live in households whose partners had never worked or to have a non-classifiable occupation (data not shown).

Breastfeeding outcomes and neighbourhood factors are displayed in Table 1. Breastfeeding decreased from the highest IMD quintile i.e. the least deprived neighbourhoods to the lowest quintile (most deprived) in a consistent stepwise fashion. Comparing the most deprived areas to the least deprived neighbourhoods, breastfeeding initiation rates were 56.0% versus 82.8%, whilst for exclusive and any breastfeeding for at least 3 and 6 months respectively, rates were 11.2% versus 25.8% and 12.7% versus 26.6%. All three breastfeeding outcomes were more favourable among mothers who reported higher levels of neighbourhood satisfaction, who perceived their neighbours as friendlier, and who said that there were places in the neighbourhood where children could play safely. For example, breastfeeding was initiated by 74.2% of mothers who were very satisfied with their neighbourhood compared to 57.7% who were very dissatisfied.

Table 2 shows breastfeeding outcomes by household and maternal characteristics. All breastfeeding outcomes were positively associated with a two-parent household, a higher household income, and a higher social class. For example, only 8.7% of mothers in families who were in semi-routine and routine occupations exclusively breastfed for at least 3 months compared to 27.4% of those in managerial and professional occupations. Breastfeeding rates were the lowest among mothers residing in the neighbourhood up to 1 year and increased proportionally with increasing maternal age. The prevalence of each breastfeeding outcome was highest among mothers who self-reported their general health as excellent, who had no longstanding illness, whose levels of education comprised NVQ levels 4 and 5, and whose ethnicity was non-White. The exception was exclusive breastfeeding for at least 3 months, for which the prevalence was slightly lower among Pakistani and Bangladeshi mothers compared to White mothers.

Breastfeeding initiation

In the bivariate analysis, the association between neighbourhood deprivation and breastfeeding initiation was inverse and significant (Model 1 in Table 3). Odds were lower for mothers living in the most deprived areas compared to those living in the least deprived neighbourhoods (OR = 0.25; 95% CI 0.20-0.32). The associations with maternal neighbourhood perceptions were positive and significant. For instance, the likelihood of breastfeeding initiation was reduced among mothers who could not express feelings about their neighbours compared to those who perceived them as friendly (OR = 0.64; 95% CI 0.49-0.82). This association, however, was not entirely linear. Results from the bivariate analyses for maternal neighbourhood perceptions are available as supplementary material.

In the multivariable analysis (Table 3), controlling for maternal age and socio-economic factors (Model 3) substantially attenuated the association between breastfeeding initiation and quintiles of IMD, which remained however statistically significant. Compared to Model 3, the association appeared to be slightly stronger after full adjustment (Model 4). The likelihood of breastfeeding initiation was 40% lower in the most deprived neighbourhoods compared to the least deprived areas (OR = 0.60; 95% CI 0.50-0.72). Very dissatisfied mothers were significantly more likely to initiate breastfeeding relative to very satisfied mothers (OR = 1.32; 95% CI 1.09-1.61). Odds for breastfeeding initiation were higher by 20% among mothers with neutral feelings about neighbours relative to those with feelings of friendliness (OR = 1.20; 95% CI 1.06-1.36). Alternatively, odds lowered by about 20% among mothers whose neighbourhoods were perceived as lacking places for children to play safely compared to those who perceived their neighbourhoods as safe to play (OR = 0.78; 95% CI 0.71-0.85).

Associations between breastfeeding initiation and maternal education and household income revealed a clear social gradient. Compared to those in the richest quintile, mothers in the poorest household income quintile were less likely to initiate breastfeeding (OR = 0.59; 95% CI 0.48-0.73). Odds were also significantly lower for families from all lower social classes compared to those in managerial and professional occupations, and for single mothers

compared to those living with a partner (OR = 0.78; 95% CI 0.68-0.88). Odds for breastfeeding initiation were significantly higher for older versus younger mothers, and for mothers from minority ethnic groups compared to White mothers (OR for Black or Black British mothers = 11.0; 95% CI 7.49-16.2). Conversely, increasing the time of residence within the neighbourhood linearly reduced the likelihood of initiating breastfeeding.

Exclusive breastfeeding for at least 3 months

In the bivariate analysis, the association between neighbourhood deprivation and exclusive breastfeeding at 3 months was inverse and significant (Model 1 in Table 4). Odds were significantly lower comparing mothers living in the most deprived areas to those living in the least deprived neighbourhoods (OR = 0.35; 95% CI 0.28-0.45). There was a positive and significant association with all three variables of maternal neighbourhood perceptions. With neighbourhood friendliness, only mothers who could not express feelings about their neighbours compared to those with feelings of friendliness had significantly reduced odds for breastfeeding exclusively at 3 months (OR = 0.67; 95% CI 0.47-0.97). Results from the bivariate analyses for maternal neighbourhood perceptions are available as supplementary material.

In the multivariable analysis (Table 4), adjusting for maternal age and socio-economic factors (Model 3), the associations with IMD and neighbourhood satisfaction were no longer statistically significant. In the full adjustment (Model 4), mothers who perceived their neighbourhood as lacking safe play areas for children were about 16% less likely to breastfeed exclusively for at least 3 months relative to mothers with the perception of safe play areas in the neighbourhood (OR = 0.84; 95% CI 0.75-0.95).

Except for families where no parent (mother or father) had ever worked, odds of exclusive breastfeeding for at least 3 months were significantly lower for families from all lower social classes compared to those in managerial and professional occupations, for all lower levels of

maternal education compared to NVQ levels 4 and 5, particularly for mothers with NVQ level 1 (OR = 0.33; 95% CI 0.25-0.42), and among single mothers compared to those living with a partner (OR = 0.70; 95% CI 0.57-0.85). Conversely, mothers reporting better general health presented significantly higher odds for exclusive breastfeeding for at least 3 months. Odds were also significantly higher per each year increase in maternal age (OR = 1.06; 95% CI 1.05-1.07), and for mothers from minority ethnic groups compared to White mothers, especially among those from Mixed ethnicity (OR = 2.46; 95% CI 1.68-3.60).

Any breastfeeding for at least 6 months

In the bivariate analysis, again as neighbourhood deprivation increased, odds for continuing breastfeeding at 6 months decreased (Model 1 in Table 5). Mothers living in the most deprived areas had significantly lower odds to breastfeed at 6 months compared to those living in the least deprived neighbourhoods (OR = 0.39; 95% CI 0.29-0.52). There was also a positive and significant association with all three variables of maternal neighbourhood perceptions. However, the association with neighbourhood satisfaction was not entirely linear. Again, results from the bivariate analyses are available as supplementary material.

In the multivariable analysis (Table 5), the full adjustment (Model 4) reveals that mothers living in the second most deprived neighbourhoods were around 20% less likely to continue with any type of breastfeeding for at least 6 months compared to those living in the least deprived areas (OR = 0.79; 95% CI 0.63-0.99). Model 4 in Table 5 also shows that mothers who perceived their neighbourhoods as lacking safe play areas for children were about 18% less likely to continue breastfeeding for at least 6 months relative to those with the perception of neighbourhoods having safe play areas (OR = 0.82; 95% CI 0.73-0.93).

Odds for continuing breastfeeding for at least 6 months were significantly higher per each year increase in maternal age, for all levels of household income compared to the highest quintile (i.e. the richest) except for the second highest, and for mothers from minority ethnic

groups compared to White mothers, especially among those from the Other ethnic group (OR = 3.96; 95% CI 2.78-5.64). Conversely, apart from families in small & self-employed and with non-classifiable occupations, odds were significantly lower for those from lower social classes compared to families in managerial and professional occupations, and for all lower levels of maternal education compared to NVQ levels 4 and 5, particularly for mothers with NVQ level 1 (OR = 0.30; 95% CI 0.23-0.39). The likelihood of breastfeeding at 6 months also lowered among single mothers compared to those living with a partner, and among mothers reporting fair or poor health compared to those in excellent health.

Discussion

To our knowledge, this is the first study in the UK that has explored the relationship between breastfeeding and neighbourhood deprivation adjusting for a comprehensive set of familial- and individual-level factors. In addition, this is the first study to explore the association between maternal neighbourhood perceptions and breastfeeding.

In the bivariate analysis, all three breastfeeding outcomes were negatively associated with neighbourhood deprivation and positively associated with more favourable maternal neighbourhood perceptions. In the full adjustment, breastfeeding initiation was independently and negatively associated with neighbourhood deprivation. There also appears to be an inverse relationship between exclusive and any breastfeeding for at least 3 and 6 months respectively and neighbourhood deprivation. All three breastfeeding outcomes were independently and positively associated with the maternal perception of the neighbourhood having safe play areas for children.

The results of this study showed that mothers living in the most deprived neighbourhoods were 40% less likely to initiate breastfeeding compared to those living in the least deprived areas. This finding was consistent with Bonet et al. (2013) who revealed that in the UK (Trent)

and Ile-de France, breastfeeding at discharge from hospital was lower in neighbourhoods presenting the highest unemployment rates than those with the lowest, and with Oakley et al. (2013) who found that outside London, Primary Care Trusts in the most deprived quintile had a 32% reduced odds of breastfeeding initiation compared to those in the least deprived quintile.

Additionally, there seems to be a negative relationship between both exclusive breastfeeding for at least 3 months and any type of breastfeeding for at least 6 months and neighbourhood deprivation. However, these relationships appeared less strong and not entirely linear once other variables were taken into account. Mothers living in the second most deprived neighbourhoods were about 20% less likely to continue breastfeeding for at least 6 months compared to those living in the least deprived areas. Findings for exclusive and any breastfeeding were consistent with Oakley et al. (2013) across non-London Primary Care Trusts, in which odds for exclusive and any breastfeeding at 6-8 weeks were negatively associated with deprivation at the area level measured by the 2010 Index of Multiple Deprivation. In their findings, however, the association with any breastfeeding at 6-8 weeks was stronger and linear.

Breastfeeding initiation appears to be indirectly influenced by neighbourhood deprivation. In poor quality neighbourhoods, for instance, there may not be the support systems in place to encourage mothers to initiate breastfeeding, and perhaps socialisation with family and friends for whom formula feeding is the norm could discourage breastfeeding. Moreover, there might be a lack of or insufficient encouragement from institutional support systems including health providers. Furthermore, education, social class, and some medical conditions such as obesity can potentially influence breastfeeding initiation (McAndrew *et al.*, 2012; Turcksin *et al.*, 2014; Jonas & Woodside, 2016). The clustering of such health conditions with low education and lower social classes appear to be a characteristic of deprived areas in England (Marmot, 2010).

Neighbourhood effects on exclusive and any type of breastfeeding are also more likely to be indirect, and particularly influenced by familial and individual levels. Proximal determinants such as self-efficacy, intention and planning appear to be paramount for exclusive breastfeeding (de Jager *et al.*, 2013; Dennis *et al.*, 2014), whilst the perception of insufficient milk supply, postpartum depression and familial support can be crucial for breastfeeding duration (Thulier & Mercer, 2009; Dias & Figueiredo, 2015). Furthermore, individual- and familial-level characteristics are likely to interact with neighbourhood characteristics (Diez Roux & Mair, 2010; Schüle & Bolte, 2015) influencing breastfeeding duration and exclusivity.

Maternal neighbourhood satisfaction was inversely associated with breastfeeding initiation. Research on the interpretation of this subjective measure of neighbourhood context seems to be scarce. However, neighbourhood satisfaction is complex as it appears to be rooted in personal, psychological, and social factors over and above the physical environment (Hur *et al.*, 2010). Moreover, neighbourhood perceptions are likely to vary by urban/rural residence (Salmon *et al.*, 2013; De Vos *et al.*, 2016). In England and Scotland, sense of belonging and area satisfaction were perceived slightly higher among rural residents (Pateman, 2011). Neighbourhood race/ethnic composition was also reported to play a significant role in individual and neighbourhood satisfaction (Swaroop & Krysan, 2011; Knies *et al.*, 2016). In this sample, specific institutional or social processes at the neighbourhood level that influenced mothers' levels of satisfaction were unknown. Perhaps, mothers might have perceived the area where they lived as unsatisfactory due to adverse physical environments such as traffic noise or living density, rather than accessibility to health facilities equipped with professionals trained in breastfeeding.

Breastfeeding initiation was higher among mothers who expressed neutral feelings about neighbours compared to those with feelings of friendship. This does not necessarily suggest that positive social support is negatively associated with breastfeeding initiation. Instead,

mothers who live in areas where they are impersonal about the relationship with their neighbours might have stronger social networks outside the immediate neighbourhood. An example might be minority ethnic groups who are more likely to breastfeed in the UK (Kelly *et al.*, 2006; Baker *et al.*, 2011). Ethnic minority mothers may have their culture and beliefs about initiation of breastfeeding preserved. Indeed, conviviality and friendship could induce mothers to the culture of using formula milk through a ‘contagious model’ (Jencks & Mayer, 1990) enabling a particular behaviour to become a norm among neighbours.

The likelihood for all breastfeeding outcomes was lowered by about 20% with the maternal perception of neighbourhoods lacking safe play areas for children. Such perceptions may indicate a more general feeling that the area is not safe. Associations between maternal perceptions of unsafe neighbourhoods with unfavourable child outcomes have been previously reported in the literature, for instance, with adverse mental health (Pettit *et al.*, 1999), obesity (Bacha, *et al.*, 2010), and asthma (Vangeepuram *et al.*, 2012). Institutional resources such as the availability of safe playgrounds and the social environment (e.g. social relationships) can be plausible pathways through which neighbourhood effects are transmitted to individuals influencing behaviour and child outcomes (Leventhal & Brooks-Gunn, 2000; Diez Roux & Mair, 2010; Christian *et al.*, 2015). In line with that, safe play areas in the neighbourhood could provide a meeting point for mothers to share positive health behaviours including breastfeeding.

This study has strengths and limitations. We used a large, nationally representative UK sample and additionally, conducted comprehensive adjustments for factors at the household and individual levels including residential mobility. Moreover, the inclusion of subjective neighbourhood measures was an important step towards exploring the role of perceived neighbourhood quality and social dimensions of breastfeeding.

However, if researchers are not explicit about the causal pathways hypothesised between neighbourhood constructs analysed and breastfeeding, inference is likely to be limited. Diez Roux (2004) explained that individual-level factors can be simultaneously mediators and confounders in neighbourhood effects on health. As noted by Diez Roux & Mair (2010), the cumulative exposure to impoverished areas early in life might reduce access to education and employment, and thus affect health later in life, whilst lack of education and low-paid jobs at the individual level may also be confounders to neighbourhood deprivation effects on health. Therefore, adjusting for individual- and familial-level factors in order to identify a direct effect of neighbourhood-level deprivation may have eliminated pathways that influenced exclusive and any breastfeeding for at least 3 and 6 months respectively.

Furthermore, the dichotomisation of breastfeeding duration could have caused a loss of potentially useful information. However, we were limited by the measurement in the MCS and additionally, a continuous variable would be subject to measurement error with peaks of reporting at the monthly intervals. Existing research demonstrates, whether for breastfeeding or unemployment spells, people find it difficult to report circumstances in continuous intervals of time. We have tested a threshold that was important at the time (i.e. 4 months) and a threshold now considered a critical point (i.e. 6 months). As previously mentioned, at the time of the first wave only 3.4% of women in the sample were breastfeeding exclusively at 4 months, therefore the threshold chosen was 3 months. Both thresholds (i.e. 3 months and 6 months) have been used in previous studies (e.g. Gore *et al.*, 2015; Hao *et al.*, 2017; O'Connor *et al.*, 2017). Ultimately, this was a cross-sectional study, and therefore causation cannot be inferred.

Our findings have potentially important policy implications. As a public health indicator, breastfeeding rates can be a good marker of social inequalities (Department of Health, 2016). Breastfeeding is determined by a range of interacting factors operating at different levels, therefore requiring a range of downstream, midstream and upstream strategies. We suggest that

policy makers should consider programmes to advocate breastfeeding more strongly in deprived neighbourhoods. The allocation of adequate resources such as the 'Baby Friendly Initiative' for deprived areas can be paramount to increase breastfeeding initiation, however, focusing solely at the time of birth could be reductionist. Therefore, multifaceted strategies designed on the social determinants of health inequalities concomitantly with individual and community empowerment are crucial (Marmot, 2010).

Policies to improve the physical and social environments of neighbourhoods along with the provision and maintenance of parks and amenities for children could indirectly result in an increase in the rates of breastfeeding initiation and duration. Nevertheless, the success of public health interventions relies upon targeting *all* determinants of breastfeeding including not only proximal factors such as professional support, but also distal factors such as public policies promoting breastfeeding in public spaces, and the regulation of marketing practice of the infant formula industry.

In conclusion, neighbourhoods and breastfeeding are both multidimensional constructs, making it challenging to provide specific recommendations. Our main finding was that breastfeeding in the UK seems to be associated with the environment over and above individual background. Therefore, multifaceted and context-led interventions seem necessary along with strategies targeting social inequalities. Future research should aim to address the issue of selection bias inherent to residential mobility using longitudinal data and causal methods of analysis, in addition to qualitative techniques to explore women's views about initiating and maintaining breastfeeding in relation to their environment.

Key messages:

- In the UK, as neighbourhood deprivation increased, odds for breastfeeding initiation decreased: odds lowered by 40% among mothers living in the most deprived areas compared to those living in the least deprived neighbourhoods.
- The likelihood of breastfeeding initiation, exclusivity at 3 months or more, and any breastfeeding for at least 6 months each reduced by about 20% with the maternal perception of neighbourhoods lacking safe play environments for children compared to neighbourhoods perceived as safe to play.
- Addressing social inequalities can be paramount in increasing the effectiveness of multifaceted and context-led interventions aiming at improving breastfeeding rates

References

- Almquist-Tangen, G., Strömberg, U., Holmén, A., Alm, B., Roswall, J., Bergman, S., et al. (2013). Influence of neighbourhood purchasing power on breastfeeding at four months of age: a Swedish population-based cohort study. *BioMed Central Public Health*, **13**, 1077.
- Bacha, J.M., Appugliese, D., Coleman, S., Kaciroti, N., Bradley, R.H., Corwyn, R.F., et al. (2010). Maternal perception of neighborhood safety as a predictor of child weight status: the moderating effect of gender and assessment of potential mediators. *International Journal of Pediatric Obesity*, **5**, 72-79.
- Baker, D., Garrow, A. & Shiels, C. (2011). Inequalities in immunisation and breast feeding in an ethnically diverse urban area: cross-sectional study in Manchester, UK. *Journal of Epidemiology and Community Health*, **65**, 346–352.
- Ballard, O. & Morrow, A.L., (2013). Human milk composition: nutrients and bioactive factors. *Pediatric Clinics of North America*, **60**, 49–74.
- Bonet, M., Smith, L., Pilkington, H., Draper, E. & Zeitlin, J. (2013). Neighbourhood deprivation and very preterm birth in an English and French cohort. *BioMed Central Pregnancy and Childbirth*, **13**, 97.
- Brown, C.R.L., Dodds, L., Attenborough, R., Bryanton, J., Rose, A.E., Flowerdew, G., et al. (2013). Rates and determinants of exclusive breastfeeding in first 6 months among women in Nova Scotia: a population-based cohort study. *Canadian Medical Association Journal Open*, **1**, e9-e17.
- Burdette, A.M. (2013). Neighborhood context and breastfeeding behaviors among urban mothers. *Journal of Human Lactation*, **29**, 597-604.
- Christian, H., Zubrick, S.R., Foster, S., Giles-Corti, B., Bull, F., Wood, L., Knuiaman, M., Brinkman, S., Houghton, S. and Boruff, B. (2015). The influence of the neighborhood physical environment on early child health and development: A review and call for research. *Health & place*, **33**, pp.25-36.
- Cubbin, C., Marchi, K., Lin, M., Bell, T., Marshall, H., Miller, C., et al. (2008). Is neighborhood deprivation independently associated with maternal and infant health? evidence from Florida and Washington. *Maternal and Child Health Journal*, **12**, 61-74.
- de Jager, E., Skouteris, H., Broadbent, J., Amir, L. & Mellor, K. (2013). Psychosocial correlates of exclusive breastfeeding: a systematic review. *Midwifery*, **29**, 506–518.
- De Vos, J., Van Acker, V. & Witlox, F. (2016). Urban sprawl: neighbourhood dissatisfaction and urban preferences. Some evidence from Flanders. *Urban Geography*, **37**, pp.839-862.
- Dennis, C., Gagnon, A., Van Hulst, A. & Dougherty, G. (2014). Predictors of breastfeeding exclusivity among migrant and Canadian-born women: results from a multi-centre study. *Maternal and Child Nutrition*, **10**, 527–544.
- Department of Health (1994). Weaning and the weaning diet. Report of the working group on the weaning diet of the Committee on Medical Aspects of Food Policy. Report on Health and Social Subjects No 45. London: HMSO.
- Department of Health (2016). Government response to the consultation refreshing the public health outcomes framework (2015). [pdf]. London: Department of Health. Available: <https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/520455/PH_OF_cons_response.pdf> [Accessed 18 August 2016].

- Dias, C.C. & Figueiredo, B. (2015). Breastfeeding and depression: a systematic review of the literature. *Journal of Affective Disorders*, **171**, 142-154.
- Diez Roux, A.V. (2004). Estimating neighborhood health effects: the challenges of causal inference in a complex world. *Social Science and Medicine*, **58**, 1953-1960.
- Diez Roux, A.V. & Mair, C. (2010). Neighborhoods and health. *Annals of the New York Academy of Sciences*, **11861**, 125-145.
- Dyson, L., Renfrew, M., McFadden, A., McCormick, F., Herbert, G. & Thomas, J. (2006). Promotion of breastfeeding initiation and duration. Evidence into practice briefing. [pdf]. London: National Institute for Health and Clinical Excellence. Available: <http://www.breastfeedingmanifesto.org.uk/doc/publication/EAB_Breastfeeding_final_version_1162237588.pdf> [Accessed 25 January 2016].
- Gore, N., Emerson, E., & Brady, S. (2015). Rates of breastfeeding and exposure to socio-economic adversity amongst children with intellectual disability. *Research in developmental disabilities*, **39**, 12-19.
- Hansen, K. (2012). Millennium Cohort Study First, Second, Third and Fourth Surveys: A Guide to the Datasets (7th ed.). London: Centre for Longitudinal Studies, Institute of Education, University of London.
- Hao, N., Jiang, H., Wu, M., Pan, T., Yan, B., Liu, J., et al., (2017). Breastfeeding initiation, duration and exclusivity in mothers with epilepsy from South West China. *Epilepsy research*, **135**, 168-175.
- Horta B.L., Loret de Mola C. & Victora C.G. (2015). Long-term consequences of breastfeeding on cholesterol, obesity, systolic blood pressure and type 2 diabetes: a systematic review and meta-analysis. *Acta Paediatrica*, **104**, 30–37.
- Hur, M., Nasar, J.L. and Chun, B. (2010). Neighborhood satisfaction, physical and perceived naturalness and openness. *Journal of Environmental Psychology*, **30**, 52-59.
- Jencks, C. & Mayer, S. (1990). The social consequences of growing up in a poor neighborhood. In: E. Lynn & M. McGahey (eds.) *Inner-City Poverty in the United States*, pp. 111-186. Washington, DC: National Academy Press.
- Jonas, W. & Woodside, B. (2016). Physiological mechanisms, behavioral and psychological factors influencing the transfer of milk from mothers to their young. *Hormones and Behavior*, **77**, 167-181.
- Kelly, Y.J., Watt, R.G. & Nazroo, J.Y. (2006). Racial/ethnic differences in breastfeeding initiation and continuation in the United Kingdom and comparison with findings in the United States. *Pediatrics*, **118**, e1428-e1435.
- Khan, J., Vesel, L., Bahl, R. & Martines, J.C. (2015). Timing of breastfeeding initiation and exclusivity of breastfeeding during the first month of life: effects on neonatal mortality and morbidity — a systematic review and meta-analysis. *Maternal and Child Health Journal*, **19**, 468-479.
- Knies, G., Nandi, A. & Platt, L., (2016). Life satisfaction, ethnicity and neighbourhoods: Is there an effect of neighbourhood ethnic composition on life satisfaction? *Social science research*, **60**, 110-124.
- Kramer, M.S. & Kakuma, R. (2012). Optimal duration of exclusive breastfeeding. *Cochrane Database of Systematic Reviews*, Aug 15 (8), CD003517.

- Lagerberg, D., Magnusson, M. & Sundelin, C. (2011). Child health and maternal stress: does neighbourhood status matter? *International Journal of Adolescent Medicine and Health*, **23**, 19-25.
- Leventhal, T. & Brooks-Gunn, J. (2000). The neighborhoods they live in: the effects of neighborhood residence on child and adolescent outcomes. *Psychological Bulletin*, **126**, 309–337.
- Marmot, M. (2010). Fair society, healthy lives: The Marmot Review. Strategic review of health inequalities in England post-2010. [pdf]. London: The Marmot Review. Available: <<http://www.instituteofhealthequity.org/Content/FileManager/pdf/fairsocietyhealthylives.pdf>> [Accessed 17 August 2016].
- McAndrew, F., Thompson, J., Fellows, L., Large, A., Speed, M. & Renfrew, M.J. (2012). Infant Feeding Survey 2010. [pdf]. Leeds: Health and Social Care Information Centre. Available: <<http://www.hscic.gov.uk/catalogue/PUB08694/Infant-Feeding-Survey-2010-Consolidated-Report.pdf>> [Accessed 17 April 2016].
- National Assembly for Wales (2005). The Welsh Index of Multiple Deprivation 2005: Local Authority Analysis. [pdf]. Available: <<http://gov.wales/docs/statistics/2005/051208wimd2005localreviseden.pdf>> [Accessed 19 June 2016].
- NISRA (2005). Northern Ireland Multiple Deprivation Measure 2005. [pdf]. Belfast: Statistics and Research Agency. Available: <<http://www.nisra.gov.uk/archive/deprivation/nimdm2005fullreport.pdf>> [Accessed 19 June 2016].
- Noble, M., McLennan, D., Wilkinson, K., Whitworth, A., Barnes, H. and Dibben, C. (2008). The English Indices of Deprivation 2007. [pdf]. London: Department for Communities and Local Government. Available: <http://geoconvert.mimas.ac.uk/help/imd-2007-manual.pdf> [Accessed 19 June 2016].
- Oakley, L.L., Renfrew, M.J., Kurinczuk, J.J. & Quigley, M.A. (2013). Factors associated with breastfeeding in England: an analysis by primary care trust. *British Medical Journal Open*, **3**, e002765.
- O'Connor, M., Allen, J., Kelly, J., Gao, Y. & Kildea, S. (2017). Predictors of breastfeeding exclusivity and duration in a hospital without Baby Friendly Hospital Initiative accreditation: A prospective cohort study. *Women and Birth*.
- ODPM (2004). The English Indices of Deprivation 2004: Revised. [pdf]. London: Office of the Deputy Prime Minister. Available: <http://www.simonpoulter.co.uk/ioid/ioidpdf/odpm_urbpol_029534.pdf> [Accessed 19 June 2016].
- Pateman, T. (2011). Rural and urban areas: comparing lives using rural/urban classifications. *Regional Trends*, **43**, pp.11-86.
- Peres, K.G., Cascaes, A.M., Nascimento, G.G. & Victora, C.G. (2015). Effect of breastfeeding on malocclusions: a systematic review and meta-analysis. *Acta Paediatrica*, **104**, 54-61.
- Pettit, G.S., Bates, J.E., Dodge, K.A. & Meece, D.W. (1999). The impact of after-school peer contact on early adolescent externalizing problems is moderated by parental monitoring perceived neighborhood safety, and prior adjustment. *Child Development*, **70**, 768-778.
- Renfrew, M.J., Pokhrel, S., Quigley, M., McCormick, F., Fox-Rushby, J., Dodds, R., et al. (2012). Preventing disease and saving resources: the potential contribution of increasing

- breastfeeding rates in the UK. [pdf]. UNICEF UK. Available: http://www.unicef.org.uk/Documents/Baby_Friendly/Research/Preventing_disease_saving_resources.pdf > [Accessed 16 April 2016].
- Rollins, N.C., Bhandari, N., Hajeebhoy, N., Horton, S., Lutter, C.K., Martines, J.C., et al. (2016). Why invest, and what it will take to improve breastfeeding practices? *The Lancet*, **387**, 491-504.
- Rose, D., Pevalin, D.J. & O'Reilly, K. (2005). The National Statistics Socio-economic Classification: origins, development and use. Basingstoke: Palgrave Macmillan.
- Salmon, J., Veitch, J., Abbott, G., ChinAPaw, M., Brug, J.J., Cleland, V., Hume, C., Crawford, D. & Ball, K. (2013). Are associations between the perceived home and neighbourhood environment and children's physical activity and sedentary behaviour moderated by urban/rural location? *Health & place*, **24**, 44-53.
- Sampson, R.J., Morenoff, J.D. & Gannon-Rowley, T. (2002). Assessing "Neighbourhood effects": social processes and new directions in research. *Annual Review of Sociology*, **28**, 443-478.
- Schafer, J. L. (1999). Multiple imputation: a primer. *Statistical methods in medical research*, **8**, 3-15.
- Schüle, S.A. & Bolte, G. (2015). Interactive and independent associations between the socioeconomic and objective built environment on the neighbourhood level and individual health: a systematic review of multilevel studies. *Public Library of Science One*, **10**, e0123456.
- Scottish Executive (2004). Using the Scottish Index of Multiple Deprivation 2004: Guidance. [pdf]. Edinburgh: Scottish Executive. Available: <http://www.gov.scot/Resource/Doc/35596/0025690.pdf> > [Accessed 19 June 2016].
- Swaroop, S. & Krysan, M., (2011). The determinants of neighborhood satisfaction: Racial proxy revisited. *Demography*, **48**, 1203-1229.
- Thulier, D. & Mercer, J. (2009). Variables associated with breastfeeding duration. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, **38**, 259-268.
- Tofani, A.A., Lamarca, G.D.A., Sheiham, A. & Vettore, M.V. (2015). The different effects of neighbourhood and individual social capital on health-compromising behaviours in women during pregnancy: a multi-level analysis. *BioMed Central Public Health*, **15**, 890.
- Turcksin, R., Bel, S., Galjaard, S. & Devlieger, R. (2014). Maternal obesity and breastfeeding intention, initiation, intensity and duration: a systematic review. *Maternal and Child Nutrition*, **10**, 166-183.
- Uchida, C.D., Swatt, M.L., Solomon, S.E., & Varano, S. (2014). Neighborhoods and Crime: Collective Efficacy and Social Cohesion in Miami-Dade County. United States Department of Justice, No. 245406.
- Vangeepuram, N., Galvez, M.P., Teitelbaum, S.L., Brenner, B. & Wolff, M.S. (2012). The association between parental perception of neighborhood safety and asthma diagnosis in ethnic minority urban children. *Journal of Urban Health*, **89**, 758-768.
- Victora, C.G., Bahl, R., Barros, A.J.D., França, G.V.A., Horton, S., Krasevec, J. et al. (2016). Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *The Lancet*, **387**, 475-490.

World Health Organization (2008). Indicators for assessing infant and young child feeding practices: Conclusions of a consensus meeting held 6–8 November 2007 in Washington D.C., USA. Geneva : WHO Press.

World Health Organization (2009). World Health Statistics. [pdf]. Geneva: World Health Organization. Available:
http://www.who.int/gho/publications/world_health_statistics/EN_WHS09_Full.pdf
 [Accessed 16 April 2016]

World Health Organization (2010). A conceptual framework for action on the social determinants of health. [pdf]. Geneva: World Health Organization. Available:
http://www.who.int/sdhconference/resources/ConceptualframeworkforactiononSDH_eng.pdf [Accessed 28 December 2015].

World Health Organization (2015). World Health Statistics. [pdf]. Geneva: World Health Organization. Available:
http://apps.who.int/iris/bitstream/10665/170250/1/9789240694439_eng.pdf [Accessed 16 April 2016].

World Health Organization (2017). The World Health Organization's infant feeding recommendation. Nutrition. [Online]. Available:
http://www.who.int/nutrition/topics/infantfeeding_recommendation/en/ [Accessed 7 November 2017].

World Health Organization (2018). Infant and young child feeding. Fact Sheet (updated July 2017). WHO Media Centre. [Online]. Available:
<http://www.who.int/mediacentre/factsheets/fs342/en/> > [Accessed 22 March 2018].

Table 1. Descriptive statistics: Breastfeeding outcomes by neighbourhood characteristics.

<i>Neighbourhood Factors</i>	<i>18,234 (N)^a</i>	<i>Breastfeeding Initiation</i>	<i>Exclusive Breastfeeding</i>	<i>Any Breastfeeding^b</i>
	<i>n</i>	<i>Yes (%)</i>	<i>≥ 3 months (%)</i>	<i>≥ 6 months (%)</i>
<i>Neighbourhood-level Deprivation</i>				
<i>IMD</i>				
Highest Quintile	2687	82.8	25.8	26.6
2nd Highest	2444	77.9	23.9	24.3
Middle quintile	3030	71.7	19.1	18.7
2nd lowest	4165	62.8	14.3	15.4
Lowest quintile	5908	56.0	11.2	12.7
<i>Maternal Neighbourhood Perceptions</i>				
<i>Neighbourhood Satisfaction</i>				

Very satisfied	7375	74.2	22.2	22.5
Fairly satisfied	7332	67.1	16.8	17.5
Neither	1344	65.6	14.1	15.1
Fairly dissatisfied	1305	65.6	13.4	16.6
Very dissatisfied	837	57.7	9.8	11.2
Missing	41	58.9	24.2	21.1
<i>Neighbourhood</i>				
<i>Friendliness</i>				
Friendly	14345	70.0	19.1	19.5
Neither	2336	70.4	16.9	18.3
Unfriendly	504	61.6	15.5	15.7
Cannot say	441	59.9	13.5	13.9
Missing	608	72.1	16.7	23.9
<i>Neighbourhood</i>				
<i>Safety (child)</i>				
Yes	10939	74.6	21.3	21.9
No	7027	60.3	13.5	14.2
Missing	268	71.3	16.0	20.4

Data from the Millennium Cohort Study wave 1: Proportions accounted for the survey design

^a Sample size after exclusion criteria and deletion of missing values on breastfeeding outcomes

^b Included babies exclusively, predominantly or partially breastfed

Table 2. Descriptive statistics: Breastfeeding outcomes by household and maternal characteristics.

<i>Familial & Individual Factors</i>	<i>18,234 (N)^a</i>	<i>Breastfeeding Initiation</i>	<i>Exclusive Breastfeeding</i>	<i>Any Breastfeeding^b</i>
	<i>n</i>	<i>Yes (%)</i>	<i>≥ 3 months (%)</i>	<i>≥ 6 months (%)</i>
<i>Household Characteristics</i>				
<i>Household Structure</i>				
Two parents	15092	73.1	20.3	20.9
Single parent	3142	49.0	8.1	9.3
<i>Household Income</i>				
Highest quintile	2907	87.0	29.9	27.3
2nd highest	3168	79.2	22.7	23.3
Middle quintile	3444	71.3	17.0	20.0
2nd lowest	4089	59.8	13.4	14.4
Lowest quintile	4565	50.6	9.3	10.8
Missing	61	66.2	28.4	28.1
<i>Household Social Class</i>				
Managerial & Professional	6878	83.4	27.4	27.6
Intermediate	2370	68.8	15.3	16.7
Small & Self-employed	1071	69.4	16.7	20.5
Low Sup. & Technical	1547	62.4	11.0	10.8
Semi-routine & Routine	4887	50.6	8.7	8.9
Never Worked	1287	50.1	9.4	11.8
Not Classifiable	194	66.8	14.7	21.6
<i>Residential Mobility</i>				
>5 years and over	4927	69.6	19.2	21.0
>1 up to 5 years	9709	71.6	19.7	19.9
up to 1 year	3558	64.0	14.2	14.8
Missing	40	56.5	17.8	23.0
<i>Maternal Characteristics</i>				
<i>Age group</i>				
12 to 19	1581	45.3	4.9	4.8
20 to 29	8557	65.1	13.3	14.0
30 to 39	7710	77.5	25.3	25.9
40 plus	383	82.9	31.2	36.9

Missing	3	43.6	0.0	0.0
---------	---	------	-----	-----

Table 2. Continued

<i>Familial & Individual Factors</i>	<i>18,234 (N)^a</i>	<i>Breastfeeding Initiation</i>	<i>Exclusive Breastfeeding</i>	<i>Any Breastfeeding^b</i>
	<i>n</i>	<i>Yes (%)</i>	<i>≥ 3 months (%)</i>	<i>≥ 6 months (%)</i>
<i>General Health</i>				
Excellent	5474	74.3	24.3	22.8
Good	9563	68.4	17.4	18.9
Fair	2669	65.0	11.4	13.5
Poor	522	62.4	8.1	13.7
Missing	6	71.8	15.2	15.2
<i>Longstanding Illness</i>				
No	14389	70.0	19.4	19.7
Yes	3838	68.1	15.3	17.5
Missing	7	73.9	35.7	35.7
<i>Education</i>				
NVQ levels 4 & 5	5278	87.9	30.5	31.9
NVQ level 3	2576	72.6	17.9	18.4
NVQ level 2	5279	62.7	12.9	12.7
NVQ level 1	1542	51.2	6.9	7.1
None	2974	46.6	8.3	9.8
Overseas qualification	554	75.5	23.0	25.8
Missing	31	58.5	29.2	24.8
<i>Ethnicity</i>				
White	15284	67.5	18.0	17.8
Mixed	188	86.9	29.5	30.3
Indian	470	85.4	22.6	27.6
Pakistani & Bangladeshi	1252	78.2	17.8	21.7
Black/Black British	665	92.9	22.8	36.3
Other ethnic group	345	92.9	27.5	45.7
Missing	30	71.7	30.5	29.7

Data from the Millennium Cohort Study wave 1: Proportions accounted for the survey design
^a Sample size after exclusion criteria and deletion of missing values on breastfeeding outcomes

^bIncluded babies exclusively, predominantly or partially breastfed

Table 3. Results of multivariable logistic regression of neighbourhood characteristics on breastfeeding initiation adjusted for familial- and individual-level factors.

	Model 1	Model 2	Model 3	Full Adjustment
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
<i>IMD</i>				
Highest quintile	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)
2nd highest	0.74 (0.61-0.89)**	0.76 (0.63-0.92)**	0.89 (0.75-1.06)	0.87 (0.73-1.04)
Middle quintile	0.52 (0.43-0.63)***	0.56 (0.46-0.67)***	0.84 (0.72-0.99)*	0.81 (0.68-0.95)*
2nd lowest	0.34 (0.28-0.41)***	0.38 (0.31-0.45)***	0.72 (0.61-0.84)***	0.66 (0.56-0.77)***
Lowest quintile	0.25 (0.20-0.32)***	0.30 (0.24-0.37)***	0.76 (0.62-0.93)**	0.60 (0.50-0.72)***
<i>Neighbourhood Satisfaction</i>				
Very satisfied		1 (Ref)	1 (Ref)	1 (Ref)
Fairly satisfied		0.94 (0.85-1.03)	1.00 (0.90-1.11)	1.01 (0.91-1.12)
Neither		1.06 (0.91-1.25)	1.11 (0.94-1.32)	1.14 (0.96-1.35)
Fairly dissatisfied		1.16 (0.97-1.37)	1.20 (1.01-1.43)*	1.27 (1.06-1.51)*
Very dissatisfied		0.99 (0.83-1.19)	1.18 (0.98-1.42)	1.32 (1.09-1.61)**
<i>Neighbourhood Friendliness</i>				
Friendly		1 (Ref)	1 (Ref)	1 (Ref)
Neither		1.14 (1.01-1.29)*	1.22 (1.08-1.39)**	1.20 (1.06-1.36)**
Unfriendly		0.85 (0.68-1.06)	1.04 (0.83-1.31)	1.01 (0.80-1.28)
Cannot say		0.77 (0.59-1.01)	0.93 (0.71-1.23)	0.88 (0.66-1.18)
<i>Neighbourhood Safety (child)</i>				
Yes		1 (Ref)	1 (Ref)	1 (Ref)
No		0.65 (0.59-0.72)***	0.75 (0.68-0.82)***	0.78 (0.71-0.85)***
<i>Maternal Age</i>				
			1.03 (1.02-1.03)***	1.02 (1.01-1.03)***
<i>Household Income</i>				
Highest quintile			1 (Ref)	1 (Ref)
2nd highest			0.79 (0.66-0.95)*	0.80 (0.66-0.97)*
Middle quintile			0.78 (0.65-0.94)*	0.78 (0.65-0.95)*
2nd lowest			0.66 (0.54-0.80)***	0.62 (0.51-0.77)***
Lowest quintile			0.59 (0.48-0.73)***	0.59 (0.48-0.74)***
<i>Household Social Class</i>				
Manage & Professional			1 (Ref)	1 (Ref)
Intermediate			0.74 (0.64-0.86)***	0.73 (0.63-0.85)***
Small & Self-employed			0.81 (0.68-0.97)*	0.77 (0.65-0.92)**
Low Sup. & Technical			0.71 (0.59-0.84)***	0.72 (0.61-0.86)***
Semi-routine & Routine			0.55 (0.48-0.64)***	0.56 (0.49-0.65)***
Never Worked			0.67 (0.51-0.87)**	0.55 (0.42-0.72)***

Not Classifiable

0.63 (0.42-0.94)*

0.59 (0.39-0.89)*

Table 3. Continued

	Model	Model	Model 3	Full Adjustment
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
<i>Maternal Education</i>				
NVQ levels 4 & 5			1 (Ref)	1 (Ref)
NVQ level 3			0.55 (0.48-0.63)***	0.57 (0.49-0.65)***
NVQ level 2			0.38 (0.33-0.43)***	0.40 (0.35-0.47)***
NVQ level 1			0.30 (0.25-0.36)***	0.33 (0.27-0.40)***
None			0.26 (0.22-0.31)***	0.25 (0.21-0.30)***
Overseas qualification			0.79 (0.58-1.07)	0.58 (0.42-0.79)**
<i>Maternal Ethnicity</i>				
White				1 (Ref)
Mixed				5.33 (3.36-8.46)***
Indian				2.91 (1.83-4.62)***
Pakistani & Bangladeshi				4.31 (3.38-5.49)***
Black/Black British				11.0 (7.49-16.2)***
Other ethnic group				6.31 (4.02-9.91)***
<i>Household Structure</i>				
Two parents				1 (Ref)
Single parent				0.78 (0.68-0.88)***
<i>Maternal General Health</i>				
Excellent				1 (Ref)
Good				0.95 (0.87-1.05)
Fair				1.02 (0.88-1.17)
Poor				0.98 (0.74-1.28)
<i>Maternal Longstanding</i>				
<i>Illness</i>				
No				1 (Ref)
Yes				1.01 (0.90-1.12)
<i>Residential Mobility</i>				
>5 years and over				1 (Ref)
>1 up to 5 years				1.11 (1.00-1.23)*
up to 1 year				1.24 (1.10-1.39)***
* p-value < 0.05				
** p-value < 0.01				
*** p-value < 0.001				

Table 4. Results from multivariable logistic regression of neighbourhood characteristics on exclusive breastfeeding ≥ 3 months adjusted for familial- and individual-level factors.

	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Full Adjustment OR (95% CI)
<i>IMD</i>				
Highest quintile	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)
2nd highest	0.91 (0.77-1.08)	0.95 (0.80-1.12)	1.07 (0.91-1.25)	1.08 (0.92-1.25)
Middle quintile	0.69 (0.56-0.84)**	0.74 (0.60-0.91)**	1.02 (0.84-1.23)	1.01 (0.83-1.22)
2nd lowest	0.47 (0.38-0.57)**	0.54 (0.44-0.67)***	0.90 (0.74-1.08)	0.88 (0.73-1.07)
Lowest quintile	0.35 (0.28-0.45)***	0.44 (0.34-0.58)***	0.96 (0.75-1.23)	0.86 (0.67-1.11)
<i>Neighbourhood Satisfaction</i>				
Very satisfied		1 (Ref)	1 (Ref)	1 (Ref)
Fairly satisfied		0.88 (0.77-1.00)*	0.94 (0.83-1.07)	0.97 (0.85-1.10)
Neither		0.83 (0.66-1.05)	0.89 (0.69-1.14)	0.95 (0.74-1.21)
Fairly dissatisfied		0.80 (0.63-1.00)	0.82 (0.65-1.04)	0.88 (0.70-1.11)
Very dissatisfied		0.66 (0.49-0.90)**	0.78 (0.57-1.07)	0.88 (0.64-1.21)
<i>Neighbourhood Friendliness</i>				
Friendly		1 (Ref)	1 (Ref)	1 (Ref)
Neither		0.97 (0.82-1.14)	1.00 (0.85-1.18)	1.03 (0.87-1.22)
Unfriendly		1.01 (0.73-1.39)	1.22 (0.87-1.73)	1.28 (0.90-1.82)
Cannot say		0.82 (0.56-1.19)	0.95 (0.64-1.41)	0.98 (0.66-1.46)
<i>Neighbourhood Safety (child)</i>				
Yes		1 (Ref)	1 (Ref)	1 (Ref)
No		0.74 (0.65-0.84)***	0.83 (0.74-0.93)**	0.84 (0.75-0.95)**
<i>Maternal Age</i>			1.06 (1.05-1.07)***	1.06 (1.05-1.07)***
<i>Household Income</i>				
Highest quintile			1 (Ref)	1 (Ref)
2nd highest			0.91 (0.78-1.06)	0.93 (0.79-1.09)
Middle quintile			0.92 (0.76-1.10)	0.96 (0.80-1.16)
2nd lowest			0.99 (0.82-1.21)	1.05 (0.85-1.29)
Lowest quintile			0.91 (0.72-1.14)	1.08 (0.85-1.37)
<i>Household Social Class</i>				
Manage & Professional			1 (Ref)	1 (Ref)
Intermediate			0.75 (0.63-0.90)**	0.77 (0.64-0.92)**
Small & Self-employed			0.78 (0.62-0.97)*	0.77 (0.61-0.96)*
Low Sup. & Technical			0.60 (0.49-0.74)***	0.64 (0.52-0.79)***
Semi-routine & Routine			0.58 (0.47-0.71)***	0.62 (0.51-0.76)***

Never Worked	0.72 (0.49-1.05)	0.70 (0.48-1.03)
Not Classifiable	0.53 (0.29-0.95)*	0.54 (0.30-0.97)*

Table 4. Continued

	Model OR (95% CI)	Model OR (95% CI)	Model 3 OR (95% CI)	Full Adjustment OR (95% CI)
<i>Maternal Education</i>				
NVQ levels 4 & 5			1 (Ref)	1 (Ref)
NVQ level 3			0.70 (0.60-0.82)***	0.73 (0.62-0.85)***
NVQ level 2			0.50 (0.44-0.58)***	0.52 (0.45-0.60)***
NVQ level 1			0.31 (0.24-0.40)***	0.33 (0.25-0.42)***
None			0.38 (0.30-0.49)***	0.38 (0.30-0.48)***
Overseas qualification			1.15 (0.82-1.61)	0.99 (0.70-1.40)
<i>Maternal Ethnicity</i>				
White				1 (Ref)
Mixed				2.46 (1.68-3.60)***
Indian				1.43 (1.03-1.98)*
Pakistani & Bangladeshi				1.96 (1.40-2.75)***
Black/Black British				1.62 (1.22-2.16)**
Other ethnic group				1.48 (1.01-2.17)*
<i>Household Structure</i>				
Two parents				1 (Ref)
Single parent				0.70 (0.57-0.85)**
<i>Maternal General Health</i>				
Excellent				1 (Ref)
Good				0.78 (0.70-0.87)***
Fair				0.55 (0.44-0.68)***
Poor				0.38 (0.25-0.57)***
<i>Maternal Longstanding</i>				
<i>Illness</i>				
No				1 (Ref)
Yes				0.95 (0.83-1.08)
<i>Residential Mobility</i>				
>5 years and over				1 (Ref)
>1 up to 5 years				1.09 (0.97-1.23)
up to 1 year				1.06 (0.88-1.28)

* p-value < 0.05
 ** p-value < 0.01
 *** p-value < 0.001

Table 5. Results from multivariable logistic regression of neighbourhood characteristics on any breastfeeding \geq 6 months adjusted for familial- and individual-level factors.

	Model 1	Model 2	Model 3	Full Adjustment
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
<i>IMD</i>				
Highest quintile	1 (Ref)	1 (Ref)	1 (Ref)	1 (Ref)
2nd highest	0.90 (0.75-1.08)	0.93 (0.78-1.11)	1.02 (0.86-1.22)	1.01 (0.85-1.20)
Middle quintile	0.63 (0.52-0.76)***	0.67 (0.55-0.82)***	0.89 (0.73-1.08)	0.85 (0.70-1.04)
2nd lowest	0.49 (0.39-0.61)***	0.55 (0.43-0.69)***	0.85 (0.69-1.06)	0.79 (0.63-0.99)*
Lowest quintile	0.39 (0.29-0.52)***	0.46 (0.34-0.63)***	0.94 (0.70-1.25)	0.77 (0.58-1.02)
<i>Neighbourhood Satisfaction</i>				
Very satisfied		1 (Ref)	1 (Ref)	1 (Ref)
Fairly satisfied		0.90 (0.80-1.02)	0.97 (0.85-1.10)	0.98 (0.86-1.11)
Neither		0.86 (0.68-1.08)	0.93 (0.73-1.17)	0.95 (0.75-1.20)
Fairly dissatisfied		1.00 (0.80-1.24)	1.05 (0.83-1.32)	1.11 (0.88-1.40)
Very dissatisfied		0.75 (0.56-1.00)	0.88 (0.66-1.17)	0.97 (0.73-1.29)
<i>Neighbourhood Friendliness</i>				
Friendly		1 (Ref)	1 (Ref)	1 (Ref)
Neither		1.01 (0.87-1.17)	1.04 (0.90-1.21)	1.06 (0.91-1.23)
Unfriendly		0.96 (0.69-1.33)	1.13 (0.80-1.59)	1.15 (0.81-1.64)
Cannot say		0.81 (0.56-1.16)	0.92 (0.64-1.34)	0.89 (0.60-1.32)
<i>Neighbourhood Safety (child)</i>				
Yes		1 (Ref)	1 (Ref)	1 (Ref)
No		0.72 (0.63-0.82)***	0.81 (0.72-0.92)**	0.82 (0.73-0.93)**
<i>Maternal Age</i>				
			1.07 (1.06-1.08)***	1.07 (1.06-1.08)***
<i>Household Income</i>				
Highest quintile			1 (Ref)	1 (Ref)
2nd highest			1.11 (0.94-1.31)	1.12 (0.95-1.32)
Middle quintile			1.46 (1.25-1.70)***	1.47 (1.26-1.72)***
2nd lowest			1.42 (1.19-1.69)***	1.40 (1.17-1.67)***
Lowest quintile			1.40 (1.14-1.72)**	1.54 (1.25-1.89)***
<i>Household Social Class</i>				

Manage & Professional	1 (Ref)	1 (Ref)
Intermediate	0.76 (0.63-0.91)**	0.77 (0.64-0.92)**
Small & Self-employed	0.93 (0.74-1.17)	0.90 (0.72-1.13)
Low Sup. & Technical	0.53 (0.41-0.67)***	0.55 (0.43-0.70)***
Semi-routine & Routine	0.54 (0.44-0.65)***	0.57 (0.47-0.69)***
Never Worked	0.79 (0.58-1.09)	0.69 (0.51-0.94)*
Not Classifiable	0.89 (0.51-1.54)	0.87 (0.51-1.49)

Table 5. Continued

	Model	Model	Model 3	Full Adjustment
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
<i>Maternal Education</i>				
NVQ levels 4 & 5			1 (Ref)	1 (Ref)
NVQ level 3			0.63 (0.54-0.74)***	0.67 (0.57-0.78)***
NVQ level 2			0.42 (0.36-0.49)***	0.45 (0.39-0.52)***
NVQ level 1			0.27 (0.21-0.35)***	0.30 (0.23-0.39)***
None			0.33 (0.26-0.43)***	0.33 (0.26-0.42)***
Overseas qualification			0.95 (0.68-1.32)	0.73 (0.52-1.02)
<i>Maternal Ethnicity</i>				
White				1 (Ref)
Mixed				2.64 (1.77-3.95)***
Indian				1.89 (1.32-2.71)**
Pakistani & Bangladeshi				2.51 (1.89-3.33)***
Black/Black British				3.18 (2.48-4.08)***
Other ethnic group				3.96 (2.78-5.64)***
<i>Household Structure</i>				
Two parents				1 (Ref)
Single parent				0.69 (0.56-0.84)***
<i>Maternal General Health</i>				
Excellent				1 (Ref)
Good				0.90 (0.81-1.00)
Fair				0.66 (0.55-0.79)***
Poor				0.66 (0.47-0.92)*
<i>Maternal Longstanding</i>				
<i>Illness</i>				
No				1 (Ref)
Yes				1.01 (0.88-1.17)

Residential Mobility

>5 years and over	1 (Ref)
>1 up to 5 years	1.03 (0.90-1.17)
up to 1 year	1.03 (0.88-1.22)

* p-value < 0.05
** p-value < 0.01
*** p-value < 0.001
