

# WINDOW consortium: A Path Towards Increased Therapy Efficacy Against Glioblastoma

Kulsoom U. Abdul<sup>1,6,7</sup>, Megan Houweling<sup>1,6,7</sup>, Fredrik Svensson<sup>2,6</sup>, Ravi S. Narayan<sup>3</sup>, Fleur M.G. Cornelissen<sup>1</sup>, Asli Küçükösmanoglu<sup>1,6</sup>, Emmanouil Metzakopian<sup>4,6</sup>, Colin Watts<sup>5,6</sup>, David Bailey<sup>2,6</sup>, Tom Wurdinger<sup>1,6</sup>, Bart A. Westerman<sup>1,6</sup>

1 Department of Neurosurgery, Neuro-oncology Research Group, Amsterdam University Medical Center, Cancer Center Amsterdam, De Boelelaan 1117, 1081 HZ Amsterdam

2 IOTA Pharmaceuticals Ltd, St Johns Innovation Centre, Cowley Road, Cambridge, CB4 0WS, UK

3 Department of Radiation Oncology, Amsterdam University Medical Center, Cancer Center Amsterdam, De Boelelaan 1117, 1081 HZ Amsterdam

4 UK Dementia Research Institute, Cambridge, CB2 0AH, UK

5 Institute of Cancer and Genomic Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT

6 The WINDOW consortium, [www.window-consortium.org](http://www.window-consortium.org)

7 These authors contributed equally

## ABSTRACT


Glioblastoma is the most common and malignant form of brain cancer, for which the standard treatment is maximal surgical resection, radiotherapy and chemotherapy. Despite these interventions, mean overall survival remains less than 15 months, during which extensive tumor infiltration throughout the brain occurs. The resulting metastasized cells in the brain are characterized by chemotherapy resistance and extensive intratumoral heterogeneity. An orthogonal approach attacking both intracellular resistance mechanisms as well as intercellular heterogeneity is necessary to halt tumor progression. For this reason, we established the WINDOW Consortium (Window for Improvement for Newly Diagnosed patients by Overcoming disease Worsening), in which we are establishing a strategy for rational selection and development of effective therapies against glioblastoma. Here, we overview the many challenges posed in treating glioblastoma, including selection of drug combinations that prevent therapy resistance, the need for drugs that have improved blood brain barrier penetration and strategies to counter heterogeneous cell populations within patients. Together, this forms the backbone of our strategy to attack glioblastoma.

## BACKGROUND AND CHALLENGES

### Classification of glioblastoma and inter-patient heterogeneity

Gliomas are the most common malignancies of the central nervous system (CNS). Most glioblastomas (GBMs) arise *de novo* without any sign of a less malignant precursor (Ohgaki and Kleihues, 2013). Primary GBMs typically occur at an average age of 62 years, rapidly progress and have an extremely poor prognosis. However, 10% of GBMs progress from low-grade diffuse astrocytomas or anaplastic astrocytomas (**Figure 1A**). These “secondary” GBMs occur at an average age of 45 years and offer a slightly better prognosis than primary lesions. According to the WHO 2016 classification, gliomas are defined based on mutations in the IDH1 gene with/without chromosomal 19q loss (summarized in **Figure 1B**, taken from Verhaak 2016). Astrocytic gliomas are classified based on histologic criteria from lower grade lesions (grades II–III) to high-grade (grade IV) malignancies (WHO 2016; previous classifications given in Louis et al, 2007; Zhu and Parada, 2002; Weller et al, 2005). Originally, the Cancer Genome Atlas Network (TCGA, Verhaak et al, 2010) defined

molecular signatures within GBM (grade IV), which were distinguishable based on expression of lineage markers. These subgroups are termed “classical”, “mesenchymal”, “proneural” and “neural”, although existence of the latter group is debated (Sideway 2017). The proneural subtype correlates with better prognosis whereas the mesenchymal and classical subtypes correspond to poor prognosis (Lin et al, 2014). These quantitative relationships, based on WHO criteria, molecular features and histology are summarized in **Figure 1**.






**Figure 1.** (A) Circos plot showing the histological classification system as used in the clinic until recently. (B) Sankey plot showing the quantitative relation of different adult gliomas (based on 1,122 patients in the TCGA database [www.cbioportal.org](http://www.cbioportal.org), see also Ceccarelli et al 2016 and Weller et al 2017), grouped by the WHO 2016 classification (IDH mutated; IDH/ATRX comutated; 1p19q codeletion); molecular profile (Mesenchymal, Classic like, LGM6-GBM\_PA [LGGs in the third methylation cluster of IDH-wild type tumours as pilocytic astrocytoma-like], GCIMP; and histological class: astrocytoma (AS), oligodendroglioma (OD) and glioblastoma (GBM). (C) Current treatment of GBM after maximal safe surgical resection. After surgery, RT is continued for 6 weeks combined with TMZ treatment. Subsequently, adjuvant TMZ five days per week is indicated for at least six months. po, per os (oral administration).

**Current clinical trials are not focused on GBM driver mutations**

The standard treatment of GBM is based on the international EORTC-study (Stupp et al. 2005), consisting of maximal safe surgical resection, followed by six weeks of radiotherapy (RT) and concomitant Temozolomide (TMZ) chemotherapy with subsequent adjuvant TMZ chemotherapy for six months, **Figure 1C**. Compared to RT alone, with a median survival time of 12.1 months, the combination of RT plus TMZ increases median overall survival time (OS) by 2.5 months to 14.6 months (Weller et al, 2017; Stupp et al, 2009; 2005). A group of patients displaying silencing of the MGMT-gene promoter via DNA methylation, have repressed DNA damage repair and is therefore more sensitive to the alkylating effect of TMZ (Hegi et al, 2005). In clinical practice, TMZ is administered regardless of the patient's promoter methylation status.

Many efforts are underway to improve the outcome of the current standard therapy, attacking driver as well as non-driver targeted therapies (graphically summarized in **Figure 2A**). Currently, there are more than 225 ongoing trials in GBM, of which over 60% include systemic therapy, comprising chemotherapy/DNA damaging compounds (23%), immunotherapy (13%), VEGF targeted therapy (8%) and PI3K targeted therapy (3%), shown in Sankey plot in **Figure 2B**.



**Figure 2.** Driver versus non-driver focussed therapies against GBM in clinical trials (ClinicalTrials.gov). (A) Circos plots showing non-driver targets and driver targets of currently applied clinical trials against GBM. Cellular location of targets is shown as extracellular, membrane bound or intracellular. Non-driver genes are classified into immune targets, receptors and receptor ligands; and cytotoxic/DNA damaging targets. The driver mutation circos plot is divided in targetable mutations, absence of druggable mutations and non-druggable driver mutations. (B) Sankey plot showing the quantitative relation of the number of trials towards each target in ongoing clinical trials. Non-driver targets (upper portion) as well as driver targets using FDA approved mutation specific small molecules (lower portion) are shown. Note that almost no approaches to driver targets are currently in clinical trials. Abbreviations: RTK, receptor tyrosine kinase; SMO, smoothened Shh receptor; Ptc, Patched Shh receptor; TGF $\beta$  Transforming growth factor beta; HGF, Hepatocyte growth factor; CRM1, chromosome region maintenance 1; COX2, Cyclooxygenase 2; PD1, Programmed cell death protein 1; EGFR, Epidermal growth factor receptor, ATRX, Alpha Thalassemia/Mental Retardation Syndrome X-Linked gene; RB1, retinoblastoma 1 protein; TP53, tumour protein p53; p15/CDK2NB, cyclin-dependent kinase inhibitor 2B; IDH1 Isocitrate dehydrogenase homolog 1; NF1, Neurofibromatosis type 1; PIK3R1 Phosphatidylinositol 3-kinase regulatory subunit alpha; PIK3CA, Phosphatidylinositol-4,5-Bisphosphate 3-Kinase Catalytic Subunit Alpha; PDGFRA, Platelet-derived growth factor receptor A; MDM2, Murine double minute 2; PTEN, Phosphatase and tensin homolog; p16/CDK2NA, cyclin-dependent kinase inhibitor 2A; CDK4, cyclin dependent kinase 4.

Although 70% of GBMs contain tumor driver mutations that are targetable by therapy (**Figure 2**, Cacarelli et al, 2016), in current clinical trials there is no focus on these driver mutations, except for EGFR- targeted therapy (11% of clinical trials, **Figure 2B**). Trials aimed at driver-mutations have been largely unsuccessful (for instance NCT00187486, NCT0062243 and NCT00671970 aimed at EGFR inhibition), possibly because of poor penetrance of drugs into the CNS through the blood brain barrier [BBB; (Oberoi et al, 2013; Tang et al, 2012; Porta et al, 2011)].

Here, we address the challenges associated with the efficient treatment of GBM through combination therapy. We see these as: (1) identification of drugs that have desired BBB penetration, (2) addressing whether combination-therapies are of benefit in targeting intratumoral heterogeneity and (3) overcoming

drug resistance. Based on this landscape, we outline the strategy adopted by the WINDOW consortium for the selection and further development of effective therapies against GBM.

**Challenge 1: The BBB and perivascular/perineural microinvasion** – The BBB is the natural barrier that prevents toxins from reaching the brain. It is a major obstacle to brain tumor therapy, preventing the delivery of most chemotherapeutic and targeted agents to the tumor location (e.g. Sminia and Westerman 2016; van Tellingen et al, 2015). Since most drugs have been developed for applications outside the brain and therefore selected for low BBB penetration to avoid neurotoxicity, most FDA approved drugs have poor target engagement in the brain.

Gliomas show perivascular and perineural microinvasion (Cuddapah et al, 2014; Gritsenko et al, 2012; Montana et al, 2011; Farin, et al, 2006). Glioma tumor cells can form multicellular networks connected through branched filamentous protrusions connecting cells, or epithelial-like linear adherent junctions between directly adjacent cells (Friedl et al, personal communication). Therefore, individual tumor cells are localized distantly from the bulk of the tumor (Sherriff et al, 2013) and might be more difficult to reach both therapeutically as well as surgically.

For many years, different strategies have been investigated to facilitate BBB penetration by chemotherapeutics, reviewed in Upadhyay, 2014 and Lu et al, 2014. These include the use of 1) non-invasive techniques, such as radiotherapy (Trnovec et al, 2016), ultrasound and microbubbles (Lamanauskas et al, 2013; Escoffre et al, 2013; Rachlin et al, 2013; Chu et al, 2016), and biological approaches via cell penetrating peptides and viral vectors (Sidaway, 2017); 2) invasive techniques, such as convection enhanced delivery (reviewed by Vogelbaum and Aghi, 2015), and 3) alternative routes such as intranasal application, bypassing the cardiovascular system (reviewed by Peterson et al, 2014).

The high prevalence of brain metastasis from the lung, breast and melanoma, makes it attractive to develop compounds that target driver mutations specific for these tumor types, leading to the development of potent kinase inhibitors that do cross the BBB. These include compounds targeting driver mutations such as Osimertinib (targeting EGFR T790M) and Lorlatinib (targeting ALK mutations), as well as the radio-sensitizing drug AZD0156 (ATM) and the PI3K/AKT/mTOR pathway inhibitor GDC-0084. GBM is rarely the main therapeutic focus, due to its relatively small market size. However, because drug discovery is a long-

term endeavour and an extremely expensive one (estimated that a single drug can take over 10 years and cost more than £500M to develop) (source ABPI, <http://www.abpi.org.uk/>), drug repurposing (i.e. the identification of new therapeutic uses for existing drugs), is the only route currently available as a solution to this challenge. To date, there has been no systematic evaluation of the already available approved drugs against GBM.


## **Challenge 2: Drug resistance and intratumoral molecular heterogeneity**

GBM is characterized by intratumoral heterogeneity in which subpopulations of cells have distinct features, largely consisting of (1) gene copy number variations as shown for PTEN, TP53 and MDM4 (Sottoriva et al, 2013; Johnson et al, 2014; Meyer et al 2015), (2) clonal ploidy differences (Stieber et al, 2014; Johnson et al. 2014; Meyer et al, 2015), (3) extrachromosomal DNA elements (deCarvalho et al, 2018; Turner et al, 2017), (4) signalling heterogeneity as a result of mosaic receptor tyrosine kinase activity (Snuderl et al, 2011; Little et al. 2012; Szerlip et al. 2012) including structural variants of EGF receptor (Francis et al, 2014; Meyer et al, 2015), and (5) lineage heterogeneity, i.e. GBM cells express markers of lineage derivatives of stem/progenitor cells such as proneural, neural as well as astroglial genes (Verhaak et al, 2010; Phillips et al, 2006; Al-Mayhany et al, 2011; Sottoriva et al. 2013; Patel et al, 2014; Kenney-Herbert et al, 2015; Piccirillo et al, 2015a; Piccirillo et al, 2015b; Wang et al, 2016, Meyer et al, 2015). The different forms of cellular heterogeneity are summarized in **Figure 3A**.

The sensitivity of individual tumors towards drugs can vary due to intrinsic resistance which is pre-existing or acquired as a result of drug therapy (Sequist et al, 2011; Garrett et al, 2011; Prahallad et al, 2012; Wilson et al, 2012). Drug resistance, intrinsic or acquired, is affected by intratumoral heterogeneity due to genetic or phenotypic heterogeneity (the mechanisms are listed in **Figure 3B**). Preclinically, intratumoral subpopulations display a differential response to therapeutics (Meyer et al, 2015; Hägerstrand et al, 2011; Heo et al, 2014; Saito et al, 2014; Lee et al, 2017; Lan et al, 2017). Moreover, a re-transplanted tumor seems to retain its intrinsic resistance, indicating that there is a population with a long term tumor initiation or drug resistance potential (Lan et al, 2017). These studies implicate intrinsic cell populations in each GBM patient exhibiting pre-existing resistance to therapy (Meyer et al, 2015). In the clinic, radiation resistance is observed frequently in relapsed GBMs (Hochberg and Pruitt, 1980; Kelley et al, 2016; Fidoamore et al, 2016). Several



alternative mechanisms have been found to underlie therapy resistance. For instance, EGFR inhibitor resistance may occur as a consequence of loss of extrachromosomal mutant EGFR DNA, promyelocytic leukemia (PML) gene expression, PTEN phosphorylation status, PDGFR upregulation, ERBB4 activation, AMPK levels or IL-6 upregulation (Nathanson et al, 2013; O'Rourke et al, 2017; Iwanami et al, 2013; Fenton et al, 2012; Akhavan et al, 2013; Donoghue et al, 2018; Guo et al, 2009; Zanca et al, 2017).



**Figure 3.** Resistance and heterogeneity mechanisms of GBM. (A) Circos plot showing different molecular mechanisms underlying the generation of cellular heterogeneity; (B) Sankey plot quantitatively summarising the causes of heterogeneity and drug resistance involving driver targets in GBM. Targets that are frequently mutated are shown in color. PML, promyelocytic leukemia; AMPK, 5' adenosine monophosphate-activated protein kinase; MAPK, Mitogen Activated Protein Kinase.

### Challenge 3 Combination therapies against heterogeneous populations

Since intratumoral heterogeneity and drug resistance can occur simultaneously, a promising approach might be to combine targeted drugs that hit several survival mechanisms at once for different cell populations. This could provide a synergistic (i.e. more than additive) effect and prevent therapy-resistance. However, identification of these synergistic combinations has thus far only been possible in an empirical setting (by experimentally testing all combinations). A major challenge is therefore to match GBM vulnerabilities to effective drug combinations.

Tumors are dependent on a limited number of molecular mechanisms for their survival and proliferation (Hanahan and Weinberg, 2011; Wang et al, 2015). Combination therapy enables simultaneous targeting of these crucial mechanisms. In addition, for personalized cancer treatment, a focus on drug/irradiation or drug/drug combinations is particularly appealing since enhanced efficacy can be assessed for each radiation-dose or drug-concentration window.

Strategies for selecting synergistic and effective drug combinations (Dancey and Chen, 2006; Day and Siu, 2016) can be based on several precepts, including: (1) maximal target inhibition, achieved by synergistic drug combinations that hit the same target, especially important in the case of oncogene addiction. Drug resistance might also be avoided by this approach as well as other novel approaches (Bar-zeev et al, 2017; Li et al, 2016; Zhitomirsky et al, 2016); (2) maximal pathway inhibition. In a similar manner, maximal pathway inhibition can be used to suppress an entire pathway by inhibiting it at multiple levels. An example of this strategy is the FDA approved combination of the BRAF inhibitor vemurafenib and the MEK inhibitor cobimetinib in BRAFV600 mutated melanoma which results in maximal pathway inhibition in cases where the tumor is dependent on the BRAFV600 mutation (Larkin et al, 2014). Also, within GBM radiotherapy and PARP-inhibition have shown synergy (Lesueur et al, 2018); (3) feedback inhibition between and within pathways. In many cases this affects upstream pathway activation through downstream targets of the initial inhibited pathway (Sun and Hobor et al, 2014; Sun and Wang et al, 2014; Rozengurt et al, 2014); and finally (4) synthetic lethality (feed forward) inhibition, in many cases affecting inhibition of two parallel pathways downstream of activated oncogenes (Croesmann et al, 2018). For feedback inhibition and synthetic lethality approaches, information on underlying therapy-resistance is scarce, and direct targets are difficult to identify, rendering the rational design of pathway combinations difficult.

## **AIMS OF THE WINDOW CONSORTIUM**

### **General aims of the consortium**

Within the WINDOW consortium, our aim is to provide a solution to these challenges by examining the links between intratumor molecular heterogeneity, prediction of *in vitro* therapeutic drug combination and testing, as well as emerging clinical data, areas which must be addressed for successful development of effective therapeutics. Our overall objective is to create a validated patient-centred platform based upon

integration of patient-derived cell systems, detailed genetic analysis of tumor cell population, together with their treatment with the most effective combination therapies. At a more immediate level, we will develop a collaborative pipeline to enable patient-stratified treatments with the most effective combinations of FDA approved drugs, thereby offering a Window for Improvement for Newly Diagnosed patients thereby Overcoming Disease Worsening (WINDOW).

### **WINDOW specific aim 1: Repurposing of clinically approved drugs**

To be able to repurpose FDA approved drugs against GBM, drugs must combine traditional qualities, such as optimal systemic absorption, distribution, metabolism, and excretion (ADME), with enhanced CNS bioavailability. Selection criteria are therefore: (1) molecular mode of action [i.e. target specificity], (2) experimental proof of efficacy, (3) ADME characteristics, (4) documented CNS penetrance and (5) toxicology including the absence of neural side-effects.

Most FDA approved drugs will never reach the brain because they were selected based on inability to cross the blood brain barrier. In addition, most drugs are substrates for efflux pumps and are actively pumped out of the brain through ABC transporters such as P-gp and ABCG2. Recent development of drugs against metastasized from the lung, skin and breast tumors have provided important information how to design drugs that have optimal characteristics to reach the brain tumor. Primary brain tumor have suffered from a lack of clinically relevant drugs targeting brain specific lesions. Nevertheless, we show here how to use data from non-primary brain tumor fields to select drugs that might be lead drugs for application against brain tumors. A first step in this process has been to curate GBM drug information in a form that is readily accessible to all preclinical GBM researchers. This has now been achieved through the establishment of the GBM Drug Bank ([www.gbmdrugbank.com](http://www.gbmdrugbank.com); Svensson et al 2018). This resource includes all FDA approved drugs suitable for repurposing that are active in GBM preclinical models.

### **WINDOW specific aim 2: Overcoming drug resistance through combination therapies**

In the WINDOW project, we have developed a novel strategy to identify effective drug-combinations using a topology-based approach which we call the drug-atlas (Narayan et al, submitted). This atlas is built from drug-response encyclopedias and can be considered as a framework of therapeutic action and therefore can be

interpreted as a drug-vulnerability landscape of cancer. The methodology is based on the finding that most tumors contain multiple independent survival/proliferation mechanisms. This novel rational and generalizable strategy opens the door to unforeseen personalized multi-drug combination approaches.

Some of the drug combinations identified with the drug atlas will show a lack of response in a number of cell lines. For this we will apply CRISPR/CAS9 technologies to elucidate the molecular basis for the lack of therapeutic response (Tzelepis et al, 2016). We will (1) identify genetic vulnerabilities of these clonally derived cell lines and to determine the mechanisms that are responsible for the lack of response therapy. Clonally derived glioblastoma cell lines that show drug resistance are infected with a CRISPR knock out library. Subsequently, next gen sequencing of the CRISPR cassettes will reveal which CRISPR constructs are lost upon exposure to the combination therapy and hence identify the genes responsible for the sensitization to the drug combination. This drop out screen methodology will reveal the sensitivities of the drug resistant cell lines, which will differ from the control cell line. Finally, drugs can be chosen that hit the identified protein or process as directly as possible. In addition, drug resistance mechanisms to monotherapy or combination therapy can be identified using CRISPR screening technologies.

Two major limitations of combination therapy with targeted drugs are the narrow time window wherein drugs can or need to be delivered, and the narrow therapeutic window between enhanced tumor kill and toxicity. Examples of drug combination-induced toxicities are shown in **Table 1**.

**Table 1**  
**Examples of accumulating toxicities**

Challenge	Drug combination	Tumour type	Clinical phase	Outcome	Reference
Target engagement & PK/PD	Sirolimus + erlotinib	GBM	Phase II	No improvement over control	Reardon et al, 2010
	Cetuximab or panitumumab to bevacizumab and chemotherapy	Colon	Phase III	No improvement over control	Hecht et al, 2009; Tol et al, 2009
Toxicity	Bevacizumab, irinotecan, temozolomide	GBM	Phase II	Hematological toxicity	Peters et al, 2015
	Bevacizumab (VEGF inhibitor), temozolomide	GBM	Phase II	Blood pressure, Hematological toxicity	Reyes Botero et al, 2018
	Onartuzumab (c-MET) bevacizumab (VEGF inhibitor)	GBM	Phase II	Peripheral Asthenia edema,	Cloughesy et al, 2017

Bevacizumab (VEGF inhibitor) + sunitinib	Renal cell carcinoma	Phase I	Vascular/hematological toxicities	Feldman et al, 2009*
Temsirolimus (mTOR inhibitor) + sunitinib	Renal cell carcinoma	Phase I	Skin/hematological toxicities	Patel et al, 2009*
Everolimus (MTORC1) combined with exemestane	Breast cancer	Phase III	Stomatitis, intestinal, hematological toxicities	Baselga et al, 2012
MEK inhibitor combined with AKT inhibitor	Various	Phase II	Intestinal, skin	Tolcher et al, 2015
Ipilimumab combined with nivolumab	Melanoma	Phase III	Intestinal, skin	Larkin et al, 2015
BRAF inhibition combined with MEK inhibition	Melanoma	Phase II	Fever	Flaherty et al, 2012

\*Taken from Day and Siu Genome Medicine (2016) 8:115

Additive toxicities can be avoided by using drug combinations that have different toxicity patterns. For example, in the combination with irradiation, the crucial issue is the scheduling and administration of targeted and radiosensitizing agents. To predict possible toxicities, we will generate a Toxicity Atlas, to enable selection of drug-combinations with complementary toxicities and prevent accumulated side effects.

## Future Perspectives

By using our WINDOW consortium approach, we aim to overcome issues that currently prevent an effective therapeutic strategy against GBM. These issues concern intratumoral heterogeneity and identification of more effective therapies targeting the heterogeneous tumor populations characteristic of GBM, initially focussing on the repurposing of FDA-approved small molecule drugs. Recognizing the fact that many of the current set of FDA-approved drugs were developed with non-CNS penetrance as an objective, the WINDOW program has the longer-term aim of re-developing approved drugs that show efficacy in preclinical models, through industry-proven lead optimization approaches.

Based on the high prevalence of brain metastases from lung cancer, breast cancer and melanoma, industry research is leading to the development of kinase inhibitors for treating brain metastases of other solid tumors, including driver-targeted compounds such as Osimertinib, AZD3759, NT113, Lorlatinib, GDC-0084, as well as the radio-sensitizing drug AZD0156. An additional focus could be on inhibitors showing

polypharmacology (Knight et al., 2010), in which either a single drug or a combination of drugs can restrict or even gridlock adaption mechanisms of the tumor.

Many of these new approaches could be tested in preclinical models recapitulating the inter- as well as intra-patient heterogeneity to enable further dissection of the molecular basis of resistance development in the context of the host-tumor interaction, as well as direct tumor toxicity. A better mechanistic understanding of the toxicity of drug combinations and/or drug polypharmacology will assist in prioritizing candidate therapies. Moreover, activation of the immune system might further enhance the effects of our strategy. The objective of the WINDOW Consortium is to combine these paradigms, to deliver new and effective therapeutic strategies for GBM.

### **Acknowledgement**

This work was made possible through grant #488097 of the Brain Tumour Charity and support from the Dutch Brain Tumor Society Stophersentumoren, grant “The high grade glioma synergy atlas”.

### **References**

1. ABPI, The Association of the British Pharmaceutical Industry, <http://www.abpi.org.uk/>Chu et al., 2016,
2. Akhavan D, Pourzia AL, Nourian AA, Williams KJ, Nathanson D, Babic I, Villa GR, Tanaka K, Nael A, Yang H, Dang J, Vinters HV, Yong WH, Flagg M, Tamanoi F, Sasayama T, James CD, Kornblum HI, Cloughesy TF, Cavenee WK, Bensinger SJ, Mischel PS. De-repression of PDGFR $\beta$  transcription promotes acquired resistance to EGFR tyrosine kinase inhibitors in glioblastoma patients. *Cancer Discov.* 2013 May;3(5):534-47.
3. Al-Mayhany MT, Grenfell R, Narita M, Piccirillo S, Kenney-Herbert E, Fawcett JW, Collins VP, Ichimura K, Watts C. NG2 expression in glioblastoma identifies an actively proliferating population with an aggressive molecular signature. *Neuro Oncol.* 2011 Aug;13(8):830-45.
4. Bar-Zeev M, Livney YD, Assaraf YG. Targeted nanomedicine for cancer therapeutics: Towards precision medicine overcoming drug resistance. *Drug Resist Updat.* 2017 Mar;31:15-30.

5. Baselga J, Campone M, Piccart M, Burris HA 3rd, Rugo HS, Sahmoud T, Noguchi S, Gnant M, Pritchard KI, Lebrun F, Beck JT, Ito Y, Yardley D, Deleu I, Perez A, Bachelot T, Vittori L, Xu Z, Mukhopadhyay P, Lebwahl D, Hortobagyi GN. Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer. *N Engl J Med*. 2012 Feb 9;366(6):520-9.
6. Baselga J, Cortés J, Kim SB, Im SA, Hegg R, Im YH, Roman L, Pedrini JL, Pienkowski T, Knott A, Clark E, Benyunes MC, Ross G, Swain SM. Pertuzumab plus trastuzumab plus docetaxel for metastatic breast cancer. *N Engl J Med*. 2012 Jan 12;366(2):109-19.
7. Ceccarelli M, Barthel FP, Malta TM, Sabedot TS, Salama SR, Murray BA, Morozova O, Newton Y, Radenbaugh A, Pagnotta SM, Anjum S, Wang J, Manyam G, Zoppoli P, Ling S, Rao AA, Grifford M, Cherniack AD, Zhang H, Poisson L, Carlotti CG Jr, Tirapelli DP, Rao A, Mikkelsen T, Lau CC, Yung WK, Rabadan R, Huse J, Brat DJ, Lehman NL, Barnholtz-Sloan JS, Zheng S, Hess K, Rao G, Meyerson M, Beroukhi R, Cooper L, Akbani R, Wrensch M, Haussler D, Aldape KD, Laird PW, Gutmann DH; TCGA Research Network, Nounmehr H, Iavarone A, Verhaak RG. Molecular Profiling Reveals Biologically Discrete Subsets and Pathways of Progression in Diffuse Glioma. *Cell*. 2016 Jan 28;164(3):550-63.
8. Cloughesy T, Finocchiaro G, Belda-Iniesta C, Recht L, Brandes AA, Pineda E, Mikkelsen T, Chinot OL, Balana C, Macdonald DR, Westphal M, Hopkins K, Weller M, Bais C, Sandmann T, Bruey JM, Koeppen H, Liu B, Verret W, Phan SC, Shames DS. Randomized, Double-Blind, Placebo-Controlled, Multicenter Phase II Study of Onartuzumab Plus Bevacizumab Versus Placebo Plus Bevacizumab in Patients With Recurrent Glioblastoma: Efficacy, Safety, and Hepatocyte Growth Factor and O6-Methylguanine-DNA Methyltransferase Biomarker Analyses. *J Clin Oncol*. 2017 Jan 20;35(3):343-351.
9. Croessmann S, Formisano L, Kinch L, Gonzalez-Ericsson PI, Sudhan DR, Nagy RJ, Mathew A, Bernicker EH, Cristofanilli M, He J, Cutler RE, Lalani AS, Miller VA, Lanman RB, Grishin N, Arteaga CL. Combined blockade of activating ERBB2 mutations and ER results in synthetic lethality of ER+/HER2 mutant breast cancer. *Clin Cancer Res*. 2018 Oct 12
10. Cuddapah, V. A., Robel, S., Watkins, S. & Sontheimer, H. A neurocentric perspective on glioma invasion. *Nature reviews. Neuroscience* 15, 455-465, (2014).

11. Dancey JE, Chen HX. Strategies for optimizing combinations of molecularly targeted anticancer agents. *Nat Rev Drug Discov*. 2006 Aug;5(8):649-59.
12. Day D, Siu LL. Approaches to modernize the combination drug development paradigm. *Genome Med*. 2016 Oct 28;8(1):115.
13. deCarvalho AC, Kim H, Poisson LM, Winn ME, Mueller C, Cherba D, Koeman J, Seth S, Protopopov A, Felicella M, Zheng S, Multani A, Jiang Y, Zhang J, Nam DH, Petricoin EF, Chin L, Mikkelsen T, Verhaak RGW. Discordant inheritance of chromosomal and extrachromosomal DNA elements contributes to dynamic disease evolution in glioblastoma. *Nat Genet*. 2018 May;50(5):708-717.
14. Donoghue JF, Kerr LT, Alexander NW, Greenall SA, Longano AB, Gottardo NG, Wang R, Tabar V, Adams TE, Mischel PS, Johns TG. Activation of ERBB4 in Glioblastoma Can Contribute to Increased Tumorigenicity and Influence Therapeutic Response. *Cancers (Basel)*. 2018 Jul 25;10(8). pii: E243.
15. Escoffre JM, Novell A, Piron J, Zeghimi A, Doinikov A, Bouakaz A. Microbubble attenuation and destruction: are they involved in sonoporation efficiency? *IEEE Trans Ultrason Ferroelectr Freq Control*. 2013 Jan;60(1):46-52.
16. Farin A, Suzuki SO, Weiker M, Goldman JE, Bruce JN, Canoll P. Transplanted glioma cells migrate and proliferate on host brain vasculature: a dynamic analysis. *Glia*. 2006 Jun;53(8):799-808.
17. Feldman DR, Baum MS, Ginsberg MS, Hassoun H, Flombaum CD, Velasco S, et al. Phase I trial of bevacizumab plus escalated doses of sunitinib in patients with metastatic renal cell carcinoma. *J Clin Oncol*. 2009;27:1432–9.
18. Feldman DR, Baum MS, Ginsberg MS, Hassoun H, Flombaum CD, Velasco S, Fischer P, Ronnen E, Ishill N, Patil S, Motzer RJ. Phase I trial of bevacizumab plus escalated doses of sunitinib in patients with metastatic renal cell carcinoma. *J Clin Oncol*. 2009 Mar 20;27(9):1432-9.
19. Fenton TR, Nathanson D, Ponte de Albuquerque C, Kuga D, Iwanami A, Dang J, Yang H, Tanaka K, Oba-Shinjo SM, Uno M, Inda MM, Wykosky J, Bachoo RM, James CD, DePinho RA, Vandenberg SR, Zhou H, Marie SK, Mischel PS, Cavenee WK, Furnari FB. Resistance to EGF receptor inhibitors in glioblastoma mediated by phosphorylation of the PTEN tumor suppressor at tyrosine 240. *Proc Natl Acad Sci U S A*. 2012 Aug 28;109(35):14164-9.



20. Fidoamore A, Cristiano L, Antonosante A, d'Angelo M, Di Giacomo E, Astarita C, Giordano A, Ippoliti R, Benedetti E, Cimini A. Glioblastoma Stem Cells Microenvironment: The Paracrine Roles of the Niche in Drug and Radioresistance. *Stem Cells Int.* 2016, 6809105 (2016).
21. Flaherty KT, Infante JR, Daud A, Gonzalez R, Kefford RF, Sosman J, Hamid O, Schuchter L, Cebon J, Ibrahim N, Kudchadkar R, Burris HA 3rd, Falchook G, Algazi A, Lewis K, Long GV, Puzanov I, Lebowitz P, Singh A, Little S, Sun P, Allred A, Ouellet D, Kim KB, Patel K, Weber J. Combined BRAF and MEK inhibition in melanoma with BRAF V600 mutations. *N Engl J Med.* 2012 Nov 1;367(18):1694-703.
22. Flaherty KT, Robert C, Hersey P, Nathan P, Garbe C, Milhem M, Demidov LV, Hassel JC, Rutkowski P, Mohr P, Dummer R, Trefzer U, Larkin JM, Utikal J, Dreno B, Nyakas M, Middleton MR, Becker JC, Casey M, Sherman LJ, Wu FS, Ouellet D, Martin AM, Patel K, Schadendorf D. Improved survival with MEK inhibition in BRAF-mutated melanoma. *N Engl J Med.* 2012 Jul 12;367(2):107-14.
23. Francis JM, Zhang CZ, Maire CL, Jung J, Manzo VE, Adalsteinsson VA, Homer H, Haidar S, Blumenstiel B, Pedamallu CS, Ligon AH, Love JC, Meyerson M, Ligon KL. EGFR variant heterogeneity in glioblastoma resolved through singlenucleus sequencing. *2014 Cancer Discov.* Aug;4(8):956-71.
24. Garrett JT, Arteaga CL. Resistance to HER2-directed antibodies and tyrosine kinase inhibitors: mechanisms and clinical implications. *Cancer Biol Ther.* 2011 May 1;11(9):793-800.
25. Gritsenko, P. G., Ilina, O. & Friedl, P. Interstitial guidance of cancer invasion. *The Journal of pathology* 226, 185-199, (2012).
26. Guo D, Hildebrandt IJ, Prins RM, Soto H, Mazzotta MM, Dang J, Czernin J, Shyy JY, Watson AD, Phelps M, Radu CG, Cloughesy TF, Mischel PS. The AMPK agonist AICAR inhibits the growth of EGFRvIII-expressing glioblastomas by inhibiting lipogenesis. *Proc Natl Acad Sci U S A.* 2009 Aug 4;106(31):12932-7.
27. Hägerstrand D, He X, Bradic Lindh M, Hoefs S, Hesselager G, Ostman A, Nistér M. Identification of a SOX2- dependent subset of tumor- and sphere-forming glioblastoma cells with a distinct tyrosine kinase inhibitor sensitivity profile. *2011 Neuro Oncol.* Nov;13(11):1178-91.
28. Hanahan D, Weinberg RA. Hallmarks of cancer: The next generation. *Cell.* 2011;144(5):646-674.

29. Hecht JR, Mitchell E, Chidiac T, Scroggin C, Hagenstad C, Spigel D, et al. A randomized phase IIIB trial of chemotherapy, bevacizumab, and panitumumab compared with chemotherapy and bevacizumab alone for metastatic colorectal cancer. *J Clin Oncol.* 2009;27:672–80.
30. Hecht JR, Mitchell E, Chidiac T, Scroggin C, Hagenstad C, Spigel D, Marshall J, Cohn A, McCollum D, Stella P, Deeter R, Shahin S, Amado RG. A randomized phase IIIB trial of chemotherapy, bevacizumab, and panitumumab compared with chemotherapy and bevacizumab alone for metastatic colorectal cancer. *J Clin Oncol.* 2009 Feb 10;27(5):672-80.
31. Hegi ME, Diserens A-C, Gorlia T, Hamou M-F, de Tribolet N, Weller M, et al. MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med.* 2005;352:997–1003.
32. Heo JC, Jung TH, Jung DY, Park WK, Cho H. Indatraline inhibits Rho- and calcium-mediated glioblastoma cell motility and angiogenesis. 2014 *Biochem Biophys Res Commun.* Jan 10;443(2):749-55.
33. Hochberg FH, Pruitt A. Assumptions in the radiotherapy of glioblastoma. *Neurology.* 1980 Sep;30(9):907-11.
34. Iwanami A, Gini B, Zanca C, Matsutani T, Assuncao A, Nael A, Dang J, Yang H, Zhu S, Kohyama J, Kitabayashi I, Cavenee WK, Cloughesy TF, Furnari FB, Nakamura M, Toyama Y, Okano H, Mischel PS. PML mediates glioblastoma resistance to mammalian target of rapamycin (mTOR)-targeted therapies. *Proc Natl Acad Sci U S A.* 2013 Mar 12;110(11):4339-44.
35. Johnson BE, Mazar T, Hong C, Barnes M, Aihara K, McLean CY, Fouse SD, Yamamoto S, Ueda H, Tatsuno K, Asthana S, Jalbert LE, Nelson SJ, Bollen AW, Gustafson WC, Charron E, Weiss WA, Smirnov IV, Song JS, Olshen AB, Cha S, Zhao Y, Moore RA, Mungall AJ, Jones SJ, Hirst M, Marra MA, Saito N, Aburatani H, Mukasa A, Berger MS, Chang SM, Taylor BS, Costello JF. Mutational analysis reveals the origin and therapy-driven evolution of recurrent glioma. 2014 *Science* 343, 189–93.
36. Kelley K, Knisely J, Symons M, Ruggieri R. Radioresistance of Brain Tumors. *Cancers (Basel).* 2016 Mar 30;8(4). pii: E42.
37. Kenney-Herbert E, Al-Mayhany T, Piccirillo SG, Fowler J, Spiteri I, Jones P, Watts C. CD15 Expression Does Not Identify a Phenotypically or Genetically Distinct Glioblastoma Population. *Stem Cells Transl Med.* 2015 Jul;4(7):822-31.

38. Knight, Z.A., Lin, H., and Shokat, K.M. (2010). Targeting the cancer kinome through polypharmacology. *Nat. Rev. Cancer* 10, 130-137.
39. Lamanaskas N, Novell A, Escoffre JM, Venslauskas M, Satkauskas S, Bouakaz A. Bleomycin delivery into cancer cells in vitro with ultrasound and SonoVue® or BR14® microbubbles. *J Drug Target.* 2013 May;21(4):407-14.
40. Lan X, Jörg DJ, Cavalli FMG, Richards LM, Nguyen LV, Vanner RJ, Guilhamon P, Lee L, Kushida MM, Pellacani D, Park NI, Coutinho FJ, Whetstone H, Selvadurai HJ, Che C, Luu B, Carles A, Moksa M, Rastegar N, Head R, Dolma S, Prinos P, Cusimano MD, Das S, Bernstein M, Arrowsmith CH, Mungall AJ, Moore RA, Ma Y, Gallo M, Lupien M, Pugh TJ, Taylor MD, Hirst M, Eaves CJ, Simons BD, Dirks PB. Fate mapping of human glioblastoma reveals an invariant stem cell hierarchy. *Nature.* 2017 Sep 14;549(7671):227-232.
41. Larkin J, Ascierto PA, Dréno B, Atkinson V, Liskay G, Maio M, Mandalà M, Demidov L, Stroyakovskiy D, Thomas L, de la Cruz-Merino L, Dutriaux C, Garbe C, Sovak MA, Chang I, Choong N, Hack SP, McArthur GA, Ribas A. Combined vemurafenib and cobimetinib in BRAF-mutated melanoma. *N Engl J Med.* 2014 Nov 13;371(20):1867-76.
42. Larkin J, Hodi FS, Wolchok JD. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med.* 2015 Sep 24;373(13):1270-1.
43. Larkin J, Hodi FS, Wolchok JD. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med.* 2015;373(13):1270-1271.
44. Lee JK, Wang J, Sa JK, Ladewig E, Lee HO, Lee IH, Kang HJ, Rosenbloom DS, Camara PG, Liu Z, van Nieuwenhuizen P, Jung SW, Choi SW, Kim J, Chen A, Kim KT, Shin S, Seo YJ, Oh JM, Shin YJ, Park CK, Kong DS, Seol HJ, Blumberg A, Lee JI, Iavarone A, Park WY, Rabadan R, Nam DH. Spatiotemporal genomic architecture informs precision oncology in glioblastoma. *Nat Genet.* 2017 Apr;49(4):594-599.
45. Lesueur P, Chevalier F, El-Habr EA, Junier MP, Chneiweiss H, Castera L, Müller E, Stefan D, Saintigny Y. Radiosensitization Effect of Talazoparib, a Parp Inhibitor, on Glioblastoma Stem Cells Exposed to Low and High Linear Energy Transfer Radiation. *Sci Rep.* 2018 Feb 26;8(1):3664.

46. Li W, Zhang H, Assaraf YG, Zhao K, Xu X, Xie J, Yang DH, Chen ZS. Overcoming ABC transporter-mediated multidrug resistance: Molecular mechanisms and novel therapeutic drug strategies. *Drug Resist Updat*. 2016 Jul;27:14-29
47. Lin F, de Gooijer MC, Roig EM, Buil LC, Christner SM, Beumer JH, Würdinger T, Beijnen JH, van Tellingen O. ABCB1, ABCG2, and PTEN determine the response of glioblastoma to temozolomide and ABT-888 therapy. *Clin Cancer Res*. 2014 May 15;20(10):2703-13.
48. Little SE, Popov S, Jury A, Bax DA, Doey L, Al-Sarraj S, Jurgensmeier JM, Jones C. Receptor tyrosine kinase genes amplified in glioblastoma exhibit a mutual exclusivity in variable proportions reflective of individual tumor heterogeneity. *2012 Cancer Res*. 72, 1614–20.
49. Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, Jouvet A, Scheithauer BW, Kleihues P. The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol*. 114, 97–109 (2007).
50. Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, Ohgaki H, Wiestler OD, Kleihues P, Ellison DW. The 2016 World Health Organization Classification of Tumors of the Central Nervous System: a summary. *Acta Neuropathol*. 2016 Jun;131(6):803-20.
51. Lu C-T, Zhao Y-Z, Wong HL, Cai J, Peng L, Tian X-Q. Current approaches to enhance CNS delivery of drugs across the brain barriers. *Int J Nanomedicine*. 2014;9:2241–57.
52. Meyer M, Reimand J, Lan X, Head R, Zhu X, Kushida M, Bayani J, Pressey JC, Lionel AC, Clarke ID, Cusimano M, Squire JA, Scherer SW, Bernstein M, Woodin MA, Bader GD, Dirks PB. Single cell-derived clonal analysis of human glioblastoma links functional and genomic heterogeneity. *2015 Proc Natl Acad Sci U S A*. Jan 20;112(3):851-6.
53. Montana, V. & Sontheimer, H. Bradykinin promotes the chemotactic invasion of primary brain tumors. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 31, 4858-4867, (2011).
54. Narayan RS, Molenaar P, Cornelissen F, Dik R, Roelofs I, Brands E, van Kuiken P, in 't Veld S, van Wieringen W, Lang FF, Verhaak R, Sulman E, Baumert BG, Stalpers LJA, Bailey D, Slotman BJ, Versteeg R, Tannous B, Noske D, Sminia P, Würdinger T, Koster J & Westerman BA, A cancer drug-atlas enables prediction of parallel drug vulnerabilities. Submitted.

55. Nathanson DA, Gini B, Mottahedeh J, Visnyei K, Koga T, Gomez G, Eskin A, Hwang K, Wang J, Masui K, Paucar A, Yang H, Ohashi M, Zhu S, Wykosky J, Reed R, Nelson SF, Cloughesy TF, James CD, Rao PN, Kornblum HI, Heath JR, Cavenee WK, Furnari FB, Mischel PS. Targeted therapy resistance mediated by dynamic regulation of extrachromosomal mutant EGFR DNA. *Science*. 2014 Jan 3;343(6166):72-6
56. Oberoi RK, Mittapalli RK, Elmquist WF. Pharmacokinetic assessment of efflux transport in sunitinib distribution to the brain. *J Pharmacol Exp Ther*. 2013 Dec;347(3):755-64.
57. Ohgaki H, Kleihues P. The definition of primary and secondary glioblastoma. *Clin Cancer Res*. 2013 Feb 15;19(4):764-72.
58. O'Rourke DM, Nasrallah MP, Desai A, Melenhorst JJ, Mansfield K, Morrisette JJD, Martinez-Lage M, Brem S, Maloney E, Shen A, Isaacs R, Mohan S, Plesa G, Lacey SF, Navenot JM, Zheng Z, Levine BL, Okada H, June CH, Brogdon JL, Maus MV. A single dose of peripherally infused EGFRvIII-directed CAR T cells mediates antigen loss and induces adaptive resistance in patients with recurrent glioblastoma. *Sci Transl Med*. 2017 Jul 19;9(399).
59. Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, Cahill DP, Nahed BV, Curry WT, Martuza RL, Louis DN, Rozenblatt-Rosen O, Suvà ML, Regev A, Bernstein BE. Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. 2014 *Science*. Jun 20;344(6190):1396-401.
60. Patel PH, Senico PL, Curiel RE, Motzer RJ. Phase I study combining treatment with temsirolimus and sunitinib malate in patients with advanced renal cell carcinoma. *Clin Genitourin Cancer*. 2009;7:24–7.
61. Patel PH1, Senico PL, Curiel RE, Motzer RJ. Phase I study combining treatment with temsirolimus and sunitinib malate in patients with advanced renal cell carcinoma. *Clin Genitourin Cancer*. 2009 Jan;7(1):24-7.
62. Peters KB, Lou E, Desjardins A, Reardon DA, Lipp ES, Miller E, Herndon JE 2nd, McSherry F, Friedman HS, Vredenburgh JJ. Phase II Trial of Upfront Bevacizumab, Irinotecan, and Temozolomide for Unresectable Glioblastoma. *Oncologist*. 2015 Jul;20(7):727-8.

63. Peters KB, Lou E, Desjardins A, Reardon DA, Lipp ES, Miller E, Herndon JE, McSherry F, Friedman HS, Vredenburgh JJ. Phase II Trial of Upfront Bevacizumab, Irinotecan, and Temozolomide for Unresectable Glioblastoma. *Oncologist*. 2015 Jul;20(7):727-8.
64. Peterson A, Bansal A, Hofman F, Chen TC, Zada G. A systematic review of inhaled intranasal therapy for central nervous system neoplasms: an emerging therapeutic option. *J Neurooncol*. 2014 Feb;116(3):437-46.
65. Phillips HS, Kharbanda S, Chen R, Forrest WF, Soriano RH, Wu TD, Misra A, Nigro JM, Colman H, Soroceanu L, Williams PM, Modrusan Z, Feuerstein BG, Aldape K. Molecular subclasses of high-grade glioma predict prognosis, delineate a pattern of disease progression, and resemble stages in neurogenesis. 2006 *Cancer Cell*. Mar;9(3):157- 73.
66. Piccirillo SG, Colman S, Potter NE, van Delft FW, Lillis S, Carnicer MJ, Kearney L, Watts C, Greaves M. Genetic and functional diversity of propagating cells in glioblastoma. *Stem Cell Reports*. 2015 Jan 13;4(1):7-15.
67. Piccirillo SG, Spiteri I, Sottoriva A, Touloumis A, Ber S, Price SJ, Heywood R, Francis NJ, Howarth KD, Collins VP, Venkitaraman AR, Curtis C, Marioni JC, Tavaré S, Watts C. Contributions to drug resistance in glioblastoma derived from malignant cells in the sub-ependymal zone. 2015 *Cancer Res*;75(1):194-202.
68. Porta R, Sánchez-Torres JM, Paz-Ares L, Massutí B, Reguart N, Mayo C, Lianes P, Queralt C, Guillem V, Salinas P, Catot S, Isla D, Pradas A, Gúrpide A, de Castro J, Polo E, Puig T, Tarón M, Colomer R, Rosell R. Brain metastases from lung cancer responding to erlotinib: the importance of EGFR mutation. *Eur Respir J*. 2011 Mar;37(3):624-31.
69. Prahallad A, Sun C, Huang S, Di Nicolantonio F, Salazar R, Zecchin D, Beijersbergen RL, Bardelli A, Bernards R. Unresponsiveness of colon cancer to BRAF(V600E) inhibition through feedback activation of EGFR. *Nature*. 2012 Jan 26;483(7387):100-3.
70. Rachlin K, Moore DH, Yount G. Infrasound sensitizes human glioblastoma cells to cisplatin-induced apoptosis. *Integr Cancer Ther*. 2013 Nov;12(6):517-27.

71. Reardon DA, Desjardins A, Vredenburgh JJ, Gururangan S, Friedman AH, Herndon JE 2nd, Marcellino J, Norfleet JA, McLendon RE, Sampson JH, Friedman HS. Phase 2 trial of erlotinib plus sirolimus in adults with recurrent glioblastoma. *J Neurooncol*. 2010 Jan;96(2):219-30.
72. Reardon DA, Desjardins A, Vredenburgh JJ, Gururangan S, Friedman AH, Herndon JE 2nd, Marcellino J, Norfleet JA, McLendon RE, Sampson JH, Friedman HS. Phase 2 trial of erlotinib plus sirolimus in adults with recurrent glioblastoma. *J Neurooncol*. 2010 Jan;96(2):219-30.
73. References
74. Reyes-Botero G, Cartalat-Carel S, Chinot OL, Barrie M, Taillandier L, Beauchesne P, Catry-Thomas I, Barrière J, Guillo JS, Fabbro M, Frappaz D, Benouaich-Amiel A, Le Rhun E, Campello C, Tennevet I, Ghiringhelli F, Tanguy ML, Mokhtari K, Honnorat J, Delattre JY. Temozolomide Plus Bevacizumab in Elderly Patients with Newly Diagnosed Glioblastoma and Poor Performance Status: An ANOCEF Phase II Trial (ATAG). *Oncologist*. 2018 May;23(5):524-e44
75. Reyes-Botero G, Cartalat-Carel S, Chinot OL, Barrie M, Taillandier L, Beauchesne P, Catry-Thomas I, Barrière J, Guillo JS, Fabbro M, Frappaz D, Benouaich-Amiel A, Le Rhun E, Campello C, Tennevet I, Ghiringhelli F, Tanguy ML, Mokhtari K, Honnorat J, Delattre JY. Temozolomide Plus Bevacizumab in Elderly Patients with Newly Diagnosed Glioblastoma and Poor Performance Status: An ANOCEF Phase II Trial (ATAG). *Oncologist*. 2018 May;23(5):524-e44.
76. Rozengurt E, Soares HP, Sinnett-Smith J. Suppression of feedback loops mediated by PI3K/mTOR induces multiple over-activation of compensatory pathways: an unintended consequence leading to drug resistance. *Mol Cancer Ther*. 2014 Nov; 13(11): 2477–2488.
77. Saito N, Fu J, Zheng S, Yao J, Wang S, Liu DD, Yuan Y, Sulman EP, Lang FF, Colman H, Verhaak RG, Yung WK, Koul D. A high Notch pathway activation predicts response to  $\gamma$  secretase inhibitors in proneural subtype of glioma tumor initiating cells. 2014 *Stem Cells*. Jan;32(1):301-12. 8.
78. Sequist LV, Waltman BA, Dias-Santagata D, Digumarthy S, Turke AB, Fidias P, Bergethon K, Shaw AT, Gettinger S, Cosper AK, Akhavanfard S, Heist RS, Temel J, Christensen JG, Wain JC, Lynch TJ, Vernovsky K, Mark EJ, Lanuti M, Iafrate AJ, Mino-Kenudson M, Engelman JA. Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors. *Sci Transl Med*. 2011 Mar 23;3(75):75ra26.

79. Sherriff J, Tamangani J, Senthil L, Cruickshank G, Spooner D, Jones B, Brookes C, Sanghera P. Patterns of relapse in glioblastoma multiforme following concomitant chemoradiotherapy with temozolomide. *Br J Radiol.* 2013 Feb;86(1022):20120414.
80. Sidaway P. CNS cancer: Glioblastoma subtypes revisited. *Nat Rev Clin Oncol.* 2017 Oct;14(10):587.
81. Sminia P, Westerman BA. Blood-brain barrier crossing and breakthroughs in glioblastoma therapy. *Br J Clin Pharmacol.* 2016;81(6):1018-1020.
82. Snuderl M, Fazlollahi L, Le LP, Nitta M, Zhelyazkova BH, Davidson CJ, Akhavanfard S, Cahill DP, Aldape KD, Betensky RA, Louis DN, Iafrate AJ. Mosaic amplification of multiple receptor tyrosine kinase genes in glioblastoma. 2011 *Cancer Cell* 20, 810–7.
83. Sottoriva A, Spiteri I, Piccirillo SG, Touloumis A, Collins VP, Marioni JC, Curtis C, Watts C, Tavaré S. Intratumor heterogeneity in human glioblastoma reflects cancer evolutionary dynamics. 2013 *Proc. Natl. Acad. Sci. U. S. A.* 110, 4009–14.
84. Stieber D, Golebiewska A, Evers L, Lenkiewicz E, Brons NH, Nicot N, Oudin A, Bougnaud S, Hertel F, Bjerkvig R, Vallar L, Barrett MT, Niclou SP. Glioblastomas are composed of genetically divergent clones with distinct tumorigenic potential and variable stem cell-associated phenotypes. *Acta Neuropathol.* 2014 Feb;127(2):203-19. doi: 10.1007/s00401-013-1196-4. Epub 2013 Oct 24.
85. Stupp R, Hegi ME, Mason WP, van den Bent MJ, Taphoorn MJ, Janzer RC, Ludwin SK, Allgeier A, Fisher B, Belanger K, Hau P, Brandes AA, Gijtenbeek J, Marosi C, Vecht CJ, Mokhtari K, Wesseling P, Villa S, Eisenhauer E, Gorlia T, Weller M, Lacombe D, Cairncross JG, Mirimanoff RO. Effects of radiotherapy with concomitant and adjuvant temozolomide versus radiotherapy alone on survival in glioblastoma in a randomised phase III study: 5-year analysis of the EORTC-NCIC trial. *Lancet Oncol.* 10, 459–466 (2009).
86. Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus Concomitant Adjuvant Temozolomide for Glioblastoma. *N. Engl. J. Med.* 987–96 (2005).
87. Sun C, Hobor S, Bertotti A, Zecchin D, Huang S, Galimi F, Cottino F, Prahallad A, Grenrum W, Tzani A, Schlicker A, Wessels LF, Smit EF, Thunnissen E, Halonen P, Liefink C, Beijersbergen RL, Di Nicolantonio F, Bardelli A, Trusolino L, Bernards R. Intrinsic resistance to MEK inhibition in KRAS



- mutant lung and colon cancer through transcriptional induction of ERBB3. *Cell Rep.* 2014 Apr 10;7(1):86-93.
88. Sun C, Wang L, Huang S, Heynen GJ, Prahallad A, Robert C, Haanen J, Blank C, Wesseling J, Willems SM, Zecchin D, Hobor S, Bajpe PK, Liefstink C, Mateus C, Vagner S, Grenrum W, Hofland I, Schlicker A, Wessels LF, Beijersbergen RL, Bardelli A, Di Nicolantonio F, Eggermont AM, Bernards R. Reversible and adaptive resistance to BRAF(V600E) inhibition in melanoma. *Nature.* 2014 Apr 3;508(7494):118-22.
89. Svensson F, Westerman BA, Wurdinger T, Bailey D. GBM Drug Bank – a new resource for glioblastoma drug discovery and informatics research. *Neuro Oncol.* 2018 Aug 20.
90. Szerlip NJ, Pedraza A, Chakravarty D, Azim M, McGuire J, Fang Y, Ozawa T, Holland EC, Huse JT, Jhanwar S, Leversha MA, Mikkelsen T, Brennan CW. Intratumoral heterogeneity of receptor tyrosine kinases EGFR and PDGFRA amplification in glioblastoma defines subpopulations with distinct growth factor response. 2012 *Proc. Natl. Acad. Sci. U. S. A.* 109, 3041–6.
91. Tang SC, Lagas JS, Lankheet NA, Poller B, Hillebrand MJ, Rosing H, Beijnen JH, Schinkel AH. Brain accumulation of sunitinib is restricted by P-glycoprotein (ABCB1) and breast cancer resistance protein (ABCG2) and can be enhanced by oral elacridar and sunitinib coadministration. *Int J Cancer.* 2012 Jan 1;130(1):223-33.
92. Tol J, Koopman M, Cats A, Rodenburg CJ, Creemers GJ, Schrama JG, Erdkamp FL, Vos AH, van Groenigen CJ, Sinnige HA, Richel DJ, Voest EE, Dijkstra JR, Vink-Börger ME, Antonini NF, Mol L, van Krieken JH, Dalesio O, Punt CJ. Chemotherapy, bevacizumab, and cetuximab in metastatic colorectal cancer. *N Engl J Med.* 2009 Feb 5;360(6):563-72.
93. Tol J, Koopman M, Cats A, Rodenburg CJ, Creemers GJ, Schrama JG. Chemotherapy, bevacizumab, and cetuximab in metastatic colorectal cancer. *N Engl J Med.* 2009;360:563–72.
94. Tolcher AW, Khan K, Ong M, Banerji U, Papadimitrakopoulou V, Gandara DR, Patnaik A, Baird RD, Olmos D, Garrett CR, Skolnik JM, Rubin EH, Smith PD, Huang P, Learoyd M, Shannon KA, Morosky A, Tetteh E, Jou YM, Papadopoulos KP, Moreno V, Kaiser B, Yap TA, Yan L, de Bono JS. Antitumor activity in RAS-driven tumors by blocking AKT and MEK. *Clin Cancer Res.* 2015 Feb 15;21(4):739-48.

95. Tolcher AW, Patnaik A, Papadopoulos KP, Rasco DW, Becerra CR, Allred AJ, Orford K, Aktan G, Ferron-Brady G, Ibrahim N, Gauvin J, Motwani M, Cornfeld M. Phase I study of the MEK inhibitor trametinib in combination with the AKT inhibitor afuresertib in patients with solid tumors and multiple myeloma. *Cancer Chemother Pharmacol*. 2015 Jan;75(1):183-9.
96. Trnovec T, Kállay Z, Bezek Š. Effects of ionizing radiation on the blood brain barrier permeability to pharmacologically active substances. *Int. J. Radiat. Oncol. Biol. Phys.* 19, 1581–1587 (2016).
97. Turner KM, Deshpande V, Beyter D, Koga T, Rusert J, Lee C, Li B, Arden K, Ren B, Nathanson DA, Kornblum HI, Taylor MD, Kaushal S, Cavenee WK, Wechsler-Reya R, Furnari FB, Vandenberg SR, Rao PN, Wahl GM, Bafna V, Mischel PS. Extrachromosomal oncogene amplification drives tumour evolution and genetic heterogeneity. *Nature*. 2017 Mar 2;543(7643):122-125.
98. Tzelepis K, Koike-Yusa H, De Braekeleer E, Li Y, Metzakopian E, Dovey OM, Mupo A, Grinkevich V, Li M, Mazan M, Gozdecka M, Ohnishi S, Cooper J, Patel M, McKerrell T, Chen B, Domingues AF, Gallipoli P, Teichmann S, Ponstingl H, McDermott U, Saez-Rodriguez J, Huntly BJ, Iorio F, Pina C, Vassiliou GS, Yusa K. A CRISPR Dropout Screen Identifies Genetic Vulnerabilities and Therapeutic Targets in Acute Myeloid Leukemia. *Cell Rep*. 2016 Oct 18;17(4):1193-1205.
99. Upadhyay RK. Drug Delivery Systems, CNS Protection, and the Blood Brain Barrier. *Biomed Res Int*. 2014;2014:1–37.
100. van Tellingen O, Yetkin-Arik B, de Gooijer MC, Wesseling P, Wurdinger T, de Vries HE. Overcoming the blood-brain tumor barrier for effective glioblastoma treatment. *Drug Resist Updat*. 2015 Mar;19:1-12.
101. Verhaak RG, Hoadley KA, Purdom E, Wang V, Qi Y, Wilkerson MD, Miller CR, Ding L, Golub T, Mesirov JP, Alexe G, Lawrence M, O'Kelly M, Tamayo P, Weir BA, Gabriel S, Winckler W, Gupta S, Jakkula L, Feiler HS, Hodgson JG, James CD, Sarkaria JN, Brennan C, Kahn A, Spellman PT, Wilson RK, Speed TP, Gray JW, Meyerson M, Getz G, Perou CM, Hayes DN. Integrated Genomic Analysis Identifies Clinically Relevant Subtypes of Glioblastoma Characterized by Abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer Cell* 17, 98–110 (2010).
102. Verhaak RG. Moving the needle: Optimizing classification for glioma. *Sci Transl Med*. 2016 Aug 3;8(350):350fs14. Review.

103. Vogelbaum MA, Aghi MK. Convection-enhanced delivery for the treatment of glioblastoma. *Neuro Oncol.* 2015 Mar;17 Suppl 2:ii3-ii8.
104. Wang J, Cazzato E, Ladewig E, et al. Clonal evolution of glioblastoma under therapy. *Nat Genet.* 2016;48(7):768-776.
105. Wang T, Birsoy K, Hughes NW, Krupczak KM, Post Y, Wei JJ, Lander ES, Sabatini DM. Identification and characterization of essential genes in the human genome. *Science* 350, 1096-1101 (2015).
106. Weller M, van den Bent M, Tonn JC, Stupp R, Preusser M, Cohen-Jonathan-Moyal E, Henriksson R, Le Rhun E, Balana C, Chinot O, Bendszus M, Reijneveld JC, Dhermain F, French P, Marosi C, Watts C, Oberg I, Pilkington G, Baumert BG, Taphoorn MJB, Hegi M, Westphal M, Reifenberger G, Soffietti R, Wick W; European Association for Neuro-Oncology (EANO) Task Force on Gliomas. European Association for Neuro-Oncology (EANO) guideline on the diagnosis and treatment of adult astrocytic and oligodendroglial gliomas. *Lancet Oncol.* 2017 Jun;18(6):e315-e329.
107. Weller M, van den Bent M, Tonn JC, Stupp R, Preusser M, Cohen-Jonathan-Moyal E, Henriksson R, Le Rhun E, Balana C, Chinot O, Bendszus M, Reijneveld JC, Dhermain F, French P, Marosi C, Watts C, Oberg I, Pilkington G, Baumert BG, Taphoorn MJB, Hegi M, Westphal M, Reifenberger G, Soffietti R, Wick W; European Association for Neuro-Oncology (EANO) Task Force on Gliomas. European Association for Neuro-Oncology (EANO) guideline on the diagnosis and treatment of adult astrocytic and oligodendroglial gliomas. *Lancet Oncol.* 2017 Jun;18(6):e315-e329.
108. Weller, M. et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *Cancer/Radiothérapie* 9, 196–197 (2005).
109. Wilson TR, Fridlyand J, Yan Y, Penuel E, Burton L, Chan E, Peng J, Lin E, Wang Y, Sosman J, Ribas A, Li J, Moffat J, Sutherlin DP, Koeppen H, Merchant M, Neve R, Settleman J. Widespread potential for growth-factor-driven resistance to anticancer kinase inhibitors. *Nature.* 2012 Jul 26;487(7408):505-9.

110. Zanca C, Villa GR, Benitez JA, Thorne AH, Koga T, D'Antonio M, Ikegami S, Ma J, Boyer AD, Banisadr A, Jameson NM, Parisian AD, Eliseeva OV, Barnabe GF, Liu F, Wu S, Yang H, Wykosky J, Frazer KA, Verkhusha VV, Isaguliantz MG, Weiss WA, Gahman TC, Shiau AK, Chen CC, Mischel PS, Cavenee WK, Furnari FB. Glioblastoma cellular cross-talk converges on NF- $\kappa$ B to attenuate EGFR inhibitor sensitivity. *Genes Dev.* 2017 Jun 15;31(12):1212-1227.
111. Zhitomirsky B, Assaraf YG. Lysosomes as mediators of drug resistance in cancer. *Drug Resist Updat.* 2016 Jan;24:23-33.
112. Zhu, Y. & Parada, L. F. The molecular and genetic basis of neurological tumours. *Nat. Rev. Cancer* 2, 616–626 (2002).