

The 2017 Report of The Lancet Countdown on Health and Climate Change

*From 25 years of inaction to a global
transformation for public health*

Nick Watts, Markus Amann, Sonja Ayeb-Karlsson, Kristine Belesova, Timothy Bouley, Maxwell Boykoff, Peter Byass, Wenjia Cai, Diarmid Campbell-Lendrum, Jonathan Chambers, Peter M Cox, Meaghan Daly, Niheer Dasandi, Michael Davies, Michael Depledge, Anneliese Depoux, Paula Dominguez-Salas, Paul Drummond, Paul Ekins, Antoine Flahault, Howard Frumkin, Lucien Georgeson, Mostafa Ghanei, Delia Grace, Hilary Graham, Rébecca Grojsman, Andy Haines, Ian Hamilton, Stella Hartinger, Anne Johnson, Ilan Kelman, Gregor Kiesewetter, Dominic Kniveton, Lu Liang, Melissa Lott, Robert Lowe, Georgina Mace, Maquins Odhiambo Sewe, Mark Maslin, Slava Mikhaylov, James Milner, Ali Mohammad Latifi, Maziar Moradi-Lakeh, Karyn Morrissey, Kris Murray, Tara Neville, Maria Nilsson, Tadj Oreszczyn, Fereidoon Owfi, David Pencheon, Steve Pye, Mahnaz Rabbaniha, Elizabeth Robinson, Joacim Rocklöv, Stefanie Schütte, Joy Shumake-Guillemot, Rebecca Steinbach, Meisam Tabatabaei, Nicola Wheeler, Paul Wilkinson, Peng Gong, Hugh Montgomery*, Anthony Costello**

* Denotes Co-Chair

[Current Word Count: 21,749

(excluding figures, captions, tables, references and executive summary)]

Table of Contents

List of Figures, Tables, and Panels	5
List of Figures	5
List of Tables	7
List of Panels	7
List of Abbreviations	9
Executive Summary	10
Introduction	14
Indicators of Progress on Health and Climate Change	14
Delivering the Paris Agreement for Better Health.....	16
1. Climate Change Impacts, Exposures and Vulnerability.....	18
Introduction	18
Indicator 1.1: Health effects of temperature change	19
Indicator 1.2: Health effects of heatwaves	20
Indicator 1.3: Change in labour capacity	22
Indicator 1.4: Lethality of weather-related disasters	24
Indicator 1.5: Global health trends in climate-sensitive diseases	26
Indicator 1.6: Climate-sensitive infectious diseases	27
Indicator 1.7: Food security and undernutrition	29
Indicator 1.7.1: Vulnerability to undernutrition	30
Indicator 1.7.2: Marine primary productivity	31
Indicator 1.8: Migration and population displacement.....	32
Conclusion.....	34
2. Adaptation Planning and Resilience for Health	36
Introduction	36
Indicator 2.1: National adaptation plans for health	36
Indicator 2.2: City-level climate change risk assessments.....	38
Indicator 2.3: Detection and early warning of, preparedness for, and response to climate related health emergencies.....	39
Indicator 2.4: Climate information services for health.....	43
Indicator 2.5: National assessments of climate change impacts, vulnerability, and adaptation for health	44
Indicator 2.6: Climate-resilient health infrastructure.....	45
Conclusion.....	46
3. Mitigation Actions and Health Co-Benefits.....	46
Introduction	47

Tracking the health co-benefits of climate change mitigation	47
Energy Supply and Demand Sectors	48
Indicator 3.1: Carbon intensity of the energy system	48
Indicator 3.2: Coal phase-out.....	49
Indicator 3.3: Zero-carbon emission electricity	51
Indicator 3.4: Access to clean energy	52
Indicator 3.5: Exposure to ambient air pollution.....	53
3.5.1: Exposure to air pollution in cities	53
3.5.2: Sectoral contributions to air pollution.....	55
3.5.3: Premature mortality from ambient air pollution by sector	57
Transport Sector	58
Indicator 3.6: Clean fuel use for transport.....	58
Indicator 3.7: Sustainable travel infrastructure and uptake.....	59
Food and agriculture.....	62
Indicator 3.8: Ruminant meat for human consumption	62
Healthcare sector	64
Indicator 3.9: Healthcare sector emissions.....	64
Conclusion.....	65
4. Finance & Economics	66
Introduction	66
Indicator 4.1: Investments in zero-carbon energy and energy efficiency	67
Indicator 4.2: Investment in coal capacity	68
Indicator 4.3: Funds divested from fossil fuels.....	69
Indicator 4.4: Economic losses due to climate-related extreme events	69
Indicator 4.5: Employment in low-carbon and high-carbon industries	72
Indicator 4.6: Fossil fuel subsidies	73
Indicator 4.7: Coverage and strength of carbon pricing	74
Indicator 4.8: Use of carbon pricing revenues.....	75
Indicator 4.9: Spending on adaptation for health and health-related activities	76
Indicator 4.10: Health adaptation funding from global climate financing mechanisms	78
Conclusion.....	79
5. Public and Political Engagement.....	81
Introduction	81
Indicator 5.1: Media coverage of health and climate change	81
5.1.1: Global newspaper reporting on health and climate change	82
5.1.2: In-depth analysis of newspaper coverage on health and climate change.....	83

Indicator 5.2: Health and climate change in scientific journals	83
Indicator 5.3: Health and climate change in the United Nations General Assembly	85
Conclusion.....	86
Conclusion - the Lancet Countdown in 2017	88
The direction of travel is set	88
References	90

List of Figures, Tables, and Panels

List of Figures

Figure 1.1 Mean summer warming from 2000 to 2016 area weighted and exposure weighted, relative to the 1986-2008 recent past average.

Figure 1.2 The change in exposure (in people aged over 65 years) to heatwaves from 2000 to 2016, relative to the heatwave exposure average from 1986-2008.

Figure 1.3 The area and exposure weighted change in mean heatwave lengths globally from 2000 to 2016 (in people aged over 65 years), relative to the 1986-2008 recent past average.

Figure 1.4 The exposure weighted labour capacity change (%) globally from 2000 to 2016, relative to the 1986-2008 recent past average.

Figure 1.5 Map of the change in labour capacity loss from 2000 to 2016, relative to the 1986-2008 recent past average.

Figure 1.6 Deaths and people affected by weather-related disasters. 1.6a) Percentage change over time in the global number of deaths, the number of those affected, and the ratio of these (measured against 1990-2009). 1.6b) Change over time in the number of people affected globally by different weather-related disasters.

Figure 1.7 Trends in mortality from selected causes of death as estimated by the Global Burden of Disease 2015, for the period 1990 to 2015, by WHO region.

Figure 1.8 Left: Academic publications reporting climate-sensitive infectious diseases by year. Right: proportion of responses reported in publications by year and direction of impact.

Figure 1.9 Average annual vectorial capacity (VC) for dengue in *Aedes aegypti* and *Aedes albopictus* for selected *Aedes*-positive countries (countries with *Aedes* present) (top panel; matrix coloured relative to country mean 1950-2015; red = relatively higher VC, blue = relatively lower VC; countries ordered by centroid latitude (north to south)). Bottom panel: average vectorial capacity (VC) for both vectors calculated globally (results shown relative to 1990 baseline).

Figure 1.10 Total number of undernourished people multiplied by regional dependency on grain production for countries.

Figure 2.1 Countries with national health climate adaptation strategies or plans.

Figure 2.2 Number of global cities undertaking climate change risk assessments by a) income grouping, and b) WHO region.

Figure 2.3 IHR Core Capacity Requirement by WHO region 2.3a) Percentage attainment of human resources available to implement the International Health Regulations Core Capacity Requirements. 2.3b) Percentage attainment of having indicator-based surveillance for early warning function for the early detection of a public health event. 2.3c) Percentage attainment for having a multi-hazard public health emergency preparedness and response plan developed and implemented. 2.3d) Percentage attainment of having a public health emergency response mechanisms established and functioning.

Figure 2.4 National Meteorological and Hydrological Services (NHMSs) of WHO member states reporting to provide targeted/tailored climate information, products and services to the health sector.

Figure 2.5 Countries with national assessment of climate change impacts, vulnerability and adaptation for health.

Figure 2.6 Countries taking measures to increase the climate resilience of health infrastructure.

Figure 3.1 Carbon intensity of Total Primary Energy Supply (TPES) for selected countries, and total CO₂ emissions (shaded area against secondary y-axis), 1971-2013.

Figure 3.2 Total primary coal supply by region, and globally (shaded area against secondary y-axis), 1990-2013.

Figure 3.3 Renewable and zero-carbon emission energy sources electricity generation a) Share of electricity generated from zero carbon sources; b) Electricity generated from zero carbon sources, TWh; c) Share of electricity generated from renewable sources (excluding hydro); d) Electricity generated from renewable sources (excl. hydro), TWh.

Figure 3.4 Proportion of population relying primarily on clean fuels and technology.

Figure 3.5 Annual mean PM_{2.5} concentration vs per capita GDP for 143 cities in the SHUE database. Colours indicate WHO regions: blue – Africa; red – Europe; green – the Americas; Lime – Eastern Mediterranean; orange – Western Pacific; purple – South East Asia. The dotted line marks the WHO recommended guidance level of 10 µg.m⁻³.

Figure 3.6 Selected primary air pollutants and their sources globally in 2015.

Figure 3.7 a) Energy related PM_{2.5} emissions in 2015 and b) NO_x emissions from transport from 1990-2010 by region.

Figure 3.8 Health impacts of exposure to ambient PM_{2.5} in terms of annual premature deaths per million inhabitants in South and East Asian countries in 2015, broken down by key sources of pollution.

Figure 3.9 Per capita fuel use by type (TJ/person) for transport sector with all fuels

Figure 3.10 Cumulative Global Electric Vehicle Sales. Note: BEV is Battery Electric Vehicle and PHEV is Plug-in Hybrid Electric Vehicle.

Figure 3.11 Modal Shares in world cities. Note: 'Other' typically includes paratransit (transport for people with disabilities) and/or electronic bikes.

Figure 3.12 Trends in modal share in selected cities. Note: Data from Santiago in 1991 represents travel on a usual day; Data from Sydney represent Weekdays only; Cycling modal share in Sydney is <1%.

Figure 3.13 The total amount of ruminant meat available for human consumption in kg/capita/year by WHO-defined regions.

Figure 3.14 The proportion of energy (kcal/capita/day) available for human consumption from ruminant meat vs from all food sources by WHO-defined regions.

Figure 4.1 Annual Investment in the Global Energy System.

Figure 4.2 Annual Investment in coal-fired power capacity.

Figure 4.3 Economic Losses from Climate-Related Events – Absolute.

Figure 4.4 Economic Losses from Climate-Related Events – Intensity.

Figure 4.5 Employment in Renewable Energy and Fossil Fuel Extraction.

Figure 4.6 Global Fossil Fuel Consumption Subsidies - 2010-2015.

Figure 4.7 Carbon Pricing Instruments implemented, scheduled for implementation and under consideration.

Figure 4.8 For the financial year 2015-2016. 4.8a) Total health and health-related adaptation spending and 4.8b) health and health-related adaptation and resilience to climate change (A&RCC) spending as a proportion of GDP. All plots are disaggregated by World Bank Income Grouping.

Figure 4.9 Year on year multilateral and bilateral funding for all adaptation projects and health adaptation projects (2003 through May 2017).

Figure 5.1 Newspaper reporting on health and climate change (for 18 newspapers) from 2007 to 2016, broken down by WHO region.

Figure 5.2 Number of scientific publications on climate change and health per year (2007-2016) from PubMed and Web of Science journals.

Figure 5.3 Political engagement with the intersection of health and climate change, represented by joint references to health and climate change in the UNGD.

Figure 5.4 Regional political engagement with the intersection of health and climate change, represented by joint references to health and climate change in the UNGD, broken down by WHO region.

List of Tables

Table 1 Thematic groups and indicators for the Lancet Countdown's 2017 report.

Table 1.1 Locations migrating now due to only climate change.

Table 4.1 Carbon Pricing - Global Coverage and Weighted Average Prices. *Global emissions coverage is based on 2012 total anthropogenic CO₂ emissions.

Table 4.2. Carbon Pricing revenues and allocation in 2016.

List of Panels

Panel 1 Developing Lancet Countdown's Indicators: An Iterative and Open Process.

Panel 1.1 Mental health and Climate Change.

Panel 2.1 WHO-UNFCCC Climate and Health Country Profiles.

Panel 2.2 The International Health Regulations.

Panel 3.1 Energy and Household Air Pollution in Peru.

Panel 4.1 International Donor Action on Climate Change and Health.

1 List of Abbreviations

- 2 A&RCC – Adaptation & Resilience to Climate
- 3 Change
- 4 AAP – Ambient Air Pollution
- 5 AUM – Assets Under Management
- 6 BEV – Battery Electric Vehicle
- 7 CDP – Carbon Disclosure Project
- 8 CFU – Climate Funds Update
- 9 CO₂ – Carbon Dioxide
- 10 COP – Conference of the Parties
- 11 COPD – Chronic Obstructive Pulmonary
- 12 Disease
- 13 CPI – Consumer Price Indices
- 14 DALYs – Disability Adjusted Life Years
- 15 DPSEEA – Driving Force-Pressure-State-
- 16 Exposure-Effect-Action
- 17 ECMWF – European Centre for Medium-
- 18 Range Weather Forecasts
- 19 EJ – Exajoule
- 20 EM-DAT – Emergency Events Database
- 21 ERA – European Research Area
- 22 ETR – Environmental Tax Reform
- 23 ETS – Emissions Trading System
- 24 EU – European Union
- 25 EU28 – 28 European Union Member States
- 26 FAO – Food and Agriculture Organization of
- 27 the United Nations
- 28 FAZ – Frankfurter Allgemeine Zeitung
- 29 FISE – Social Inclusion Energy Fund
- 30 GBD – Global Burden of Disease
- 31 GDP – Gross Domestic Product
- 32 GHG – Greenhouse Gas
- 33 GtCO₂ – Gigatons of Carbon Dioxide
- 34 GW – Gigawatt
- 35 GWP – Gross World Product
- 36 HAB – Harmful Algal Blooms
- 37 HIC – High Income Countries
- 38 ICS – Improved Cook Stove
- 39 IEA – International Energy Agency
- 40 IHR – International Health Regulations
- 41 IPC – Infection Prevention and Control
- 42 IPCC - Intergovernmental Panel on Climate
- 43 Change
- 44 IRENA - International Renewable Energy
- 45 Agency
- 46 LMICs – Low and Middle Income Countries
- 47 LPG – Liquefied Petroleum Gas
- 48 Mt – Megaton
- 49 MtCO_{2e} – Metric Tons of Carbon Dioxide
- 50 Equivalent
- 51 NAP – National Adaptation Plan
- 52 NDCs = Nationally Determined Contributions
- 53 NHMSs – National Meteorological and
- 54 Hydrological Services
- 55 NHS- National Health Service
- 56 NO_x – Nitrogen Oxide
- 57 OECD – Organization for Economic
- 58 Cooperation and Development
- 59 PHEV – Plug-in Hybrid Electric Vehicle
- 60 PM_{2.5}– Fine Particulate Matter
- 61 PV – Photovoltaic
- 62 SDG – Sustainable Development Goal
- 63 SDU – Sustainable Development Unit
- 64 SHUE – Sustainable Healthy Urban
- 65 Environments
- 66 SO₂ – Sulphur Dioxide
- 67 SSS – Sea Surface Salinity
- 68 SST – Sea Surface Temperature
- 69 tCO₂ – Tons of Carbon Dioxide
- 70 tCO₂/TJ – Total Carbon Dioxide per Terajoule
- 71 TJ – Terajoule
- 72 TPES – Total Primary Energy Supply
- 73 TWh – Terawatt Hours
- 74 UN – United Nations
- 75 UNFCCC – United Nations Framework
- 76 Convention on Climate Change
- 77 UNGA – United Nations General Assembly
- 78 UNGD – United Nations General Debate
- 79 VC – Vectorial Capacity
- 80 WHO – World Health Organization
- 81 WMO – World Meteorological Organization

82

83 Executive Summary

84

85 The Lancet Countdown tracks progress on the relationships between human health and climate
86 change, providing an independent assessment of global progress to implement the Paris Agreement,
87 and the health implications of these actions.

88 It follows on from the work of the 2015 Lancet Commission, which concluded that anthropogenic
89 climate change threatens to undermine the last 50 years of gains in public health, and conversely,
90 that a comprehensive response to climate change could be “the greatest global health opportunity
91 of the 21st century”.

92 The Lancet Countdown exists as a collaboration between 24 academic institutions and inter-
93 governmental organisations, based in every continent, and with representation from a wide range of
94 disciplines, including: climate scientists, ecologists, economists, engineers, experts in energy, food
95 and transport systems, geographers, mathematicians, social and political scientists, public health
96 professionals, and physicians. The collaboration reports annual indicators across five domains:
97 climate change impacts, exposures and vulnerability; adaptation planning and resilience for health;
98 mitigation actions and health co-benefits; economics and finance; and public and political
99 engagement.

100 The 2017 key messages from its 40 indicators in its first annual report are summarised below.

101

102 **The human symptoms of climate change are unequivocal and potentially irreversible – affecting**
103 **the health of populations around the world, today. Whilst these effects will disproportionately**
104 **impact the most vulnerable in society, every community will be affected.**

105 The impacts of climate change are disproportionately affecting the health of vulnerable populations,
106 and those in low- and middle-income countries. By undermining the social and environmental
107 determinants that underpin good health, it exacerbates social, economic and demographic
108 inequalities with the effects eventually felt by all populations.

109 The evidence is clear that exposure to more frequent and intense heatwaves are increasing, with an
110 estimated 125 million additional vulnerable adults exposed to heatwaves from 2000 to 2016
111 (Indicator 1.2). Higher ambient temperatures have resulted in estimated reduction of 5.3% in labour
112 productivity, globally, from 2000 to 2016 (Indicator 1.3). Taken as a whole, a 44% increase in
113 weather-related disasters has been observed since 2000, with no clear upward or downward trend
114 in the lethality of these extreme events (Indicator 1.4), potentially suggesting the beginning of an
115 adaptive response to climate change. Yet, the impacts of climate change are projected to worsen
116 over time, with current levels of adaptation becoming insufficient in the future. The total value of
117 economic losses that resulted from climate-related events has been increasing since 1990, and
118 totalled \$129 billion in 2016, with 99% of these losses in low-income countries uninsured (Indicator
119 4.4). Additionally, over the longer-term, altered climatic conditions are contributing to growing
120 vectorial capacity for the transmission of dengue fever by *Aedes aegypti*, reflecting an estimated
121 9.4% increase since 1950 (Indicator 1.6).

122 If governments and the global health community do not learn from the past experience of HIV/AIDS
123 and the recent outbreaks of Ebola and Zika virus, another slow response will result in an irreversible
124 and unacceptable cost to human health.

125

126 **The delayed response to climate change over the past 25 years has jeopardised human life and**
127 **livelihoods.**

128 Since the UN Framework Convention on Climate Change (UNFCCC) commenced global efforts to
129 tackle climate change in 1992, most of the indicators tracked by the Lancet Countdown have either
130 shown limited progress, particularly with regards to adaptation, or moved in the wrong direction,
131 particularly in relation to mitigation. Most fundamentally, carbon emissions, and global
132 temperatures, have continued to rise..

133 A growing number of countries are assessing their vulnerabilities to climate change, and are
134 increasingly developing adaptation and emergency preparedness plans, and providing climate
135 information to health services (Indicators 2.1, 2.3-2.6). The same is seen at the city-level, with over
136 449 cities around the world reporting having undertaken a climate change risk assessment (Indicator
137 2.2). However, the coverage and adequacy of such measures in protecting against the growing risks
138 of climate change to health remains uncertain. Indeed, health and health-related adaptation funding
139 accounts for 4.6% and 13.3% of total global adaptation spending, respectively (Indicator 4.9).

140 Whilst there has been some recent progress in strengthening health resilience to climate impacts, it
141 is clear that adaptation to new climatic conditions can only protect up to a point; an analogy to
142 human physiology is useful here. The human body can adapt to insults caused by a self-limiting
143 minor illness with relative ease. However, where disease steadily worsens, positive feedback cycles
144 and limits to adaptation are quickly reached. This is particularly true when many systems are
145 affected, and where the failure of one system may impact on the function of another, as is the case
146 for 'multi-organ system failure', or where the body has already been weakened through repeated
147 previous diseases or exposures. The same is true for the health consequences of climate change. It
148 acts as a threat multiplier, compounding many of the issues communities already face, and
149 strengthening the correlation between multiple health risks, making them more likely to occur
150 simultaneously. Indeed, it is not a 'single system disease', instead, often acting to compound existing
151 pressures on housing, food and water security, poverty, and many of the determinants of good
152 health. Adaptation has limits, and prevention is better than cure to prevent potentially irreversible
153 effects of climate change.

154 Progress in mitigating climate change since the signing of the UNFCCC has been limited across all
155 sectors, with only modest improvements in carbon emission reduction from electricity generation.
156 Whilst there are increasing levels of sustainable travel in Europe and some evidence of decline in
157 dependence on private motor vehicles in cities in the USA and Australia, the situation is generally
158 less favourable in cities in emerging economies (Indicator 3.7). This, and a slow transition away from
159 highly-polluting forms of electricity generation, has yielded a modest improvement in air pollution in
160 some urban centres. However, global population-weighted PM_{2.5} exposure has increased by 11.2%
161 since 1990 and some 71.2% of the 2971 cities in the WHO air pollution database exceed
162 recommendations of annual fine particulate matter exposure (Indicator 3.5). The strength and
163 coverage of carbon pricing covers only 13.1% of global anthropogenic CO₂ emissions, with the
164 weighted average carbon price of these instruments at 8.81USD/tCO₂e in 2017 (Indicator 4.7).
165 Furthermore, responses to climate change have yet to fully take advantage of the health co-benefits

166 of mitigation and adaptation interventions, with action taken to-date only yielding modest
167 improvements in human wellbeing. In part, this reflects a need for further evidence and research on
168 these ancillary effects and the cost-savings available. However, it also reflects a need for more
169 joined-up policymaking across health and non-health ministries of national governments.

170 This delayed mitigation response puts the world on a 'high-end' emissions trajectory, resulting in
171 global warming of between 2.6°C and 4.8°C of warming by the end of the century.

172

173 **The voice of the health profession is essential in driving forward progress on climate change and**
174 **realising the health benefits of this response.**

175 This report, and previous Lancet Commissions, have argued that the health profession has not just
176 the ability but the responsibility to act as public health advocates, communicating the threats and
177 opportunities to the public and policymakers, and ensuring climate change is understood as being
178 central to human wellbeing.

179 There is evidence of growing attention to health and climate change in the media and in academic
180 publications, with global newspaper coverage of the issue increasing 78% and the number of
181 scientific papers more than tripling, since 2007 (Indicator 5.1.1 and 5.2). However, despite these
182 positive examples, the 2017 indicators make it clear that further progress is urgently required.

183

184 **Whilst progress has historically been slow, the last five years have seen an accelerated response,**
185 **and the transition to low-carbon electricity generation now appears inevitable, suggesting the**
186 **beginning of a broader transformation. In 2017, momentum is building across a number of sectors,**
187 **and the direction of travel is set, with clear and unprecedented opportunities for public health.**

188 In 2015, the Lancet Commission made 10 recommendations to governments, to accelerate action
189 over the following five years. The Lancet Countdown's 2017 indicators track against these 2015
190 recommendations, with results suggesting that discernible progress has been made in many of these
191 areas, breathing life into previously stagnant mitigation and adaptation efforts. Alongside the Paris
192 Agreement, these provide reason to believe that a broader transformation is under way.

193 *Recommendation 1) Invest in climate change and public health research:* since 2007, the number of
194 scientific papers on health and climate change has more than trebled (Indicator 5.2).

195

196 *Recommendation 2) Scale-up financing for climate-resilient health systems:* spending on health
197 adaptation is currently at 4.63% (16.46 billion USD) of global adaptation spend; and in 2017, health
198 adaptation from global development and climate financing mechanisms is at an all-time high –
199 although absolute figures remain low (Indicators 4.9 and 4.10).

200

201 *Recommendation 3) Phase-out coal-fired power:* In 2015, more renewable energy capacity (150GW)
202 than fossil fuel capacity was added to the global energy mix. Overall, annual installed renewable
203 generation capacity (almost 2000 GW) exceeds that for coal, with about 80% of this recently added
204 renewable capacity located in China (Indicator 3.2). Whilst investment in coal capacity has increased
205 since 2006, in 2016 this turned and declined substantially (Indicator 4.1) and several countries have
206 now committed to phasing-out coal.

207

208 *Recommendation 4) Encourage a city-level low-carbon transition, reducing levels of urban pollution:*

209 Despite historically modest progress over the last two decades, the transport sector is approaching a
210 new threshold, with electric vehicles expected to reach cost-parity with their non-electric
211 counterparts by 2018 – a phenomenon that was not expected to occur until 2030 (Indicator 3.6).

212
213 *Recommendation 6) Rapidly expand access to renewable energy, unlocking the substantial economic*
214 *gains available from this transition:* Every year since 2015, more renewable energy has been added
215 to the global energy mix than all other sources, and in 2016, global employment in renewable energy
216 reached 9.8 million, over one million more than are employed in fossil fuel extraction. The transition
217 has become inevitable. However, in the same year, 1.2 billion people still did not have access to
218 electricity, with 2.7 billion people relying on the burning of unsafe and unsustainable solid fuels
219 (Indicators 3.3, 4.6 and 3.4).

220
221 *Recommendation 9) Agree and implement an international treaty which facilitates the transition to a*
222 *low-carbon economy:* In December 2015, 195 countries signed the Paris Agreement, which provides
223 a framework for enhanced mitigation and adaptation, and pledges to keep the global mean
224 temperature rise to “well below 2°C”. Going forward, a formal Health Work Programme within the
225 UNFCCC would provide a clear and essential entry point for health professionals at the national
226 level, ensuring that the implementation of the Paris Agreement maximises the health opportunities
227 for populations around the world.

228
229 Following the United States government’s announced intention to withdraw from the Paris
230 Agreement, the global community has demonstrated overwhelming support for enhanced action on
231 climate change, affirming clear political will and ambition to reach the treaty’s targets. The
232 mitigation and adaptation interventions committed to under the Paris Agreement have
233 overwhelmingly positive short- and long-term health benefits, but greater ambition is now essential.
234 Whilst progress has been historically slow, there is evidence of a recent turning point, with
235 transitions in sectors crucial to public health accelerating towards a low-carbon world. Whilst these
236 efforts must be greatly accelerated and sustained over the coming decades in order meet these
237 commitments, recent policy changes and the indicators presented here suggest that the direction of
238 travel is set.

239 From 2017 until 2030, the Lancet Countdown: Tracking Progress on Health and Climate Change will
240 continue its work, reporting annually on progress implementing the commitments of the Paris
241 Agreement, future commitments that build on them, and the health benefits that result.

242 Introduction

243 Climate change has serious implications for our health, wellbeing, livelihoods and the structure of
244 organised society. Its direct effects result from rising temperatures, and changes in the frequency
245 and strength of storms, floods, droughts, and heatwaves – with physical and mental health
246 consequences. Its impacts will also be mediated through less direct pathways, including changes in
247 crop yields, the burden and distribution of infectious disease, and in climate-induced population
248 displacement and violent conflict.¹⁻³ Whilst many of these effects are already being experienced,
249 their progression in the absence of climate change mitigation will greatly amplify existing global
250 health challenges and inequalities.⁴ It threatens to undermine many of the social, economic and
251 environmental drivers of health, which have contributed greatly to human progress.

252 Urgent and substantial climate change mitigation will help to protect human health from the worst
253 of these impacts, with a comprehensive and ambitious response to climate change potentially
254 transforming the health of the world's populations.⁴ The potential benefits and opportunities are
255 enormous, including cleaning up the air of polluted cities, delivering more nutritious diets, ensuring
256 energy, food and water security, and alleviating poverty and social and economic inequalities.

257 Monitoring this transition – from threat to opportunity – is the central role of the Lancet
258 Countdown: Tracking Progress on Health and Climate Change.⁵ The collaboration exists as a
259 partnership of 24 academic institutions from every continent, and brings together individuals with a
260 broad range of expertise across disciplines (including climate scientists, ecologists, mathematicians,
261 geographers, engineers, energy, food, and transport experts, economists, social and political
262 scientists, public health professionals, and physicians). The Lancet Countdown aims to track a series
263 of indicators of progress, publishing an annual 'health check', from now until 2030, on the state of
264 the climate, progress made in meeting global commitments under the Paris Agreement, and
265 adapting and mitigating to climate change (Panel 1). The initiative was formed following the 2015
266 Lancet Commission, which concluded that "tackling climate change could be the greatest global
267 health opportunity of the 21st century".⁴ It builds on, and reinforces, the work of the expanding
268 group of researchers, health practitioners, national governments, and the World Health Organization
269 (WHO), who are working to ensure that this opportunity becomes a reality.

270

271 Indicators of Progress on Health and Climate Change

272 In 2016, the Lancet Countdown proposed a set of potential indicators to be monitored, launching a
273 global consultation to define a conclusive set for 2017.⁵ A number of factors determined the
274 selection of indicators, including: (i) their relevance to public health, both in terms of the impacts of
275 climate change on health, and the health effects of the response to climate change; (ii) their
276 relevance to the main anthropogenic drivers of climate change; (iii) their geographical coverage and
277 relevance to a broad range of countries and income-groups; (iv) data availability; and (v) resource
278 and timing constraints. Table 1 divides these into broad themes, aligned with the global action
279 agenda on climate change and health, agreed at the Second WHO Global Conference on Health and
280 Climate, Paris, July 2016: climate change impacts, exposures, and vulnerabilities; adaptation
281 planning and resilience for health; mitigation actions and health co-benefits; economics and finance;
282 and public and political engagement.⁶

283 **Panel 1 Developing Lancet Countdown's Indicators: An Iterative and Open Process.**

284 The development of the Lancet Countdown's indicators took a pragmatic approach, taking in to
285 account the considerable limitations in data availability, resources, and time. Consequently, the

286 indicators presented here represent what is feasible for 2017 and will evolve over time in response
 287 to feedback and data improvements.

288 The purpose of this collaboration is to track progress on the links between public health and climate
 289 change, and yet, much of the data analysed here was originally collected for purposes not directly
 290 relevant to health. Initial analysis therefore principally captures changes in exposure, states, or
 291 processes, as proxies for health outcomes – the ultimate goal. Employing new methodologies to
 292 improve attribution to climate change is a particular priority. Subsequent reports will see the Lancet
 293 Countdown set 2030 targets for its indicators which align more directly with the Paris Agreement,
 294 allowing an assessment of its implementation over the course of the next 13 years.

295 The indicators presented thus far are the beginning of an ongoing, iterative and open process, which
 296 will work to continuously improve as capacity, data quality, and methods evolve. The objectives of
 297 the Lancet Countdown are both ambitious and essential, requiring support from a broad range of
 298 actors. To this end, the collaboration welcomes support from academic institutions and technical
 299 experts able to provide new analytical methods and novel data sets with appropriate geographical
 300 coverage. Appendix 1 provides a short overview of several parallel and complementary processes
 301 currently underway.

302 Throughout this report, the results and analysis of each indicator are presented alongside a brief
 303 description of the data sources and methods. A more complete account of each indicator can be
 304 found in the corresponding appendices. For a number of areas – such as the mental health impacts
 305 of climate change, or hydrological mapping of flood exposure – a robust methodology for an annual
 306 indicator has not been reported, reflecting the complexity of the topic and the paucity of data,
 307 rather than its lack of importance. Table 1 provides a summary of the 2017 indicators, with a more
 308 complete overview of these indicators provided in the supplementary online material. The thematic
 309 groups and indicator titles provide an overview of the domain being tracked, allowing for the growth
 310 and development of these metrics – for example, to more directly capture health outcomes – in
 311 subsequent years.

312

Thematic Group	Indicators	
1. Climate Change Impacts, Exposures and Vulnerability	1.1. Health effects of temperature change	
	1.2. Health effects of heatwaves	
	1.3. Change in labour capacity	
	1.4. Lethality of weather-related disasters	
	1.5. Global health trends in climate-sensitive diseases	
	1.6. Climate-sensitive infectious diseases	
	1.7. Food security and undernutrition	1.7.1. Vulnerability to undernutrition 1.7.2. Marine primary productivity
	1.8. Migration and population displacement	
2. Adaptation Planning and Resilience for Health	2.1. National adaptation plans for health	
	2.2. City-level climate change risk assessments	
	2.3. Detection and early warning of, preparedness for, and response to health emergencies	
	2.4. Climate information services for health	
	2.5. National assessment of vulnerability, impacts and adaptation for health	
	2.6. Climate-resilient health infrastructure	
3. Mitigation Actions and Health Co-Benefits	3.1. Carbon intensity of the energy system	
	3.2. Coal phase-out	
	3.3. Zero-carbon emission electricity	

	3.4. Access to clean energy	
	3.5. Exposure to ambient air pollution	3.5.1. Exposure to air pollution in cities
		3.5.2. Sectoral contributions to air pollution
		3.5.3. Premature mortality from ambient air pollution by sector
	3.6. Clean fuel use for transport	
	3.7. Sustainable travel infrastructure and uptake	
	3.8. Ruminant meat for human consumption	
	3.9. Healthcare sector emissions	
4. Economics and Finance	4.1. Investments in zero-carbon energy and energy efficiency	
	4.2. Investment in coal capacity	
	4.3. Funds divested from fossil fuels	
	4.4. Economic losses due to climate-related extreme events	
	4.5. Employment in low-carbon and high-carbon industries	
	4.6. Fossil fuel subsidies	
	4.7. Coverage and strength of carbon pricing	
	4.8. Use of carbon pricing revenues	
	4.9. Spending on adaptation for health and health-related activities	
	4.10. Health adaptation funding from global climate financing mechanisms	
5. Public and Political Engagement	5.1. Media coverage of health and climate change	5.1.1. Global newspaper reporting on health and climate change
		5.1.2. In-depth analysis of newspaper coverage on health and climate change
	5.2. Health and climate change in scientific journals	
	5.3. Health and climate change in the United Nations General Assembly	

313 Table 1 Thematic groups and indicators for the Lancet Countdown's 2017 report.

314

315

316 [Delivering the Paris Agreement for Better Health](#)

317 The Paris Agreement has been ratified at the national level by 153 of 197 parties to the UNFCCC, and
 318 currently covers 84.7% of greenhouse gas (GHG) emissions. It set out a commitment of ambitious
 319 GHG emissions reduction to limit climate change to well below a global average temperature rise of
 320 2°C above pre-industrial levels, with an aim to limit temperature increases to 1.5°C.⁷

321 Most countries (187) have committed to near-term GHG emission reduction actions up to 2030,
 322 through their Nationally Determined Contributions (NDCs). Article 4 paragraph 2 of the Paris
 323 Agreement states that each signatory “shall prepare, communicate and maintain successive
 324 nationally determined contributions that it intends to achieve”.⁷ However, the NDCs of the 153
 325 parties that have ratified the agreement currently fall short of the necessary reductions by 2030 to
 326 meet the 2°C pathway.⁸

327 The Lancet Countdown's indicators place national decisions within a broader context. They highlight
 328 the fact that globally, total power capacity of ‘pre-construction’ coal (commitments for new coal
 329 power plants) has halved from 2016 to 2017 alone; that every year since 2015, more renewable
 330 energy has been added to the global energy mix than all other sources combined; its installed costs
 331 continue to fall (with solar photovoltaic (PV) electricity generation now being cheaper than
 332 conventional fossil fuels in an ever growing number of countries); electric vehicles are poised to
 333 reach cost-parity with their petrol-based counterparts; and in 2016 global employment in renewable
 334 energy reached 9.8 million, over one million greater than that in fossil fuel extraction.

335 These positive examples in recent years must not mask the dangerous consequences of failing to
336 meet the Paris Agreement, the past two decades of relative inaction, the economies and sectors
337 currently lagging behind, and the enormity of the task ahead, which leave achieving the Agreement's
338 aims in a precarious position. Indeed, much of the data presented should serve as a wake-up call to
339 national governments, businesses, civil society, and the health profession.

340 However, as this report demonstrates, the world has already begun to embark on a path to a low-
341 carbon and healthier world. Whilst the pace of action must greatly accelerate, the direction of travel
342 is set.

343 1. Climate Change Impacts, Exposures and Vulnerability

344

345 Introduction

346 This section provides a set of indicators that track health impacts related to anthropogenic climate
347 change. Such impacts are dependent upon the nature and scale of the hazard, the extent and nature
348 of human exposure to them, and the underlying vulnerability of the exposed population.⁹ Thus,
349 these indicators aim to measure exposure to climatic hazards and vulnerabilities of people to them,
350 and over time, quantify the health impacts of climate change. These, in turn, inform protective
351 adaptation and mitigation interventions (sections two and three), the economic and financial tools
352 available to enable such responses (section four), and the public and political engagement that
353 facilitates them (section five).

354 Climate change affects human health primarily through three pathways: direct; ecosystem-
355 mediated; and human-institution-mediated.¹⁰ Direct effects are diverse, being mediated, for
356 instance, by increases in the frequency, intensity, and duration of extreme heat, and by rises in
357 average annual temperature experienced (leading to, for instance, increased heat-related mortality).
358 Rising incidence of other extremes of weather, such as flood and storms, increase the risk of
359 drowning and injury, damage to human settlements, the spread of water-borne disease, and mental
360 health sequelae.¹⁰ Ecosystem-mediated impacts include changes in the distribution and burden of
361 vector-borne diseases (such as malaria and dengue) and food and water-borne infectious disease.
362 Human undernutrition from crop failure, population displacement from sea-level rise, and
363 occupational health risks are examples of human-institution-mediated impacts.

364 Whilst the literature, and indeed some of the data presented here has traditionally focused on
365 impacts such as the spread of infectious diseases and mortality from extremes of weather, the
366 health effects from non-communicable diseases are just as important. Mediated through a variety of
367 pathways, they take the form of cardiovascular disease and acute and chronic respiratory disease
368 from worsening air pollution and aero-allergens, or the often-unseen mental health effects of
369 extreme weather events, or of population displacement.^{11,12} Indeed, emerging evidence is exploring
370 links between a rising incidence of chronic kidney disease, dehydration, and climate change.^{13,14}

371 Eight indicators were selected and developed for this section:

- 372 1.1 Health effects of temperature change
- 373 1.2 Health effects of heatwaves
- 374 1.3 Change in labour capacity
- 375 1.4 Lethality of weather-related disasters
- 376 1.5 Global health trends in climate-sensitive diseases
- 377 1.6 Exposure to climate-sensitive infectious diseases
- 378 1.7 Food security and undernutrition
- 379 1.8 Migration and population displacement

380

381 Appendix 2 provides a more detailed discussion on the data and methods used, as well as the
382 limitations and challenges encountered in the selection of each indicator. The indirect indicators (1.5
383 to 1.8) each provide a 'proof of concept', rather than being fully comprehensive, focusing variably on
384 a specific diseases, populations, or locations. Additionally, future iterations of the Lancet
385 Countdown's work will seek to capture indicators of the links between climate change and air
386 pollution, and with mental ill-health.

387 Indicator 1.1: Health effects of temperature change

388 **Headline Finding:** *People experience far more than the global mean temperature rise. Between 2000*
389 *and 2016, human exposure to warming was about 0.9°C - more than double the global area average*
390 *temperature rise over the same period.*


391 Rising temperatures can exacerbate existing health problems among populations and also introduce
392 new health threats (including cardiovascular disease and chronic kidney disease). The extent to
393 which human populations are exposed to this change, and thus the health implications of
394 temperature change, depend on the detailed spatial-temporal trends of population and temperature
395 over time.

396 Temperature anomalies were calculated relative to 1986 to 2008, from the European Research Area
397 (ERA) produced by the European Centre for Medium-Range Weather Forecasts (ECMWF).¹⁵ This
398 dataset uses climate reanalysis to give a description of recent climate, produced by combining
399 models with observations.¹⁶ The time series shown in Figure 1.1 are global mean temperatures
400 calculated from the gridded data, weighted by area (to avoid bias from measurements near the
401 poles) and by population (to show the number of people exposed); these are described as “area
402 weighted” and “exposure weighted”, respectively.

403 Changes in population were obtained per country and the data projected onto the gridded
404 population.¹⁷ Figure 1.1 shows area- (yellow lines) and exposure-weighted (blue lines) changes in
405 mean summer temperatures since 2000. Exposure-weighted warming from 2000 to 2016 (0.9°C) is
406 much higher than the area-weighted warming (0.4°C) over the same period. Hence, mean exposure
407 to warming is more than double the global warming since 2000.

408 The increase in exposure relative to the global average is driven partly by growing population
409 densities in India, parts of China and Sub-Saharan Africa. Accounting for population when assessing
410 temperature change provides a vital insight into how human wellbeing is likely to be affected by
411 temperature change, with the analysis here showing that temperature change where people are
412 living is much higher than average global warming. Details of the global distribution of this warming
413 can be found in Appendix 2.

414



415

416 Figure 1.1 Mean summer warming from 2000 to 2016 area weighted and exposure weighted, relative to the
 417 1986-2008 recent past average.

418

419 [Indicator 1.2: Health effects of heatwaves](#)


420 **Headline Finding:** *Between 2000 and 2016, the number of vulnerable people exposed to heatwave*
 421 *events has increased by approximately 125 million, with a record 175 million more people exposed to*
 422 *heatwaves in 2015.*

423 The health impacts of extremes of heat range from direct heat stress and heat stroke, through to
 424 exacerbations of pre-existing heart failure, and even an increased incidence of acute kidney injury
 425 resulting from dehydration in vulnerable populations. The elderly, children under the age of 12
 426 months, and people with chronic cardiovascular and renal disease are particularly sensitive to these
 427 changes.¹⁰


428 Here, a heatwave is defined as a period of more than 3 days where the minimum temperature is
 429 greater than the 99th percentile of the historical minima (1986-2008 average).¹⁸ This metric
 430 therefore focuses on periods of high night-time temperatures, which are critical in denying
 431 vulnerable people vital recuperation between hot days. Heatwave data were calculated against the
 432 historical period 1986-2008. The population for the exposure calculations was limited to people over
 433 the age of 65 (as this age group is most vulnerable to the health impacts of heatwaves), which was
 434 obtained on a per-country basis from the UN World Population Prospects archives for each year
 435 considered.

436 Figure 1.2 shows the increase in total exposure to heatwaves over the 2000-2016 period (one
 437 heatwave experienced by one person). In 2015, the highest number of exposure events was
 438 recorded, with approximately 175 million additional people exposed to heatwaves. Figure 1.3 shows
 439 how the mean number of heatwave days experienced by people during any one heatwave
 440 (exposure-weighted) increases at a much faster rate than the global mean (area-weighted) number

441 of heatwave days per heatwave; this is due to high populations densities in areas where heatwaves
442 have occurred.
443



444
445 Figure 1.2 The change in exposure (in people aged over 65 years) to heatwaves from 2000 to 2016, relative to
446 the heatwave exposure average from 1986-2008.
447



448

449 Figure 1.3 The area and exposure weighted change in mean heatwave lengths globally from 2000 to 2016 (in
 450 people aged over 65 years), relative to the 1986-2008 recent past average.


451

452 **Indicator 1.3: Change in labour capacity**

453 **Headline Finding:** Global labour capacity in populations exposed to temperature change is estimated
 454 to have decreased by 5.3% from 2000 to 2016.

455 Higher temperatures pose significant threats to occupational health and labour productivity,
 456 particularly for those undertaking manual labour outside in hot areas. This indicator shows the
 457 change in labour capacity (and thus productivity) globally and specifically for rural regions, weighted
 458 by population (see Appendix 2 for details). Reductions in labour capacity have important
 459 implications for the livelihoods of individuals, families, and communities, with particular impacts on
 460 those relying on subsistence farming.


461 Labour capacity was estimated in the manner documented by Watts et al. (2015), based on wet bulb
 462 globe temperatures.⁴ Figure 1.4 shows the estimated change in outdoor labour productivity
 463 represented as a percentage relative to the reference period (1986-2008), with 0% implying no
 464 change. Labour capacity is estimated to have decreased by 5.3% between 2000 and 2016, with a
 465 dramatic decrease of over 2% between 2015 and 2016. Although there are some peaks of increased
 466 labour capacity (notably 2000, 2004 and 2008), the overwhelming trend is one of reduced capacity
 467 (Figure 1.4). These effects are most notable in some of the most vulnerable countries in the world
 468 (Figure 1.5).



469

470 Figure 1.4 The exposure weighted labour capacity change (%) globally from 2000 to 2016, relative to the recent
 471 past (1986-2008) average

472



473

474 Figure 1.5 Map of the change in labour capacity loss from 2000 to 2016, relative to the recent past (1986-2008)
 475 average.

476

477 This indicator currently only captures the effects of heat on rural labour capacity. The Lancet
 478 Countdown will work to expand this metric in the future to capture impacts on labour capacity in
 479 other sectors, including manufacturing, construction, transportation, tourism and agriculture.
 480 Through collaboration with HEAT-SHIELD, the Lancet Countdown will work to develop this process
 481 going forward, providing more detailed analysis of labour capacity loss and the health implications of
 482 heat and heatwaves, globally.^{19,20}


483 Indicator 1.4: Lethality of weather-related disasters

484 **Headline Finding:** *Despite a 46% increase in annual weather-related disasters from 2007 to 2016,*
485 *compared with the 1990-1999 average, there has been no accompanying increase in the number of*
486 *deaths, nor in those affected by disasters, nor in the ratio of these two outcomes.*


487 Weather-related events have been associated with over 90% of all disasters worldwide over the last
488 twenty years. As expected, considering its population and area, the continent most affected by
489 weather-related disasters is Asia, with some 2,843 events between 1990-2016 affecting 4.8 billion
490 people and killing 505,013. Deaths from natural hazard-related disasters are largely concentrated in
491 poorer countries.²¹ Crucially, this must be understood in the context of potentially overwhelming
492 health impacts of future climate change, worsening significantly over the coming years. Indeed, the
493 2015 Lancet Commission estimated an additional 1.4 billion drought exposure events, and 2.3 billion
494 flood exposure events occurring by the end of the century – demonstrating clear public health limits
495 to adaptation.⁴

496 Disaster impact is a function of hazard and vulnerability, with vulnerability from a climate change
497 perspective sometimes defined as a function of exposure, sensitivity, and adaptive capacity.²² This
498 indicator measures the ratio of the number of deaths, to the number of people affected by weather-
499 related disasters. Weather-related disasters included are: droughts, floods, extreme temperature
500 events, storms and wildfires. The health impacts of weather-related disasters expand beyond
501 mortality alone, including injuries, mental health impacts, spread of disease, and food and water
502 insecurity. Data for the calculations for this indicator come from the Emergency Events Database
503 (EM-DAT).^{23,24} Here, in line with the EM-DAT data used for analysis, a disaster is defined as either: 1)
504 10 or more people reported killed, 2) 100 or more people affected, 3) a declaration of a state of
505 emergency, or 4) a call for international assistance.

506 Between 1994 and 2013, the frequency of reported weather-related events (mainly floods and
507 storms) increased significantly. However, this trend may be partially accounted for by information
508 systems having improved in the last 35 years, and statistical data are now more available as a result
509 of increased socio-cultural sensitivity to disaster consequences and occurrence.²⁵ From 2007 to
510 2016, EM-DAT recorded an average of 306 weather-related disasters per annum, up 46% from the
511 1990-1999 average. However, owing to impressive poverty reduction and health adaptation efforts,
512 this has not yet been accompanied by any discernible trend in number of deaths, nor in those
513 affected by disasters, nor in the ratio of these two (Figure 1.6a). Indeed, separating out the disasters
514 by the type of climate and weather hazard associated with the disaster (Figure 1.6b) shows there has
515 been a statistically significant global decrease in the numbers affected by floods, equating to a
516 decrease of 3 million people annually. Importantly, best available estimates and projections expect a
517 sharp reversal in these trends over the coming decades, and it is notable that a number of countries
518 have experienced increases in deaths associated with weather-related disasters, with many of these
519 being high-income countries, illustrating that no country is immune to the impacts of climate change
520 (see Appendix 2 for more details).A



521 a)



522 b)

523 Figure 1.6 Deaths and people affected by weather-related disasters. 1.6a) Percentage change over time in the
 524 global number of deaths, the number of those affected, and the ratio of these (measured against 1990-2009).
 525 1.6b) Change over time in the number of people affected globally by different weather-related disasters.

526


527 The relative stability of the number of deaths in a disaster as a proportion of those affected, despite
 528 an increase in the number of disasters, could be interpreted in a number of ways. One plausible
 529 conclusion is that this represents an increase in health service provision and risk reduction. However,
 530 although weather-related disasters have increased in number over the past three to four decades,
 531 the data here does not capture the severity of such events – a factor directly relevant to a country’s
 532 vulnerability and ability to adapt.²²It is also important to note the difficulties in discerning overall
 533 trends, owing to the stochastic nature of the data and the relatively short time series. This poses

534 limitation on the significance of findings that can be drawn from analysis to date. Improving the
535 validity of this indicator will be a focus going forward.

536 **Indicator 1.5: Global health trends in climate-sensitive diseases**

537 **Headline Finding:** Global health initiatives have overwhelmingly decreased deaths associated with
538 climate-sensitive diseases since 1990, owing to important economic and public health advances over
539 the last three decades.

540 Disease occurrence is determined by a complex composite of social and environmental conditions
541 and health service provision, all of which vary geographically. Nonetheless, some diseases are
542 particularly sensitive to variations in climate and weather, and may thus be expected to vary with
543 both longer-term climate change and shorter-term extreme weather events.¹⁰ This indicator draws
544 from Global Burden of Disease (GBD) mortality estimates to show trends in deaths associated with
545 seven climate-sensitive diseases since 1990 (Figure 1.7).²⁷



546 Figure 1.7 Trends in mortality from selected causes of death as estimated by the Global Burden of Disease
547 2015, for the period 1990 to 2015, by WHO region.²⁷ (Created using Global Burden of Disease, 2016 data).
548

549 The disease trends above reveal global increases in dengue mortality, particularly in the Asia-Pacific
550 and Latin America and Caribbean regions, with some peak years (including 1998) known to be
551 associated with El Niño conditions.²⁸ Beyond climate, likely drivers of dengue mortality include trade,
552 urbanization, global and local mobility and climate variability; the association between increased
553 dengue mortality and climate change is therefore complex.²⁹ It naturally follows that an increase
554 spread of the disease resulting from climate change will be a significant contributing factor in the
555 increased likelihood of an associated increase in mortality. Malignant melanoma is a distinctive
556 example of a non-communicable disease with a clear link to ultraviolet exposure, with mortality
557 increasing steadily despite advances in surveillance and treatment; although it is important to
558 recognise that increased exposures also occur as a result of changing lifestyles (for example, a rise in
559 sun tanning). Heat and cold exposure is a potentially important aspect of climate-influenced
560 mortality, although the underlying attribution of deaths to these causes in the estimates is

561 uncertain.³⁰⁻³⁵ Deaths directly related to forces of nature have been adjusted for the effects of the
562 most severe seismic events. Of the ten highest country-year mortality estimates due to forces of
563 nature, seven were directly due to specific seismic activity, and these have been discounted by
564 replacing with the same countries' force of nature mortality for the following year. The remaining
565 major peaks relate to three extreme weather events (Bangladesh cyclone of 1991, Venezuela floods
566 and mudslides of 1999 and Myanmar cyclone of 2008), which accounted for over 300,000 deaths.

567 Overall, the findings here highlight the effectiveness and success of global health initiatives since
568 1990, in largely reducing deaths associated with these diseases. Furthermore, these trends provide a
569 proxy for the global health profile of climate-sensitive diseases and thus to some degree, indication
570 of existing vulnerabilities and exposures to them.

571 [Indicator 1.6: Climate-sensitive infectious diseases](#)


572 **Headline Finding:** *Vectorial capacity for the transmission of dengue by the mosquito vectors *Aedes**
573 *aegypti and *Aedes albopictus* in regions where these vectors are currently present has increased*
574 *globally due to climate trends by an average of 3% and 5.9%, respectively, compared to 1990 levels,*
575 *and by 9.4% and 11.1%, respectively, compared to 1950s levels.*

576 Despite a declining overall trend, infectious diseases still account for around 20% of the global
577 burden of disease and underpin more than 80% of international health hazards as classified by the
578 World Health Organization (WHO).^{36,37} Climatic factors are routinely implicated in the epidemiology
579 of infectious diseases, and they often interact with other factors, including behavioural,
580 demographic, socio-economic, topographic and other environmental factors, to influence infectious
581 disease emergence, distribution, incidence and burden.^{2,38} Understanding the contribution of
582 climate change to infectious disease risk is thus complex, but necessary for advancing climate
583 change mitigation and adaptation policies.¹⁴ This indicator is split into two components: a systematic
584 literature review of the links between climate change and infectious diseases, and a vectorial
585 capacity model for the transmission of dengue virus by the climate-sensitive vectors.

586 For the first component, a systematic review of the climate change infectious disease literature was
587 performed (see Appendix 2 for details), in which trends in the evolution of knowledge and direction
588 of impact of climate change disease risk associations were measured (Figure 1.8). The number of
589 new publications fitting the search criteria in 2016 (n=89) was the highest yet reported, almost
590 double the number published in 2015 (n=50) and more than triple the number published in 2014
591 (n=25) (Figure 1.8, left). Over this period, the complexity of interactions between climate change and
592 infectious disease has been increasingly recognised and understood (Figure 1.8, right).

593

594




595

596 Figure 1.8 Left: Academic publications reporting climate-sensitive infectious diseases by year. Right: proportion
597 of responses reported in publications by year and direction of impact.

598

599 Trends in the global potential for dengue virus transmission (as represented by vectorial capacity
600 (VC) in the mosquito vectors *Aedes aegypti* and *Aedes albopictus*) are presented. VC is “the rate
601 (usually daily) at which a bloodsucking insect population generates new inoculations from a
602 currently infectious case”.³⁹ A global, mechanistic investigation was conducted of changes in annual
603 transmission potential for a model, high burden, climate-sensitive vector-borne disease, dengue
604 fever (Figure 1.9). For both vectors, VC in locations where these vectors are currently present
605 reached its highest or equal highest average level in 2015 over the period considered (Figure 1.9,
606 bottom panel). This consolidates a clear and significant increase in VC starting in the late 1970s
607 (+3.0% and +6.0% compared to 1990 levels for *A. aegypti* and *A. albopictus*, respectively). Nearly all
608 *Aedes*-positive countries showed relative increases in VC for both vectors over the period considered
609 (Figure 1.9, top panel). Annual numbers of cases of dengue have doubled every decade since 1990,
610 with 58.4 million (23.6 million–121.9 million) apparent cases in 2013, accounting for over 10,000
611 deaths and 1.14 million (0.73 million–1.98 million) disability-adjusted life-years.⁴⁰ Climate change has
612 been suggested as one potential contributor to this increase in burden.⁴¹ *Aedes aegypti* and *Aedes*
613 *albopictus*, the principal vectors of dengue, also carry other important emerging or re-emerging
614 arboviruses, including Yellow Fever, Chikungunya, Mayaro and Zika viruses, which are likely similarly
615 responsive to climate change.



616

617 Figure 1.9 Average annual vectorial capacity (VC) for dengue in *Aedes aegypti* and *Aedes albopictus* for
 618 selected *Aedes*-positive countries (countries with *Aedes* present) (top panel; matrix coloured relative to
 619 country mean 1950-2015; red = relatively higher VC, blue = relatively lower VC; countries ordered by centroid
 620 latitude (north to south)). Bottom panel: average vectorial capacity (VC) for both vectors calculated globally
 621 (results shown relative to 1990 baseline).

622

623 [Indicator 1.7: Food security and undernutrition](#)

624 Isolating the impact of climate change on health through the indirect impacts on food security is
 625 complicated, as policies, institutions, and the actions of individuals, organisations, and countries,
 626 strongly influence the extent to which food systems are resilient to climate hazards or can adapt to
 627 climate change, and whether individual households are able to access and afford sufficient nutritious
 628 food. For example, with respect to undernourishment, vulnerability has been shown to be more
 629 dependent on adaptive capacity (such as infrastructure and markets) and sensitivity (such as forest
 630 cover and rain-fed agriculture) than exposure (such as temperature change, droughts, floods,
 631 storms).⁴² Given the role of human systems in mediating the links between climate, food, and health,
 632 the chosen indicators focus on abiotic and biotic indicators and current population vulnerabilities,
 633 considering both terrestrial and marine ecosystems. Undernutrition has been identified as the
 634 largest health impact of climate change in the 21st century.^{10,43-46}

635

636 Indicator 1.7.1: Vulnerability to undernutrition

637 **Headline Finding:** *The number of undernourished people in the 30 countries located in Africa and*
638 *Southern Asia with the highest prevalence (>15%) has increased from 398 million in 1990 to 422*
639 *million in 2016. These are countries located in regions which are highly dependent on regional*
640 *production for their food needs and where climate change is predicted to have the greatest negative*
641 *impact on yields.*


642 The purpose of this indicator is to track the extent to which health will be compromised by climate
643 change in countries where both current dependence on domestic production of food, and current
644 level of undernourishment (which is strongly related to undernutrition) is already high. Climate
645 change could further compromise health through changes in localised temperature and
646 precipitation, manifested in falling yields.

647 Food markets are increasingly globalised, and food security is increasingly driven by human systems.
648 In response to falling yields caused by temperature increases, governments, communities, and
649 organisations can and will undertake adaptation activities that might variously include breeding
650 programmes, expansion of farmland, increased irrigation, or switching crops. However, the greater
651 the loss of yield potential due to temperature increases, the more difficult adaptation becomes for
652 populations dependent upon domestic food supply.

653 Rising temperatures have been shown to reduce global wheat production, which has been estimated
654 to fall 6% for each degree Celsius of additional temperature increase.⁴⁷⁻⁴⁹ Rice yields are sensitive to
655 higher night temperatures, with each 1°C increase in growing-season minimum temperature in the
656 dry season resulting in a fall in rice grain yield of 10%.⁵⁰ Higher temperatures have been
657 demonstrated rigorously to have a negative impact on crop yields in lower-latitude countries.⁵¹⁻⁵³
658 Moreover, agriculture in lower-latitudes tends to be more marginal, and more people are food
659 insecure.

660 This indicator, using data from the Food and Agriculture Organization of the United Nations (FAO),
661 focuses on vulnerability to undernutrition.⁵⁴ Countries are selected for inclusion based on three
662 criteria: the presence of moderate or high level of undernourishment, reflecting vulnerability; their
663 physical location, focusing on geographies where a changing climate is predicted with high
664 confidence to have a negative impact on the yields to staples produced; and dependence on regional
665 production for at least half of its cereal consumption, reflecting high exposure to localised climate
666 hazards. Based on these criteria, 30 countries, all located in Africa or Southern Asia, are included.
667 Figure 1.10 presents the aggregated indicators, which shows the total number within the population
668 undernourished in these 30 countries, multiplied by total dependence on regional production of
669 grains. This gives a measure of how exposed already undernourished populations, who are highly
670 dependent on regionally produced grains, are to localized climate hazards.

671



672

673 Figure 1.10 Total number of undernourished people multiplied by regional dependency on grain production for
674 countries.

675 The regions with the highest vulnerability to undernutrition also coincide with areas where yield
676 losses due to warming are predicted to be relatively high, thus increasing the vulnerability of these
677 populations to the negative health consequences of undernutrition. High dependence on one crop
678 increases the vulnerability of individual countries further. For example, Kenya, which has a domestic
679 production dependency for cereals of almost 80%, 69% dependent on maize, is experiencing high
680 levels of undernutrition, and is particularly vulnerable to climate-related yield losses. Going forward,
681 these data will be refined through country-level exploration, incorporation of the predicted impact
682 of warming on yield losses, and incorporation of key temperature indicators such as 'growing degree
683 days' above critical crop-specific thresholds.^{55,56}

684

685 [Indicator 1.7.2: Marine primary productivity](#)

686 Declining fish consumption provides an indication of food insecurity, especially in local shoreline
687 communities dependent upon marine sources for food, and hence are especially vulnerable to any
688 declines in marine primary productivity affecting fish stocks.⁵⁷ This is particularly concerning for the
689 1 billion people around the world who rely on fish as their principal source of protein, placing them
690 at increased risk of stunting (prevented from growing or developing properly) and malnutrition from
691 food insecurity.⁵⁸ In addition, fish are important for providing micronutrients, such as zinc, iron,
692 vitamin A, vitamin B12, and Omega-3 fatty acids. If current fish declines continue, as many as 1.4
693 billion people are estimated to become deficient and at elevated risk of certain diseases, particularly
694 those associated with the cardiovascular system.^{59,60}

695 Marine primary productivity is determined by abiotic and biotic factors; measuring these globally
696 and identifying relevant marine basins is complex. Factors such as sea surface temperature (SST), sea
697 surface salinity (SSS), coral bleaching and phytoplankton numbers are key determinants of marine

698 primary productivity. Other local determinants have particularly strong influences on marine primary
699 productivity. For example, harmful algal blooms (HAB) occur as a result of uncontrolled algal growth
700 producing deadly toxins. The consumption of seafood contaminated with the toxins of harmful algal
701 blooms, such as those produced by *Alexandrium tamarense*, is often very dangerous to human
702 health, and potentially fatal.⁶¹

703 Changes in SST and SSS from 1985 to present, for twelve fishery locations essential for aquatic food
704 security are presented here. Data was obtained from NASA’s Earth Observatory Databank, and
705 mapped across to the significant basins outlined in Appendix 2. From 1985 to 2016, a 1°C increase in
706 SST (from an annual average of 22.74°C to 23.73°C) was recorded in these locations.⁶² This indicator
707 requires significant further work to draw out the attribution to climate change and the health
708 outcomes that may result. A case study on food security and fish stocks in the Persian Gulf is
709 presented in Appendix 2.

710

711 [Indicator 1.8: Migration and population displacement](#)

712 **Headline Finding:** *Climate change is the sole contributing factor for at least 4,400 people already*
713 *being forced to migrate, globally. The total number for which climate change is a significant or*
714 *deciding factor is significantly higher.*

715 Climate change-induced migration may occur through a variety of different social and political
716 pathways, ranging from sea level rise and coastal erosion, through to changes in extremes and
717 averages of precipitation and temperature decreasing the arability of land and exacerbating food
718 and water security issues. Estimates of future “climate change migrants” up to 2050 vary widely,
719 from 25 million to 1 billion.⁶³ Such variation indicates the complexity of the multi-factorial nature of
720 human migration, which depends on an interaction of local environmental, social, economic, and
721 political factors. For example, in Syria, many attribute the initial and continued conflict to the rural-
722 to-urban migration that resulted from a climate change-induced drought.^{64,65} However, the factors
723 leading to the violence are wide-ranging and complex, with clear quantifiable attribution particularly
724 challenging. Indeed, climate change is often thought of as playing an important role in exacerbating
725 the likelihood of conflict, and as a threat multiplier and an accelerant of instability. Nonetheless,
726 migration driven by climate change has potentially severe impacts on mental and physical health,
727 both directly and through the disruption of essential health and social services.⁶⁶

728 Despite the methodological difficulties in proving a direct causal relationship between climate
729 change and population displacement, there are areas where this is methodologically possible. This
730 indicator focuses on these situations, attempting to isolate instances (as exemplars) where climate
731 change is the sole contributory factor in migration decisions. Sea level rise provides the clearest
732 example of this, although other examples exist as shown in Table 1.1. Estimating the number of
733 people who have involuntarily migrated (both internally and internationally) as a result of climate
734 change alone helps overcome the complexity of accounting for other societal, economic and
735 environmental factors that also influence migration.

736 Based on data derived from peer-reviewed academic publications (see Appendix 2 for full details). A
737 minimum of 4,400 people have been forced to migrate due solely to climate change (Table 1.1). This
738 will be an underestimate, as it excludes cases where more than one factor may be contributing to a
739 migration decision – such as a combination of both climate-related sea level rise and coastal erosion
740 not associated with climate change (possibly such as the village of Vunidogola, relocated by the

741 Fijian Government in 2014 for such reasons, and the planned relocation of the Fijian village of
 742 Narikoso by 2018).⁶⁷⁻⁶⁹
 743

Location	Population	Citation	Notes on causes
Carteret Islands, PNG	1,200	Connell (2016) ⁷⁰ Strauss (2012) ⁷¹	Migrating due to sea-level rise
Alaska (need to migrate as soon as possible)*		Bronen and Chapin III (2013) ⁷² Shearer (2012) ⁷³	Migrating due to changing ice conditions leading to coastal erosion and due to permafrost melt, destabilising infrastructure
Kivalina	398-400		
Newtok	353		
Shaktoolik	214		
Shismaref	609		
Alaska (need to migrate gradually)*			
Allakaket	95		
Golovin	167		
Hughes	76		
Huslia	255		
Koyukuk	89		
Nulato	274		
Teller	256		
Unalakleet	724		
Isle de Jean Charles, Louisiana	25 homes		Coastal erosion, wetland loss, reduced accretion, barrier island erosion, subsidence, and saltwater intrusion were caused by dredging, dikes, levees, controlling the Mississippi River, and agricultural practices. Climate change is now bringing sea-level rise

744 Table 1.1 Locations migrating now due to only climate change. *The village names and populations are sourced
 745 from the US Government Accountability Office’s report, “Alaska Native Villages: Limited Progress Has Been
 746 Made on Relocating Villages Threatened by Flooding and Erosion”.⁷⁰⁻⁷³

747

748 Over the long-term, human exposure and vulnerability to ice sheet collapse is increasing, as the
 749 number of people living close to the coast and at elevations close to sea level are also increasing. In
 750 1990, 450 million people lived within 20 km of the coast and less than 20 metres above sea level.⁷⁴
 751 In 2000, 634 million (~10% of the global population), of whom 360 million are urban, lived below 10
 752 metres above sea level, (the highest vertical resolution investigated).⁷⁵ With 2000 as a baseline, the
 753 population living below 10 metres above sea level will rise from 634 million to 1,005-1,091 million by
 754 2050 and 830-1,184 million by 2100.⁷⁶ From 2100 and beyond, without mitigation and adaptation

755 interventions, over one billion people may need to migrate due to sea level rise caused by any ice
756 sheet collapse which occurs.^{76,77}

757 Whilst this indicator is not yet able to capture the true number of people being forced to migrate
758 due to climate change, that at least 4,400 people are already being forced to migrate as a result of
759 climate change only is concerning and demonstrates that there are limits to adaptation. The fact
760 that this is a significant underestimate further highlights the need to mitigate climate change and
761 improve the adaptive capacity of populations to reduce future forced migration. Significantly, only
762 instances of migration where climate change is isolated as the only factor are captured. Moving
763 forward, new approaches will be required to more accurately reflect the number of people forced to
764 migrate due to climate change, looking to capture situations where climate change plays an
765 important contributory role alongside other social and economic considerations.

766

767 Conclusion

768 Climate change impacts health through diverse direct and indirect mechanisms. The indicators
769 captured here provide an overview of a number of these effects, capturing exposure, impact, and
770 underlying vulnerabilities. Going forward, indicators will be developed to better measure direct
771 health outcome from climate change, in addition to exposure and vulnerabilities.

772 The indicators presented here will be continuously developed over time in order to more directly
773 capture mortality and morbidity outcomes from communicable and non-communicable diseases.
774 Indeed, work is already underway to produce new indicators to capture these concepts for
775 subsequent reports. Panel 1.1 and Appendix 2 describe one such ongoing process focused on mental
776 health and climate change.

777 Adaptation pathways can help to minimise some of the negative health impacts of global warming,
778 especially for the lower range of projected average temperature rises. However, there are powerful
779 limits to adaptation, and this section has drawn attention to the non-linearity and the spatial
780 distribution of the health impacts of climate change. The indicators presented here demonstrate
781 clearly that these impacts are being experienced across the world today, and provide a strong
782 imperative for both adaptation and mitigation interventions to protect and promote public health.

783

784 Panel 1.1 Mental Health and Climate Change

785 Measuring progress in the effects of climate change on mental health and wellbeing is difficult.
786 Whilst this is partly due to problems of attribution, the main measurement difficulty lies in the
787 inherently complicated nature of mental health, which embraces a diverse array of outcomes (for
788 instance, anxiety and mood disorders), many of which co-occur and all of which vary over contexts
789 and lifetimes. They are products of long and complex causal pathways, many of which can be traced
790 back to distal but potent root causes, such as famine, war and poverty, of which climate change is
791 both an example and an accelerator.⁷⁸

792 Mental health, with its inherent intricacy, is a field where *systems thinking* is likely to be particularly
793 valuable. A first step, therefore, in tracking progress on mental health and climate change is to build
794 a conceptual framework using systems thinking. Initial work in partnership with the University of
795 Sydney has begun to trace through the many direct and indirect causal pathways, in order to aid the
796 identification of indicators. A number of challenges (e.g. how to gather and interpret highly

797 subjective measures across cultures and income settings) are immediately apparent. Whilst further
798 work, and engagement with other partners will be required, potential indicators may focus on a
799 range of issues, including: national and local mental health emergency response capacity to climate-
800 related extreme events; the extent to which climate change is considered within national mental
801 health strategies; or the social and psychological impact of uninsured economic losses that result
802 from extreme weather events.

803 2. Adaptation Planning and Resilience for Health

804

805 Introduction

806

807 Climate change adaptation is defined by the IPCC as the “adjustment in natural or human systems in
808 response to actual or expected climatic stimuli or their effects, which moderates harm or exploits
809 beneficial opportunities”.⁸⁰ With respect to health, adaptation consists of efforts to reduce injury,
810 illness, disability, and suffering from climate-related causes. Resilience has been defined as “the
811 capacity of individuals, communities and systems to survive, adapt, and grow in the face of stress
812 and shocks, and even transform when conditions require it”.⁸¹ In the context of climate change and
813 health, resilience is an attribute of individuals, communities, and health care systems; resilience at
814 all levels can reduce adverse health outcomes of climate change and should be a goal of adaptation
815 planning.

816 Indicators of resilience and adaptation are challenging to identify. Resilience is related to
817 preparedness, response, resource management and coordination capacity, but it is not synonymous
818 with them. Understanding the current resilience of a population’s health and health systems
819 provides some indication of resilience to climate change, although direct indicators measuring this
820 have not yet been developed by the Lancet Countdown. The indicators presented here are
821 predominantly process-based, focusing on health adaptation planning, capacity, and response.
822 Whilst the underlying resilience of communities is present to some extent in all of the indicators in
823 this section, it is currently only captured directly for health systems, and hence most indicators that
824 follow will focus more specifically on health adaptation.

825

826 The indicators presented here are:

827 2.1 National adaptation plans for health

828 2.2 City-level climate change risk assessments

829 2.3 Detection and early warning of, preparedness for, and response to health emergencies

830 2.4 Climate information services for health

831 2.5 National assessment of vulnerability, impacts and adaptation for health

832 2.6 Climate-resilience health infrastructure

833

834 Corresponding Appendix 3 provides more detailed discussion of the data and methods used.

835

836 Indicator 2.1: National adaptation plans for health

837 **Headline finding:** *30 out of 40 responding countries have a national health adaptation plan or*
838 *strategy approved by the relevant national health authority.*

839 Effective national responses to climate risks require that the health sector identify strategic goals in
840 response to anticipated – and unanticipated – threats. A critical step in achieving these strategic
841 goals is the development of a national health adaptation plan, outlining priority actions, resource
842 requirements and a specific timeline and process for implementation. This indicator tracks the policy
843 commitments of national governments for health and climate change adaptation. Data are drawn
844 from the recent WHO Climate and Health Country Survey (Panel 2.1).

845 Of the 40 countries responding to this baseline survey, 30 reported having a national adaptation
846 strategy for health, approved by their Ministry of Health or relevant health authority (Figure 2.1).
847 This number includes countries with a health component of their National Adaptation Plan (NAPs),
848 which was established by the UNFCCC to help nations identify medium- and long-term adaptation
849 needs and develop and implement programmes to address those needs.⁸² There is a need for
850 caution in extrapolating the results to global level, as many of the respondent countries have
851 received support from WHO in developing and implementing their plans.^{83,84} Nonetheless, with 75%
852 of respondents in the survey having an approved national health adaptation plan there is evidence
853 of the recognition of the need to adapt to climate change. Countries with national health adaptation
854 plans are found across all regions and, perhaps most significantly, among some of the most
855 vulnerable countries across Africa, South East Asia and South America. In future iterations of the
856 survey, data will be gathered on the content and quality of these adaptation plans, their level of
857 implementation, the main priorities for health adaptation, internal monitoring and review processes,
858 and the level of funding available to support policy interventions.

859
860
861



862
863 Figure 2.1 Countries with national health climate adaptation strategies or plans.

864

865 **Panel 2.1: WHO-UNFCCC Climate and Health Country Profiles.**
866 The WHO-UNFCCC Climate and Health Country Profile Project forms the foundation of WHO's
867 national level provision of information, and monitoring of progress, in this field. The profiles,
868 developed in collaboration with ministries of health and other health determining sectors, support
869 evidence-based decision making to strengthen the climate resilience of health systems and promote

870 actions that improve health while reducing carbon emissions. In part, the data used in the
871 development of the climate and health country profiles is collected through a biennial WHO Climate
872 and Health Country Survey. Data from this survey is reported on for indicators 2.1, 2.5 and 2.6

873 The 2015 baseline survey findings for 40 responding nations are presented in this report (for a
874 complete list of country respondents, see Appendix 3). The findings include countries from all WHO
875 regions (high, middle and low income groups) and with varying levels of risks and vulnerabilities to
876 the health impacts of climate change. The 2015 survey data were validated as part of the national
877 consultation process seeking input on respective WHO UNFCCC Climate and Health Country Profiles
878 from key in-country stakeholders, including representatives of the Ministry of Health, Ministry of
879 Environment, meteorological services and WHO country and regional technical officers.

880 The validated data presented in this report tended to include a high number of countries that are
881 actively working on climate and health with WHO; as such, the results here are indicative and are
882 not meant to be inferred as an exact indicator of global status. The number of country respondents
883 is expected to double in subsequent iterations of the survey. As such, the results presented here
884 represent the beginning of the development of a more comprehensive survey, presenting results
885 available at the start of this process.

886

887 [Indicator 2.2: City-level climate change risk assessments](#)


888 **Headline Finding:** *Of the 449 self-reporting cities, 45% have climate change risk assessments in*
889 *place.*

890 Globally, 54.5% of people live in cities, where key health infrastructure is often concentrated.⁸⁵
891 These urban centres are increasingly at risk from climate change, with negative impacts predicted
892 for human health and health services. These risks require city-level responses to complement NAPs,
893 in order to improve cities' ability to adapt to climate change. Indeed, cities have a unique
894 opportunity to provide adaptation measures that help improve the resilience of urban populations,
895 whilst also helping mitigate the impacts of climate change on public health.⁸⁶

896 Data for this indicator comes from the 2016 global survey of the Compact of Mayors and the Carbon
897 Disclosure Project (CDP).^{87 88} Of the 449 cities with public responses (533 cities responded overall),
898 45% reported to "have undertaken a climate change risk or vulnerability assessment for [their] local
899 government" (Figure 2.2).⁸⁹

900 The highest number of cities with climate change risk assessments are in high income countries
901 (HICs) (118 cities), with only 42 cities in low-income countries. This partly reflects the fact that more
902 cities in HICs were surveyed, and partly the fact that these cities have a greater capacity to develop
903 such plans. There were a higher number of respondents from cities in HICs compared with low
904 income (236 versus 61).

905 European cities in this survey have the highest number of climate change risk assessments (56
906 cities), representing 83% of European cities surveyed. Conversely, only 28% of surveyed African cities
907 have climate change risk assessments. This has serious implications for the adaptive capacity of
908 some of the most vulnerable populations to climate change in low income countries. A concerted
909 effort must be made to increase the number of climate change risk assessment in cities in low-
910 income countries, in order to better understand their vulnerability to climate change impacts and
911 implement adaptation actions.



912

913 Figure 2.2 Number of global cities undertaking climate change risk assessments by a) income grouping, and b)
 914 WHO region.

915

916 **Indicator 2.3: Detection and early warning of, preparedness for, and response to climate**
 917 **related health emergencies**

918 **Headline Finding:** *Due to focused investment in the implementation of the International Health*
 919 *Regulations (2005), national capacities relevant to climate adaptation and resilience, including*
 920 *disease surveillance and early detection, multi-hazard public health emergency preparedness and*
 921 *response, and the associated human resources to perform these public health functions, have*
 922 *increased markedly from 2010 to 2016 in all world regions.*

923 Many initiatives at community, national, regional and global levels support strengthening country
 924 capacities for health emergency and disaster risk management and complement the implementation
 925 of the Sendai Framework for Disaster Risk Reduction, Sustainable Development Goal 3D, the Paris
 926 Agreement on Climate Change and the International Health Regulations (2005). Under the
 927 International Health Regulations (IHR (2005)), all States Parties should report to the World Health
 928 Assembly annually on the implementation of IHR (2005).^{91,92} In order to facilitate this process, WHO
 929 developed an IHR Monitoring questionnaire, interpreting the Core Capacity Requirements in Annex 1

930 of IHR (2005) into 20 indicators for 13 capacities (Panel 2.2). These metrics can serve as important
931 proxies of health system adaptive capacity and system resilience, since they measure the extent to
932 which health systems demonstrate a range of attributes necessary to detect, prepare for and
933 respond to public health emergencies, some of which are climate sensitive. Four capacities reflecting
934 seven indicators from IHR Monitoring questionnaire are reported here: surveillance, preparedness,
935 response, and human resources. Additional details of all four of these IHR Capacities can be found in
936 Appendix 3.

937 **Panel 2.2: The International Health Regulations (2005).**

938 The current IHR (2005), which entered into force in 2007, is legally binding on 196 States Parties,
939 including all WHO member states. It requires States Parties to detect, assess, notify and report, and
940 respond promptly and effectively to public health risks and public health emergencies of
941 international concern (IHR Article 5, 13) and to develop, strengthen and maintain the capacity to
942 perform these functions (IHR Article 5). Examples of required core capacities include national
943 legislation, policy and financing; public health surveillance; preparedness and response; risk
944 communication; human resources; and laboratory services. Under the International Health
945 Regulations (IHR (2005)), all States Parties should report to the World Health Assembly annually on
946 the implementation of IHR (2005). In order to facilitate this process, WHO developed an IHR
947 Monitoring questionnaire.⁹³ The method of estimation calculates the proportion/percentage of
948 attributes (a set of specific elements or functions that reflect the performance or development of a
949 specific indicator) reported to be in place in a country. Since 2010, 195 States Parties have submitted
950 self-reports at least once. Indicator 2.3 is drawn from the results of these questionnaires to which
951 129 of 196 States Parties responded in 2016.⁹⁴


952

953 The first of these capacities is human resources, which reflects a single indicator: ‘human resources
954 available to implement the International Health Regulations Core Capacities’. This is a useful proxy in
955 lieu of an indicator that looks at specific capacity for health adaptation to climate change (Figure
956 2.3a). In 2010, capacity scores ranged from 25% in Africa to 57% in Western Pacific. Human resource
957 capacity has improved markedly by 2016, where on the average the capacity score is 67% (with the
958 lowest score in the Africa region reporting 51% and the highest in the Western Pacific Region 89%).

959 Secondly, surveillance capacity, summarizes two indicators in the IHR questionnaire ‘Indicator-based
960 surveillance includes an early warning function for early detection of a public health event’, and
961 ‘Event-Based Surveillance is established and functioning’. This capacity score is used as a proxy for a
962 health system’s ability to anticipate and identify outbreaks and changing patterns of climate-
963 sensitive infectious diseases, such as zoonosis and food-related outbreaks. Globally, 129 reporting
964 States Parties scored 88% for this capacity in 2016 (Figure 2.3b). This proportion has increased
965 steadily since 2010 (average score of 63%), indicating that health systems have increasing capacity
966 for early detection of public health events.


967 Thirdly, preparedness capacity reflects ‘Multi-hazard National Public Health Emergency
968 Preparedness and Response Plan is developed and implemented’, comprised of the presence of a
969 plan, the implementation of the plan, and the ability for this plan to operate under unexpected
970 stress, and ‘priority public health risks and resources are mapped and utilized’. Of responding
971 countries, progress can be seen in all world regions from 49% in 2010 to a 2016 global average of
972 76% (Figure 4.3c).

973 Finally, response capacity, reflects the availability and functioning of public health emergency
 974 response mechanisms, and Infection Prevention and Control (IPC) at national and hospital levels.
 975 This capacity is an important proxy for the ability of the health system to mobilize effective
 976 responses when shocks or stresses are detected. All countries demonstrate between 73-91%
 977 response capacity in 2016, with notable progress seen in Africa between 2010 (47%) and 2016 (73%)
 978 (Figure 2.3d).



979 a)


IHR Core Capacities implementation status: Surveillance



980


b)

IHR Core Capacities implementation status: Preparedness



981

c)



982

d)

983 Figure 2.3: IHR capacity scores by WHO region. 2.3a) Human Resources capacity score. 2.3b) Surveillance
 984 capacity score. 2.3c) Preparedness capacity score. 2.3d) Response capacity score.

985 There are some limitations to considering these capacities. Most importantly, IHR survey responses
 986 are self-reported; although national-level external verification has begun it currently remains
 987 relatively limited. Additionally, these findings capture potential capacity – not action. Finally, the
 988 quality of surveillance for early detection and warning is not shown, nor is the impact of that
 989 surveillance on public health. Response systems have been inadequate in numerous public health
 990 emergencies and thus the presence of such plans is not a proxy for their effectiveness.

991


992 [Indicator 2.4: Climate information services for health](#)

993 **Headline Finding:** *Out of the 100 WHO Member States responding to the WMO Survey, 73% report*
 994 *providing climate information to the health sector in their country.*

995 This indicator measures the proportion of countries whose Meteorological and Hydrological services
 996 self-reported to the World Meteorological Organization (WMO), providing tailored climate
 997 information, products and services to their national public health sector.⁹⁵ Response rates for the
 998 2015 WMO survey were: 71% in the African region, 67% in the Eastern Mediterranean Region, 79%
 999 in the European Region, 81% in the Region of the Americas, 67% in the South-East Asia Region and
 1000 44% in the Western Pacific Region.

1001 Taking into account the total number of WHO members (respondent and non-respondent) per WHO
 1002 region, only between 14.8 % and 51.4% are known to provide climate information to the health
 1003 sector (Figure 2.4) and between 18% and 55% did not provide information.

1004



1005

1006 Figure 2.4: National Meteorological and Hydrological Services (NHMSs) of WHO member states reporting to
 1007 provide targeted/tailored climate information, products and services to the health sector.

1008 However, it is important to note that this sample is not representative of all countries (49% non-
 1009 response rate) and these are self-reported results. Crucially, this indicator does not capture the type
 1010 of climate products made available, quality of the data provided, the ways in which the health sector
 1011 makes use of this data (if at all), and whether the data is presented in a format and timely fashion
 1012 relevant to public health. Future WMO surveys will aim to provide greater insight to the specific
 1013 applications of climate information. See Appendix 3 for more information.

1014


1015 **Indicator 2.5: National assessments of climate change impacts, vulnerability, and adaptation**
 1016 **for health**

1017 **Headline Finding:** Over two thirds of responding countries report having conducted a national
 1018 assessment of climate change impacts, vulnerability, and adaptation for health.

1019 National assessments of climate change impacts, vulnerability, and adaptation for health allow
 1020 governments to understand more accurately the extent and magnitude of potential threats to health
 1021 from climate change, the effectiveness of current adaptation and mitigation policies and future
 1022 policy and programme requirements. Although national assessments may vary in scope between
 1023 countries, the number of countries that have conducted a national assessment of climate change
 1024 impacts, vulnerability, and adaptation for health is a key indicator to monitor the global availability
 1025 of information required for adequate management of health services, infrastructure and capacities
 1026 to address climate change. This indicator tracks the number of countries that have conducted
 1027 national assessments, based on responses to the 2015 WHO Climate and Health Country Survey
 1028 (Panel 2.1).

1029 Over two-thirds of countries sampled (27 out of 40) reported having conducted a national
 1030 assessment of impacts vulnerability, and adaptation for health (Figure 2.5). These countries cover all
 1031 regions and include countries that are particularly vulnerable; for instance, of the nine responding
 1032 countries in the South-East Asia Region, eight countries (Bangladesh, Bhutan, Indonesia, Maldives,
 1033 Nepal, Sri Lanka, Thailand and Timor-Leste) reported having national assessments of impacts,

1034 vulnerability, and adaptation for health. Increasing global coverage of countries with national
1035 vulnerability and adaptation assessments for health is the result of WHO's support to countries
1036 through projects and technical guidance.⁹⁶




1037
1038 Figure 2.5 Countries with national assessment of climate change impacts, vulnerability and adaptation for
1039 health.

1040
1041 **Indicator 2.6: Climate-resilient health infrastructure**
1042 **Headline Finding:** Only 40% (16 out of 40) of responding countries reported implementing activities
1043 to increase the climate resilience of their health infrastructure.

1044 Functioning health infrastructure is essential during emergencies. Climate-related events, such as
1045 severe storms and flooding, may compromise electrical and water supplies, interrupt supply chains,
1046 disable transportation links, and disrupt communications and IT networks, contributing to reduced
1047 capacity to provide medical care. This indicator measures efforts by countries to increase the climate
1048 resilience of health infrastructure. The climate resiliency of health infrastructure reflects the extent
1049 to which these systems can prepare for and adapt to changes in climate impacting the system. Data
1050 is drawn from the WHO Climate and Health Country Survey (Panel 2.1). Only 40% of countries (16
1051 out of 40) reported having taken measures to increase the climate resilience of their health
1052 infrastructure (Figure 2.6). These results suggest widespread vulnerability of health system
1053 infrastructure to climate change. For example, only two out of nine responding countries in the
1054 African Region report efforts to improve the climate resiliency of health infrastructure. Similar trends
1055 were found across other WHO regions.

1056



1057

1058 Figure 2.6 Countries taking measures to increase the climate resilience of health infrastructure.

1059

1060 This indicator does not capture the quality or effectiveness of efforts to build climate-resilient health
 1061 system infrastructure. Nonetheless, it highlights the importance of ensuring that countries work to
 1062 implement climate-resilient health infrastructure, as these findings suggest this is generally lacking.

1063

1064 **Conclusion**

1065 This section has presented indicators across a range of areas relevant to health adaptation and
 1066 resilience. It is clear that the public, and the health systems they depend upon, are ill-prepared to
 1067 manage the health impacts of climate change.

1068 In many cases, the data and methods available provide only a starting-point for an eventual suite of
 1069 indicators that capture health-specific adaptation, and include both process-and outcome-based
 1070 indicators. New indicators will also be required to better capture important indicators of resilience.

1071

1072

1073

1074 **3. Mitigation Actions and Health Co-Benefits**

1075

1076 Introduction

1077 Sections one and two have covered the health impacts of climate change, the adaptation available
1078 and currently being implemented, and the limits to this adaptation.¹⁰ This third section presents a
1079 series of indicators relevant to the near-term health co-benefits of climate mitigation policies.
1080 Accounting for this enables a more complete consideration of the total cost and benefits of such
1081 policies, and is essential in maximising the cumulative health benefits of climate change mitigation.

1082 The health co-benefits of meeting commitments under the Paris Agreement are potentially
1083 immense, reducing the burden of disease for many of the greatest global health challenges faced
1084 today and in the future.⁹⁷ The indicators presented in this section describe a clear and urgent need
1085 to increase the scope of mitigation ambition if the world is to keep global average temperatures
1086 “well below 2°C”.⁷

1087 Countries are accelerating their response to climate change, with Finland, the UK, China, France,
1088 Canada and the Netherlands making strong commitments to phase-out or dramatically reduce their
1089 dependence on coal.⁹⁸⁻¹⁰¹ By 2017, electric vehicles are poised to be cost-competitive with their
1090 petroleum equivalents, a phenomenon that was not expected until 2030. Globally, more renewable
1091 energy capacity is being built every year than all other sources combined.^{101,102} Consequently,
1092 renewable energy is now broadly cost-competitive with fossil fuels, with electricity from low-latitude
1093 solar PV being cheaper than natural gas.¹⁰¹⁻¹⁰³

1094

1095 Tracking the health co-benefits of climate change mitigation

1096 Meeting the Paris Agreement will require global GHG emissions to peak within the next few years
1097 and undergo rapid reduction thereafter, implying near-term actions and medium- and long-term
1098 cuts through country-level activities.⁸ Global CO₂ emissions from fossil fuels and industry were 36.3
1099 GtCO₂ in 2015 (60% higher than in 1990), while emissions from land use change – which is
1100 intrinsically difficult to estimate – was approximately 4.8 GtCO₂. In the same year, 41% of the total
1101 fossil fuel and industry emissions were estimated to come from coal, 34% from oil, 19% from gas,
1102 and 6% from cement.¹⁰⁴ In 2015, the largest emitters of CO₂ were China (29%), the USA (15%), the
1103 European Union’s (EU) 28 member states ((EU28); 10%) and India (6.3%). However, per capita
1104 emissions of CO₂ belie the disparity driven by consumption, with global mean emissions at 4.8 tCO₂
1105 per person per year compared to 16.8 in the USA, 7.7 in China, 7.0 in EU28, and 1.8 in India.¹⁰⁴

1106 The actions needed to embark on rapid decarbonisation include avoiding the ‘lock-in’ of carbon
1107 intensive infrastructure and energy systems, reducing the cost of ‘scaling-up’ low-carbon systems,
1108 minimising reliance on unproven technologies, and realising opportunities of near-term co-benefits
1109 for health, security, and the environment.⁸ These actions will need to also be cost-effective and
1110 supported by non-state actors and industry.

1111 Indicators in this section are broadly considered within the framework of Driving Force-Pressure-
1112 State-Exposure-Effect-Action (DPSEEA). The DPSEEA framework is recognized as being suitable for
1113 the development of environmental health indicators, and identification of entry points for policy
1114 intervention.¹⁰⁵ An adaptation of the framework for examination of the health co-benefits of climate
1115 change mitigation is explained in Appendix 4.

1116 Here, health co-benefit indicators are captured for four sectors: 1) energy, 2) transport, 3) food, and
1117 4) healthcare. Appendix 4 provides more detailed discussion of the data and methods used.

1118 Energy Supply and Demand Sectors

1119 Fossil fuel burning comprises the largest single source of GHG emissions globally, producing an
1120 estimated 72% of all GHG emissions resulting from human activities.^{106,107} The majority (66%) of
1121 these emissions arise in the energy sector from the production of thermal and electric power for
1122 consumption across a range of sectors including industry, commercial, residential and transport.

1123 To meet the climate change mitigation ambitions of the Paris Agreement, it is widely accepted that
1124 the energy system will need to largely complete the transition towards near zero-carbon emissions
1125 by, or soon after, 2050, and then to negative emissions in the latter part of the century.^{108,109} Recent
1126 analysis has framed the necessary action as a halving of CO₂ emissions every decade.¹¹⁰

1127 The potential short-term health benefits of such strategies are substantial, with significant
1128 improvements from a reduction in indoor and outdoor air pollution; more equitable access to
1129 reliable energy for health facilities and communities; and lower costs of basic energy services for
1130 heating, cooking, and lighting to support higher quality of life.

1131

1132 Indicator 3.1: Carbon intensity of the energy system


1133 **Headline Finding:** *Globally, the carbon intensity of total primary energy supply (TPES) has remained*
1134 *stable since 1990, between 55-56 tCO₂/TJ, reflecting the significant global challenge of energy*
1135 *system decarbonisation. This has occurred because countries, which have achieved a reduction in*
1136 *carbon intensity (USA, UK, Germany), have been offset by those which have increased the carbon*
1137 *intensity of their energy supply (India and China).*

1138 To achieve the 2°C target (at a 66% probability), the global energy sector must reduce CO₂ emissions
1139 to more than 70% below current levels by 2050. This means a large reduction in the carbon intensity
1140 of the global energy system, which can be measured as the tonnes of CO₂ for each unit of total
1141 primary energy supplied (tCO₂/TJ). TPES reflects the total amount of primary energy used in a
1142 specific country, accounting for the flow of energy imports and exports.¹¹¹ Commitments under the
1143 Paris Agreement should begin to lower the overall carbon intensity of TPES, with the aim of reducing
1144 to near-zero by 2050.

1145 Drawing on data from the International Energy Agency (IEA), this indicator shows that globally, since
1146 the 1990s, the carbon intensity of primary energy supply has remained between 55-56 tCO₂/TJ.¹¹²
1147 However, a 53% growth in energy demand over the period has meant that global CO₂ emissions have
1148 grown significantly. Rapidly, low and middle income countries (LMICs) have seen an increase in
1149 carbon intensity since the 1970s, driven by increased coal use (Figure 3.1). For example, India's TPES
1150 has almost tripled since 1980, with the share of coal in the mix doubling (from 22% to 44%). Over the
1151 same period, 1980-2014, a fourfold increase in China's TPES, combined with increasing carbon
1152 intensity due to the coal share of TPES increasing from 52% to 66%, has led to strong growth in
1153 emissions.

1154 High-income countries have seen carbon intensity fall since the 1970s (for example, the USA and
1155 Germany in Figure 3.1). This decrease has resulted from a move away from coal use in energy
1156 production and use, reduced heavy industrial output, and increased use of lower carbon fuels,
1157 notably moving from coal to natural gas in the power sector and the use of renewable energy.

1158



1159

1160 Figure 3.1 Carbon intensity of Total Primary Energy Supply (TPES) for selected countries, and total CO₂
 1161 emissions (shaded area against secondary y-axis), 1971-2013.

1162

1163 **Indicator 3.2: Coal phase-out**

1164 **Headline Finding:** Globally, total primary coal supply has increased from 92 EJ in 1990, to 160 EJ in
 1165 2015. However, the 2015 supply level represents a reduction from the high point of 164 EJ in 2013,
 1166 providing an encouraging indication that global coal consumption has peaked and is now in decline.

1167

1168 The primary means of reducing carbon intensity of the energy system within necessary timescales
 1169 will be the phase-out of coal. Worldwide, coal supplies 30% of energy use and is the source of 44%
 1170 of global CO₂ emissions. The dirtiest form of coal produces almost twice the carbon per unit of
 1171 primary energy than the least carbon intensive fossil fuel – natural gas.¹¹² Given that a large share of
 1172 coal is used for power generation, it is an important sector of focus, both to reduce CO₂ emissions
 1173 and mitigate a major source of air pollution.¹¹²

1174 This indicator of coal phase-out is the total primary coal supply (EJ) in the energy system (Figure 3.2),
 1175 which makes use of recent data from the IEA.¹¹²

1176 Globally, coal use has increased by just under 60% since 1990. This is due to strong growth in global
 1177 energy demand, and an increasing share of TPES coming from coal, rising from 26% to 29% between
 1178 1990 and 2014.¹¹² This growth has largely been driven by China's increasing use of coal in industry
 1179 and for electricity production, particularly in the 2000s (see East Asia trend in Figure 3.2). Crucially,
 1180 growth in coal use has plateaued and reduced since 2013, in large part due to a recognition of the
 1181 health effects of air pollution, slower growth and structural changes in China's economy, and a
 1182 slowing in energy sector expansion.¹¹³ India has also seen significant growth in coal use, with the
 1183 share of coal in TPES increasing from 31% in 1990 to 46% in 2015. The other large coal consuming


1184 regions are the USA and Europe. The USA has had a stable level of consumption since the 1990s, but
 1185 experienced a recent fall in use, particularly in energy production and use, due to the cost-
 1186 competitiveness of shale gas. Europe has seen a steady decline in coal use since the 1990s, again
 1187 through a move to gas in economies such as the UK, although this overall downward trend has
 1188 transitioned to a plateau in recent years.

1189 Today, China and India both have similar shares of electricity generate by coal, at around 75% of
 1190 total generation. Whilst this trend is plateauing in China, this is not observed in other parts of Asia,
 1191 and the rapidly-emerging economies of Indonesia, Vietnam, Malaysia, and the Philippines see strong
 1192 growth from coal.¹¹²

1193 Meeting the IEA’s 2°C pathway and the Paris Agreement requires that no new coal-fired plants be
 1194 built (beyond those with construction currently underway), with a complete phase-out of unabated
 1195 plants (not fitted with carbon capture and storage) occurring by 2040. Crucially, such a transition
 1196 may have started, with the amount of coal power capacity in pre-construction planning at 570
 1197 gigawatts (GW) in January 2017, compared to 1,090 GW in January 2016.¹¹⁴ There are a range of
 1198 reasons for this large reduction, including decreasing planned capacity expansion, a desire to tackle
 1199 air pollution, and active efforts to expand renewable investment.

1200

1201



1202

1203 Figure 3.2 Total primary coal supply by country or region, and globally (shaded area against secondary y-axis),
 1204 1990-2015.

1205

1206

1207 Indicator 3.3: Zero-carbon emission electricity

1208 **Headline Finding:** Globally, renewable electricity as a share of total generation has increased by over
1209 20% from 1990 to 2013. In 2015, renewable energy capacity added exceeded that of new fossil fuel
1210 capacity, with 80% of recently added global renewable energy capacity currently located in China.
1211 Where renewables displace fossil fuels, in particular coal, it represents the beginning of reductions in
1212 morbidity and mortality from air pollution, and a potentially remarkable success for global health.


1213 As coal is phased out of the energy system, in particular in electricity production, the rapid scaling up
1214 of zero-carbon energy production and use will be crucial. To remain on a 2°C pathway, renewables-
1215 based capacity additions will need to be sustained over the next 35 years, reaching 400 GW per year
1216 by 2050, which is two and a half times the current level. Critical renewable technologies for
1217 achieving this will be solar, wind and hydroelectric.

1218 Indicator 3.3 draws on IEA data, and considers both renewable and other zero-carbon electricity.¹¹²
1219 Conversely, renewable energy refers to “all forms of energy produced from renewable sources in a
1220 sustainable manner, which include: bioenergy, geothermal, hydropower, ocean energy (tidal, wave,
1221 thermal), solar energy and wind energy”.¹¹⁵ By comparison, zero-carbon energy means no GHG
1222 emissions (i.e. zero-carbon and carbon equivalent) at the point of energy production and use, which
1223 therefore also includes nuclear-powered electricity, but excludes biomass.


1224 Both displace the use of fossil fuels (although notably fossil capacity tends to have annual higher
1225 load factors than renewables), reducing air pollution and GHG emissions, and so are important
1226 indicators for climate change and for health. One caveat is that the combustion of solid biomass
1227 fuels such as wood, sometimes promoted for climate change mitigation purposes, may increase fine
1228 particulate air pollution exposure and may not be carbon-neutral.¹¹⁶

1229 As a share of total generation, renewable energy has increased by over 20% from 1990 to 2013.
1230 Renewable energy continues to grow rapidly, mainly from increasing wind and solar PV investment,
1231 most notably in the USA, China and Europe (Figure 3.3). In 2015, more renewable energy capacity
1232 (150GW) was added than fossil fuel plant capacity added globally. Overall, there is now more added
1233 renewable generation capacity installed globally (almost 2000 GW) than coal, with about 80% of this
1234 newly installed capacity located in China.¹¹²

1235



1236



1237

1238 Figure 3.3 Renewable and zero-carbon emission electricity generation a) Share of electricity generated from
 1239 zero carbon sources; b) Electricity generated from zero carbon sources, TWh; c) Share of electricity generated
 1240 from renewable sources (excluding hydro); d) Electricity generated from renewable sources (excl. hydro), TWh.

1241


1242 **Indicator 3.4: Access to clean energy**

1243 **Headline Finding:** In 2016, it was reported that 1.2 billion people did not have access to electricity,
 1244 with 2.7 billion people relying on the burning of unsafe, unsustainable, and inefficient solid fuels.

1245 Increased access to clean fuels and clean energy technologies will have the dual benefit of reducing
 1246 indoor air pollution exposure, and reducing GHG emissions by displacing fossil fuels.¹¹⁷ The use of
 1247 clean energy for heating, cooling, cooking and lighting plays an important role in improving global
 1248 health and wellbeing, economic productivity, and reducing the risk of harm from living in energy
 1249 poverty.¹¹⁸

1250 It is estimated that globally, 1.2 billion people do not currently have access to electricity and 2.7
 1251 billion people rely on burning unsustainable and inefficient solid fuels, which contributes to poor
 1252 indoor air quality (see Panel 3.1), estimated to result in 4.3 million premature deaths related to
 1253 pneumonia, stroke, lung cancer, heart disease, and chronic obstructive pulmonary disease (COPD)
 1254 each year.^{119,120} Access to electricity, an energy source that emits no direct airborne particles
 1255 (though particles may be emitted indirectly through the fuel used to generate the electrical power),
 1256 is currently 85.3% globally but varies widely among countries and urban and rural settings.

1257 This indicator draws on and aligns with the proposed Sustainable Development Goal (SDG) indicator
 1258 7.1.2, defining ‘clean energy’ in terms of emission rate targets and specific fuel recommendations
 1259 (i.e. against unprocessed coal and kerosene) included in the WHO normative guidance.¹²¹ It
 1260 estimates the proportion of the population who primarily rely on clean fuels (including liquefied
 1261 petroleum gas, which, while still a fossil fuel, is cleaner than many solid fuels) and technologies for
 1262 cooking, heating and lighting compared to all people accessing those services. The data used for this
 1263 indicator comes from estimates of fuel use from WHO household survey data from roughly 800
 1264 nationally representative surveys and censuses, and is modelled to estimate the proportion of their
 1265 reliance on clean fuels (Figure 3.4).¹²²



1266
 1267 Figure 3.4 Proportion of population relying primarily on clean fuels and technology.

1268

1269 **Indicator 3.5: Exposure to ambient air pollution**

1270 **Headline Finding:** 71% of the 2,971 cities in the WHO’s database do not satisfy WHO annual fine
 1271 particulate matter exposure recommendations.

1272 Air pollutants directly harmful to health are emitted by combustion processes that also contribute to
 1273 emissions of GHGs. As such, properly designed actions to reduce GHG emissions will lead to
 1274 improvements in ambient air quality, with associated benefits for human wellbeing.¹²³ Current
 1275 estimates suggest that global population-weighted fine particulate matter (PM_{2.5}) exposure has
 1276 increased by 11.2% since 1990.^{123,124} To represent levels of exposure to air pollution, this indicator
 1277 collects information on annual average urban background concentrations of PM_{2.5} in urban settings
 1278 across the world.

1279

1280 **3.5.1: Exposure to air pollution in cities**


1281 The data for this indicator makes use of the WHO’s Urban Ambient Air Pollution Database, which
 1282 compiles information from a range of public sources, including national and subnational reports and
 1283 websites, regional networks, intergovernmental agencies, and academic publications.¹²⁵ The air
 1284 pollution measurements are taken from monitoring stations located in urban background,

1285 residential, commercial, and mixed areas. The annual average density of emission sources in urban
1286 areas and the proximity of populations to those sources led the Lancet Countdown to focus on
1287 exposure in cities.

1288 For this indicator, the Lancet Countdown has combined the WHO database with the Sustainable
1289 Healthy Urban Environments (SHUE) database, presenting data on 246 randomly sampled cities
1290 across the world (stratified by national wealth, population size, and Bailey's Ecoregion) (Figure
1291 3.5).¹²⁶

1292

1293



1294

1295 Figure 3.5 Annual mean PM_{2.5} concentration vs per capita GDP for 246 cities in the SHUE database. Colours
1296 indicate WHO regions: blue – Africa; red – Europe; green – the Americas; Lime – Eastern Mediterranean;
1297 orange – Western Pacific; purple – South East Asia. The dotted line marks the WHO recommended guidance
1298 level of 10 $\mu\text{g}\cdot\text{m}^{-3}$.

1299

1300 PM_{2.5} levels in the majority of global cities are currently well above the WHO's annual guideline level
1301 of 10 $\mu\text{g}\cdot\text{m}^{-3}$, with particularly high levels in cities in central, South and East Asia. Of almost 3,000
1302 cities in the WHO database, levels in 71.2% are above the guideline level. However, since monitoring
1303 is more common in high income settings, this is likely to represent an underestimation; for
1304 randomly-selected cities in the SHUE database, 87.3% of cities are above the guideline. The data
1305 suggests that air pollution levels have generally decreased in high income settings over recent
1306 decades, although it has marginally increased, globally.¹²⁷

1307

Panel 3.1. Energy and Household Air Pollution in Peru.

1308 Universal access to energy is a major challenge in most LMICs and access to clean energy or energy
 1309 sources that do not adversely affect health is a considerable problem. In Peru, low-income families
 1310 spend a higher percentage (5%-18%) of average monthly income on energy services than those with
 1311 higher-incomes.¹²⁸ Furthermore, a large portion of Peru’s rural population (83%) use firewood, dung,
 1312 or coal for cooking, making indoor air pollution one of the main environmental risk factors
 1313 experienced.¹²⁹


1314 Since the 1990s, the Peruvian government and various NGOs have promoted programmes and
 1315 policies oriented towards addressing the problem of solid fuels’ use for lighting, cooking and heating
 1316 and lack of access to energy sources in low-income sectors. In 2009, legislative changes enabled sub-
 1317 national governments to invest up to 2.5% of the national mining revenues in improved cook stove
 1318 (ICS) deployment, resulting in more than 280,000 ICS installed nationwide (52% public and 43%
 1319 private) as part of the multi-sectorial campaign “Half Million ICS for a Smokeless Peru”. This
 1320 campaign to help improve quality of life and health through the instalment of certified ICS.
 1321 Studies show that well-kept and certified ICS can reduce personal exposure to particulate matter
 1322 (PM_{2.5}).

1323 Peru released its 2010-2040 National Energy Policy in 2010. Of the nine goals, two discuss access to
 1324 energy services to low-income sectors. Special programmes have been developed in rural high
 1325 altitude and Amazonian regions in Peru to address energy access issues. In 2012, programmes were
 1326 established to substitute kerosene and other contaminating stoves with liquefied petroleum gas
 1327 (LPG) and ICS; and the Social Inclusion Energy Fund (FISE) was established, promoting access to LPG
 1328 for the most vulnerable populations through subsidies. By 2015, according to FISE, more than 1.3
 1329 million families had received an LPG stove, mitigating 91% of their CO₂ emissions and leading to a
 1330 corresponding reduction of 553,000 tons of CO₂ in using cleaner sources of energy.^{130,131}

1331

1332 3.5.2: Sectoral contributions to air pollution

1333 The energy sector –both production and use - is the single largest source of man-made air pollution
 1334 emissions, producing 85% of particulate matter and almost all of the sulphur oxides and nitrogen
 1335 oxides emitted around the world (Figure 3.6).¹¹²



1336


1337 Figure 3.6 Selected primary air pollutants and their sources globally in 2015.¹¹² (Source: IEA, 2016)

1338


1339 Of this, coal power is responsible for three-quarters of the energy production and use sector's
 1340 Sulphur Dioxide (SO₂) emissions, 70% of its Nitrogen Oxide (NO_x) emissions and more than 90% of its
 1341 PM_{2.5} emissions.¹¹² However, over the past decade, these emissions have largely decoupled from
 1342 increases in coal-fired generation in several geographies, due to the introduction of emission
 1343 standards for coal power plants.^{132,133}

1344 In 2015, manufacturing and other industries (for example, refining and mining) were responsible for
 1345 about half of global energy-related emissions of SO₂ as well as 30% of both NO_x (28 Mt) and PM_{2.5}.¹¹²
 1346 Furthermore, transport was responsible for around half of all energy-related NO_x emissions in 2015
 1347 as well as 10% of PM_{2.5}. Within this sector, road vehicles were by far the largest source of the
 1348 sector's NO_x and PM_{2.5} emissions (58% and 73%, respectively), while the largest portion of SO₂
 1349 emissions came from shipping.¹¹² Trends in NO_x emissions from the transport sector (1990 to 2010)
 1350 are shown in Figure 3.7.

1351



1352 a) This map is without prejudice to the status of or sovereignty over any territory, to the delimitation of international frontiers and boundaries, and to the name of any territory, city or area.



1353 b)

1354 Figure 3.7 a) Energy related PM_{2.5} emissions in 2015 and b) NO_x emissions from transport from 1990-2010 by
 1355 region.¹¹² (Created using IEA, 2016 data)

1356


1357 **3.5.3: Premature mortality from ambient air pollution by sector**

1358 The extent to which emissions of different pollutants from different sectors contribute to ambient
 1359 PM_{2.5} levels depends on atmospheric processes, such as the dispersion of primary particles and the
 1360 formation of secondary aerosols from precursor emissions. Sources with low stack heights located
 1361 close to populations, such as household combustion for cooking and heating as well as road vehicles,
 1362 typically play a disproportionately larger role for total population exposure in relation to their
 1363 absolute emissions.

1364 Long-term exposure to ambient PM_{2.5} is associated with increased mortality and morbidity from
 1365 cardiovascular and pulmonary diseases.¹³⁴⁻¹³⁶ A recent WHO assessment estimated that ambient air
 1366 pollution (AAP) is responsible for roughly three million premature deaths worldwide every year.¹³⁷
 1367 As the sources of air pollution and greenhouse gases are overlapping in many cases, greenhouse gas
 1368 mitigation measures can have large co-benefits for human health.

1369 Figure 3.8 shows an attribution of estimated premature mortality from AAP to the sources of
 1370 pollution as calculated in the GAINS model for the year 2015 in a set of South and East Asian
 1371 countries, using emissions data as published by the IEA.¹³⁸ Here, the contributions of individual
 1372 source sectors to ambient PM_{2.5} concentrations have been calculated using linearized relationships
 1373 based on full atmospheric chemistry transport model simulations, and premature deaths are
 1374 calculated following the methodology used by the WHO and the GBD 2013 study.^{136,137}

1375 In some countries, such as China, North Korea and the Republic of Korea, agriculture is a large
 1376 contributor to premature deaths. Significant direct benefits for human health can therefore be
 1377 expected if these emission sources are addressed by climate policies. Significant benefits could also
 1378 be available if, for instance, coal fired power plants were replaced by wind and solar.
 1379 Replacement of household combustion of coal, for example in China, would result in health benefits
 1380 not only from ambient (outdoor) but also household (indoor) exposure to air pollution.



1381

1382 Figure 3.8 Health impacts of exposure to ambient PM_{2.5} in terms of annual premature deaths per million
1383 inhabitants in South and East Asian countries in 2015, broken down by key sources of pollution.

1384

1385

1386 Transport Sector

1387 Transportation systems – including road vehicles, rail, shipping, and aviation – are a key source of
1388 GHG emissions, contributing 14% of global emissions in 2010.^{111,112} In order to meet the 2°C target,
1389 the global transport sector must reduce its total GHG emissions by more than 20% below current
1390 levels, by 2050, and to be on a trajectory to zero carbon emissions in the second half of the
1391 century.¹³⁹ Compared to other energy demand sectors, key sub-sectors of transportation (urban
1392 personal and freight transport, long distance road transport, shipping, short haul aviation, and long
1393 haul aviation) are more difficult to decarbonise because of the high energy density of fossil fuels,
1394 thus emissions reductions targets are lower for transport than the energy sector as a whole.

1395 The transport sector is also a major source of air pollutants, including particulate matter, nitrogen
1396 oxides, sulphur dioxide, carbon monoxide, volatile organic compounds, and indirectly, ozone.
1397 Furthermore, exposure to air pollution from road transport is particularly challenging in cities where
1398 vehicles emit street-level air pollution. In turn, significant opportunities for health exist through the
1399 reduction of GHG emissions from transport systems, both in the near-term through cleaner air and
1400 increased physical activity, and the long-term through the mitigation of climate change.

1401

1402 Indicator 3.6: Clean fuel use for transport


1403 **Headline Finding:** *Global transport fuel use (TJ) has increased by almost 24% since 1990 on a per*
1404 *capita basis. While petrol and diesel continue to dominate, non-conventional fuels have been rapidly*
1405 *expanding, with more than 2 million electric vehicles being sold between 2010 and 2016.*

1406 Fuels used for transport produce more than half the nitrogen oxides emitted globally and a
1407 significant proportion of particulate matter.^{111,112} Switching to low-emission transport systems is an
1408 important component of climate change mitigation and will help to reduce concentrations of most
1409 ambient air pollutants. However, the transport sector's extremely high reliance on petroleum-based
1410 fuels makes this transition particularly challenging.

1411 This indicator focuses on monitoring global trends in levels of fuel efficiency, and on the transition
1412 away from the most polluting and carbon intensive transport fuels. More specifically, this indicator
1413 follows the metric of fuel use for transportation on a per capita basis (TJ/person) by type of fuel. To
1414 develop this indicator, the Lancet Countdown draws on transport fuel data from the IEA and
1415 population data from the World Bank.¹¹²

1416 While some transition away from carbon-intensive fuel use, towards increasing levels of fuel
1417 efficiency has occurred in select countries, transport is still heavily dominated by gasoline and diesel.
1418 Global transport fuel use has increased by almost 65% since 1970 on a per capita basis (Tj/person)
1419 (Figure 3.9). However, non-conventional fuels (for example, electricity, biofuels, and natural gas)
1420 have been rapidly gaining traction since the 2000s, with more than two million electric vehicles
1421 having been sold around the globe since 2010, mostly in the US, China, Japan and some European
1422 countries (Figure 3.10).¹⁴⁰ These figures remain modest when compared to the overall number of
1423 cars sold per year, 77 million in 2017, and the total global fleet of 1.2 billion cars.


1424



1425

1426 Figure 3.9 Per capita fuel use by type (TJ/person) for transport sector with all fuels.

1427



1428

1429 Figure 3.10 Cumulative Global Electric Vehicle Sales. Note: BEV is Battery Electric Vehicle and PHEV is Plug-in
1430 Hybrid Electric Vehicle.^{141,142} (Source: IEA, 2017)

1431

1432 Indicator 3.7: Sustainable travel infrastructure and uptake


1433 **Headline Finding:** Levels of sustainable travel appear to be increasing in many European cities, but
1434 cities in emerging economies are facing sustainable mobility challenges. While levels of private
1435 transport use remain high in many cities in the USA and Australia, evidence suggests that they are
1436 starting to decline.

1437 Global trends of population growth and increasing urbanization suggests that demand for mobility in
1438 urban areas will increase. Moving from private motorized transport to more sustainable modes of
1439 travel (such as public transport, walking and cycling) in urban areas not only helps to reduce
1440 emissions from vehicles, but also has several health co-benefits. This indicator tracks trends in
1441 sustainable travel infrastructure and uptake in urban areas.

1442 Whilst this indicator would ideally track the proportion and distance of journeys undertaken by
1443 different modes of transport over time, data availability for city-level trends in modal share is
1444 particularly scarce. Therefore, the Lancet Countdown will instead present data for selected locations,

1445 across a limited time-scale. Figure 3.11 presents data on current modal shares (i.e. recent year
 1446 estimates of the proportion of trips by different modes of transport) in world cities (see Appendix 4
 1447 for details). The data, collated by the Land Transport Authority come from travel surveys of
 1448 individual cities and national census data (see Appendix 4 for details).¹⁴³

1449



1450


1451 Figure 3.11 Modal Shares in world cities. Note: 'Other' typically includes paratransit (transport for people with
 1452 disabilities) and/or electric bikes.

1453

1454 Figure 3.12 collates data on trends in modal share in select cities, where data from at least three
 1455 time points (including one pre-2000 time point) is available. While many cities have started to collect
 1456 this information in the past decade, there is a paucity of data on trends from before 2000, with
 1457 particularly wide gaps in data availability from cities in Asia, Africa and South America.¹⁴⁴

1458 In Berlin, London and Tokyo, the proportion of trips by privatised motor transport has slowly
 1459 declined since the late 1990s, while levels have remained high in Vancouver and Sydney and appear
 1460 to be increasing in Santiago. Levels of cycling are generally low, but appear to be increasing in many
 1461 cities.

1462 Public transport in emerging cities is often insufficient, inefficient and in poor condition, potentially
 1463 leading to further declines in sustainable travel in many rapidly growing cities in the future. ¹⁴⁵ As
 1464 this transition occurs, ensuring the mistakes made in Organization for Economic Cooperation and
 1465 Development (OECD) countries are not repeated will be vital. In particular, it is critical to improve
 1466 walking and cycling environments, in order to both make these modes attractive choices and protect
 1467 road users from injury. Recent United Nations (UN) guidance recommends devoting 20% of
 1468 transport budgets to funding non-motorized transport at national and local levels in low- and
 1469 middle-income countries.¹⁴⁶



1470
1471
1472
1473
1474
1475
1476
1477

Figure 3.12 Trends in modal share in selected cities. Note: Data from Santiago in 1991 represents travel on a usual day; Data from Sydney represent Weekdays only; Cycling modal share in Sydney is <1%.¹⁴⁷⁻¹⁵⁶ (Figure created using data from the following sources: Institute for Mobility Research (2016); Transport for London (2016); NSW Department of Transport (1996); NSW Department of Transport (2003); NSW Department of Transport (2009); NSW Department of Transport (2017); Translink (2012); Dictuc S.A. (1992); Rode et al (2015); and City of Berlin (2013))

1478 **Food and agriculture**

1479 The availability of food is central to human health. Its production, however, is also a major
1480 contributor to climate change, with the agricultural sector alone contributing 19-29% of
1481 anthropogenic GHG emissions globally.^{10,157}

1482 Dietary choices determine food energy and nutrient intake, which are essential for human health,
1483 with inadequate and unhealthy diets associated with malnutrition and health outcomes including
1484 diabetes, cardiovascular diseases, and some cancers. Globally, dietary risk factors were estimated to
1485 account for over 10% of all Disability Adjusted Life Years (DALYs) lost in 2013.¹⁵⁸ A transition to
1486 healthier diets, with reduced red and processed meat consumption, and higher consumption of
1487 locally and seasonally produced fruits and vegetables, could provide significant emissions savings.¹⁵⁹


1488 Tracking progress towards more sustainable diets requires consistent and continuous data on food
1489 consumption, and related GHG emissions throughout food product life cycles. This would require
1490 annual nationally representative dietary survey data on food consumption. However, due to the
1491 complexity and cost of such data collection, dietary surveys are available for a limited number of
1492 countries and years only.¹⁶⁰ Although efforts to compile data and ensure comparability are under
1493 way, their current format is not suitable for global monitoring of progress towards optimal dietary
1494 patterns in terms of health benefits of climate change mitigation.^{161,162}

1495

1496 **Indicator 3.8: Ruminant meat for human consumption**

1497 **Headline Finding:** *Globally, the amount of ruminant meat available for human consumption has*
1498 *declined slightly from 12.09 kg/capita/year in 1990 to 11.23 in 2013; the proportion of energy*
1499 *(kcal/capita/day) available for human consumption from ruminant meat as opposed to other sources*
1500 *has declined marginally from 1.86% in 1990 to 1.65% in 2013.*


1501 This indicator focuses on ruminants because the production of ruminant meat, in particular cattle,
1502 dominates GHG emissions from the livestock sector (estimated at 5.6-7.5 GtCO₂e per year), and
1503 consumption of red meat has known associations with adverse health outcomes.¹⁶³ It measures the
1504 total amount of ruminant meat available for consumption, and the ratio of ruminant meat energy
1505 supply to total energy supply. Together, these reflect the relative amount of high GHG emission
1506 foods in the system (Figure 3.13).¹⁶⁴⁻¹⁶⁶ Assuming correlation between ruminant meat supply and
1507 consumption, the indicator therefore also provides information on variations in certain diet-related
1508 health outcomes (such as colorectal cancer and heart disease).^{167,168} This indicator should be viewed
1509 in the context of the specific setting where this trend is examined (in some populations, meat
1510 consumption is a main source of food energy and provides essential micronutrients, as well as
1511 livelihoods). Data was constructed using data from the FAO food balance sheets, which comprises
1512 national supply and utilisation accounts of primary foods and processed commodities.¹⁶⁹



1513

1514 Figure 3.13. The total amount of ruminant meat available for human consumption in kg/capita/year by WHO-
 1515 defined regions.

1516 The amount of ruminant meat available for consumption is high in the Americas and has remained
 1517 relatively stable across 1990-2013. In Europe, the amount of ruminant meat was relatively high in
 1518 1990, declined rapidly from 1990-2000 and has remained stable from 2000-2013. Amounts are more
 1519 moderate in Africa and the Eastern Mediterranean and have remained reasonably constant over
 1520 time; South East Asia and Western Pacific have low amounts but have been slowly increasing in the
 1521 Western Pacific since 1990.



1522

1523 Figure 3.14 The proportion of energy (kcal/capita/day) available for human consumption from ruminant meat
 1524 vs from all food sources by WHO-defined regions.

1525 The proportion of energy supply from ruminant meat has been markedly higher in the Americas than
 1526 other regions since the 1990s, although the trend has been decreasing over time (Figure 3.14). In
 1527 Europe, the proportion of energy from ruminant meat rapidly declined from 1990-2000 and has
 1528 continued to slowly decline. By contrast, the trend has been increasing in the Western Pacific,
 1529 possibly reflecting the increasing trend in beef consumption in China (16% annually).¹⁷⁰

1530 Healthcare sector

1531 The healthcare sector is a considerable contributor to GHG emissions, and has both a responsibility
 1532 and an appreciable opportunity to lead by example in reducing its carbon footprint. In 2013, the
 1533 estimated US healthcare sector emissions were 655 MtCO₂e, which exceeded emissions of the entire
 1534 UK.¹⁷¹ GHG emissions in the healthcare sector illustrate an obvious externality which contributes to
 1535 climate change, contradicting the sector's aim of improving population health.

1536 The World Bank estimates that a 25% reduction from existing healthcare emissions in Argentina,
 1537 Brazil, China, India, Nepal, Philippines, and South Africa would equate to 116-194 million metric tons
 1538 of CO₂e emission reduction, in other terms equal to decommissioning of 34-56 coal fired power
 1539 plants or removing 24-41 million passenger vehicles from the road.¹⁷¹

1540

1541 Indicator 3.9: Healthcare sector emissions

1542 **Headline Finding:** Whilst no systematic global standard for measuring the greenhouse gas emissions
 1543 of the healthcare sector currently exists, a number of healthcare systems in the UK, US, and around
 1544 the world are working to reduce their contribution to climate change.

1545 Several health sector emission reduction targets can be highlighted as positive examples. The
1546 National Health Service (NHS) in the UK set an ambitious target of 34% health-system wide GHG
1547 emission reduction by 2020; Kaiser Permanente in the U.S. has set 2025 as a target to become net
1548 carbon positive; the Western Cape Government health system in South Africa committed to 10%
1549 emission reduction by 2020 and 30% by 2050 in government hospitals; and Albert Einstein Hospital
1550 in Sao Paulo, Brazil, has reduced its annual emissions by 41%.¹⁷¹

1551 In the UK, comprehensive GHG emissions reporting was facilitated by the centralized structure of the
1552 NHS. The Sustainable Development Unit (SDU) of the NHS has been monitoring GHG emissions from
1553 a 1992 baseline, including major contributions from procurement of pharmaceuticals and other
1554 products. NHS emissions reduced by 11% from 2007 to 2015, despite an 18% increase in activity.¹⁷²
1555 Mitigation efforts from the healthcare sector provide remarkable examples of hospitals and health
1556 care systems leading by example, yielding impressive financial savings and health benefits for their
1557 patients. To this end, the efforts of the hospitals, governments, and civil society organisations driving
1558 this work forward must be supported and redoubled, ensuring a full transition to a healthier, more
1559 sustainable model of climate-smart, and increasingly carbon neutral healthcare.¹⁷¹

1560 Monitoring healthcare system emissions is an essential step towards accounting for the externality
1561 of these emissions. Comprehensive national GHG emissions reporting by the healthcare system is
1562 currently only routinely performed in the UK. Elsewhere, select healthcare organisations, facilities,
1563 and companies provide self-reported estimates of emissions, however this is rarely standardized
1564 across sites. The Lancet Countdown will continue to work on developing a standardised indicator on
1565 health sector emissions for subsequent reports.

1566

1567 Conclusion

1568 The indicators presented in this section have provided an overview of activities relevant to public
1569 health for the energy, transport, food and healthcare sectors' mitigation. They have been selected
1570 for their relevance to both climate change and human health and wellbeing.

1571 A number of areas show remarkable promise – each of which should yield impressive benefits for
1572 human health. However, these positive examples must not distract from the enormity of the task at
1573 hand. The indicators presented in this section serve as a reminder of the scale and scope of
1574 increased ambition required to meet commitments under the Paris Agreement. They demonstrate a
1575 world which is only just beginning to respond to climate change, and hence only just unlocking the
1576 opportunities available for better health.

1577

1578

1579 4. Finance & Economics

1580

1581 Introduction

1582 Interventions to protect human health from climate change risks have been presented above. This
1583 section focuses on the economic and financial mechanisms necessary for them to be implemented,
1584 and their implications. Some the indicators here do not have an explicit link to human health, and yet,
1585 investment in renewable energy and a declining investment in coal capacity, for instance, is essential
1586 in displacing fossil fuels and reducing their two principal externalities – the social cost of climate
1587 change and the health costs from air pollution. Other indicators, such as economic and social losses
1588 from extreme weather events, have more explicit links to human wellbeing.

1589 The 2006 Stern Review on the Economics of Climate Change estimated that the impacts of climate
1590 change would cost the equivalent of reducing annual global Gross World Product (GWP) – the sum
1591 of global economic output – by “5-20% now, and forever”, compared to a world without climate
1592 change.¹⁷³ The Intergovernmental Panel on Climate Change’s (IPCC) AR5 estimates an aggregate loss
1593 of up to 2% GWP even if the rise in global mean temperatures is limited to 2.5°C above pre-industrial
1594 levels.²² However, such estimates depend on numerous assumptions, such as the rate at which
1595 future costs and benefits are discounted. Further, existing analytical approaches are poorly suited to
1596 producing estimates of the economic impact of climate change, and hence their magnitude is likely
1597 greatly underestimated.^{174 175} In the presence of such uncertainty, with potentially catastrophic
1598 outcomes, risk minimisation through stringent emissions reduction seems the sensible course of
1599 action.

1600 The indicators in this section, which seek to track flows of finance and impacts on the economy and
1601 social welfare resulting from (in)action on climate change, fall into four broad themes: investing in a
1602 low-carbon economy; the economic benefits of tackling climate change; pricing GHG emissions from
1603 fossil fuels; and adaptation financing. The indicator presented are:

- 1604 4.1 Investments in zero-carbon energy and energy efficiency
- 1605 4.2 Investment in coal capacity
- 1606 4.3 Funds divested from fossil fuels
- 1607 4.4 Economic losses due to climate-related extreme events
- 1608 4.5 Employment in low-carbon and high-carbon industries
- 1609 4.6 Fossil fuel subsidies
- 1610 4.7 Coverage and strength of carbon pricing
- 1611 4.8 Use of carbon pricing revenues
- 1612 4.9 Spending on adaptation for health and health-related activities
- 1613 4.10 Health adaptation funding from global climate financing mechanisms

1614

1615 Appendix 5 provides more detailed discussion of the data and methods used.

1616


1617

1618 Indicator 4.1: Investments in zero-carbon energy and energy efficiency

1619 **Headline Finding:** Proportional investment in renewable energy and energy efficiency increased in
 1620 2016, whilst absolute and proportional investment in fossil fuels decreased, and crucially, ceased to
 1621 account for the majority of annual investments in the global energy system.

1622 This indicator tracks the level of global investment in zero-carbon energy and energy efficiency in
 1623 absolute terms, and as a proportion of total energy system investment. Figure 4.1 illustrates the data
 1624 for 2015 and 2016; the data for this indicator is sourced from the IEA.^{176,177}

1625



1626 Figure 4.1 Annual Investment in the Global Energy System.

1627

1628 In 2015, total investment in the energy system was around \$1.83 trillion (in US\$2016), accounting
 1629 for 2.4% of GWP. Renewables and nuclear comprised 19% of this investment, and energy efficiency
 1630 12%. Most investment (54%) was in fossil fuel infrastructure. Electricity networks accounted for the
 1631 remaining 15%. In 2016, total investment in the energy system reduced to around \$1.68 trillion,
 1632 accounting for 2.2% of GWP. Although the absolute value of investment in renewables and nuclear
 1633 energy reduced slightly in absolute (real) terms, its proportional contribution increased to 20%.
 1634 Investment in energy efficiency increased in both absolute and proportional terms to 14%. Fossil fuel
 1635 infrastructure suffered a significant reduction in investment, ceasing to account for the majority of
 1636 investment (at 49%). Such trends broadly represent a continuation of the trends experienced
 1637 between 2014 and 2015.¹⁷⁸

1638 Investment in renewables and nuclear is driven by renewable electricity capacity (with over 87% of
 1639 investment by value in this category in 2016). This, in turn, is largely driven by investments in solar
 1640 PV and onshore wind. Solar PV capacity additions in 2016 were 50% higher than 2015 (reaching
 1641 record levels of 73GW), driven by new capacity in China, the USA and India. However, this was
 1642 coupled with just a 20% increase in investment, resulting from a 20% reduction in the cost of solar
 1643 PV units. By contrast, investments in onshore wind reduced by around 20% between 2015 and 2016,
 1644 largely driven by changes to incentive schemes and elevated wind power curtailment rates in China.
 1645 The increase in energy efficiency investment was driven by policies that shifted markets towards
 1646 more energy efficient goods (such as appliances and lighting) and buildings (along with the

1647 expansion of the construction industry), and an increase in the sales of energy efficient (and low-
 1648 carbon) vehicles. Europe accounted for the largest proportion of spending on energy efficiency
 1649 (30%), followed by China (27%), driven by efficiency investments in the buildings and transport
 1650 sectors.¹⁷⁷

1651 The substantial reduction in fossil fuel infrastructure investment, both upstream (such as mining,
 1652 drilling and pipelines, which dominate fossil fuel investment) and downstream (such as fossil fuel
 1653 power plants) is driven by a combination of low (and reducing) fossil fuel prices and cost reductions
 1654 (particularly upstream, which have on average reduced by 30% since 2014).¹⁷⁷

1655

1656 In order to hold a 66% probability of remaining within 2°C of warming, it is estimated that average
 1657 annual investments in the energy system between 2016 and 2050 must reach \$3.5 trillion, with
 1658 renewable energy investments increasing by over 150%, and energy efficiency increasing by around
 1659 a factor of ten.¹⁷⁹


1660

1661 **Indicator 4.2: Investment in coal capacity**

1662 **Headline Finding:** *Although investment in coal capacity has increased since 2006, in 2016 this trend*
 1663 *turned and declined substantially.*

1664 The combustion of coal is the most CO₂-intensive method of generating of electricity..¹⁸⁰This
 1665 indicator tracks annual investment in coal-fired power capacity. Figure 4.2 presents an index of
 1666 global annual investment in coal power generation capacity from 2006 to 2016, using IEA data.¹⁷⁷

1667



1668 Figure 4.2. Annual Investment in coal-fired power capacity.

1669

1670 It is clear that global investment in coal-fired electricity capacity generally increased from 2006 to
 1671 2012, before returning to 2006 levels in 2013-14, and rebounding significantly to over 40% above
 1672 this level in 2015. This rapid growth was driven principally by China, which increased investment in

1673 coal-fired power capacity by 60% from 2014, representing half of all new global coal capacity in 2015
1674 (with investment in India and other non-OECD Asia countries also remaining high).¹⁷⁸ The
1675 subsequent reduction in investment in 2016 was similarly driven by reduced investment in China,
1676 due to overcapacity in generation, concerns about local air pollution and new government measures
1677 to reduce new capacity additions and halt the construction of some plants already in progress.¹⁷⁷

1678

1679 [Indicator 4.3: Funds divested from fossil fuels](#)

1680 **Headline Finding:** *Global Value of Funds Committing to Divestment in 2016 was \$1.24 trillion, of*
1681 *which Health Institutions represent \$2.4 billion; this represents a cumulative sum of \$5.45 trillion*
1682 *(with health accounting for \$30.3 billion).*

1683 The fossil fuel divestment movement seeks to encourage institutions and investors to divest
1684 themselves of assets involved in the extraction of fossil fuels. ‘Divestment’ is defined relatively
1685 broadly, ranging from an organisation that has made a binding commitment to divest from coal
1686 companies only, to those who have fully divested from any investments in fossil fuel companies and
1687 have committed to avoiding such investments in future. Proponents cite divestment as embodying
1688 both a moral purpose (for example, reducing the fossil fuel industry’s ‘social licence to operate’), and
1689 an economic risk reduction strategy (for example, through reducing the investor’s exposure to the
1690 risk of ‘stranded assets’). However, others believe active engagement between investors and fossil
1691 fuel businesses is a more appropriate course of action (for instance, encouraging diversification into
1692 less carbon-intensive assets, through stakeholder resolutions).¹⁸¹

1693 This indicator tracks the global total value of funds committing to divestment in 2016, and the value
1694 of funds committed to divestment by health institutions in 2016, which was \$1.24 trillion, and \$2.4
1695 billion respectively. The values presented above are calculated from data collected and provided by
1696 350.org. They represent the total assets (or assets under management (AUM)) for institutions that
1697 have committed to divest in 2016, and thus do not directly represent the sums divested from fossil
1698 fuel companies. It also includes only those institutions for which such information is publicly
1699 available (or provided by the institution itself), with non-US\$ values converted using the market
1700 exchange rate when the commitment was made.

1701 By the end of 2016, a total of 694 organisations with cumulative assets worth at least \$5.45 trillion,
1702 including 13 health organisations with assets of at least \$30.3 billion, had committed to divestment.
1703 From the start of January 2017 to the end of March 2017, a further 12 organisations with assets
1704 worth \$46.87 billion joined this total (including Australia’s Hospitals Contribution Fund – HCF – with
1705 assets of \$1.45 billion).

1706

1707 [Indicator 4.4: Economic losses due to climate-related extreme events](#)

1708 **Headline Finding:** *In 2016, a total of 797 events resulted in \$129 billion in overall economic losses,*
1709 *with 99% of losses in low-income countries uninsured.*


1710 Climate change will continue to increase the frequency and severity of meteorological (tropical
1711 storms), climatological (droughts) and hydrological (flooding) phenomena, across the world. As
1712 demonstrated by indicator 1.4, the number of weather-related disasters has increased in recent
1713 years. The number of people affected and the economic costs associated with this increase is
1714 expected to have risen. This indicator tracks the number of events and the total economic losses
1715 (insured and uninsured) resulting from such events. In addition to the health impacts of these

1716 events, economic losses (particularly uninsured losses) have potentially devastating impacts on
 1717 wellbeing and mental health.¹⁸²

1718 The data upon which this indicator is based is sourced from Munich Re.¹⁸³ Economic losses (insured
 1719 and uninsured) refer to the value of physical assets, and do not include the economic value of loss of
 1720 life or ill health, or health and casualty insurance. Values are first denominated in local currency,
 1721 converted to US\$ using the market exchange rate in the month the event occurred, and inflated to
 1722 US\$2016 using country-specific Consumer Price Indices (CPI). This indicator and underlying data does
 1723 not seek to attribute events and economic losses to climate change *per se*, but may plausibly be
 1724 interpreted as showing how climate change is changing the frequency and severity of these events.

1725 Figure 4.3 presents insured and uninsured economic losses resulting from all significant
 1726 meteorological, climatological and hydrological events across the world, from 2010 to 2016, by
 1727 country income group. An annual average of 700 events resulted in an annual average of \$127 billion
 1728 in overall economic losses per year over this timeframe. Upper-middle and high-income countries
 1729 experienced around two-thirds of the recorded events and around 90% of economic losses, with
 1730 <1% attributable to those of low-income. The same ratios for the number of events and economic
 1731 losses between income groups is present in the data for the period 1990-2016, despite an increasing
 1732 trend in the total global number of events and associated total value of economic losses over this
 1733 period.

1734



1735


1736 Figure 4.3 Economic Losses from Climate-Related Events – Absolute.

1737

1738 However, the data in Figure **Error! Reference source not found.3** does not indicate the relative scale
 1739 of impacts across different income groups. For example, although the majority of economic losses
 1740 have occurred in upper-middle and high-income countries, these countries are among the most

1741 populous, with more economically valuable property and infrastructure (in absolute terms). A rather
 1742 different picture emerges in Figure 4.4, which presents the data in terms of ‘intensity’ – insured and
 1743 uninsured economic losses per \$1000 GDP (in US\$2016).

1744



1745 Figure 4.4 Economic Losses from Climate-Related Events - Intensity.

1746

1747 Between 2010 and 2016, high and upper-middle income countries experienced the least average
 1748 annual economic loss as a proportion of GDP (\$1.45/\$1000 GDP and \$1.95/\$1000 GDP, respectively),
 1749 with low and lower-middle income countries subject to somewhat higher values (\$2.65/\$1000 GDP
 1750 and \$2.3/\$1000 GDP, respectively). Economic losses in low-income countries were more than three
 1751 times as high in 2016 than in 2010. However, for 1990-2016, average annual values vary significantly
 1752 (see Appendix 5 for the full dataset). Whilst high and upper-middle income countries maintain
 1753 relatively similar values (\$1.60/\$1000 GDP and \$2.9/\$1000 GDP, respectively), average annual
 1754 economic losses experienced by (particularly) low and lower-middle income countries increase
 1755 substantially (to \$10.95/\$1000 GDP and \$4.22/\$1000 GDP, respectively).

1756 It is clear that, on average, lower income countries experience greater economic loss as a proportion
 1757 of GDP as a result of climate-related events than higher-income countries. However, a more striking
 1758 result is the difference in the proportion of economic losses that are uninsured. In high-income
 1759 countries, on average around half of economic losses experienced are insured. This share drops
 1760 rapidly to under 10% in upper-middle income countries, and to well under 1% in low-income
 1761 countries. Over the period 1990-2016, uninsured losses in low-income countries were on average
 1762 equivalent to over 1.5% of their GDP. For contrast, expenditure on healthcare in low-income
 1763 countries on average for the period 1995-2015 was equivalent to 5.3% of GDP.¹⁸⁴


1764 **Indicator 4.5: Employment in low-carbon and high-carbon industries**

1765 **Headline Finding:** In 2016, global employment in renewable energy reached 9.8 million, with
 1766 employment in fossil fuel extraction trending down, to 8.6 million.

1767 The generation and presence of employment opportunities in low- and high-carbon industries have
 1768 important health implications, both in terms of the safety of the work environment itself and
 1769 financial security for individuals and communities. As the low-carbon transition gathers pace, high-
 1770 carbon industries and jobs will decline. A clear example is seen in fossil fuel extraction. Some fossil
 1771 fuel extraction activities, such as coal mining, have substantial impacts on human health. Coal mining
 1772 accidents led to over 1,000 deaths in 2008 in China alone (a rapid decline from nearly 5,000 in 2003),
 1773 with exposure to particulate matter and harmful pollutants responsible for elevated levels of
 1774 cardiovascular, respiratory and kidney disease, in coal mining areas.¹⁸⁵⁻¹⁸⁸ The low-carbon transition
 1775 is also likely to stimulate the growth of new industries and employment opportunities. With
 1776 appropriate planning and policy, the transition from employment in high-carbon to low-carbon
 1777 industries will yield positive consequences for human health.

1778 This indicator tracks global employment levels in fossil fuel extraction industries (coal mining and oil
 1779 and gas exploration and production), and in renewable energy. Figure 4.5 presents these values for
 1780 2012-2016. The data for this indicator is sourced from International Renewable Energy Agency
 1781 (IRENA) (renewables), and IBIS World (fossil fuel extraction).¹⁸⁹⁻¹⁹¹

1782



1783

1784 Figure 4.5 Employment in Renewable Energy and Fossil Fuel Extraction.

1785

1786 From a peak of 9.1 million in 2014, jobs in the global fossil fuel extraction industry reduced by
 1787 around 500,000 to 8.6 million in 2016. Reductions in the coal mining industry largely drove this
 1788 change, which was the result of a range of factors, including its substitution by lower-cost natural
 1789 gas in the power sector in many countries, reducing the demand for coal and leading to
 1790 overcapacity, industry consolidation, and the rising automation of extractive activities.¹⁹¹

1791 By contrast, employment in the renewable energy industry increased rapidly from over 7.1 million
 1792 jobs in 2012 to over 9.3 million in 2014, and reaching 9.8 million in 2016. This growth has largely
 1793 been driven by the solar PV industry, which added over 1.7 million jobs between 2012 and 2016.
 1794 Solar PV is now the largest renewable energy employer, overtaking bioenergy, which has
 1795 experienced a reduction of 250,000 jobs since 2012.

1796


1797 **Indicator 4.6: Fossil fuel subsidies**

1798 **Headline Finding:** In 2015, fossil fuel consumption subsidies followed a trend seen since 2012,
 1799 decreasing markedly to \$327 billion, principally as a result of declining global oil prices.

1800 The combustion of fossil fuels results in a variety of harmful consequences for human health, and
 1801 the presence of subsidies for fossil fuels, either for its production (such as fossil fuel extraction) or
 1802 consumption (such as regulated gasoline prices), artificially lowers prices, promoting
 1803 overconsumption. This indicator tracks the global value of fossil fuel consumption subsidies. Figure
 1804 4.6 illustrates the value of fossil fuel consumption subsidies for 2010-2016 using IEA data.^{178,192}

1805

1806



1807 Figure 4.6 Global Fossil Fuel Consumption Subsidies - 2010-2015.

1808

1809 Despite rising from \$444 billion in 2010 to a peak of \$571 billion in 2012, fossil fuel consumption
 1810 subsidies have decreased markedly to \$327 billion in 2015 (in US\$2016). The principal driver for this
 1811 is the doubling in oil price between 2010 and 2012, after which it plateaued, before falling rapidly to
 1812 below 2010 levels from mid-2014. Fossil fuel consumption subsidies are typically applied in order to
 1813 moderate energy costs for low-income consumers (although in practice, 65% of such subsidies in
 1814 LMICs benefit the wealthiest 40% of the population).¹⁹³ As such, rising oil (and other fossil fuel)
 1815 prices tend to increase subsidy levels, as the differences between market and regulated consumer
 1816 prices increase, and governments take further action to mitigate the impact on citizens. When fossil

1817 fuel prices decrease, the gap between market and regulated prices reduces, and governments can
1818 reform fossil fuel subsidies whilst keeping overall prices relatively constant.

1819 Between 2014 and 2015, several countries took advantage of this opportunity, particularly regarding
1820 oil-based fuels, which accounted for over 60% of the reduction in total fossil fuel subsidies between
1821 2012 and 2015 (followed by natural gas at around 25%). This included India, which in deregulating
1822 diesel prices accounted for a \$19 billion subsidy reduction between 2014 and 2015 (~13% of the
1823 global total reduction), and the major oil and natural gas producing nations (including Angola,
1824 Algeria, Indonesia, Iran, Qatar, Saudi Arabia and Venezuela), in which reduced hydrocarbon revenue
1825 created pressure for fiscal consolidation, and in turn for consumption subsidy reform.¹⁷⁸ To
1826 encourage the low-carbon transition, fossil fuel subsidies should be phased out as soon as possible.
1827 The commitment made by the G7 in 2016 to achieve this goal by 2025 should be extended to all
1828 OECD countries, and globally by 2030.¹⁹⁴

1829

1830 [Indicator 4.7: Coverage and strength of carbon pricing](#)

1831 **Headline Finding:** *So far in 2017, various carbon pricing mechanisms covered 13.1% of global*
1832 *anthropogenic CO₂ emissions, up from 12.1% in 2016. This reflects a doubling in the number of*
1833 *national and sub-national jurisdictions with a carbon pricing mechanism over the last decade.*

1834 This indicator tracks the extent to which carbon pricing instruments are applied around the world as
1835 a proportion of total GHG emissions, and the weighted average carbon price such instruments
1836 provide (Table 4.1).

1837

	2016	2017
Global Emissions Coverage*	12.1%	13.1%
Weighted Average Carbon Price of Instruments (current prices, US\$)	\$7.79	\$8.81
Global Weighted Average Carbon Price (current prices, US\$)	\$0.94	\$1.12


1838 Table 4.1 Carbon Pricing - Global Coverage and Weighted Average Prices per tCO₂e. *Global emissions
1839 coverage is based on 2012 total anthropogenic GHG emissions.¹⁹⁵ (Source: World Bank, 2017)

1840

1841 Between 2016 and 2017, the proportion of global emissions covered by carbon pricing instruments,
1842 and the weighted average price of these instruments (and thus the global weighted average price for
1843 all anthropogenic GHG emissions), increased. This is due to the introduction of four new instruments
1844 in 2017 (note, this data runs up to 1 April 2017) - the carbon taxes in Alberta, Chile and Colombia,
1845 and an Emissions Trading System (ETS) in Ontario. As such, over 40 national and 25 sub-national
1846 jurisdictions now put a price on at least some of their GHG emissions (with substantially varying
1847 prices, from less than \$1/tCO₂e in Chongqing, to over \$126/tCO₂e in Sweden). The last decade has
1848 seen a rapid increase in the number of carbon pricing instruments around the world, with the
1849 number of jurisdictions introducing them doubling.¹⁹⁶ Over 75% of the GHG emissions covered by
1850 carbon pricing instruments are in HICs, with the majority of the remainder covered by the 8 pilot
1851 pricing instruments in China (Figure 4.7).

1852 The World Bank provides the data for this indicator.^{195,196} Prices for 2016 and 2017 are those as of 1
 1853 August 2016 and 1 April 2017, respectively. For 2017, the indicator includes only instruments that
 1854 had been introduced by 1 April 2017. Instruments without price data are excluded.

1855



1856

1857 Figure 4.7 Carbon Pricing Instruments implemented, scheduled for implementation and
 1858 under consideration.¹⁹⁶ (Source: World Bank, 2017)

1859

1860 In total, a further 21 carbon pricing instruments are either scheduled for implementation, or are
 1861 under consideration. This includes the commencement of a national ETS in China expected in the
 1862 second half of 2017. Although this would replace the 8 pilot schemes currently in place in China, it
 1863 could expand their emissions coverage fourfold, surpassing the European ETS to become the largest
 1864 carbon pricing instrument in the world.¹⁹⁶

1865

1866 Indicator 4.8: Use of carbon pricing revenues

1867 **Headline Finding:** 40% of government revenues generated from carbon pricing are spent on climate
 1868 change mitigation, totalling US\$9 billion.

1869 Carbon pricing instruments require those responsible for producing the emissions concerned to pay
 1870 for their emissions, in one form or another. In most cases this generates revenue for the
 1871 governments or authorities responsible for introducing the instrument. Such revenue may be put to
 1872 a range of uses, including investment in climate change mitigation or adaptation or environmental
 1873 tax reform (ETR), which involves shifting the burden of tax from negative activities, such as the
 1874 generation of pollution, to positive activities, such as labour or environmentally beneficial products
 1875 or activities. Such options may produce a ‘double dividend’ of environmental improvement with
 1876 social and economic benefits.¹⁹⁷ This indicator tracks the total government revenue from carbon
 1877 pricing instruments, and how such income is allocated.

	Mitigation	Adaptation	Environmental Tax Reform (ETR)	General Funds	Total Revenue (US\$2016)
Proportion (%)	40.4%	4%	19.5%	36.1%	\$22.31 Billion
Value (US\$2016)	\$9.01 Billion	\$0.9 Billion	\$4.34 Billion	\$8.06 Billion	

1878

1879 Table 4.2. Carbon Pricing revenues and allocation in 2016.¹⁹⁵ (Source: World Bank, 2017)

1880

1881 Tale 4.2 presents total government revenue generated by carbon pricing instruments in 2016, and
 1882 four categories of expenditure for this revenue. The largest expenditure category is climate change
 1883 mitigation, which is in receipt of over \$9 billion annually in funds. Despite this, less than half of
 1884 revenue-generating instruments allocate revenue for mitigation.

1885 ETR policies accounted for around 20% of revenue allocation in 2016. Just two instruments (the
 1886 Portuguese and British Colombia Carbon Taxes) allocate all their revenue to allowing revenue-neutral
 1887 reduction in other (for example, income) taxes, with another four allocating part of their revenue to
 1888 this purpose. By contrast, only four instruments do not have any revenue allocated to general
 1889 government funds (The British Colombian, Swiss, Japanese and Portuguese carbon taxes), with 11
 1890 instruments allocating all revenues to this category (reaching €8 billion – or more than a third – of
 1891 revenues generated in 2016). Data for individual carbon pricing instruments may be found in Appendix
 1892 5.

1893 Data on revenue generated is provided by the World Bank, with revenue allocation information
 1894 obtained from various sources (see Appendix 5).¹⁹⁵ Only instruments with revenue estimates, and only
 1895 revenue received by the administering authority before redistribution, are considered. Revenue must
 1896 be explicitly allocated to climate change mitigation or adaptation, or for ETR, to be considered in these
 1897 categories. If such explicit earmarking is not present, or no data is available, then revenue is assumed
 1898 to be allocated to general funds.

1899

1900 [Indicator 4.9: Spending on adaptation for health and health-related activities](#)

1901 **Headline finding:** *Out of the world's total adaptation spend just 4.63% (\$16.46 billion USD) is on*
 1902 *health and 13.3% (\$47.29 billion USD) on health-related adaptation.*

1903 This indicator reports estimates of spending on health and health-related climate change adaptation
 1904 and resilience. Many adaptation activities within and beyond the formal health sector yield health


1905 co-benefits, which are important to understand and capture. Here, estimates of the total health and
 1906 health-related adaptation spending were derived from the Adaptation & Resilience to Climate
 1907 Change (A&RCC) dataset produced by kMatrix. This global dataset, covering financial transactions
 1908 relevant to climate change adaptation, was compiled from a relevant subset of over 27,000
 1909 independent databases and sources (such as public disclosures and reports from insurance
 1910 companies, the financial sector, and governments).¹⁹⁸ In this case, entries were triangulated
 1911 between at least seven independent sources before being included.

1912 Examples of transactions captured here range from the procurement of goods or services (for
 1913 example, purchasing sandbags for flood levees) through to spending on research and development
 1914 (for example, for vulnerability and adaptation assessments) or staff training.¹⁹⁸ Each of these
 1915 ‘adaptation activities’ are grouped in to eleven sectors: Agriculture and Forestry, Built Environment,
 1916 Disaster-Preparedness, Energy, Health, ICT, Natural Environment, Professional Services, Transport,
 1917 Waste, and Water. Whilst adaptation spending relevant directly to the formal health sector is clearly
 1918 important (the ‘health’ category), interventions outside of the healthcare system will also yield
 1919 important benefits for health and wellbeing. ‘Health-related adaptation spending’ was defined as
 1920 that which additionally included adaptation spending from the agricultural sector (due to the
 1921 centrality of food and nutrition to health) and disaster preparedness sector (due to the direct public
 1922 health benefits that often result from these efforts).

1923 This data from the A&RCC dataset is reported here, showing health and health-related adaptation
 1924 spending for 180 countries for the 2015-2016 financial year. Global health adaptation spending for
 1925 the financial year 2015-2016, calculated in this way, totalled 16.46 billion USD, representing 4.63% of
 1926 the global aggregate adaptation spend. Health-related adaptation spending totalled 47.29 billion
 1927 USD, or 13.3% of the global total adaptation spend (Figure 4.8).

1928 Health-related adaptation and resilience spending, both national totals and per capita levels, is
 1929 extremely low in low-income countries, and increase across the continuum towards high-income
 1930 countries. Interestingly, health and health-related adaptation spending as a proportion of total
 1931 adaptation spending is relatively constant across income groups.

1932



1933

1934 Figure 4.8 For the financial year 2015-2016. 4.8a) Total health and health-related adaptation spending and
 1935 4.8b) health and health-related adaptation and resilience to climate change (A&RCC) spending as a proportion
 1936 of GDP. All plots are disaggregated by World Bank Income Grouping.

1937
1938
1939
1940
1941
1942
1943
1944
1945

1946

1947
1948
1949

1950
1951
1952
1953
1954
1955
1956

1957
1958
1959
1960
1961
1962

1963


It is important to note that further work is required to more completely determine what should be considered as ‘health-related adaptation spending’. Spending for agriculture and disaster preparedness were included here, however other forms of adaptation spending clearly have important health implications. Second, only economic data relating to the financial year 2015-2016 was available, precluding time trend analysis. Third, since public sector transactions may not leave a sufficient ‘footprint’ to be picked up by this methodology, adaptation spending data here may exclude some public-sector spending.

Indicator 4.10: Health adaptation funding from global climate financing mechanisms

Headline Finding: *Between 2003 and 2017, 0.96% of total adaptation funding for development, flowing through global climate change financing mechanisms, was dedicated to health adaptation.*

The final indicator in this section is designed in parallel with indicator 4.9, and aims to capture development funds available for climate change adaptation. It reports global financial flows dedicated to health adaptation to climate change, moving through established global climate financing mechanisms. Data was drawn from the Climate Funds Update (CFU), an independent source which aggregates funding data from multilateral and bilateral development agencies since 2003.^{16,199} CFU data is presented in four categories (pledged, deposited, approved, and disbursed); this indicator uses data designated as ‘approved’.

Between 2003 and 2017, only 0.96% of approved adaptation funding was allocated to health adaptation, corresponding with a cumulative total of 39.55 million USD (Figure 4.9). Total global adaptation funding peaked in 2013 at 910.36 million USD and declined thereafter. However, health-related adaptation funding reached its highest level in early 2017, resulting in the near-doubling in the proportion of adaptation funding allocated to health. Panel 4.1 provides a brief overview of growing interest in health and climate change from the international donor community.



■ Total Approved Adaptation Funding (USD millions)
 ■ Total Approved Adaptation Funding 2017 to date (USD millions)
 ■ Total Approved Health Adaptation Funding (USD millions)
 ■ Total Approved Health Adaptation Funding 2017 to date (USD millions)

Figure 4.9 Year on year multilateral and bilateral funding for all adaptation projects and health adaptation projects (2003 through May 2017).

Panel 4.1 International Donor Action on Climate Change and Health.

In 2017, the World Bank released three independent reports on climate change and health, articulating (i) a new action plan for climate change and health, (ii) geographic focus areas, and (iii) new strategy for climate-smart healthcare. In addition to training staff and increasing government capacity, the World Bank outlines an approach to ensuring that at least 20% of new World Bank health investments are climate-smart by 2020, corresponding to as much as \$1bn in new climate-smart health finance for countries. Other development institutions and foundations are also getting involved. Two separate, major gatherings of public and private funders occurred in 2016 (May, Helsinki) and 2017 (May, Chicago) toward establishing new channels for health and climate finance, and a third is planned for late 2017 (October, Washington, DC).

Conclusion

The indicators presented in this section seek to highlight the status of the economics and finance associated with climate change and health across four themes; investing in a low-carbon economy, economic benefits of tackling climate change, pricing the GHG emissions from fossil fuels, and adaptation financing.

Many of the trends show positive change over time, notably global investment in zero-carbon energy supply, energy efficiency, new coal-fired electricity capacity, employment in renewable energy, and

1985 divestment in fossil fuels. However, the rate of change is relatively slow, and must accelerate rapidly
1986 to meet the objectives of the Paris Agreement.

1987 5. Public and Political Engagement

1988

1989 Introduction

1990 So far, this report has presented indicators on the health impacts of climate hazards; resilience and
1991 adaptation to climate change; health co-benefits of climate change mitigation; and economics and
1992 finance mechanisms that facilitate a transition to a low-carbon economy.

1993 Policy change requires public support and government action. This is particularly true of policies with
1994 the reach and impact to enable societies to transition to a low-carbon future.²⁰⁰ The overarching
1995 theme of this section is therefore the importance of public and political engagement in addressing
1996 health and climate change, and the consequent need for indicators that track engagement in the
1997 public and political domains.

1998 The aim is to track engagement with health and climate change in the public and political domains
1999 and identify trends since 2007. In selecting indicators, priority has been given to high-level
2000 indicators, which can be measured globally, tracked over time and provide a platform for more
2001 detailed analysis in future Lancet Countdown reports. The indicators relate to coverage of health and
2002 climate change in the media, science, and government. Search terms for the indicators are aligned
2003 and a common time-period was selected for all indicators (2007-2016). The period runs from before
2004 the resolution on health and climate change by the 2008 World Health Assembly, which marked a
2005 watershed in global engagement in health and climate change; for the first time, member states of
2006 the UN made a multilateral commitment to protect human health from climate change.²⁰¹

2007 The indicators presented are:

2008 5.1. Media coverage of health and climate change

2009 5.2. Health and climate change in scientific journals

2010 5.3. Health and climate change in the United Nations General Assembly

2011

2012 Corresponding Appendix 6 provide more detailed discussion of the data and methods used.

2013

2014 Indicator 5.1: Media coverage of health and climate change

2015 **Headline Finding:** *Global newspaper coverage of health and climate change has increased 78% since*
2016 *2007, with marked spikes in 2009 and 2015, coinciding with the 15th and 21st Conference of the*
2017 *Parties (COP).*


2018 Media plays a crucial role in communicating risks associated with climate change.²⁰² Knowledge
2019 about climate change is related to perceptions of risk and intentions to act.^{203,204} Public perceptions
2020 of a nation's values and identity are also an important influence on public support for national
2021 action.²⁰⁵ Indicator 5.1 therefore tracks media coverage of health and climate change, with a global
2022 indicator on newspaper coverage on health and climate change (5.1.1), complemented by an in-
2023 depth analysis of newspaper coverage on health and climate change for two national newspapers
2024 (5.1.2).

2025

2026 5.1.1: Global newspaper reporting on health and climate change
 2027 Focusing on English-language and Spanish-language newspapers, this indicator tracks global
 2028 coverage of health and climate change in high-circulation national newspapers from 2007 to 2016.
 2029 Using 18 high-circulation ‘tracker’ newspapers, global trends are shown and disaggregated regionally
 2030 to provide a global indicator of public exposure to news coverage of health and climate change.

2031 Since 2007, newspaper coverage of health and climate change has risen globally by 78% (Figure 5.1).
 2032 However, this trend is largely driven by South-East Asian newspapers. Although mostly due to the
 2033 higher number of South-East Asian newspapers included in this analysis, the South-East Asian
 2034 newspapers here did have a higher than average coverage of health and climate change than other
 2035 regions, particularly among Indian sources (see Appendix 6). This generally high volume of coverage
 2036 in the Indian press can be attributed to the centrality of newspapers as communication channels for
 2037 elite-level discourse in India and to relatively high levels of climate change coverage throughout
 2038 Asia.²⁰⁶⁻²⁰⁸ For the Eastern Mediterranean, Americas, and Western Pacific, there is not a strong trend
 2039 in the media reporting. Some spikes are notable in 2009 in Europe, which is largely maintained for
 2040 the rest of the time series, and in the Americas, which drops until a secondary spike between 2012
 2041 and 2014. The first major spike globally was in 2009, coinciding with COP15 (Conference of the
 2042 Parties) in Copenhagen, for which there was high expectation. Newspaper reporting then dropped
 2043 around 2010, but since 2011 has been rising overall globally.

2044



2045

2046 Figure 5.1 Newspaper reporting on health and climate change (for 18 newspapers) from 2007 to 2016, broken
 2047 down by WHO region.

2048

2049 Data was assembled by accessing archives through the Lexis Nexis, Proquest and Factiva databases.
 2050 These sources were selected through the weighting of four main factors: geographical diversity
 2051 (favouring a greater geographical range), circulation (favouring higher circulating publications),
 2052 national sources (rather than local/regional), and reliable access to archives over time (favouring
 2053 those accessible consistently for longer periods). Search terms were aligned to those used for the
 2054 indicators of scientific and political engagement and searches, with Boolean searches done in English
 2055 and Spanish.

2056

2057 [5.1.2: In-depth analysis of newspaper coverage on health and climate change](#)
2058 The second part of this indicator provides an analysis of two national newspapers; Le Monde
2059 (France) and Frankfurter Allgemeine Zeitung (FAZ) (Germany). Le Monde and FAZ were chosen for
2060 this analysis, as these are leading newspapers in France and Germany; two countries with political
2061 weight in Europe. Both newspapers continue to set the tone of public debates in France and
2062 Germany.^{209,210}


2063 Only a small proportion of articles on climate change mentioned the links between health and
2064 climate change: 5% in Le Monde and 2% in FAZ. The analysis also pointed to important national
2065 differences in reporting on health and climate change. For example, in France, 70% of articles
2066 referring to health and climate change represented the health-climate change nexus as an
2067 environmental issue, whereas in Germany articles had a broader range of references: the economy
2068 (23%), local news (20%) and politics (17%). The recommended policy responses also differed; in Le
2069 Monde, the emphasis was on adaptation (41% of articles), while FAZ put more emphasis on
2070 mitigation (40% of articles). The co-benefits that public health policies can represent for mitigation
2071 were mentioned by 17% of Le Monde articles and 9% of FAZ articles. Overall, the analysis points to
2072 the marked differences in media reporting of health and climate change, and therefore in the
2073 information and perspectives to which the public is exposed (see Appendix 6 for details).

2074

2075 [Indicator 5.2: Health and climate change in scientific journals](#)

2076 **Headline Finding:** *Since 2007, the number of scientific papers on health and climate change has more*
2077 *than trebled.*

2078 Science is critical to increasing public and political understanding of the links between climate
2079 change and health; informing mitigation strategies; and accelerating the transition to low-carbon
2080 societies.^{211,212} This indicator, showing scientific engagement with health and climate change, tracks
2081 the volume of peer-reviewed publications in English-language journals from PubMed and Web of
2082 Science (see Appendix 6 for details). The results show there has been a marked increase in published
2083 research on health and climate change in the last decade, from 94 papers in 2007 to over 275
2084 published in both 2015 and 2016. Within this overall upward trend, the volume of scientific papers
2085 increased particularly rapidly from 2007-2009 and from 2012, with a plateauing between these
2086 periods (Figure 5.2).



2087

2088 Figure 5.2 Number of scientific publications on climate change and health per year (2007-2016) from PubMed
 2089 and Web of Science journals.

2090

2091 The two periods of growth in scientific outputs coincided with the run-up to the UNFCCC COPs held
 2092 in Copenhagen in 2009 (COP15) and in Paris in 2015 (COP21). This pattern suggests that scientific
 2093 and political engagement in health and climate change are closely linked, with the scientific
 2094 community responding quickly to the global climate change agenda and the need for evidence.

2095 Most publications focus on the impacts of climate change and health in Europe and North America.
 2096 Overall, more than 2000 scientific articles were identified, of which 30% of papers focussed on
 2097 Europe, followed by 29% on the Americas. Within the Americas, the large majority (72%) of the
 2098 papers related to health and climate change in North America (see Figure S5.1 in Appendix 6). By
 2099 contrast, only 10% of published articles had a focus on Africa or the Eastern Mediterranean Region,
 2100 demonstrating a marked global inequality in the science of health and climate change (see Figures
 2101 S5.1 and S5.2 in Appendix 6).

2102 Among the journals in the analysis, infectious diseases, particularly dengue fever and other
 2103 mosquito-transmitted infections, are the most frequently investigated health outcomes;
 2104 approximately 30% of selected papers covered these health-related issues. Important gaps in the
 2105 scientific evidence base were identified, including migration and mental ill-health.


2106 For this indicator, a scoping review of peer-reviewed articles on health and climate change,
 2107 published in English between 2007 and 2016, was conducted; an appropriate approach for broad
 2108 and inter-disciplinary research fields.²¹³ Two databases were used, PubMed and Web of Science, to
 2109 identify papers through a bibliometric analysis using keyword searches (see Appendix 6 for
 2110 details).²¹⁴ Inclusion and exclusion criteria were applied to capture the most relevant literature on
 2111 the human health impacts of climate change within the chosen timeframe and papers were
 2112 independently reviewed and screened three times to identify relevant publications.²¹⁵

2113

2114 [Indicator 5.3: Health and climate change in the United Nations General Assembly](#)
2115 **Headline Finding:** *There is no overall trend in United Nations General Debate (UNGD) references to*
2116 *health and climate change, but two significant peaks occurred in 2009 and 2014.*

2117 The General Debate (GD) takes place every September at the start of each new session of the United
2118 Nations General Assembly (UNGA). Governments use their annual statements to present their
2119 perspective on events and issues they consider the most important in global politics, and to call for
2120 greater action from the international community. All UN Member States can address the UNGA, free
2121 from external constraints. Therefore, GD statements provide an ideal data source on political
2122 engagement with health and climate change, which is comparable spatially and temporally. This
2123 indicator focuses on the extent to which governments refer to linkages between health and climate
2124 change issues in their annual statements in the GD, with one reference representing one ‘hit’.

2125 Health and climate change are issues frequently raised in UNGD statements (see Figures S5.3-S5.5 in
2126 Appendix 6). However, statements less frequently link health and climate change together. Between
2127 2007 and 2016, linked references to health and climate change in the annual UNGD ranged from 44
2128 to 124 (Figure 5.3). The comparable figures for references to climate change alone were 378 and
2129 989. It was found that there is no overall trend in conjoint references to health and climate change
2130 across the period.




2131
2132 Figure 5.3 Political engagement with the intersection of health and climate change, represented by joint
2133 references to health and climate change in the UNGD.

2134
2135 While no overall trend is apparent, there are two distinct peaks between 2009 and 2011 and in
2136 2014. In both 2009 and 2014, there were 124 references linking health and climate change in the GD
2137 statements. The 2009 peak occurred after the 2008 World Health Day, which focussed on health and
2138 climate change, and in the build-up to COP15 in Copenhagen in 2009. The 2014 peak is indicative of
2139 the influence of the large UNGA on climate change in 2014 and the lead up to COP21 in Paris in
2140 2015.

2141 The 2015 UNGA, which focused on the Sustainable Development Goals, made relatively limited
 2142 reference to climate change, and, after the 2014 peak, conjoint references to health and climate
 2143 change declined. This irregular pattern points to the importance of key events in the global
 2144 governance of health and climate change in driving high-level political engagement.

2145 There are country-level differences in the attention given to health and climate change in UNGD
 2146 statements (Figure 5.4). More frequent reference is made to the issue by countries in the Western
 2147 Pacific, particularly by the SIDS in these regions. In contrast, governments in the East Mediterranean,
 2148 the Americas and South-East Asia tend to make fewer references to health and climate change.



2149
 2150 Figure 5.4 Regional political engagement with the intersection of health and climate change, represented by
 2151 joint references to health and climate change in the UNGD, broken down by WHO region.

2152
 2153 This indicator is based on the application of keyword searches in the text corpus of debates. A new
 2154 dataset of GD statements was used (UNGD corpus), in which the annual UNGD statements have
 2155 been pre-processed and prepared for use in quantitative text analysis (see Appendix 6 for details).²¹⁶

2156
 2157 **Conclusion**

2158 The indicators in this section have demonstrated the importance of global governance in mobilising
 2159 public and political engagement in health and climate change. The UN (and particularly the annual
 2160 COPs) have a significant role here, clearly influencing media, scientific and political engagement with
 2161 health and climate change.

2162 To further improve understanding of public and political engagement, indicators relating to national
 2163 governments' health and climate change legislation, private sector engagement, the inclusion of
 2164 climate change in professional health education, and the prominence given to health in UNFCCC
 2165 negotiations are proposed for future analysis. The previous sections in this report have presented
 2166 findings on the impacts of climate hazards, adaptation and resilience, co-benefits of mitigation, and

2167 finance and economics. All of these hinge upon policy, which in turn is dependent upon public and
2168 political engagement.

2169 Conclusion - the Lancet Countdown in 2017

2170 In June 2015, the Lancet Commission laid the groundwork for its global monitoring platform,
2171 designed to systematically track progress on health and climate change, and hold governments to
2172 account for their commitments under the then to-be-finalised Paris Agreement.⁴ The Lancet
2173 Countdown will continue this work, reporting annually on the indicators presented in this report and
2174 on new indicators in future.

2175

2176 The direction of travel is set

2177 The data and analysis presented in this 2017 report cover a wide range of topics and themes from
2178 the lethality of weather-related disasters, to the phase-out of coal-fired power. The report begins
2179 with an indicator set dedicated to tracking the health effects of climate change and climate hazards.
2180 The analysis here demonstrates that the symptoms of climate change have been clear for a number
2181 of years, with the health impacts far worse than previously understood. These effects have been
2182 spread unequally, with a 9.4% increase in vectorial capacity of the dengue fever carrying *Aedes*
2183 *aegypti* predominantly spreading to low- and middle-income countries since 1950; and India
2184 disproportionately affected by the additional 75 million exposure events to potentially fatal
2185 heatwaves since 2000.

2186 These indicators also suggest that populations are beginning to adapt, with improvements in the
2187 world's overall health profile strengthening its resilient capacity, and national governments
2188 beginning to invest in health adaptation planning for climate change. This is supported by some
2189 \$47.29 billion USD spent annually on health-related adaptation (some 13.3% of global total
2190 adaptation spend). However, the academic literature and past experience make it clear that there
2191 are very real and immediate technological, financial, and political barriers to adaptation.¹⁰

2192 The indicators in the third section track health-relevant mitigation trends across four sectors, with an
2193 ultimate focus of keeping temperature rise "well below 2°C" and meeting the Paris Agreement. At an
2194 aggregate level, the past two decades have seen limited progress here, with many of the trends and
2195 indicators remaining flat or moving strongly in the opposite direction. More recently, trends in the
2196 electricity generation (deployment of renewable energy and a dramatic slow-down in coal-fired
2197 power) and transport sectors (soon-to-be cost parity of electric vehicles with their petrol-based
2198 equivalents) provide cause for optimism, which, if sustained, could reflect the beginning of system-
2199 wide transformation.

2200 Indicators in the fourth and fifth sections underpin and drive forward this transition. Again, trends
2201 across the last two decades reflect concerning levels of inaction, with accelerated investment and
2202 intervention seen in more recent years. They reflect record levels of employment in the renewable
2203 energy sector to overtake those in fossil fuel extraction, and a global reduction in fossil fuel
2204 consumption subsidies. Carbon pricing mechanisms are slowly widening and now cover some 13.1%
2205 of global CO₂ emissions. The final section considers the degree to which the public, political and
2206 academic communities have engaged with the links between climate change and health. It points to
2207 uneven patterns of engagement and the vital role of global institutions, and the UN particularly, in
2208 driving forward public, political and scientific support for enhanced mitigation and adaptation
2209 policies.

2210 Overall, the trends elucidated in the Lancet Countdown's 2017 report provide cause for deep
2211 concern, highlighting the immediate health threats from climate change and the relative inaction
2212 seen across the world over the past two decades. However, they also point to more recent trends

2213 over the last five years demonstrating a rapid increase in action, which was solidified in the Paris
2214 Agreement. These ‘glimmers of progress’ are encouraging, and reflect a growing political consensus
2215 and ambition, which was seen in full-force in response to the US’s departure from the 2015 climate
2216 change treaty. Whilst action needs to increase rapidly, taken together, this provides the clearest
2217 signal to-date that the world is beginning to transition to a low-carbon world, that no one country or
2218 head of state can halt this progress, and that from today until 2030, the direction of travel is set.

2219

2220

2221 **Contributors**

2222 The Lancet Countdown: Tracking Progress on Health and Climate Change is an international
2223 academic collaboration which builds off the work of the 2015 Lancet Commission on Health and
2224 Climate Change, convened by The Lancet. The Lancet Countdown’s work for this paper was
2225 conducted by its five working groups, each of which were responsible for the design, drafting, and
2226 review of their individual indicators and sections. All authors contributed to the overall paper
2227 structure and concepts, and provided input and expertise to the relevant sections. Authors
2228 contributing to Working Group 1: Jonathan Chambers; Peter M Cox; Mostafa Ghanei; Ilan Kelman; Lu
2229 Liang; Ali Mohammad Latifi; Maziar Moradi-Lakeh; Kris Murray; Fereidoon Owfi; Mahnaz Rabbaniha;
2230 Elizabeth Robinson; Meisam Tabatabaei. Authors contributing to Working Group 2: Sonja Ayeb-
2231 Karlsson; Peter Byass; Diarmid Campbell-Lendrum; Michael Depledge; , Paula Dominguez-Salas;
2232 Howard Frumkin; Lucien Georgeson; Delia Grace; Anne Johnson; Dominic Kniveton; Georgina Mace;
2233 Maquins Odhiambo Sewe; Mark Maslin; Maria Nilsson; Tara Neville; Karyn Morrissey; Joacim
2234 Rocklöv; Joy Shumake-Guillemot. Authors contributing to Working Group 3: Markus Amann; Kristine
2235 Belesova; Wenjia Cai; Michael Davies; Andy Haines; Ian Hamilton; Stella Hartinger; Gregor
2236 Kiesewetter; Melissa Lott, Robert Lowe; James Milner; Tadj Oreszczyn; David Pencheon, Steve Pye;
2237 Rebecca Steinbach; Paul Wilkinson. Authors contributing to Working Group 4: Timothy Bouley; Paul
2238 Drummond; Paul Ekins. Authors Contributing to Working Group 5: Maxwell Boykoff; Meaghan Daly;
2239 Niheer Dasandi; Anneliese Depoux; Antoine Flahault; Hilary Graham; Rébecca Grojsman; Slava
2240 Mikhaylov; Stefanie Schütte. The coordination, strategic direction, and editorial support for this
2241 paper was provided by Anthony Costello (Co-Chair), Hugh Montgomery (Co-Chair), Peng Gong (Co-
2242 Chair), Nick Watts (Executive Director), and Nicola Wheeler (Programme Officer). The findings and
2243 conclusions in this article are those of the authors and do not necessarily represent the official
2244 position of World Health Organization, the World Bank, or the World Meteorological Organization.

2245

2246 **Declarations of Interest**

2247 The Lancet Countdown’s work is supported by an unrestricted grant from the Wellcome Trust (ref:
2248 200890/Z/16/Z)). The Lancet Countdown covered travel costs for meetings related to the
2249 development of the paper. Seven of the authors (NWA, NWH, ML, PD, MB, MD and JC) were
2250 compensated for their time while working on the Lancet Countdown’s drafting and development.
2251 HM is a board member of the UK Climate and Health Council, an Advisory Board member of the
2252 Energy and Climate Intelligence Unit, and is developing an air pollution mask (which represents no
2253 conflict of interest). NWA reports being the Director of the UK Health Alliance on Climate Change. AJ
2254 is a Governor of the Wellcome Trust, and a member of the Adaptation Sub-Committee of the
2255 Committee on Climate Change. MA, SAK, KB, TB, PB, WC, DCL, AC, PC, ND, MDa, MDe, AD, PDS, PE,
2256 AF, HF, LG, MG, PG, DG, HG, RG, AH, IH, SH, IK, GK, DK, LL, RL, GM, MM, SM, JM, AML, MML, KMo,

2257 KMu, TN, MN, TO, FO, DP, SP, MR, ER, JR, SS, MS, JSG, RS, MT, and PW declare no conflicts of
2258 interest.

2259

2260 **Acknowledgements**

2261 The Lancet Countdown would like to thank the Wellcome Trust, in particular Saskia Heijnen, Sarah
2262 Molten and Sophie Tunstall-Behrens, for its financial and strategic support—without which, this
2263 research collaboration would not be possible. While carrying out its work, the Lancet Countdown
2264 received invaluable technical advice and input from a number of individuals, including Neil Adger
2265 (University of Exeter), Kevin Andrews (University of Colorado Boulder), Nigel Arnell (University of
2266 Reading), Rob Bailey (Chatham House), John Balbus (National Institute of Environmental Health
2267 Sciences), Simon Bennet (International Energy Agency), Helen Berry (Australiana National
2268 University), Kathryn Brown (Climate Change Committee), Yossi Cadan (350.org), Tony Capon
2269 (University of Sydney), Carbon Disclosure Project (CDP), Michelle Chan (Universidad Peruana
2270 Cayetano Heredia), Lucia Fernandez (World Health Organization), Lauren Gifford (University of
2271 Colorado Boulder), Francesca Harris (London School of Hygiene & Tropical Medicine), Mathieu
2272 Hemon (Centre Virchow-Villermé), Niamh Herlihy (Centre Virchow-Villermé), Richard King
2273 (Chatham House), Tord Kjellstrom (Australian National University), Noemie Klein (Ecofys), Long Lam
2274 (Ecofys), Seline Lo (The Lancet), Rachel Lowe (London School of Hygiene & Tropical Medicine), Gesa
2275 Luedecke (University of Colorado Boulder), Lucy McAllister (University of Colorado Boulder), Marisa
2276 McNatt (University of Colorado Boulder), Jonathan Patz (University of Wisconsin-Madison), Sonia
2277 Roschnik (Sustainable Health Solutions), Osman Sankoh (INDEPTH), Ami Nacu-Schmidt (University of
2278 Colorado Boulder), Pauline Scheelbeek (London School of Hygiene & Tropical Medicine), Jan
2279 Semenza (European Centre for Disease Prevention and Control), Imogen Tennison (National Health
2280 Service), Hanna Tuomisto (London School of Hygiene and Tropical Medicine), Armando Valdes
2281 Valasquez (Universidad Peruana Cayetano Heredia) and Shelagh Whitley (Overseas Development
2282 Institute). Administrative and communications support was provided by Richard Black (Energy and
2283 Climate Intelligence Unit), Pete Chalkley (Energy and Climate Intelligence Unit), Tan Copsey (Climate
2284 Nexus), Tom Fern, Jack Fisher (University College London), Sarah Hurtes (European Climate
2285 Foundation), Paige Knappenberger (Climate Nexus) and George Smeeton (Energy and Climate
2286 Intelligence Unit). Mr Georgeson wishes to express gratitude for funding from the Economic and
2287 Social Research Council and the Natural Environment Research Council (grant number
2288 ES/J500185/1).

2289 The Lancet Countdown is funded through an unrestricted grant from the Wellcome Trust
2290 (200890/Z/16/Z).

2291

2292

2293

2294

2295 **References**

2296

- 2297 1. Kang Y, Khan S, Ma X. Climate change impacts on crop yield, crop water productivity
2298 and food security – A review. *Progress in Natural Science* 2009; **19**(12): 1665–74.
- 2299 2. Lindgren E, Andersson, Y, Suk, J.E, Sudre, B. and Semenza, J.C. Monitoring EU
2300 Emerging Infectious Disease Risk Due to Climate Change. *Science* 2012; **336**(6080): 418-9.
- 2301 3. Reuveny R. Climate change-induced migration and violent conflict. *Political*
2302 *Geography* 2007; **26**(6): 656–73.
- 2303 4. Watts N, et al. Health and climate change: policy responses to protect public health.
2304 *The Lancet* 2015; **386**(10006): 1861–914.
- 2305 5. Watts N, et al. The Lancet Countdown: tracking progress on health and climate
2306 change. *The Lancet* 2016; **389**(10074): 1151–64.
- 2307 6. World Health Organization. Second Global Conference: Health and Climate (Paris 7-8
2308 July 2016), 2016.
- 2309 7. United Nations Framework Convention on Change. Paris Agreement. In: United
2310 Nations, editor. Paris, France; 2015.
- 2311 8. United Nations Environment Program. The Emissions Gap Report 2016. In: United
2312 Nations, editor. Nairobi, Kenya; 2016.
- 2313 9. Intergovernmental Panel on Climate Change. Climate Change 2014: Impacts,
2314 Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working
2315 Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.
2316 Cambridge, United Kingdom and New York, NY, USA, 2014.
- 2317 10. Smith KR, Woodward A, Campbell-Lendrum D, et al. Human health: Impacts,
2318 adaptation, and co-benefits. In: Field CB, Barros VR, Dokken DJ, et al., eds. Climate Change
2319 2014: Impacts, Adaptation, and Vulnerability Part A: Global and Sectoral Aspects
2320 Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental
2321 Panel of Climate Change. Cambridge and New York: Cambridge University Press; 2014: 709-
2322 54.
- 2323 11. Berry HL, Bowen K, Kjellstrom T. Climate change and mental health: a causal
2324 pathways framework. *International Journal of Public Health* 2010; **55**(2): 123-32.
- 2325 12. Reinmuth-Selzle K, et al. Air Pollution and Climate Change Effects on Allergies in the
2326 Anthropocene: Abundance, Interaction, and Modification of Allergens and Adjuvants.
2327 *Environmental Science and Technology* 2017; **51**(8): 4119-41.
- 2328 13. Glaser J, et al. Climate Change and the Emergent Epidemic of CKD from Heat Stress
2329 in Rural Communities: The Case for Heat Stress Nephropathy. *Clinical Journal of the*
2330 *American Society of Nephrology* 2016; **11**(8): 1472–83.
- 2331 14. McMichael AJ. Globalization, Climate Change, and Human Health. *The New England*
2332 *Journal of Medicine* 2013; **368**: 1335-43.

- 2333 15. European Centre for Medium-Range Weather Forecasts (ECMWF). 2017.
2334 <https://www.ecmwf.int/>.
- 2335 16. European Centre for Medium-Range Weather Forecasts (ECMWF). Climate
2336 Reanalysis. 2017. <https://www.ecmwf.int/en/research/climate-reanalysis>.
- 2337 17. NASA. Gridded Population of the World (GPW), v4. 2017.
- 2338 18. Jacob D, Petersen J, Eggert B, et al. EURO-CORDEX: new high-resolution climate
2339 change projections for European impact research. *Regional Environmental Change* 2014;
2340 **14**(2): 563-78.
- 2341 19. HEAT-SHIELD. 2017. <https://www.heat-shield.eu/>.
- 2342 20. Kjellstrom T, Briggs D, Freyberg C, Lemke B, Otto M, Hyatt O. Heat, Human
2343 Performance, and Occupational Health: A Key Issue for the Assessment of Global Climate
2344 Change Impacts. *Annual Review of Public Health* 2016; **37**: 97–112.
- 2345 21. Ijaz K, Kasowski E, Arthur RR, Angulo FJ, Dowell SF. International Health
2346 Regulations—What Gets Measured Gets Done. *Emerging Infectious Diseases* 2012; **18**(7):
2347 1054-7.
- 2348 22. Intergovernmental Panel on Climate Change. Climate Change 2014: Synthesis
2349 Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the
2350 Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A.
2351 Meyer (eds.)]. Geneva, Switzerland: IPCC, 2014.
- 2352 23. Emergency Events Database. The Human Cost of Weather-Related Disasters 1995-
2353 2015: Centre for Research on the Epidemiology of Disasters (CRED), 2015
- 2354 24. Emergency Events Database. The International Disaster Database - Centre for
2355 Research on the Epidemiology of Disasters. 2017.
- 2356 25. International Federation of Red Cross and Red Crescent Societies. World Disasters
2357 Report 2014: Focus on Culture and Risks, 2014.
- 2358 26. Centre for Research on the Epidemiology of Disasters. The Human Cost of Natural
2359 Disasters: A Global Perspective. Brussels: CRED (Centre for Research on the Epidemiology of
2360 Disasters), 2015.
- 2361 27. Global Burden of Disease. Global Burden of Disease Study 2015. Global Burden of
2362 Disease Study 2015 (GBD 2015) Results. In: Institute for Health Metrics and Evaluation
2363 (IHME), editor. Seattle, United States; 2016.
- 2364 28. Liyanage P, Tissera H, Sewe M, et al. A Spatial Hierarchical Analysis of the Temporal
2365 Influences of the El Nino-Southern Oscillation and Weather on Dengue in Kalutara District,
2366 Sri Lanka. *International journal of environmental research and public health* 2016; **13**(11).
- 2367 29. Wilder-Smith A, Byass P. The elusive global burden of dengue. *The Lancet Infectious*
2368 *diseases* 2016; **16**(6): 629-31.

- 2369 30. Mitchell D, Heaviside C, Vardoulakis S, et al. Attributing human mortality during
2370 extreme heat waves to anthropogenic climate change. *Environmental Research Letters*
2371 2016; **11**(7): 074006.
- 2372 31. Zanobetti A, Schwartz J. Temperature and mortality in nine US cities. *Epidemiology*
2373 (*Cambridge, Mass*) 2008; **19**(4): 563-70.
- 2374 32. Gasparrini A, Guo Y, Hashizume M, et al. Temporal Variation in Heat-Mortality
2375 Associations: A Multicountry Study. *Environmental health perspectives* 2015; **123**(11): 1200-
2376 7.
- 2377 33. Shi L, Kloog I, Zanobetti A, Liu P, Schwartz JD. Impacts of Temperature and its
2378 Variability on Mortality in New England. *Nature climate change* 2015; **5**: 988-91.
- 2379 34. Guo Y, Gasparrini A, Armstrong BG, et al. Temperature Variability and Mortality: A
2380 Multi-Country Study. *Environmental health perspectives* 2016.
- 2381 35. Wang Y, Shi L, Zanobetti A, Schwartz JD. Estimating and projecting the effect of cold
2382 waves on mortality in 209 US cities. *Environment international* 2016; **94**: 141-9.
- 2383 36. Murray CJL, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries
2384 in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study
2385 2010. *The Lancet* 2013; **380**(9859): 2197–223.
- 2386 37. Dye C. After 2015: infectious diseases in a new era of health and development. *Royal*
2387 *Philosophical Transactions of The Royal Society B* 2014; **369**(1645).
- 2388 38. World Health Organization. Using climate to predict infectious disease epidemics.
2389 Geneva, Switzerland, 2005.
- 2390 39. World Health Organization. The Health and Environment Lexicon: Multi-Language
2391 Glossary of Health and Environment Terminology. 2017.
2392 <http://apps.who.int/thelexicon/entry.php>.
- 2393 40. Stanaway JD, et al. The global burden of dengue: an analysis from the Global Burden
2394 of Disease Study 2013. *The Lancet: Infectious Diseases* 2016; **16**(6): 712–23.
- 2395 41. Hales S, de Wet, N, Maindonald, J. and Woodward, A. Potential effect of population
2396 and climate changes on global distribution of dengue fever: an empirical model. *The Lancet*
2397 2002; **360**(9336): 830-4.
- 2398 42. Krishnamurthy PK, et al. A methodological framework for rapidly assessing the
2399 impacts of climate risk on national-level food security through a vulnerability index. *Global*
2400 *Environmental Change* 2014; **24**: 121–32.
- 2401 43. Nelson GC, Rosegrant MW, Palazzo A, et al. Food Security Farming and Climate
2402 Change to 2050: scenarios, results, policy options: International Food Policy Research
2403 Institute, 2010.

- 2404 44. Schmidhuber J, Tubiello FN. Global food security under climate change. *Proceedings*
2405 *of the National Academy of Sciences of the United States of America* 2007; **104**(50): 19703-8.
- 2406 45. Campbell-Lendrum D, Woodruff R. Comparative risk assessment of the burden of
2407 disease from climate change. *Environmental health perspectives* 2006; **114**: 1935-41.
- 2408 46. Campbell-Lendrum DH, Corvalán CF, Prüss-Ustün A. How Much Disease Could
2409 Climate Change Cause? Geneva: The World Health Organization, 2003.
- 2410 47. Naylor RL, Falcon WP. Food security in an era of economic volatility. *Population and*
2411 *development review* 2010; **36**(4): 693-723.
- 2412 48. Headey D, Fan S. Reflections on the global food crisis: how did it happen? how has it
2413 hurt? and how can we prevent the next one?, 2010.
- 2414 49. Asseng S, Ewert F, Martre P, et al. Rising temperatures reduce global wheat
2415 production. *Nature climate change* 2015; **5**(2): 143-7.
- 2416 50. Peng S, Huang, J., Sheehy, J.E., Laza, R.C., Visperas, R.M., Zhong, X., Centeno, G.S.,
2417 Khush, G.S. and Cassman, K.G. Rice yields decline with higher night temperature from global
2418 warming. *Proceedings of the National academy of Sciences of the United States of America*
2419 2004; **101**(27): 9971-5.
- 2420 51. Lobell DB, Burke, M.B., Tebaldi, C., Mastrandrea, M.D., Falcon, W.P. and Naylor, R.L.
2421 Prioritizing climate change adaptation needs for food security in 2030. *Science* 2007;
2422 **319**(5863): 607-10.
- 2423 52. Lobell DB, Schlenker, W. and Costa-Roberts, J. Climate trends and global crop
2424 production since 1980. *Science* 2011; **333**(6042): 616-20.
- 2425 53. Gornall J, et al. Implications of climate change for agricultural productivity in the
2426 early twenty-first century. *Philosophical Transactions of the Royal Society B Biological*
2427 *Sciences* 2010; **365**(1554).
- 2428 54. FAO. The FAO Hunger Map 2015. In: Food and Agriculture Organization of the United
2429 Nations, editor.; 2015.
- 2430 55. Jones PG, Thornton PK. The potential impacts of climate change on maize production
2431 in Africa and Latin America in 2055. *Global Environmental Change* 2003; **13**(1): 51-9.
- 2432 56. Lobell DB, Bänziger M, Magorokosho C, Vivek B. Nonlinear heat effects on African
2433 maize as evidenced by historical yield trials. *Nature climate change* 2011; **1**(1): 42-5.
- 2434 57. High Level Panel of Experts on Food Security and Nutrition of the Committee on
2435 World Food Security. Sustainable Fisheries and Aquaculture for Food Security and Nutrition.
2436 Rome, Italy, 2014.
- 2437 58. World Health Organization. Availability and consumption of fish. 2017.
2438 http://www.who.int/nutrition/topics/3_foodconsumption/en/index5.html (accessed July
2439 2017.

- 2440 59. Djoue L, et al. Fish consumption, omega-3 fatty acids and risk of heart failure: a
2441 meta-analysis. *Clinical Nutrition* 2012; **31**(6): 846-53.
- 2442 60. Golden Cea. Fall in fish catch threatens human health. *Nature* 2016; **534**(317).
- 2443 61. Bushaw-Newton KL, Sellner KG. Harmful Algal Blooms: National Oceanic and
2444 Atmospheric Administration, 1999.
- 2445 62. Armstrong EM, Vazquez-Cuervo J. A New Global Satellite-Based Sea Surface
2446 Temperature Climatology. *Geophysical Research Letters* 2001; **28**(22): 4199-202.
- 2447 63. International Organization for Migration. Migration and Climate Change, 2008.
- 2448 64. Gleick PH. Water, Drought, Climate Change, and Conflict in Syria. *American
2449 Meteorological Society* 2014.
- 2450 65. Kelley CP, Mohtadib S, Canec MA, Seagerc R, Kushnirc Y. Climate change in the
2451 Fertile Crescent and implications of the recent Syrian drought. *Proceedings of the National
2452 Academy of Sciences of the United States of America* 2015; **112**(11): 3241–6.
- 2453 66. McMichael C, Barnett J, McMichael AJ. An Ill Wind? Climate Change, Migration, and
2454 Health. *Environmental health perspectives* 2012; **120**(5): 646-54.
- 2455 67. Green M. Contested territory. *Nature climate change* 2016; **6**: 817–20.
- 2456 68. United Nations Office for the Coordination of Humanitarian Affairs. Fiji: Building
2457 resilience in the face of climate change, 2014.
- 2458 69. EU-GIZ Adapting to Climate Change and Sustainable Energy Programme. Planned
2459 Relocation Project. 2017.
- 2460 70. Connell J. Last days in the Carteret Islands? Climate change, livelihoods and
2461 migration on coral atolls. *Asia Pacific Viewpoint* 2016; **57**(1): 3-15.
- 2462 71. Strauss S. Are cultures endangered by climate change? Yes, but ... *WIREs Climate
2463 Change* 2012; **3**(4): 371-7.
- 2464 72. Bronen R, Chapin I. Adaptive governance and institutional strategies for climate-
2465 induced community relocations in Alaska. *PNAS, Proceedings of the National Academy of
2466 Sciences* 2013; **110**(23): 9320-5.
- 2467 73. Shearer C. The political ecology of climate adaptation assistance: Alaska Natives,
2468 displacement, and relocation. *Journal of Political Ecology* 2012; **19**: 174-83.
- 2469 74. Small C, Nicholls RJ. A Global Analysis of Human Settlement in Coastal Zones.
2470 *Journal of Coastal Research* 2003; **19**(3): 584-99.
- 2471 75. McGranahan G, Balk, D. and Anderson, B. The rising tide: assessing the risks of
2472 climate change and human settlements in low elevation coastal zones. *Environment &
2473 Urbanization* 2007; **19**(1): 17-37.

- 2474 76. Merkens JL, Reimann, L, Hinkel, J. and Vafeidis, A.T. Gridded population projections
2475 for the coastal zone under the Shared Socioeconomic Pathways. *Global and Planetary*
2476 *Change* 2016; **145**: 57-66.
- 2477 77. Gregory J. Projections of sea level rise: Working Group I contribution to the IPCC
2478 Fifth Assessment Report - Climate Change 2013: The Physical Science Basis, 2013.
- 2479 78. Collins PY, et al. Grand challenges in global mental health: A consortium of
2480 researchers, advocates and clinicians announces here research priorities for improving the
2481 lives of people with mental illness around the world, and calls for urgent action and
2482 investment. *Nature* 2011; **475**: 27-30.
- 2483 79. Vins H, Bell J, Saha S, Hess J. The mental health outcomes of drought: A systematic
2484 review and causal process diagram. *International journal of environmental research and*
2485 *public health* 2015; **12**: 13251.
- 2486 80. Intergovernmental Panel on Climate Change. Climate Change 2007: Working Group
2487 II: Impacts, Adaptation and Vulnerability, 2001.
- 2488 81. Rockefeller Foundation. Resilience. 2017.
2489 <https://www.rockefellerfoundation.org/our-work/topics/resilience/> (accessed 02 Jul 2017).
- 2490 82. United Nations Framework Convention on Change. National Adaptation Plans. 2017.
- 2491 83. World Health Organization. WHO guidance to protect health from climate change
2492 through health adaptation planning. Switzerland, 2014.
- 2493 84. World Health Organization. Operational framework for building climate resilient
2494 health systems. Geneva, Switzerland, 2015.
- 2495 85. United Nations. The World's Cities in 2016, 2016.
- 2496 86. Doherty M, Klima K, Hellmann JJ. Climate change in the urban environment:
2497 Advancing, measuring and achieving resiliency. *Environmental Science & Policy* 2016; **66**:
2498 310-3.
- 2499 87. Compact of Mayors. 2017. <https://www.compactofmayors.org/>.
- 2500 88. Carbon Disclosure Project. Carbon Disclosure Project Data. 2017.
- 2501 89. Carbon Disclosure Project. CDP Cities 2016 Information Request, 2016.
- 2502 90. Sustainable Development Knowledge Platform. Sustainable Development Goal 3 -
2503 Ensure healthy lives and promote well-being for all at all ages. 2017.
2504 <https://sustainabledevelopment.un.org/sdg3>.
- 2505 91. World Health Organization. International Health Regulations (2005) Second Edition,
2506 2008.
- 2507 92. World Health Assembly. World Health Assembly resolution WHA 62.1 2008.

- 2508 93. International Health Regulations. IHR Core Capacity Monitoring Framework:
2509 Questionnaire for monitoring progress in the implementation of IHR core capacities in states
2510 parties: World Health Organization, 2005.
- 2511 94. World Health Organization. International Health Regulations (2005). IHR Core
2512 Capacity Monitoring Framework: Questionnaire for Monitoring Progress in the
2513 Implementation of IHR Core Capacities in States Parties, 2017.
- 2514 95. World Meteorological Organization. Monitoring and Evaluation. 2017.
- 2515 96. World Health Organization. Protecting Health from Climate Change: Vulnerability
2516 and Adaptation Assessment, 2013.
- 2517 97. Lim SS, Allen K, Bhutta ZA, et al. Measuring the health-related Sustainable
2518 Development Goals in 188 countries: a baseline analysis from the Global Burden of Disease
2519 Study 2015. *The Lancet* 2016; **16**: 31467-2.
- 2520 98. Ministry of Economic Affairs and Employment. National Energy and Climate Strategy.
2521 In: Government of Finland, editor.; 2016.
- 2522 99. Department for Business Energy and Industrial Strategy. Coal Generation in Great
2523 Britain. The pathway to a low-carbon future: consultation document In: HM Government,
2524 editor.; 2016.
- 2525 100. Mason J. In latest move, China halts over 100 coal power projects. 2017.
2526 <http://uk.reuters.com/article/us-china-coal-idUKKBN151090> (accessed July 2017).
- 2527 101. UBS. UBS Evidence Lab Electric Car Teardown – Disruption Ahead?, 2017.
- 2528 102. International Energy Agency. Medium-Term Renewable Energy Market Report 2016,
2529 2016.
- 2530 103. Dezem V. Solar Sold in Chile at Lowest Ever, Half Price of Coal. 2016.
2531 [https://www.bloomberg.com/news/articles/2016-08-19/solar-sells-in-chile-for-cheapest-](https://www.bloomberg.com/news/articles/2016-08-19/solar-sells-in-chile-for-cheapest-ever-at-half-the-price-of-coal)
2532 [ever-at-half-the-price-of-coal.](https://www.bloomberg.com/news/articles/2016-08-19/solar-sells-in-chile-for-cheapest-ever-at-half-the-price-of-coal)
- 2533 104. Le Quéré C, Andrew, R.M, Canadell, J.G, et al. Global Carbon Budget 2016. *Earth*
2534 *System Science Data* 2016; **8**: 605-49.
- 2535 105. Schirnding YV. Framework for Linkages between Health, Environment and
2536 Development, 2002.
- 2537 106. Intergovernmental Panel on Climate Change. Climate Change 2014: Mitigation of
2538 Climate Change. Contribution of Working Group III to the Fifth Assessment
2539 Report of the Intergovernmental Panel on Climate Change. Cambridge, United Kingdom and
2540 New York, NY, USA, 2014.
- 2541 107. International Energy Agency. Energy and Climate Change, 2015.

- 2542 108. Rogelj J, Schaeffer, M, Meinshausen, M, et al. Zero emission targets as long-term
2543 global goals for climate protection. *Environmental Research Letters* 2015; **10**(105007).
- 2544 109. Pye S, Li FGN, Price J, Fais B. Achieving net-zero emissions through the reframing of
2545 UK national targets in the post-Paris Agreement era. *Nature Energy* 2017; **2**.
- 2546 110. Rockström J, Gaffney, O, Rogelj, J, Meinshausen, M, Nakicenovic, N. and
2547 Schellnhuber, H.J. A roadmap for rapid decarbonization. *Science* 2017; **80**(355): 1269–71.
- 2548 111. International Energy Agency. World Energy Outlook. Paris, France, 2016.
- 2549 112. International Energy Agency. Energy and Air Pollution: World Energy Outlook Special
2550 Report. Paris, France, 2016.
- 2551 113. Green F, Stern N. China’s changing economy: implications for its carbon dioxide
2552 emissions. *Climate Policy* 2017; **17**: 423–42.
- 2553 114. Shearer C, Ghio, N, Myllyvirta, L, Yu, A. and Nace, T. Boom and Bust 2017 - Tracking
2554 the global coal plant pipeline, 2017.
- 2555 115. IRENA. Conference on the establishment of the international renewable energy
2556 agency. Bonn, Germany, 2009.
- 2557 116. World Health Organization. The European health report 2015: Targets and beyond –
2558 reaching new frontiers in evidence, 2015.
- 2559 117. World Health Organization. Proportion of population with primary reliance on clean
2560 fuels and technology. Geneva, Switzerland, 2017.
- 2561 118. International Energy Agency, World Bank. Sustainable Energy for All 2017—Progress
2562 toward Sustainable Energy. Washington, D.C., 2017.
- 2563 119. World Energy Outlook. WEO 2016 Biomass Database. 2016.
- 2564 120. World Energy Outlook. WEO 2016 Electricity Access Database. 2016.
- 2565 121. World Health Organization. Household Fuel Combustion: WHO guidelines for indoor
2566 air quality, 2014.
- 2567 122. Bonjour S, Adair-Rohani, H, Wolf, J. et al. Solid Fuel Use for Household Cooking:
2568 Country and Regional Estimates for 1980–2010. *Environmental health perspectives* 2013;
2569 **121**: 784–90.
- 2570 123. West JJ, Smith, S.J, Silva, R.A. et al. Co-benefits of Global Greenhouse Gas Mitigation
2571 for Future Air Quality and Human Health. *Nature climate change* 2013; **3**: 885–9.
- 2572 124. Institute for Health Metrics and Global Burden of Disease. State of Global Air - 2017:
2573 A special report on global exposure to air pollution and its disease burden, 2017.

- 2574 125. World Health Organization. WHO's Urban Ambient Air Pollution database – Update
2575 2016. Geneva, Switzerland, 2017.
- 2576 126. Milner J, Taylor, J, Barreto, M.L. et al. Environmental risks of cities in the European
2577 region: analyses of the Sustainable Healthy Urban Environments (SHUE) database. *Public*
2578 *Health Panorama* 2017; **3**: 300-9.
- 2579 127. World Health Organization. Air pollution levels rising in many of the world's poorest
2580 cities, 2016.
- 2581 128. European Commission. Europe, Latin America and The Caribbean: sharing
2582 experiences in regional development policies, 2015.
- 2583 129. Instituto Nacional de Estadística e Informática. 2017.
- 2584 130. Fondo De Inclusion Social Energetico. Memoria Anual De Gestion Fise 2015, 2015.
- 2585 131. Fondo De Inclusion Social Energetico. Memoria Anual De Gestion Fise 2014, 2014.
- 2586 132. European Commission. Directive 2001/80/EC of the European Parliament and of the
2587 Council on the limitation of emissions of certain pollutants into the air from large
2588 combustion plants. OJEC L 309/1. 2001.
- 2589 133. EU Parliament and Council. Directive 2010/75/EU of the European Parliament and of
2590 the Council of 24 November 2010 on industrial emissions (integrated pollution prevention
2591 and control). OJEC L 334/17. 2010.
- 2592 134. Pope CAea. Lung Cancer, Cardiopulmonary Mortality and Long-Term Exposure to
2593 Fine Particulate Air Pollution. *Journal of the American Medical Association* 2002; **287** (9):
2594 1132–41.
- 2595 135. Lim SSea. A Comparative Risk Assessment of Burden of Disease and Injury
2596 Attributable to 67 Risk Factors and Risk Factor Clusters in 21 Regions, 1990–2010: A
2597 Systematic Analysis for the Global Burden of Disease Study 2010. *The Lancet* 2012;
2598 **380**(9859): 2224–60.
- 2599 136. Forouzanfar MHea. Global, Regional, and National Comparative Risk Assessment of
2600 79 Behavioural, Environmental and Occupational, and Metabolic Risks or Clusters of Risks in
2601 188 Countries, 1990–2013: A Systematic Analysis for the Global Burden of Disease Study
2602 2013. *The Lancet* 2015; **386**(10010): 2287–323.
- 2603 137. World Health Organization. Ambient Air Pollution: A Global Assessment of Exposure
2604 and Burden of Disease. Geneva, Switzerland, 2016.
- 2605 138. Amann M, I. Bertok, J. Borken-Kleefeld, J. Cofala, C. Heyes, L. Höglund-Isaksson, Z.
2606 Klimont, et al. Cost-Effective Control of Air Quality and Greenhouse Gases in Europe:
2607 Modeling and Policy Applications. *Environmental Modelling & Software* 2011; **26**(2): 1489–
2608 501.
- 2609 139. International Energy Agency. Energy Technology Perspectives. Paris, France, 2016.

- 2610 140. International Energy Agency. Global EV Outlook 2016: Beyond on million electric
2611 cars. Paris, France, 2016.
- 2612 141. International Energy Agency. Global EV Outlook 2017: Two Million and Counting,
2613 2017.
- 2614 142. Electric Vehicle Initiative. Global Electric Vehicle Outlook. 2016.
- 2615 143. Land Transport Authority. Passenger Transport Mode Shares in World Cities, 2014.
- 2616 144. Salon D, Gulyani S. Mobility, Poverty, and Gender: Travel 'Choices' of Slum Residents
2617 in Nairobi, Kenya. *Transport Reviews* 2010; **30**: 641–57.
- 2618 145. Sims R. RS, F. Creutzig, X. Cruz-Núñez, M. D'Agosto, D. Dimitriu, M.J. Figueroa Meza,
2619 L. Fulton, S., Kobayashi, O. Lah, A. McKinnon, P. Newman, M. Ouyang, J.J. Schauer, D.
2620 Sperling, and G. Tiwari. Transport, 2014.
- 2621 146. United Nations Environment Program. Global Outlook on Walking and Cycling:
2622 United Nations, 2016.
- 2623 147. Institute for Mobility Research. Mobility trends in cutting edge cities, 2016.
- 2624 148. Transport for London. Travel in London Report 9, 2016.
- 2625 149. NSW Department of Transport. Public transport travel patterns in the greater Sydney
2626 Metropolitan area 1981-1991. 1996.
- 2627 150. NSW Department of Transport. Household Travel Survey Summary Report 2002.
2628 2003.
- 2629 151. NSW Department of Transport. Household Travel Survey Summary Report 2007.
2630 2009.
- 2631 152. NSW Department of Transport. Key Transport Indicators- How do people travel.
2632 2017.
- 2633 153. Translink. Metro Vancouver Regional Trip Diary Survey Briefing Paper #1. 2012.
- 2634 154. S.A. D. Estudio Encuesta origen destino de viajes del Gran Santiago, 1991, 1992.
- 2635 155. Rode P, Hoffmann C, Kandt J, Smith D, Graff A. Toward New Urban. Mobility: The
2636 case of London and Berlin. London: London School of Economics and Political Science, 2015.
- 2637 156. City of Berlin. Mobility in the City: Berlin Traffic in Figures, 2013.
- 2638 157. Vermeulen SJ, Campbell BM, Ingram JSI. Climate change and food systems. *Annual
2639 Review of Environment and Resources* 2012; **37**: 195-222.
- 2640 158. Lim SS, Vos, T, Flaxman, A.D. et al. Global, regional, and national comparative risk
2641 assessment of 79 behavioural, environmental and occupational, and metabolic risks or

- 2642 clusters of risks in 188 countries, 1990-2013: a systematic analysis for the Global Burden of
2643 Disease Study 2013. *The Lancet* 2016; **380**: 2224–60.
- 2644 159. Springmann Mea. Analysis and valuation of the health and climate change cobenefits
2645 of dietary change. *PNAS, Proceedings of the National Academy of Sciences* 2016; **15**: 4146-
2646 51.
- 2647 160. Hawkesworth S, Dangour, A.D, Johnston, D. et al. Feeding the world healthily: the
2648 challenge of measuring the effects of agriculture on health. *Philosophical Transactions of the*
2649 *Royal Society B Biological Sciences* 2010; **365**: 3083–97.
- 2650 161. Smith MR, Micha, R, Golden, C.D. et al. Global Expanded Nutrient Supply (GENUS)
2651 Model: A New Method for Estimating the Global Dietary Supply of Nutrients. *PLoS One*
2652 2016; **11**.
- 2653 162. Gobbo LCD, Khatibzadeh, S, Imamura, F. et al. Assessing global dietary habits : a
2654 comparison of national estimates from the FAO and the Global Dietary Database 1 – 4,
2655 2015.
- 2656 163. Herrero Mea. Greenhouse gas mitigation potentials in the livestock sector. *Nature*
2657 *climate change* 2016; **6**: 452–61.
- 2658 164. O’Mara FP. The significance of livestock as a contributor to global greenhouse gas
2659 emissions today and in the near future. *Animal Feed Science and Technology* 2011; **166–167**:
2660 7–15.
- 2661 165. Herrero M, Havlík P, Valin H, et al. Biomass use, production, feed efficiencies, and
2662 greenhouse gas emissions from global livestock systems. *Proceedings of the National*
2663 *Academy of Sciences of the United States of America* 2013; **110**: 20888–93.
- 2664 166. Carlsson-Kanyama A, González AD. Potential contributions of food consumption
2665 patterns to climate change. *American Journal of Clinical Nutrition* 2009; **89**: 1704S–9S.
- 2666 167. Larsson SC, Wolk A. Meat consumption and risk of colorectal cancer: A meta-analysis
2667 of prospective studies. *International Journal of Cancer* 2006; **119**: 2657–64.
- 2668 168. Norat T, Lukanova, A, Ferrari, P. and Riboli, E. Meat consumption and colorectal
2669 cancer risk: Dose-response meta-analysis of epidemiological studies. *International Journal of*
2670 *Cancer*; **98**: 241–56.
- 2671 169. FAOSTAT. Food Balance Sheets, 2017.
- 2672 170. Ng M, Fleming T, Robinson M, al. e. Global, regional, and national prevalence of
2673 overweight and obesity in children and adults during 1980-2013: A systematic analysis for
2674 the Global Burden of Disease Study 2013. *The Lancet* 2013; **384**: 766–81.
- 2675 171. World Bank. Climate-Smart Healthcare: Low-Carbon and Resilience Strategies for the
2676 Health Sector. Washington DC, 2017.
- 2677 172. NHS Sustainable Development Unit. NHS carbon footprint, 2016.

- 2678 173. Stern N. Stern Review on the Economics of Climate Change. In: Kingdom GotU,
2679 editor. London: Blackwell Publishing; 2006. p. 5.
- 2680 174. Weitzmann. M. Fat-Tailed Uncertainty in the Economics of Catastrophic Climate
2681 Change. *Review of Environmental Economics and Policy* 2011; **5**(2): 17.
- 2682 175. Stern N. The Structure of Economic Modeling of the Potential Impacts of Climate
2683 Change: Grafting Gross Underestimation of Risk onto Already Narrow Science Models.
2684 *Journal of Economic Literature* 2013; **51**(3): 21.
- 2685 176. International Energy Agency. World Energy Investment 2016. Paris, 2016.
- 2686 177. International Energy Agency. World Energy Investment 2017. Paris, 2017.
- 2687 178. International Energy Agency. World Energy Outlook 2016. Paris, France, 2016.
- 2688 179. International Energy Agency, International Renewable Energy Agency. Perspectives
2689 for the energy transition: Investment needs for a low-carbon energy system. Berlin,
2690 Germany, 2017.
- 2691 180. Olivier JGJ, Janssens-Maenhout G, Muntean M, Peters JAHW. Trends in Global CO₂
2692 Emissions: 2016 Report: PBL Netherlands Environmental Assessment Agency, The Hague,
2693 2016.
- 2694 181. Ansar A, Caldecott B, Tilbury J. Stranded assets and the fossil fuel divestment
2695 campaign: what does divestment mean for the valuation of fossil fuel assets?: Smith School
2696 of Enterprise and the Environment, 2013.
- 2697 182. North CS. Disaster Mental Health Epidemiology: Methodological Review and
2698 Interpretation of Research Findings. *Psychiatry* 2016; **79**(2): 16.
- 2699 183. Munich Re. NatCatSERVICE. In: Re M, editor.; 2017.
- 2700 184. World Health Organization. Global Health Observatory (GHO) data. In: World Health
2701 Organization, editor. Geneva; 2017.
- 2702 185. Ming-Xiao. W, Tao. Z, Miao-Rong. X, Bin. Z, Ming-Qiu. J. Analysis Of National Coal-
2703 mining Accident data In China, 2001–2008. *Public Health Rep* 2011; **126**(2): 5.
- 2704 186. Hendryx M, Ahern MM. Relations Between Health Indicators and Residential
2705 Proximity to Coal Mining in West Virginia. *American Journal of Public Health* 2008; **98**(4): 2.
- 2706 187. Zullig. KJ, Hendryx. M. A Comparative Analysis of Health-Related Quality of Life for
2707 Residents of U.S. Counties with and without Coal Mining. *Public Health Reports* 2010;
2708 **125**(4): 7.
- 2709 188. Hendryx M. Mortality from heart, respiratory, and kidney disease in coal mining
2710 areas of Appalachia. *International Archives of Occupational and Environmental Health* 2009;
2711 **82**(2): 6.

- 2712 189. IBIS World. Global Coal Mining: Market Research Report, 2016.
- 2713 190. IBIS World. Global Oil & Gas Exploration & Production: Market Research Report,
2714 2017.
- 2715 191. IRENA. Renewable Energy and Jobs: Annual Review 2017. Abu Dhabi: International
2716 Renewable Energy Agency, 2017.
- 2717 192. International Energy Agency. World Energy Outlook 2012. Paris, France, 2012.
- 2718 193. Granado JA, Coady D, Gillingham R. The Unequal Benefits of Fuel Subsidies: A Review
2719 of Evidence for Developing Countries, 2010.
- 2720 194. General Secretariat of the Council. G7 Ise-Shima Leaders' Declaration. Brussels,
2721 Belgium; 2016. p. 1.
- 2722 195. World Bank. Carbon Pricing Dashboard. 2017.
2723 <http://carbonpricingdashboard.worldbank.org> (accessed 06.06.2017 2017).
- 2724 196. World Bank, Ecofys. Carbon Pricing Watch 2017. Washington DC, USA, 2017.
- 2725 197. Patuelli R, Nijkamp P, Pels E. Environmental tax reform and the double dividend: a
2726 meta-analytical performance assessment. *Ecological Economics* 2005; **55**: 564-83.
- 2727 198. Georgeson L, al. e. Global Disparity in the Supply of Commercial Weather and
2728 Climate Information Services. *Science Advances* 2017; **3**(5).
- 2729 199. Climate Funds Update. Climate Funds Update: The Data. 2017.
2730 <http://www.climatefundsupdate.org/>.
- 2731 200. Crompton T. Common Values: The Case for Working with our Cultural Values: WWF,
2732 2010.
- 2733 201. World Health Assembly. Sixty-first World Health Assembly WHA61.19. 2008.
- 2734 202. Boykoff MT, Goodman MK, Curtis I. Cultural Politics of Climate Change: Interactions
2735 in Everyday Spaces, 2009.
- 2736 203. Lee TM, Markowitz EM, Howe PD, al e. Predictors of public climate change
2737 awareness and risk perception around the world *Nature climate change* 2015; **5**: 1014–20.
- 2738 204. Boykoff MT. Media and scientific communication: a case of climate change
2739 *Geological Society* 2008; **305**: 11-8.
- 2740 205. Steentjes K, al e. European Perceptions of Climate Change: Topline findings of a
2741 survey conducted in four European countries in 2016, 2017.
- 2742 206. Billett S. Dividing climate change: global warming in the Indian mass media. *Climatic
2743 Change* 2010; **99**(1/2): 525-37.

- 2744 207. Bhatta SNA. Coverage of Climate Change Issues in Indian Newspapers and Policy
2745 Implications. *Current Science* 2015; **108**(11): 1972-3.
- 2746 208. Andrews K, Boykoff, M., Daly, M., Gifford, L., Luedecke, G., McAllister, L., and Nacu-
2747 Schmidt, A. . World Newspaper Coverage of Climate Change or Global Warming, 2004-2017:
2748 Center for Science and Technology Policy Research, Cooperative Institute for Research in
2749 Environmental Sciences, University of Colorado, 2017.
- 2750 209. Schütte S, Depoux A, Vigil S, al. e. The influence of health concerns in scientific and
2751 policy debates on climate change. 2015; **Journal of Epidemiology and Community Health**.
- 2752 210. Depoux A, Hémono M, Puig-Malet S, Pédrón R, Flahault A. Communicating climate
2753 change and health in the media. *Public Health Reviews* 2017; **38**(1): 7.
- 2754 211. Hosking J, Campbell-Lendrum D. How well does climate change and human health
2755 research match the demands of policymakers? A scoping review. *Environmental health*
2756 *perspectives* 2012; **120**(8): 1076-82.
- 2757 212. Campbell-Lendrum D, Bertollini R, Neira M, Ebi K, McMichael A. Health and climate
2758 change: a roadmap for applied research. *The Lancet* 2009; **373**(9676): 1663-5.
- 2759 213. Arksey H, O'Malley L. Scoping studies: towards a methodological framework.
2760 *International Journal of Social Research Methodology* 2005; **8**(1): 19-32.
- 2761 214. Janssen MA, Schoon, M.L, Ke, W, and Börner, K. Scholarly networks on resilience,
2762 vulnerability and adaptation within the human dimensions of global environmental change.
2763 *Global Environmental Change* 2006; **16**(3): 240-52.
- 2764 215. Herlihy N, Bar-Hen A, Verner G, et al. Climate change and human health: what are
2765 the research trends? A scoping review protocol. *British Medical Journal* 2016; **6**.
- 2766 216. Baturo A, Dasandi N, Mikhaylov S. Understanding State Preferences with Text as
2767 Data: Introducing the UN General Debate Corpus. *Research and Politics* 2017; **4**(2).
2768