

## A Novel ELISA for the Measurement of Cerebrospinal Fluid SNAP-25 in Patients with Alzheimer's Disease

Annika Öhrfelt,<sup>a\*</sup> Ann Brinkmalm,<sup>a,b</sup> Julien Dumurgier,<sup>c</sup> Henrik Zetterberg,<sup>a,b,d,e</sup> Elodie Bouaziz-Amar,<sup>f</sup> Jacques Hugon,<sup>c</sup> Claire Paquet<sup>c</sup> and Kaj Blennow<sup>a,b</sup>

<sup>a</sup> Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, The Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden

<sup>b</sup> Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden

<sup>c</sup> Centre de Neurologie Cognitive CMRR Paris Nord Ile de France, INSERM UMR-S942, Groupe Hospitalier Lariboisière Fernand-Widal Saint-Louis, Paris, France

<sup>d</sup> Department of Molecular Neuroscience, UCL Institute of Neurology, London, United Kingdom

<sup>e</sup> UK Dementia Research Institute, London, United Kingdom

<sup>f</sup> Service de Biochimie, Groupe Hospitalier Lariboisière FW Saint-Louis, APHP, Université Paris Diderot, 75010 Paris, France

**Abstract**—Synaptic degeneration is central in Alzheimer's disease (AD) pathogenesis and biomarkers to monitor this pathophysiology in living patients are warranted. We developed a novel sandwich enzyme-linked immunosorbent assay (ELISA) for the measurement of the pre-synaptic protein SNAP-25 in cerebrospinal fluid (CSF) and evaluated it as a biomarker for AD. CSF samples included a pilot study consisting of AD ( $N = 26$ ) and controls ( $N = 26$ ), and two independent clinical cohorts of AD patients and controls. Cohort I included CSF samples from patients with dementia due to AD ( $N = 17$ ), patients with mild cognitive impairment (MCI) due to AD ( $N = 5$ ) and controls ( $N = 17$ ), and cohort II CSF samples from patients with dementia due to AD ( $N = 24$ ), patients with MCI due to AD ( $N = 18$ ) and controls ( $N = 36$ ). CSF levels of SNAP-25 were significantly increased in patients with AD compared with controls ( $P \leq 0.00001$ ). In both clinical cohorts, CSF levels of SNAP-25 were significantly increased in patients with MCI due to AD ( $P < 0.0001$ ). SNAP-25 could differentiate dementia due to AD ( $N = 41$ ) from controls ( $N = 52$ ) and MCI due to AD ( $N = 23$ ) from controls ( $N = 52$ ) with areas under the curve of 0.967 ( $P < 0.0001$ ) and 0.948 ( $P < 0.0001$ ), respectively. CSF SNAP-25 is a promising AD biomarker that differentiates AD patients in different clinical stages of the disease from controls with excellent diagnostic accuracy. Future studies should address the specificity of the CSF SNAP-25 against common differential diagnoses to AD, as well as how the biomarker changes in response to treatment with disease-modifying drug candidates.

*This article is part of a Special Issue entitled: SNARE proteins.* © 2018 The Authors. Published by Elsevier Ltd on behalf of IBRO. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

**Key words:** Alzheimer's disease, biomarker, cerebrospinal fluid, ELISA, mild cognitive impairment, SNAP-25.

### INTRODUCTION

Alzheimer's disease is characterized of extra-cellular accumulation of aggregated amyloid  $\beta$ , intra-cellular

neurofibrillary tangles, synaptic degeneration and neuronal degeneration (Blennow et al., 2006). Several cerebrospinal fluid (CSF) biomarkers for Alzheimer's disease are accessible, including total tau (T-tau) and phosphorylated tau protein (P-tau), mirroring tau pathology and neurodegeneration, respectively, and amyloid- $\beta_{1-42}$  ( $A\beta_{1-42}$ ), mirroring aggregation of the peptide into plaques (Blennow et al., 2010; Olsson et al., 2016). Numerous studies have consistently shown a reduction in  $A\beta_{1-42}$  attended by a marked increase in CSF T-tau and P-tau in Alzheimer's disease, and also in the mild cognitive impairment (MCI) stage of the disease (Blennow et al., 2010; Olsson et al., 2016), while there not yet is a conventional CSF biomarker for synaptic dysfunction. Synaptic degeneration of the most vulnerable brain regions is an early key characteristic of Alzheimer's disease (Davies

\*Correspondence to: Annika Öhrfelt, Clinical Neurochemistry Laboratory, Inst. of Neuroscience and Physiology, Dept. of Psychiatry and Neurochemistry, Sahlgrenska Academy at the University of Gothenburg, Sahlgrenska University Hospital, Mölndal, SE-431 80 Mölndal, Sweden.

E-mail addresses: [annika.ohrfelt@neuro.gu.se](mailto:annika.ohrfelt@neuro.gu.se) (A. Öhrfelt), [ann.brinkmalm@neuro.gu.se](mailto:ann.brinkmalm@neuro.gu.se) (A. Brinkmalm), [henrik.zetterberg@clinchem.gu.se](mailto:henrik.zetterberg@clinchem.gu.se) (H. Zetterberg), [elodie.amar@lrp.aphp.fr](mailto:elodie.amar@lrp.aphp.fr) (E. Bouaziz-Amar), [jacques.hugon@inserm.fr](mailto:jacques.hugon@inserm.fr) (J. Hugon), [claire.paquet@inserm.fr](mailto:claire.paquet@inserm.fr) (C. Paquet), [kaj.blennow@neuro.gu.se](mailto:kaj.blennow@neuro.gu.se) (K. Blennow).

Abbreviations: CV, coefficients of variation; MCI, mild cognitive impairment; MMSE, mini-mental state examination; PBS, phosphate-buffered saline; ROC, receiver operating characteristic; SNAP-25, synaptosomal-associated protein 25.

et al., 1987; Masliah et al., 2001; Scheff et al., 2007). Earlier post-mortem studies suggested that synaptic dysfunction in Alzheimer's disease is related to cognitive decline (DeKosky and Scheff, 1990; Blennow et al., 1996) and that synaptic loss occurs early in the disease (Davies et al., 1987; Masliah et al., 2001), with disturbances in presynaptic terminals (Masliah et al., 1991) and reductions in synaptic protein levels (DeKosky and Scheff, 1990; Blennow et al., 1996). Thus, it is evident that reliable CSF biomarkers to monitor synaptic dysfunction and degeneration directly in Alzheimer's disease patients would be very useful.

In recent years, there are promising results for some synaptic biomarkers in CSF, including the pre-synaptic proteins synaptosomal-associated protein 25 (SNAP-25) (Brinkmalm et al., 2014a; b) and synaptotagmin (Öhrfelt et al., 2016), as well as the post-synaptic protein neurogranin (Kvartsberg et al., 2015a,b; Sanfilippo et al., 2016; Wellington et al., 2016). A marked increase of these synaptic CSF markers were found in dementia due to Alzheimer's disease and already in MCI due to Alzheimer's disease (Brinkmalm et al., 2014a,b; Kvartsberg et al., 2015a,b; Öhrfelt et al., 2016; Sanfilippo et al., 2016; Wellington et al., 2016), with higher CSF levels correlating with more marked future cognitive decline among MCI patients (Kvartsberg et al., 2015a,b).

The pre-synaptic protein SNAP-25 is one of the major proteins involved in the formation of the SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) complexes (Sollner et al., 1993a; Sollner et al., 1993b; Jahn et al., 2003). This protein assembly is a crucial step in neurotransmitter release and modifications of any of the SNARE proteins could alter the apposition of them, which could influence calcium-dependent exocytosis of neuro-transmitters (Sollner et al., 1993a; Sollner et al., 1993b; Jahn et al., 2003; Sudhof 2004). The central function of SNAP-25 in the regulation of neuro-transmitter release along with the recently suggested post-synaptic impact on receptor trafficking, spine morphogenesis and plasticity (Antonucci et al., 2013; Antonucci et al., 2016), makes it as a potential biomarker candidate reflecting synaptic dysfunction and degeneration in Alzheimer's disease. We have previously shown that a N-terminal fragment of SNAP-25 is a promising biomarker by utilizing an approach of affinity purification and mass spectrometry (Brinkmalm et al., 2014a,b), and up to now, no enzyme-linked immunosorbent assay (ELISA) for assessment of SNAP-25 in CSF samples has been available. One advantage of the ELISA technology is the ease with which it can be performed in a high-through-put format. The feasibility and the accessibility that the ELISA offers would be required in future studies for assessment of synaptic proteins in large patient cohorts.

In this study, we report a novel ELISA for measurements of the pre-synaptic protein SNAP-25 in CSF. The utility of the novel SNAP-25 ELISA was initially verified in brain tissue extracts and from patients with Alzheimer's disease and age-matched controls, followed by a pilot study of CSF samples. Then, CSF SNAP-25 was assessed in two independent clinical cohorts, with the main finding being markedly higher

levels in patients with MCI due to Alzheimer's disease and dementia due to Alzheimer's disease.

## EXPERIMENTAL PROCEDURES

### Human brain tissue samples

All brain tissues, from the superior parietal gyrus, were obtained from the Netherlands Brain Bank. The clinical and demographic characteristics autopsy-confirmed patients with Alzheimer's disease ( $N = 15$ ) and age-matched controls ( $N = 15$ ) have previously been published (Brinkmalm et al., 2014a,b). In our study, all Alzheimer's disease patients fulfilled Braak stages 5 or 6, i.e. late stages of disease, while the controls fulfilled Braak stages 0 or 1 (Braak and Braak, 1991). The brain extraction procedure was performed as described by Brinkmalm et al. (2014a,b). In the present study, brain homogenates from the Tris fractions (soluble proteins) were analyzed.

### Quality control (QC) CSF samples

The repeatability of the novel SNAP-25 ELISA was examined on decoded CSF samples supplied by the clinical routine section at the Clinical Neurochemistry Laboratory, The Sahlgrenska University Hospital, Mölndal, Sweden. The procedure making pools of left-over CSF aliquots were approved by the Ethics Committee at University of Gothenburg. The quality control CSF pool 1 (QC1 sample) had an  $A\beta_{1-42}$  of 446 ng/L, a T-tau level of 332 ng/L and a P-tau level of 46 ng/L. The QC2 sample had an  $A\beta_{1-42}$  level of 405 ng/L, a T-tau level below 561 ng/L and a P-tau level of 50 ng/L.

### CSF samples in the pilot study

An initial pilot study was performed using de-identified CSF samples supplied by the Clinical Neurochemistry Laboratory, Sahlgrenska University, Mölndal, following procedures approved by the Ethics Committee at University of Gothenburg. Patients were designated as control or Alzheimer's disease according to CSF Alzheimer's disease core biomarker levels using in-house optimized cut-off levels for Alzheimer's disease (Hansson et al., 2006):  $A\beta_{1-42} < 550$  ng/L, T-tau  $> 400$  ng/L, and P-tau  $> 50$  ng/L. The subjects were older than 55 years. The age-matched test material included 26 patients with an Alzheimer's disease biomarker profile and 26 subjects with a control biomarker profile (Fig. 2).

### CSF samples in the clinical studies

In this study, SNAP-25 levels in CSF were measured in two independent clinical patient cohorts. The clinical and demographic characteristics have been reported previously (Öhrfelt et al., 2016). To facilitate for the reader essential parts used for diagnosing the patients and selecting the CSF are briefly given below (Öhrfelt et al., 2016). At the Center of Cognitive at Lariboisière Fernand-Widal University Hospital APHP, patients underwent a thorough clinical examination involving personal

151 medical and family histories, neurological examination,  
 152 neuropsychological assessment, lumbar puncture with  
 153 CSF biomarker analysis, and a brain structural imaging  
 154 study with MRI. The diagnosis for each patient was made  
 155 by neurologists considering CSF results and according to  
 156 validated clinical diagnostic criteria for dementia due to  
 157 Alzheimer's disease (McKhann et al., 2011), MCI due to  
 158 Alzheimer's disease (Albert et al., 2011; Dubois et al.,  
 159 2014), subjective cognitive impairment (Sperling et al.,  
 160 2011), psychiatric disorder (DSM-IV). The CSF samples  
 161 of the study were selected after a second validation step  
 162 by a neurologist (CP) and a biochemist (EAB). Patients  
 163 were not included in the study, without a consensus diag-  
 164 nosis or in case of disagreement about the final diagnosis.  
 165 This procedure resulted in selection of CSF samples from  
 166 subject with MCI due to Alzheimer's disease, dementia  
 167 due to Alzheimer's disease, and neurological controls  
 168 (no neurodegenerative disorders). The Alzheimer's dis-  
 169 ease core CSF biomarkers have been included in the  
 170 research criteria for the diagnosis of both early and man-  
 171 ifest Alzheimer's disease by the International Working  
 172 Group (Dubois et al., 2014) and in the diagnostic guideli-  
 173 nes from the National Institute on Aging-Alzheimer's  
 174 Association (McKhann et al., 2011), respectively. The fol-  
 175 lowing cut-off values were used to define a biochemical  
 176 Alzheimer's disease signature as supportive criteria for  
 177 dementia due to Alzheimer's disease (McKhann et al.,  
 178 2011):  $A\beta_{1-42}$  ( $< 550$  ng/L), T-tau ( $> 400$  ng/L), and P-  
 179 tau ( $> 50$  ng/L). CSF was obtained by lumbar puncture  
 180 between the L3/L4 or L4/L5 intervertebral space, and  
 181 samples were immediately centrifuged at 1800g for  
 182 10 min at  $+4$  °C, and stored at  $-80$  °C pending analysis.

### 183 Demographics of the clinical CSF studies

184 The demographic characteristics and the biomarker CSF  
 185 levels of the Alzheimer's disease core biomarkers for the  
 186 cohorts have been reported previously (Öhrfelt et al.,  
 187 2016). Briefly, cohort I consisted of five patients with  
 188 MCI due to Alzheimer's disease (one man and four  
 189 women, 62–88 years), 17 patients with dementia due to  
 190 Alzheimer's disease (five men and 12 women,  
 191 52–86 years), and 17 neurological controls (seven men  
 192 and ten women, 41–82 years) (Öhrfelt et al., 2016). The  
 193 replication sample set (cohort II) consisted of 18 patients  
 194 with MCI due to Alzheimer's disease (five men and 13  
 195 women, 58–83 years), 24 patients with dementia due to  
 196 Alzheimer's disease (seven men and 17 females,  
 197 52–84 years) and 36 neurological controls (13 men and  
 198 23 women, 43–80 years) (Öhrfelt et al., 2016). In cohort  
 199 I, the patients with MCI due to Alzheimer's disease were  
 200 older than the controls. Both patients with MCI due to  
 201 Alzheimer's disease and dementia due to Alzheimer's dis-  
 202 ease were slightly but significantly older than the controls  
 203 in cohort II (Öhrfelt et al., 2016).

### 204 Analysis of CSF biomarkers

205  $A\beta_{1-42}$ , T-tau, and tau phosphorylated at threonine 181  
 206 (P-tau) protein measurements were performed using  
 207 commercially available assays from Fujirebio  
 208 (INNOTEST®  $\beta$ -AMYLOID<sub>(1-42)</sub>, INNOTEST® hTAU Ag

and INNOTEST® PHOSPHO-TAU(181P) according to 209  
 the manufacturer's instructions. 210

### Synthetic peptides of SNAP-25 and antibodies 211

212 The synthetic peptide of N-terminal acetylated SNAP-25  
 213 (Ac-2-47 SNAP-25) was bought from CASLO Aps  
 214 (Lyngby, Denmark). The monoclonal mouse antibody  
 215 clone 71.1 recognizing the N-terminal portion of SNAP-  
 216 25 (aa 20–40) was purchased from Synaptic Systems  
 217 (Göttingen, Germany). Polyclonal chicken IgY antibody  
 218 was produced by immunization with Ac-2-47 SNAP-25  
 219 and the subsequent antigen affinity purification of the  
 220 total IgY extract was conducted by Getica AB  
 221 (Gothenburg, Sweden). Biotinylation of the Ac-2-47  
 222 SNAP-25 purified chicken IgY antibody was performed  
 223 accordingly to the manual, Simoa Homebrew Detector  
 224 Biotinylation Protocol, provided by Quanterix (Lexington,  
 225 MA, USA). A ratio of biotin to antibody of 40:1 was  
 226 applied.

### A novel sandwich ELISA method for SNAP-25 227

228 F16 Maxisorp Loose Nunc-Immuno plates (Thermo  
 229 Fisher Scientific Nunc A/S, Roskilde, Denmark) were  
 230 coated with 100  $\mu$ L of monoclonal mouse antibody clone  
 231 71.1 (1 g/L) diluted 1:400 in 50 mM carbonate buffer, pH  
 232 9.6 and incubated over night or up to three nights at  
 233  $+2$ – $8$  °C. The plates were washed with 385  $\mu$ L of  
 234 phosphate-buffered saline PBS-Tween20 (0.05%) (PBS-  
 235 T). The same washing procedure was repeated  
 236 between every following incubation step. After the  
 237 coating and washing steps, the plates were blocked with  
 238 300  $\mu$ L Roti®-Block (Carl Roth, Germany) diluted 1:10 in  
 239 PBS-T for one hour at room temperature. All standards  
 240 and samples were analyzed in duplicate. The standards  
 241 of Ac-2-47 SNAP-25 were diluted in assay buffer, i.e.  
 242 Roti®-Block diluted 1:100 in PBS-T, to providing a final  
 243 concentration range of 4000–62.5 ng/L or 1000–7.8 ng/L  
 244 for brain samples and CSF samples, respectively. Brain  
 245 tissue homogenates were diluted 1:15 in assay buffer,  
 246 while neat CSF samples were added to the plates.  
 247 Samples and standards (50  $\mu$ L) were incubated over  
 248 night at  $+2$ – $8$  °C, simultaneously with 50  $\mu$ L biotinylated  
 249 affinity Ac-2-47 SNAP-25 purified chicken IgY antibody  
 250 (1 g/L) diluted 1:500 in assay buffer. Enhanced  
 251 Streptavidin-HRP conjugate (0.01 g/L) (Kem-En-Tec  
 252 Diagnostics, Taastrup, Denmark), pre-diluted 1:100 in  
 253 Uni-Stabil Plus (Kem-En-Tec Diagnostics) (stored at  
 254  $+2$ – $8$  °C pending analysis), was then diluted 1:200 in  
 255 assay buffer, and was incubated for 30 min at room  
 256 temperature. Then, 100  $\mu$ L TMB ONE™, ready-to-use  
 257 substrate (KE-MEN-TEC Diagnostics) were added. The  
 258 reaction was quenched with 100  $\mu$ L of  $H_2SO_4$  (0.2 M).  
 259 The absorbance was measured at 450 nm. The  
 260 concentrations of SNAP-25 in samples were calculated  
 261 from the four parameter standard curve. For each brain  
 262 sample a ratio was calculated where the SNAP-25 level  
 263 was divided with the total protein concentration.



## Assay performance

264 The within-day precision (repeatability) and the between-  
265 day repeatability (intermediate precision) were  
266 determined using two QC samples (QC1 and QC2)  
267 analyzing them at three different days ( $N = 5$  or  $N = 6$ ).  
268 Lower limit of quantification (LLOQ) was calculated  
269 according to Andreasson et al. (2015).  
270

## Statistical analysis

271 Because most of the analytes were not normally  
272 distributed (Shapiro-Wilk test,  $P < 0.05$ ), non-parametric  
273 statistics were used for analysis. Data are given as  
274 median (inter-quartile range). Differences between more  
275 than two groups were assessed with Kruskal–Wallis  
276 test. Statistically significant results ( $P < 0.05$ ) were  
277 followed by Mann–Whitney  $U$ -tests to investigate group  
278 differences. Receiver operating characteristic (ROC)  
279 curves were performed on each subject group on the  
280 levels of SNAP-25 in order to assess its diagnostic  
281 value. The area under the curve (AUC) and a 95%  
282 confidence interval (CI) was calculated for SNAP-25  
283 using GraphPad Prism 7.02. The correlation coefficients  
284 ( $\rho$ ) were calculated using the Spearman two-tailed  
285 correlation test. SPSS 24 was employed for most of the  
286 statistical analyzes.  
287

## RESULTS

### Assay performance

288 The novel ELISA is directed against the N-terminal of  
289 SNAP-25, that measure both partially degraded N-  
290 terminal SNAP-25 fragments as well as the possible full-  
291 length protein. Within-day repeatability was 9.6% for QC  
292 sample 1 and 15% for QC sample 2. Between-day  
293 repeatability was 13% (QC1) and 16% (QC2). The  
294 repeatability was within acceptable ranges, i.e. within-  
295 day  $\leq 15$  and between-day  $\leq 20$  (Lee and Hall (2009)).  
296 LLOQ was 15.7 ng/L.  
297  
298

### Human brain and the pilot CSF study

299 Initially, we tested the novel SNAP-25 ELISA on brain  
300 tissue homogenates from age-matched patients with  
301 Alzheimer's disease and controls. We found that SNAP-  
302 25 levels were significantly decreased in patients with  
303 later stages of Alzheimer's disease compared with the  
304 controls (Fig. 1). In the pilot CSF study, the levels of  
305 SNAP-25 were significantly increased in the group with  
306 an Alzheimer's disease biomarker profile ( $N = 26$ ) than  
307 in the group with a control biomarker profile ( $N = 26$ )  
308 (Fig. 2).  
309

### CSF SNAP-25 in the clinical cohorts

310 CSF levels of the SNAP-25 were significantly higher in  
311 patients with MCI due to Alzheimer's disease (cohort I,  
312 II and all samples), and in dementia due to Alzheimer's  
313 disease compared with controls (cohort I, II and all  
314 samples) (Fig. 3). SNAP-25 could differentiate MCI due  
315 to Alzheimer's disease from controls in both cohorts and  
316 in the entire set of samples, with AUCs (confidence  
317





Fig. 1. SNAP-25 in brain tissue in Alzheimer's disease (AD) and controls. The figure shows the individual values SNAP-25 (displayed as the ratio SNAP-25/total protein) in the soluble protein fraction in the superior parietal gyrus from controls (green) and patients with AD (violet). The lower, upper and middle lines of the error bars correspond to the 25th and 75th percentiles and medians, respectively.




	Control biomarker profile	AD biomarker profile
Subjects, N (men/women)	26 (14/12)	26 (7/19)
Age (years)	71 (68-75)	74 (70-77)
A $\beta$ <sub>1-42</sub> (ng/L)	748 (669-900)	440 (375-489) <sup>a</sup>
T-tau (ng/L)	234 (188-325)	642 (514-870) <sup>a</sup>
P-tau (ng/L)	34 (31-44)	82 (70-91) <sup>a</sup>


Fig. 2. Individual values for SNAP-25 (A) and demographic data including Alzheimer's disease (AD) core biomarker levels (B) from the pilot study for the patients with AD (violet) and controls (green) based on the biomarker profile. The lower, upper and middle lines of the error bars correspond to the 25th and 75th percentiles and medians, respectively (A).

interval (CI) of 1 (1-1) ( $P = 0.001$ ) (cohort I), 0.975 (0.943–1.008) ( $P < 0.0001$ ) (cohort II) and 0.948 (0.964–1.004) ( $P < 0.0001$ ) (all samples) (Fig. 4A, C). SNAP-25 could also differentiate dementia due to Alzheimer's disease from controls with AUCs (CI) of 0.982 (0.946–1.017) ( $P < 0.0001$ ) (cohort I), 0.970 (0.935–1.005) ( $P < 0.0001$ ) (cohort II) and 0.967 (0.938–0.996) ( $P < 0.0001$ ) (all samples) (Fig. 4B, C).

318  
319  
320  
321  
322  
323  
324  
325



**Fig. 3.** Individual values for SNAP-25 in CSF samples within cohort I (A), cohort II (B) and for the entire set of samples (C) from subjects with dementia due to Alzheimer's disease (AD) (violet), mild cognitive impairment due to Alzheimer's disease (MCI-AD) (orange) and control (green) individuals. The lower, upper and middle lines of the error bars correspond to the 25th and 75th percentiles and medians, respectively.



	Control versus MCI-AD	Control versus AD
<b>Cohort I</b>	N=16 and N=5	N=16 and N=17
SNAP-25	1 (1-1), P=0.001	0.982 (0.946-1.017), P<0.0001
<b>Cohort II</b>	N=36 and N=18	N=36 and N=24
SNAP-25	0.975 (0.943-1.008), P<0.0001	0.970 (0.935-1.005), P<0.0001
<b>All samples</b>	N=52 and N=23	N=52 and N=41
SNAP-25	0.984 (0.964-1.004), P<0.0001	0.967 (0.938-0.996), P<0.0001

**Fig. 4.** ROC curve analysis for SNAP-25 in CSF for differentiation of MCI due to Alzheimer's disease (MCI-AD) from controls in cohort I (violet), cohort II (green) and in the entire set of samples (black) (A). ROC curve analysis for SNAP-25 in CSF for differentiation of dementia due to Alzheimer's disease (AD) from controls in cohort I (violet), cohort II (green) and in the entire set of samples (black) (B). The area under the curve (95% confidence interval) is shown in the included table (C).

326 There was a correlation between the CSF levels of  
327 SNAP-25 and the age in patients with dementia due to  
328 Alzheimer's disease (cohort I), while there were no  
329 statistically significant correlations between SNAP-25  
330 and age in any other of the investigated groups

(Table 1). There were no statistically significant correlations between CSF SNAP-25 and minimal state examination (MMSE) scores in any group.

The CSF levels of SNAP-25 correlated with the levels of T-tau and P-tau in both the control group and in patients with dementia due to Alzheimer's disease (Table 1). Additionally, the CSF levels of SNAP-25 correlated with the levels of T-tau and P-tau in patients with MCI due to Alzheimer's disease within the entire set of samples, but only with the levels of P-tau within cohort II (Table 1). SNAP-25 correlated positively with  $A\beta_{1-42}$  in the control group of cohort II and for the entire set of samples, while there were no correlations within other investigated groups (Table 1).

## DISCUSSION

We developed a novel ELISA for assessment of the pre-synaptic protein SNAP-25 in CSF samples. In one pilot study and both investigated clinical cohorts, we found that the CSF levels of SNAP-25 were significantly higher in patients with dementia due to Alzheimer's disease than in controls. There was also a consistent increase in early disease (i.e. MCI due to Alzheimer's disease) as compared to controls.

Synaptic dysfunction and degeneration predict cognitive decline in Alzheimer's disease (Davies et al., 1987; Masliah et al., 2001). The pre-synaptic protein SNAP-25 is one of the prominent proteins involved in the regulation of synaptic transmission (Sollner et al., 1993a,b; Sudhof, 2004), and therefore could possibly be a biomarker candidate that mirrors synaptic degeneration and dysfunction in Alzheimer's disease. We found that the CSF levels of SNAP-25 were consistently elevated in patients with dementia due to Alzheimer's disease compared with controls in two separate clinical cohorts, as well as in a group having an Alzheimer's disease biomarker profile compared to a group with a control biomarker profile. Addition-

**Table 1.** Correlation between cerebrospinal fluid SNAP-25, age, MMSE and biomarker levels for the diagnostic groups in the clinical cohorts<sup>a</sup>

	SNAP-25	SNAP-25	SNAP-25
<i>Cohort I</i>	<i>Control (N = 17)</i>	<i>MCI-AD (N = 18)</i>	<i>AD (N = 17)</i>
Age	N.S.		rho = -0.503, P = 0.04
MMSE	N.S.		N.S.
Amyloid- $\beta_{1-42}$	N.S.		N.S.
Total tau	rho = 0.805, P = 0.0002		rho = 0.738, P = 0.001
Phosphorylated tau	rho = 0.715, P = 0.002		rho = 0.830, P = 0.00004
<i>Cohort II</i>	<i>Control (N = 36)</i>	<i>MCI-AD (N = 18)</i>	<i>AD (N = 24)</i>
Age	N.S.	N.S.	N.S.
MMSE	N.S.	N.S.	N.S.
Amyloid- $\beta_{1-42}$	rho = 0.363, P = 0.03	N.S.	N.S.
Total tau	rho = 0.743, P < 0.00001	N.S.	rho = 0.663, P = 0.0004
Phosphorylated tau	rho = 0.618, P = 0.00008	rho = 0.513, P = 0.03	rho = 0.604, P = 0.002
<i>All samples</i>	<i>Control (N = 53)</i>	<i>MCI-AD (N = 23)</i>	<i>AD (N = 41)</i>
Age	N.S.	N.S.	N.S.
MMSE	N.S.	N.S.	N.S.
Amyloid- $\beta_{1-42}$	rho = 0.325, P = 0.02	N.S.	N.S.
Total tau	rho = 0.744, P < 0.00001	rho = 0.453, P = 0.03	rho = 0.726, P < 0.00001
Phosphorylated tau	rho = 0.639, P < 0.00001	rho = 0.637, P = 0.001	rho = 0.736, P < 0.00001

<sup>a</sup> Correlations presented by the Spearman's rank correlation coefficient (rho). Non-significant (N.S., P > 0.05) correlations were not reported.

ally, the level of SNAP-25 was increased already in the MCI stage of Alzheimer's disease, supporting the notion that this pre-synaptic protein might be an early marker for Alzheimer's disease (Brinkmalm et al., 2014a,b). There is evidence suggesting that pre-synaptic dysfunction may occur early in the pathogenesis of dementia (Masliah et al., 2001), and that compensatory post-synaptic alterations may occur in response to pre-synaptic discrepancies (DeKosky and Scheff, 1990). These results are altogether in agreement with our earlier studies of the synaptic proteins SNAP-25 (Brinkmalm et al., 2014a; b), synaptotagmin (Öhrfelt et al., 2016) and neurogranin (Kvartsberg et al., 2015a,b).

We present a sensitive ELISA, which showed reproducibility and intermediate precision not exceeding %CV of 15 and 16, respectively. SNAP-25 exists in two isoforms in the brain, SNAP-25A and SNAP-25B (Bark and Wilson, 1994). These isoforms differ only in nine alternate amino acids 58, 60, 65, 69, 79, 84 and 88–89, which are located beyond the potential cleavage site of SNAP-25, all of which can be measured using the novel ELISA. The design of the novel ELISA is based on our previous finding of numerous N-terminally acetylated soluble SNAP-25 fragments in both human brain tissue and CSF from subjects with Alzheimer's disease and controls (Brinkmalm et al., 2014a,b). In the previous study, we applied affinity purification (immunoprecipitation) against the N-terminal of SNAP-25 and mass spectrometry analyzed for subsequently quantification of tryptic peptides in CSF (Brinkmalm et al., 2014a,b). The most prominent result was that the tryptic peptide furthest away from the targeted N-terminal provided the best differential diagnostic biomarker of Alzheimer's disease (Brinkmalm et al., 2014a,b), which might correspond to a truncated SNAP-25 fragment ending after amino acid 47 (Ac-2–47) (Brinkmalm et al., 2014a,b). In the present study, we confirm that CSF SNAP-25 can discriminate both patients

with dementia due to Alzheimer's disease and patients with MCI due to Alzheimer's disease from controls with high diagnostic accuracy in ROC curve analyzes (Brinkmalm et al., 2014a,b). In agreement, we also found that the CSF levels of SNAP-25 were significantly elevated in Alzheimer's disease (Brinkmalm et al., 2014a, b). The novel ELISA does not exclusively target the Ac-2–47, and possibly longer N-terminal forms of SNAP-25 might also be analyzed. Interestingly, truncated N-terminal fragments of SNAP-25 might be created by calpain cleavage (Ando et al., 2005; Grumelli et al., 2008), and the activity of calpain is increased in Alzheimer's disease brain (Kurbatskaya et al., 2016). The cleavage of SNAP-25 by calpain might regulate synaptic transmission by suppressing the neuro-transmitter release (Ando et al., 2005).

In agreement with the majority of previous reports summarized by Honer (2003), we found that the SNAP-25 levels in brain were significantly decreased in later stages of Alzheimer's disease compared with the controls (Gabriel et al., 1997; Mukaetova-Ladinska et al., 2000; Brinkmalm et al., 2014a,b). The lower levels of SNAP-25 might reflect the synaptic degeneration known to occur in disease-affected regions of the brain in Alzheimer's disease (DeKosky and Scheff, 1990). Intra-cellular SNAP-25 is anchored to the pre-synaptic membrane by palmitoylation of a central cysteine-rich region (amino acids 85, 88, 90 and 92) (Veit et al., 1996). Since the palmitoylation is a reversible reaction, SNAP-25 could possibly reside free in the pre-synaptic cytoplasm. However, the mechanism of liberation of SNAP-25 into CSF and what it reflects are unknown. Herein, we found that SNAP-25 correlated with the levels of T-tau and P-tau in both the control group and in patients with dementia due to Alzheimer's disease in all examined sample sets. CSF T-tau has previously been suggested to be a general marker of damage to cortical non-myelinated neurons (Blennow et al., 2010). In con-



464 trast, P-tau might be a more specific marker for Alzheimer's  
 465 disease (Blennow et al., 2010), since high CSF  
 466 levels of P-tau have been found to correlate to the accu-  
 467 mulation of cortical neurofibrillary tangles (Buerger  
 468 et al., 2006; Tapiola et al., 2009). Altogether, these find-  
 469 ings suggest that SNAP-25 is a sensitive Alzheimer's dis-  
 470 ease biomarker that to some extent mirrors general  
 471 neurodegeneration, which is in agreement with our first  
 472 pilot study (Brinkmalm et al., 2014a,b). The result that  
 473 the levels of SNAP-25 correlated well with T-tau and P-  
 474 tau, imply that SNAP-25 might be a valuable surrogate  
 475 biomarker in future clinical treatment studies with tau-  
 476 based- modifying drugs (Panza et al., 2016).

477 Marked synaptic degeneration and loss are the main  
 478 pathological features of Alzheimer's disease that  
 479 correlate with cognitive decline. Since SNAP-25 is  
 480 directly involved in the maintenance of synaptic function  
 481 (Sollner et al., 1993a,b; Sudhof, 2004), CSF SNAP-25  
 482 could be a potential biomarker to follow progression of  
 483 clinical symptoms. In the present study, there were no  
 484 correlations between the MMSE score, *i.e.*, the severity  
 485 of cognitive impairment, and SNAP-25 in any of the exam-  
 486 ined groups. Although we did not find correlation  
 487 between cognition and SNAP-25, previous studies sup-  
 488 port that SNAP-25 single nucleotide polymorphisms are  
 489 associated with cognitive decline (Gosso et al., 2008;  
 490 Guerini et al., 2014). Further studies using a larger set  
 491 of clinical samples are warranted to investigate if SNAP-  
 492 25 in CSF could be used for assessment of future rate  
 493 of cognitive decline. The relationship of CSF SNAP-25  
 494 with neuroimaging markers (positron emission tomogra-  
 495 phy and magnetic resonance imaging) would also be  
 496 important to evaluate. For instance, changes in glucose  
 497 utilization identified with fluorodeoxyglucose positron  
 498 emission tomography could possibly reflect neurodegen-  
 499 eration/synaptic dysfunction (Petrie et al., 2009), and  
 500 the cortical glucose metabolism would therefore be inter-  
 501 esting to study together with CSF SNAP-25.

502 The strengths of our study are that we present a novel  
 503 ELISA for assessment of the CSF levels of SNAP-25 and  
 504 that consistent findings were shown in one pilot set and  
 505 two independent replication cohorts of CSF samples.  
 506 One drawback is the cross-sectional design that  
 507 complicates the investigation of possible association  
 508 between CSF SNAP-25 and synaptic degeneration over  
 509 time.

510 In summary, we present a novel ELISA for  
 511 measurement of the pre-synaptic protein SNAP-25 in  
 512 CSF samples. CSF SNAP-25 levels were increased in  
 513 patients with MCI due to Alzheimer's disease and  
 514 dementia due to Alzheimer's disease compared with  
 515 controls, which are in agreement with our previous  
 516 findings, and supports the notion that SNAP-25 could be  
 517 a valuable biomarker both in early Alzheimer's disease  
 518 and in manifest Alzheimer's disease dementia. Future  
 519 studies should examine the ability to monitor cognitive  
 520 decline, the specificity of the biomarker against non-  
 521 Alzheimer's disease dementias, as well as how it  
 522 changes in response to treatment with novel disease-  
 523 modifying drug candidates.

## DECLARATIONS

### Ethical approval and consent to participate

The study was approved by the Ethics Committee of Paris  
 Diderot University Hospital (Bichat Hospital). All patients  
 or caregivers gave their written informed consents for  
 research, which was conducted in accordance with the  
 Helsinki Declaration. The use of de-identified leftover  
 samples for method development and validation studies  
 was approved by the Regional Ethical Review Board at  
 University of Gothenburg (08-11-14).

### CONSENT FOR PUBLICATION

Not applicable.

### AVAILABILITY OF DATA AND MATERIALS

The datasets used and/or analyzed during the present  
 study are available from the corresponding author on  
 reasonable request.

### COMPETING INTERESTS

KB has served at advisory boards or as a consultant for  
 Alzheon, BioArctic, Biogen, Eli Lilly, Fujirebio Europe,  
 IBL International, Pfizer, and Roche Diagnostics, and is  
 a co-founder of Brain Biomarker Solutions in  
 Gothenburg AB, a GU Ventures-based platform  
 company at the University of Gothenburg. HZ is another  
 co-founder of this company. The other authors declare  
 that they have no competing interests.

### FUNDING

The work was supported by grants from the Swedish  
 Brain Power Consortium, the Swedish Alzheimer  
 Foundation (#AF-553101 and # AF-646211), the  
 Research Council, Sweden (project #14002), the Brain  
 Foundation, Sweden (project # FO2015-0021), LUA/ALF  
 project, Västra Götalandsregionen, Sweden (project #  
 ALFGBG-139671), European Research Council, the  
 Knut and Alice Wallenberg Foundation, Demensfonden,  
 Eivind och Elsa K:son Sylvans stiftelse, the Wolfson  
 Foundation, Märtha och Gustaf Ågrens stiftelse,  
 Stohnes stiftelse, Stiftelsen Gamla Tjänarinnor, Magn.  
 Bergvalls stiftelse, Svenska Läkaresällskapet, the  
 Torsten Söderberg Foundation at the Royal Swedish  
 Academy of Sciences, Åhlén-stiftelsen, and BMBF  
 BIOMARK-APD (DLR 01ED1203 J).

### AUTHORS' CONTRIBUTIONS

AÖ and KB performed the study design, interpretation of  
 the results, and writing of the manuscript draft. AB, JD,  
 HZ, EB-A, JH and CP contributed to the study concept  
 and design and/or to critical revision of the manuscript  
 for important intellectual content. AÖ performed the  
 experiments, analyzed and compiled data. All authors  
 read and approved the final manuscript.

## ACKNOWLEDGMENTS

We are grateful to Åsa Källén and Sara Skoglar for their technical assistance.

## REFERENCES

- Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, Gamst A, Holtzman DM, Jagust WJ, Petersen RC, Snyder PJ, Carrillo MC, Thies B, Phelps CH (2011) The diagnosis of mild cognitive impairment due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 7(3):270–279.
- Ando K, Kudo Y, Takahashi M (2005) Negative regulation of neurotransmitter release by calpain: a possible involvement of specific SNAP-25 cleavage. *J Neurochem* 94(3):651–658.
- Andreasson U, Perret-Liaudet A, van Waalwijk van Doorn LJ, Blennow K, Chiasserini D, Engelborghs S, Fladby T, Genc S, Kruse N, Kuiperij HB, Kulic L, Lewczuk P, Mollenhauer B, Mroczko B, Parnetti L, Vanmechelen E, Verbeek MM, Winblad B, Zetterberg H (2015) A practical guide to immunoassay method validation. *Front Neurol* 6:179.
- Antonucci F, Corradini I, Morini R, Fossati G, Menna E, Pozzi D, Pacioni S, Verderio C, Bacci A, Matteoli M (2013) Reduced SNAP-25 alters short-term plasticity at developing glutamatergic synapses. *EMBO Rep* 14(7):645–651.
- Antonucci F, Corradini I, Fossati G, Tomasoni R, Menna E, Matteoli M (2016) SNAP-25, a known presynaptic protein with emerging postsynaptic functions. *Front Synaptic Neurosci* 8:7.
- Bark IC, Wilson MC (1994) Human cDNA clones encoding two different isoforms of the nerve terminal protein SNAP-25. *Gene* 139(2):291–292.
- Blennow K, Bogdanovic N, Alafuzoff I, Ekman R, Davidsson P (1996) Synaptic pathology in Alzheimer's disease: relation to severity of dementia, but not to senile plaques, neurofibrillary tangles, or the ApoE4 allele. *J Neural Transm (Vienna)* 103(5):603–618.
- Blennow K, de Leon MJ, Zetterberg H (2006) Alzheimer's disease. *Lancet* 368(9533):387–403.
- Blennow K, Hampel H, Weiner M, Zetterberg H (2010) Cerebrospinal fluid and plasma biomarkers in Alzheimer disease. *Nat Rev Neurol* 6(3):131–144.
- Braak H, Braak E (1991) Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol* 82(4):239–259.
- Brinkmalm A, Brinkmalm G, Honer WG, Moreno JA, Jakobsson J, Mallucci GR, Zetterberg H, Blennow K, Öhrfelt A (2014b) Targeting synaptic pathology with a novel affinity mass spectrometry approach. *Mol Cell Proteomics* 13(10):2584–2592.
- Brinkmalm A, Brinkmalm G, Honer WG, Frölich L, Hausner L, Minthon L, Hansson O, Wallin A, Zetterberg H, Blennow K, Öhrfelt A (2014a) SNAP-25 is a promising novel cerebrospinal fluid biomarker for synapse degeneration in Alzheimer's disease. *Mol Neurodegener* 9:53.
- Buerger K, Ewers M, Pirttilä T, Zinkowski R, Alafuzoff I, Teipel SJ, DeBernardis J, Kerkman D, McCulloch C, Soininen H, Hampel H (2006) CSF phosphorylated tau protein correlates with neocortical neurofibrillary pathology in Alzheimer's disease. *Brain* 129(Pt 11):3035–3041.
- Davies CA, Mann DM, Sumpter PQ, Yates PO (1987) A quantitative morphometric analysis of the neuronal and synaptic content of the frontal and temporal cortex in patients with Alzheimer's disease. *J Neurol Sci* 78(2):151–164.
- DeKosky ST, Scheff SW (1990) Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann Neurol* 27(5):457–464.
- Dubois B, Feldman HH, Jacova C, Hampel H, Molinuevo JL, Blennow K, DeKosky ST, Gauthier S, Selkoe D, Bateman R, Cappa S, Crutch S, Engelborghs S, Frisoni GB, Fox NC, Galasko D, Habert MO, Jicha GA, Nordberg A, Pasquier F, Rabinovici G, Robert P, Rowe C, Salloway S, Sarazin M, Epelbaum S, de Souza LC, Vellas B, Visser PJ, Schneider L, Stern Y, Scheltens P, Cummings JL (2014) Advancing research diagnostic criteria for Alzheimer's disease: the IWG-2 criteria. *Lancet Neurol* 13(6):614–629.
- Gabriel SM, Haroutunian V, Powchik P, Honer WG, Davidson M, Davies P, Davis KL (1997) Increased concentrations of presynaptic proteins in the cingulate cortex of subjects with schizophrenia. *Arch Gen Psychiatry* 54(6):559–566.
- Gosso MF, de Geus EJ, Polderman TJ, Boomsma DI, Heutink P, Posthuma D (2008) Common variants underlying cognitive ability: further evidence for association between the SNAP-25 gene and cognition using a family-based study in two independent Dutch cohorts. *Genes Brain Behav* 7(3):355–364.
- Grumelli C, Berghuis P, Pozzi D, Caleo M, Antonucci F, Bonanno G, Carmignoto G, Dobszay MB, Harkany T, Matteoli M, Verderio C (2008) Calpain activity contributes to the control of SNAP-25 levels in neurons. *Mol Cell Neurosci* 39(3):314–323.
- Guerini FR, Agliardi C, Sironi M, Arosio B, Calabrese E, Zanzottera M, Bolognesi E, Ricci C, Costa AS, Galimberti D, Griffanti L, Bianchi A, Savazzi F, Mari D, Scarpini E, Baglio F, Nemni R, Clerici M (2014) Possible association between SNAP-25 single nucleotide polymorphisms and alterations of categorical fluency and functional MRI parameters in Alzheimer's disease. *J Alzheimers Dis* 42(3):1015–1028.
- Hansson O, Zetterberg H, Buchhave P, Londos E, Blennow K, Minthon L (2006) Association between CSF biomarkers and incipient Alzheimer's disease in patients with mild cognitive impairment: a follow-up study. *Lancet Neurol* 5(3):228–234.
- Honer WG (2003) Pathology of presynaptic proteins in Alzheimer's disease: more than simple loss of terminals. *Neurobiol Aging* 24(8):1047–1062.
- Jahn R, Lang T, Sudhof TC (2003) Membrane fusion. *Cell* 112(4):519–533.
- Kurbatskaya K, Phillips EC, Croft CL, Dentoni G, Hughes MM, Wade MA, Al-Sarraj S, Troakes C, O'Neill MJ, Perez-Nievas BG, Hanger DP, Noble W (2016) Upregulation of calpain activity precedes tau phosphorylation and loss of synaptic proteins in Alzheimer's disease brain. *Acta Neuropathol Commun* 4:34.
- Kvartsberg H, Duits FH, Ingelsson M, Andreassen N, Öhrfelt A, Andersson K, Brinkmalm G, Lannfelt L, Minthon L, Hansson O, Andreasson U, Teunissen CE, Scheltens P, Van der Flier WM, Zetterberg H, Portelius E, Blennow K (2015a) Cerebrospinal fluid levels of the synaptic protein neurogranin correlates with cognitive decline in prodromal Alzheimer's disease. *Alzheimers Dement* 11(10):1180–1190.
- Kvartsberg H, Portelius E, Andreasson U, Brinkmalm G, Hellwig K, Lelental N, Kornhuber J, Hansson O, Minthon L, Spitzer P, Maler JM, Zetterberg H, Blennow K, Lewczuk P (2015b) Characterization of the postsynaptic protein neurogranin in paired cerebrospinal fluid and plasma samples from Alzheimer's disease patients and healthy controls. *Alzheimers Res Ther* 7(1):40.
- Lee JW, Hall M (2009) Method validation of protein biomarkers in support of drug development or clinical diagnosis/prognosis. *J Chromatogr B Analyt Technol Biomed Life Sci* 877(13):1259–1271.
- Masliah E, Hansen L, Albricht T, Mallory M, Terry RD (1991) Immunoelectron microscopic study of synaptic pathology in Alzheimer's disease. *Acta Neuropathol* 81(4):428–433.
- Masliah E, Mallory M, Alford M, DeTeresa R, Hansen LA, McKeel Jr DW, Morris JC (2001) Altered expression of synaptic proteins occurs early during progression of Alzheimer's disease. *Neurology* 56(1):127–129.
- McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack Jr CR, Kawas CH, Klunk WE, Koroshetz WJ, Manly JJ, Mayeux R, Mohs RC, Morris JC, Rossor MN, Scheltens P, Carrillo MC, Thies B, Weintraub S, Phelps CH (2011) The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* 7(3):263–269.



- 711 Mukaetova-Ladinska EB, Garcia-Siera F, Hurt J, Gertz HJ, Xuereb  
712 JH, Hills R, Brayne C, Huppert FA, Paykel ES, McGee M, Jakes  
713 R, Honer WG, Harrington CR, Wischik CM (2000) Staging of  
714 cytoskeletal and beta-amyloid changes in human isocortex  
715 reveals biphasic synaptic protein response during progression of  
716 Alzheimer's disease. *Am J Pathol* 157(2):623–636.
- 717 Öhrfelt A, Brinkmalm A, Dumurgier J, Brinkmalm G, Hansson O,  
718 Zetterberg H, Bouaziz-Amar E, Hugon J, Paquet C, Blennow K  
719 (2016) The pre-synaptic vesicle protein synaptotagmin is a novel  
720 biomarker for Alzheimer's disease. *Alzheimers Res Ther* 8(1):41.
- 721 Olsson B, Lautner R, Andreasson U, Öhrfelt A, Portelius E, Bjerke M,  
722 Holtta M, Rosen C, Olsson C, Strobel G, Wu E, Dakin K, Petzold  
723 M, Blennow K, Zetterberg H (2016) CSF and blood biomarkers for  
724 the diagnosis of Alzheimer's disease: a systematic review and  
725 meta-analysis. *Lancet Neurol* 15(7):673–684.
- 726 Panza F, Solfrizzi V, Seripa D, Imbimbo BP, Lozupone M, Santamato  
727 A, Tortelli R, Galizia I, Prete C, Daniele A, Pilotto A, Greco A,  
728 Logroscino G (2016) Tau-based therapeutics for Alzheimer's  
729 disease: active and passive immunotherapy. *Immunotherapy* 8  
730 (9):1119–1134.
- 731 Petrie EC, Cross DJ, Galasko D, Schellenberg GD, Raskind MA,  
732 Peskind ER, Minoshima S (2009) Preclinical evidence of  
733 Alzheimer changes: convergent cerebrospinal fluid biomarker  
734 and fluorodeoxyglucose positron emission tomography findings.  
735 *Arch Neurol* 66(5):632–637.
- 736 Sanfilippo C, Forlenza O, Zetterberg H, Blennow K (2016) Increased  
737 neurogranin concentrations in cerebrospinal fluid of Alzheimer's  
738 disease and in mild cognitive impairment due to AD. *J Neural  
739 Transm (Vienna)* 123(12):1443–1447.
- 740 Scheff SW, Price DA, Schmitt FA, DeKosky ST, Mufson EJ (2007)  
741 Synaptic alterations in CA1 in mild Alzheimer disease and mild  
742 cognitive impairment. *Neurology* 68(18):1501–1508.
- 743 Sollner T, Bennett MK, Whiteheart SW, Scheller RH, Rothman JE  
744 (1993a) A protein assembly-disassembly pathway in vitro that  
745 may correspond to sequential steps of synaptic vesicle docking,  
746 activation, and fusion. *Cell* 75(3):409–418.
- 747 Sollner T, Whiteheart SW, Brunner M, Erdjument-Bromage H,  
748 Geromanos S, Tempst P, Rothman JE (1993b) SNAP receptors  
749 implicated in vesicle targeting and fusion. *Nature* 362  
750 (6418):318–324.
- 751 Sperling RA, Aisen PS, Beckett LA, Bennett DA, Craft S, Fagan AM,  
752 Iwatsubo T, Jack Jr CR, Kaye J, Montine TJ, Park DC, Reiman  
753 EM, Rowe CC, Siemers E, Stern Y, Yaffe K, Carrillo MC, Thies B,  
754 Morrison-Bogorad M, Wagster MV, Phelps CH (2011) Toward  
755 defining the preclinical stages of Alzheimer's disease:  
756 recommendations from the National Institute on Aging-  
757 Alzheimer's Association workgroups on diagnostic guidelines for  
758 Alzheimer's disease. *Alzheimers Dement* 7(3):280–292.
- 759 Sudhof TC (2004) The synaptic vesicle cycle. *Annu Rev Neurosci*  
760 27:509–547.
- 761 Tapiola T, Alafuzoff I, Herukka SK, Parkkinen L, Hartikainen P,  
762 Soininen H, Pirttilä T (2009) Cerebrospinal fluid {beta}-amyloid 42  
763 and tau proteins as biomarkers of Alzheimer-type pathologic  
764 changes in the brain. *Arch Neurol* 66(3):382–389.
- 765 Veit M, Sollner TH, Rothman JE (1996) Multiple palmitoylation of  
766 synaptotagmin and the t-SNARE SNAP-25. *FEBS Lett* 385(1–  
767 2):119–123.
- 768 Wellington H, Paterson RW, Portelius E, Tornqvist U, Magdalinou N,  
769 Fox NC, Blennow K, Schott JM, Zetterberg H (2016) Increased  
770 CSF neurogranin concentration is specific to Alzheimer disease.  
771 *Neurology* 86(9):829–835.

(Received 27 April 2018, Accepted 28 November 2018)  
(Available online xxxx)