

Exploration and Development of a C–H-Activated Route to Access the [1,2]Dithiolo[4,3-*b*]indole-3(4*H*)-thione Core and Related Derivatives

Christopher R. M. Asquith *^{a,1} Lidia S. Konstantinova^{b,c}Graham J. Tizzard^dTuomo Laitinen^eSimon J. Coles^dOleg A. Rakitin^{b,c}Stephen T. Hilton *^a

^a School of Pharmacy, Faculty of Life Sciences, University College London, London, WC1N 1AX, UK
ucnvcrm@ucl.ac.uk
s.hilton@ucl.ac.uk

^b N.D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, Moscow, 119991, Russia

^c Nanotechnology Education and Research Center, South Ural State University, Lenina Ave. 76, Chelyabinsk, Russia

^d School of Chemistry, University of Southampton, Highfield, Southampton SO17 1BJ, UK

^e School of Pharmacy, Faculty of Health Sciences, University of Eastern Finland, Kuopio, 70211, Finland

Received: 08.10.2018

Accepted after revision: 19.11.2018

Published online: 19.12.2018

DOI: 10.1055/s-0037-1611692; Art ID: st-2018-v0647-l

Abstract A robust procedure for the production of [1,2]dithiolo[4,3-*b*]indole-3(4*H*)-thione analogues using a DABCO/S₂Cl₂ complex as a sulfur source via a C–H activated approach.

Key words indole, indole sulfides, sulfur-nitrogen heterocycles, fused 3*H*-1,2-dithiole-3-thiones, C–H activation, disulfur dichloride

The privileged indole structure has been shown to be a favorable biological scaffold, and despite a plethora of examples of known drugs containing indole only limited indole polysulfide research has been carried out.² This is despite the fact that examples of sulfur-containing indoles such as **1–4** show various biological activities including tyrosine kinase inhibition,³ anti-inflammation,⁴ cytoprotectant activity,⁵ antifungal activity,⁶ antiviral properties,⁷ cannabinoid receptor binding,⁸ and ion channel modulation.⁹ (Figure 1). As part of a wider research program within our group, we became interested in the electronics of the ring system attached to a thione functionality⁶ and its potential biological applications, along with development of a robust mild synthetic route to access them.

There are several examples in the literature of synthetic methodologies towards indole polysulfides. Initial work by Snyder *et al.* and Wawzonek *et al.* involved heating elemental sulfur with indole and cyclic amines, respectively, producing a number of sulfides, with little control of the products produced.^{10,11} Several methods have since been employed to control the sulfur addition and improve the yields including use of thioureas,¹² sulfur,¹³ thiols,¹⁴ sulphenyl






Figure 1 Examples of biologically active indole sulfides

chlorides,¹⁵ disulfides,¹⁶ and more recently sulfonyl hydrazides¹⁷ and carbon disulfide/iodine.¹⁸ Despite a number of methods and sulfur sources, the chemistry described has limited scope and suffers from a lack of full exploration. Broad, general methods have yet to be developed.

This trend continues when looking into the indole attached to a cyclized ring systems such as the pentathiepin^{6,7,19} and the 3*H*-1,2-dithiole-3-thione. The latter caught our interest due in part to its unusual electronic properties which provide a potentially mutually beneficial stabilizing electronic effect for both the indole and thione parts of the molecule. We sought to develop a robust syn-

thetic pathway to access these molecules in order to open this class of compound up to the wider medicinal chemistry applications.

There are limited examples of [1,2]dithiolo[4,3-*b*]indole-3(4*H*)-thione in the literature with synthesis tending to lack substrate scope, likely due in part to the complex mechanistic pathways required for these reactions to proceed. There are two potential isomers of the indole thione, the *alpha* (**5**) and *Pri* (**6**). Previous routes to access these compounds proceed using phosphorous sulfide to form **5** from **7**, and have limited scope (Scheme 1).²⁰ Other examples include a two-step phosphorous sulfide, carbon disulfide combination to produce **6** from **9**,²¹ and a method using carbon disulfide produces compounds **6** from **10** in a three-step protocol.²² These methods produce **5** and **6** in low yields, consistent with more recent reports on analogous ring systems.¹⁹ However utilizing the disulfur dichloride/DABCO complex²³ enabled the 3*H*-1,2-dithiole-3-thione to be accessed in a respectable yield in one step.²⁴



Scheme 1 Previously reported routes to fused 3*H*-1,2-dithiole-3-thiones **6** and **9**

To date there has been little exploration to produce a route that can reliably and reproducibly generate a library of compounds under mild conditions in high yield. We now present a robust one-pot route to synthesize a broad range of cyclized indole sulfide derivatives. We confirm their structures with X-ray crystallography.²⁵ We first treated 2-methyl indoles **11a,b** with a series of alkyl halides using mild conditions (method A: powdered potassium hydroxide in DMF at room temperature, 48 h). In some cases (**12**

a–p) that were sluggish or proceeded in low yields we utilized Finkelstein-type conditions (conditions B) utilizing sodium hydride and potassium iodide (Table 1). A head to head comparison of the two methods showed that method B can increase the yield by a factor of three in certain cases. The yields tended to be higher with the *meta*-substituted derivatives (**12f** and **12k**), which also corresponded to a milder reaction requirement.

Table 1 Results of Coupling Reactions on 2-Methyl-Indoles

11a,b		KOH or NaH/KI, R'-Br DMF or THF 48 or 96 h, 20 °C	12a–q	
Name	R	R'	Method	Yield (%) ^a
12a	H	methyl	A	81
12b	H	ethyl	A	74
12c	H	isopropyl	A	51
12d	H	(2-methyl)propyl	A	52
12e	H	2-fluorobenzyl	B	63
12f	H	3-fluorobenzyl	A	72
12g	H	4-fluorobenzyl	B	81
12h	H	3,4-difluorobenzyl	A	56
12i	H	propylbenzene	A	91
12j	H	2-cyanobenzyl	B	52
12k	H	3-cyanobenzyl	A	74
12l	H	4-cyanobenzyl	A	25
12l	H	4-cyanobenzyl	B	64
12m	H	benzyl	A	65
12n	5-F	benzyl	A	67
12o	H	2-methoxybenzyl	A	83
12p	H	3-methoxybenzyl	A	56

^a Yield of isolated product.

2-Methyl indole (**11a**) was also treated with corresponding aryl bromides using standard Ullman conditions at 180 °C for 45 min in the microwave to afford **13a,b** in good yield (Table 2). This allowed for testing the versatility of the incorporation of sulfur on indole N-aryl systems.²⁶

Table 2 Ullman Coupling Reactions on 2-Methyl-Indole (**11a**)

11a		Cs ₂ CO ₃ , R-Br, CuI DMF, 180 °C, 45 min microwave	13a,b	
Name	R	Method	Yield (%) ^a	
13a	3,4-dimethoxybenzene	C	82	
13b	3-fluoro-4-methoxybenzene	C	81	

^a Yield of isolated product.

With the substituted indoles (**11a**, **12a–q** and **13a,b**) in hand we treated them with the disulfur dichloride/DABCO complex in an operationally simple three-step, one-pot protocol.²⁴ The reaction can be followed through a series of visual transitions between a set of previous proposed intermediates.²⁴ We were able to access the [1,2]dithiolo[4,3-*b*]indole-3(4*H*)-thione products **14a–s** in moderate to excellent yields (Table 3). In keeping with our previous observation, the *meta*-substituted indoles (**12F** and **12K**) had the highest yields. We did, however, notice a drop-off in yield in the production of **14r,s**; this is likely due to the potential steric and electronic interference of the directly bound phenyl ring system. Overall this method is operationally simple, robust, and amenable to scale up and array synthesis.

Table 3 [1,2]Dithiolo[4,3-*b*]indole-3(4*H*)-thione Results

Name	R	R'		Yield (%) ^a
14a	H	N-H		27
14b	H	N-methyl		94 ^b
14c	H	N-ethyl		56 ^b
14d	H	N-isopropyl		66 ^b
14e	H	N-(2-methyl)propyl		52
14f	H	2-fluoro-N-benzyl		64
14g	H	3-fluoro-N-benzyl		79
14h	H	4-fluoro-N-benzyl		71
14i	H	3,4-difluoro-N-benzyl		39
14j	H	N-propylbenzene		62
14k	H	2-cyano-N-benzyl		74
14l	H	3-cyano-N-benzyl		82
14m	H	4-cyano-N-benzyl		73
14n	H	N-benzyl		70 ^b
14o	5-F	N-benzyl		63
14p	H	2-methoxy-N-benzyl		58
14q	H	3-methoxy-N-benzyl		86
14r	H	3,4-dimethoxy-N-benzene		34
14s	H	3-fluoro-4-methoxy-N-benzene		24

^a Yield of isolated product.

^b Consistent with previous report.²⁴

The disulfide bridges contained within these compounds have never been investigated and we wondered if the stabilizing effect of the indole would lengthen or shorten the thio carbonyl or the disulfide bridge head. The typical bond length for a S-S bond is 2.07 Å and C=S is 1.63 Å.²⁶ We observed a slight lengthening of the C=S bond by 0.01–

0.04 Å in all structures except **14q** and the minor disorder component of **14h**. The only significant deviation from the typical S-S bond length was observed in the minor disorder component of **14h** (Figure 2, Table 4).




Figure 2 *ortho*-Cyano (**14k**)/*meta*-cyano (**14l**)/*para*-cyano (**14m**) crystal structures showing orthogonal ring system.

Table 4 [1,2]Dithiolo[4,3-*b*]indole-3(4*H*)-thione **14a–t** Crystal Structure Results Showing S–S and S–C Bonds

Name	S–S	S–C	Crystal structure	
14a	2.073(2)	1.664(4)	triclinic	plate
14b	2.0755(9)	1.670(3)	monoclinic	block
14c	–	–	–	–
14d	–	–	–	–
14e	2.0668(13)	1.653(3)	monoclinic	block
14f	2.0717(10)	1.651(3)	trigonal	block
14g	2.0734(6)	1.6700(17)	monoclinic	block
14h	2.0745(17) 2.001(16)	1.655(4) 1.604(16)	monoclinic	plate
14i	2.065(2) 2.067(2)	1.669(6) 1.669(5)	triclinic	plate
14j	2.0772(13)	1.656(3)	orthorhombic	blade
14k	2.0701(12)	1.647(3)	orthorhombic	block
14l	2.0751(7)	1.6620(19)	monoclinic	block
14m	2.0752(7)	1.6612(19)	monoclinic	block
14n	–	–	–	–
14o	2.0710(6) 2.0746(6)	1.6549(18) 1.6532(18)	monoclinic	rod
14p	2.0669(19)	1.554(5)	triclinic	plate
14q	2.0766(6)	1.6643(16)	monoclinic	block
14r	2.0634(9)	1.657(3)	monoclinic	block
14s	2.0799(19) 2.079(2)	1.664(6) 1.663(6)	orthorhombic	needle

There are no hydrogen-bond donors or acceptors present in these compounds (aside from the indole N–H in **14a**) so the crystal structures arise from simple close-packing. The only exception to this is the higher symmetry trigonal structure of **14f** which is solvent mediated (Table 4). The smaller R' residues of **14a** and **14b** result in structures comprising of simple stacks of alternately orientated molecules whilst the larger R' residues of **14e–s** give rise to more complex stacking arrangements of molecules (Table 4).

The observed lengthening of S–S bonds in X-ray structures led us to study how well commonly used electronic-structure calculations are able to reproduce these X-ray results. We used crystal-form conformation for the compound **14l** and calculated standard geometry optimizations using standard DFT geometry optimization (B3LYP/6-31G*) using Gaussian09 software.²⁸ Results reveal that this level of theory, which usually produces reliable molecular geometries, is actually overestimating the S–S bond lengths of [1,2]dithiolo system; giving a 2.13 Å length for S–S bond in the case of **14l**, which has an actual length of 2.08 by X-ray. Whereas with a length of 1.67 Å for C=S bond, this is more consistent with our experimental data. Optimization using higher level basis sets B3LYP/6-311+G** and B3LYP/cc-PVTZ did not improve the performance of DFT while both unconstrained optimizations resulted in a calculated 2.14 Å bond length for the S–S bond in question.²⁸

This C–H activation to produce the 3*H*-1,2-dithiole-3-thiones has been observed as a thermodynamic product of a reaction of methyl indole with elemental sulfur.²⁹ To our knowledge this is the first report of crystallization, full crystallographic structural determination, and characterization of the [1,2]dithiolo[4,3-*b*]indole-3(4*H*)-thione core functionality.³⁰ This work has demonstrated an efficient synthetic route to 3*H*-1,2-dithiole-3-thiones with broad substrate scope which we hope to utilize in further investigation of the potential of disulfide bridge networks in medicinal and organic chemistry.

Acknowledgment

The authors are grateful to Bloomsbury Colleges – University of London; University College London; Russian Science Foundation (grant no. 15-13-10022), and the Biocenter Finland/DDCB for financial support towards the goals of our work. We would also like to thank the EPSRC UK National Mass Spectrometry Facility at the Swansea University for spectroscopic services and the EPSRC UK National Crystallography Service at the University of Southampton for the collection of the crystallographic data. In addition, we thank CSC – IT Center for Science Ltd. for software licenses and computational resources and Prof. David H. Drewry and Prof. Timothy M. Wilson (University of North Carolina – Chapel Hill, NC, USA) for their insights in preparation of this manuscript.

References and Notes

- (1) Current address - Department of Pharmacology, School of Medicine, University of North Carolina at Chapel Hill, NC 27599, USA
- (2) de Sá Alves, F. R.; Barreiro, E. J.; Fraga, C. A. *Mini-Rev. Med. Chem.* **2009**, *9*, 782.
- (3) (a) Rewcastle, G. W.; Denny, W. A. *Heterocycles* **1994**, *37*, 701. (b) Palmer, B. D.; Rewcastle, G. W.; Thompson, A. M.; Boyd, M.; Showalter, H. D.; Sercel, A. D.; Fry, D. W.; Kraker, A. J.; Denny, W. A. *J. Med. Chem.* **1995**, *38*, 58. (c) Dobrusin, E. M.; Showalter, H. D. H.; Denny, W. A.; Rewcastle, G. W.; Tercel, M.; Thompson, A. M. US 5464861, **1994**.
- (4) Greenhouse, R. J.; Muchowski, J. M. US 4654360, **1987**.
- (5) Hosoi, M. K.; Fumio, N. O.; Ichikawa, H. T.; Ichikawa, K. N.; Okazaki, S. N. US 4743609, **1988**.
- (6) Asquith, C. R. M.; Meli, M. L.; Konstantinova, L. S.; Laitinen, T.; Poso, A.; Rakitin, O. A.; Hofmann-Lehmann, R.; Allenspach, K.; Hilton, S. T. *Bioorg. Med. Chem. Lett.* **2015**, *25*, 1352.
- (7) Rewcastle, G. W.; Janosik, T.; Bergman, J. *Tetrahedron* **2001**, *57*, 7185.
- (8) (a) Friary, R. J.; Zhou, G.; Shih, N.; Chen, L.; Shu, Y.; Tong, L.; Lavey, B. J.; Wong, M. K. C.; Kozlowski, J. A.; Shankar, B. B. US 200410013, **2004**. (b) Shankar, B. B.; Gilbert, E.; Huang, C.; McCombie, S.; Shih, N.; Kozlowski, J. A.; Rizvi, R. K. WO 20062133, **2006**.
- (9) Gao, D.; Sand, R.; Fu, H.; Sharmin, N.; Gallin, W. J.; Hall, D. G. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 5503.
- (10) Carpenter, W.; Grant, M. S.; Snyder, H. R. *J. Am. Chem. Soc.* **1960**, *82*, 2739; and references within.
- (11) Wawzonek, S.; Hansen, G. R. *J. Org. Chem.* **1966**, *31*, 3580.
- (12) Harris, R. L. N. *Aust J Chem.* **1970**, *23*, 1199.
- (13) Kobayashi, K.; Kobayashi, A.; Ezaki, K. *Tetrahedron* **2013**, *69*, 7936.
- (14) Zhang, H.; Bao, X.; Song, Y.; Qu, J.; Wang, B. *Tetrahedron* **2015**, *71*, 8885.
- (15) (a) Hamel, P.; Prévile, P. *J. Org. Chem.* **1996**, *61*, 1573. (b) Hamel, P. *J. Org. Chem.* **2002**, *67*, 2854. (c) Plate, R.; Nivard, R. J. F.; Ottenheijm, H. C. J. *Tetrahedron* **1986**, *42*, 4503.
- (16) (a) Prasad, C. D.; Kumar, S.; Sattar, M.; Adhikary, A.; Kumar, S. *Org. Biomol. Chem.* **2013**, *11*, 8036. (b) Silveira, C. C.; Mendes, S. R.; Wolf, L.; Martins, G. M.; Mühlen, L. V. *Tetrahedron* **2012**, *68*, 10464. (c) Hamel, P.; Zajac, N.; Atkinson, J. G.; Girard, Y. *J. Org. Chem.* **1994**, *59*, 6372. (d) Hamel, P.; Girard, Y.; Atkinson, J. G. *J. Org. Chem.* **1992**, *57*, 2694. (e) Anzai, K. *J. Heterocycl. Chem.* **1979**, *16*, 567.
- (17) Yang, F. L.; Tian, S. K. *Angew. Chem. Int. Ed.* **2013**, *18*, 4929.
- (18) Halimehjani, A. Z.; Shokrgozar, S.; Beier, P. *J. Org. Chem.* **2018**, *83*, 5778.
- (19) (a) Konstantinova, L. S.; Rakitin, O. A.; Rees, C. W. *Chem. Commun.* **2002**, *11*, 1204. (b) Amelichev, S. A.; Konstantinova, L. S.; Lyssenko, K. A.; Rakitin, O. A.; Rees, C. W. *Org. Biomol. Chem.* **2005**, *19*, 3496. (c) Amelichev, S. A.; Aysin, R. R.; Konstantinova, L. S.; Obruchnikova, N. V.; Rakitin, O. A.; Rees, C. W. *Org. Lett.* **2005**, *25*, 5725. (d) Konstantinova, L. S.; Amelichev, S. A.; Rakitin, O. A. *Russ. Chem. Bull.* **2006**, *55*, 2081. (e) Janosik, T.; Bergman, J.; Stensland, B.; Stalhandske, C. *J. Chem. Soc., Perkin Trans. 1.* **2002**, *330*. (f) Bergman, J.; Stalhandske, C. *Tetrahedron Lett.* **1994**, *5279*. (g) Konstantinova, L. S.; Amelichev, S. A.; Rakitin, O. A. *Russ. Chem. Rev.* **2007**, *76*, 195.
- (20) Jardine, R. V.; Brown, R. K. *Can. J. Chem.* **1965**, *43*, 1293.
- (21) Kobayashi, G.; Furukawa, S.; Matsuda, Y.; Natsuki, R. *Yakugaku Zasshi* **1970**, *90*, 132.
- (22) Rewcastle, G. W.; Denny, W. A. *Heterocycles* **1994**, *37*, 701.

- (23) (a) Konstantinova, L. S.; Rakitin, O. A.; Rees, C. W.; Amelichev, S. A. *Mendeleev Commun.* **2004**, 91. (b) Konstantinova, L. S.; Rakitin, O. A. *Mendeleev Commun.* **2009**, 55.
- (24) Konstantinova, L. S.; Lysov, K. A.; Amelichev, S. A.; Obruchnikova, N. V.; Rakitin, O. A. *Tetrahedron* **2009**, 65, 2178.
- (25) CCDC 1517984–1517999 contain the supplementary crystallographic data for **14a,b,e–m** and **14o–s**. The data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/getstructures
- (26) Sambiagio, C.; Marsden, S. P.; Blacker, A. J.; McGowan, P. C. *Chem. Soc. Rev.* **2014**, Recent Ullman Coupling review: 10, 3525.
- (27) Cremlyn, R. J. *An Introduction to Organosulfur Chemistry*; John Wiley and Sons: Chichester, **1996**.
- (28) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A. Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Liyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. *Gaussian 09, Revision E.01*; Gaussian, Inc: Wallingford, CT, **2009**.
- (29) Schmitt, J.; Lespagnol, A. *Bull. Soc. Chim. Fr.* **1950**, 459.
- (30) **General Procedures**

Method A

2-Methylindole (1.31 g, 10.0 mmol) in dry DMF (15 mL) was treated with powdered KOH (0.67 g, 12.0 mmol) and the mixture stirred for 30 min, whereupon 3-(bromomethyl)benzonitrile (1.99 g, 10.0 mmol) was added in one portion. The reaction became exothermic and was cooled in an ice bath and after stirring for 96 h at r.t., the mixture was concentrated under vacuum and the residue taken up in EtOAc. The solution was washed with water, brine, dried (MgSO_4), and concentrated to dryness *in vacuo*. Purification *via* flash column chromatography on silica gel (hexanes/EtOAc, 1:1) to afford 3-[(2-methyl-1H-indol-1-yl)methyl]benzonitrile (**12k**, 1.95 g, 7.4 mmol, 74% yield) as colorless crystals; mp 120–121 °C. HMRS: m/z calcd for $\text{C}_{11}\text{H}_{14}\text{N}_2$ [M + H]⁺: 247.1235; found: 247.1231. IR (film): ν_{\max} = 3019 m (C–H), 2945 m (C–H), 2915 m (C–H), 2224 s (C≡N), 1552 m (C=C), 1457 m, 1426 w, 1395 w, 1136 w, 783 m 744 s, 686 m cm^{-1} . ¹H NMR (400 MHz, CDCl_3): δ = 7.60–7.57 (1 H, m, CH^{Ar}), 7.54 (1 H, d, J = 8.1 Hz, CH^{Ar}), 7.38 (1 H, t, J = 7.8 Hz, CH^{Ar}), 7.28 (1 H, s, CH^{Ar}), 7.17 (1 H, dq, J = 1.0, 8.1 Hz, CH^{Ar}), 7.14–7.11 (3 H, m, CH^{Ar}), 6.38 (1 H, s, CH^{Ar}), 5.43 (2 H, s, CH_2), 2.37 (3 H, s, CH_3).

¹³C NMR (100 MHz, CDCl_3): δ = 139.6 (C^{Ar}), 136.9 (C^{Ar}), 136.1 (C^{Ar}), 131.2 (CH^{Ar}), 130.4 (CH^{Ar}), 129.7 (CH^{Ar}), 129.5 (CH^{Ar}), 128.3 (C^{Ar}), 121.2 (CH^{Ar}), 120.1 (CH^{Ar}), 120.0 (CH^{Ar}), 118.5 (C^{Ar}), 113.1 (C≡N), 108.8 (CH^{Ar}), 101.3 (CH^{Ar}), 45.7 (CH_2), 12.7 (CH_3).

Method B

Sodium hydride (0.5 g of a 60% dispersion in mineral oil, 12.5 mmol) was added portionwise to a solution of 2-methylindole (1.31 g, 10.0 mmol) in dry THF (20 mL). The mixture was stirred for 30 min, and then 2-(bromomethyl)benzonitrile (1.96 g, 10.0 mmol) was added in one portion, in addition to NaI (1.50 g, 10.0 mmol). The reaction became exothermic and was cooled in an ice bath and after stirring for 48 h at r.t., it was concentrated *in vacuo* and the residue taken up in EtOAc. The solution was washed with water, brine, dried (MgSO_4), and concentrated to dryness *in vacuo*. Purification *via* flash column chromatography on silica gel (hexanes/EtOAc, 1:1) to afford 2-[(2-methyl-1H-indol-1-yl)methyl]benzonitrile (**12j**, 1.28 g, 5.2 mmol, 52% yield) as colorless crystals; mp 124–125 °C. HMRS: m/z calcd for $\text{C}_{11}\text{H}_{15}\text{N}_2$ [M + H]⁺: 247.1235; found: 247.1233. IR (film): ν_{\max} = 3055 m (C–H), 2920 m (C–H), 2851 m (C–H), 2220 s (C≡N), 1460 m (C=C), 1396 w, 747 s, 703 s cm^{-1} . ¹H NMR (400 MHz, CDCl_3): δ = 7.74 (1 H, dd, J = 1.2, 7.0 Hz, CH^{Ar}), 7.62–7.59 (1 H, m, CH^{Ar}), 7.40–7.34 (2 H, m, CH^{Ar}), 7.16–7.12 (3 H, m, CH^{Ar}), 6.52 (1 H, d, J = 6.3, CH^{Ar}), 6.41 (1 H, s, CH^{Ar}), 5.55 (2 H, s, CH_2), 2.39 (3 H, s, CH_3). ¹³C NMR (100 MHz, CDCl_3): δ = 141.7 (C^{Ar}), 137.0 (C^{Ar}), 136.4 (C^{Ar}), 133.5 (CH^{Ar}), 132.9 (CH^{Ar}), 128.3 (C^{Ar}), 127.8 (CH^{Ar}), 126.5 (CH^{Ar}), 121.2 (CH^{Ar}), 120.0 (CH^{Ar}), 117.1 (C^{Ar}), 110.1 (C≡N), 108.9 (CH^{Ar}), 101.2 (CH^{Ar}), 44.7(CH_2), 12.6 (CH_3).

Method C

Disulfur dichloride (0.4 mL, 5 mmol) was added dropwise at –35 °C to a stirred solution of DABCO (1.12 g, 10 mmol) in chloroform (25 mL) under nitrogen. The mixture was stirred at r.t. for 1 h. 3-[(2-Methyl-1H-indol-1-yl)methyl]benzonitrile (**12j**, 0.239 g, 1 mmol) in chloroform (5 mL) was added, and the mixture was stirred at r.t. for 48 h under nitrogen. Triethylamine (1.4 mL, 10 mmol) was then added to the resultant mixture at 0 °C, the mixture stirred at r.t. for 2 h, heated at reflux for 3 h, filtered, and solvents evaporated. Purification *via* flash column chromatography on silica gel (hexanes/ CH_2Cl_2) to afford 3-[(3-thioxo-[1,2]dithiolo[4,3-*b*]indol-4(3H)-yl)methyl]benzonitrile (**14l**) (0.277 g, 0.82 mmol, 82% yield) as orange crystals; mp 190–191 °C. HMRS: m/z calcd for $\text{C}_{17}\text{H}_{11}\text{N}_2\text{S}_3$ [M + H]⁺: 339.0079; found: 339.0077. IR (film): ν_{\max} = 3054 m (C–H), 2952 m (C–H), 2921 m (C–H), 2872 w (C–H), 2851 w (C–H), 2226 s (C≡N), 1472 s (C=C), 1330 s, 1258 w, 1122 w, 1060 s, 685 m cm^{-1} . (400 MHz, CDCl_3): δ = 7.77 (1 H, dt, J = 1.0, 8.0 Hz, CH^{Ar}), 7.46 (2 H, ddd, J = 1.0, 4.8, 5.9 Hz, CH^{Ar}), 7.40–7.32 (3 H, m, CH^{Ar}), 7.24–7.19 (2 H, m, CH^{Ar}), 6.15 (2 H, s, CH^{Ar}). ¹³C NMR (100 MHz, CDCl_3): δ = 195.9 (C=S), 146.9 (C^{Ar}), 142.2 (C^{Ar}), 149.4 (C^{Ar}), 139.0 (C^{Ar}), 131.4 (CH^{Ar}), 131.3 (C^{Ar}), 130.3 (CH^{Ar}), 129.6 (CH^{Ar}), 125.5 (C^{Ar}), 121.9 (CH^{Ar}), 121.7 (CH^{Ar}), 120.6 (CH^{Ar}), 114.3 (CH^{Ar}), 113.0 (C≡N), 111.8 (CH^{Ar}), 44.0 (CH_2).