

## Hemostatic spray powder TC-325 in the primary endoscopic treatment of peptic ulcer related bleeding: multicentre international registry

Mohamed Hussein<sup>1</sup>, Durayd Alzoubaidi<sup>1</sup>, Miguel-Fraile Lopez<sup>2</sup>, Michael Weaver<sup>3</sup>, Jacobo Ortiz-Fernandez-Sordo<sup>2</sup>, Paul Bassett<sup>4</sup>, Johannes W Rey<sup>5</sup>, Bu'Hussain Hayee<sup>6</sup>, Edward Despott<sup>7</sup>, Alberto Murino<sup>7</sup>, Sulleman Moreea<sup>8</sup>, Philip Boger<sup>9</sup>, Jason Dunn<sup>10</sup>, Inder Mainie<sup>11</sup>, David Graham<sup>12</sup>, Daniel K Mullady<sup>3</sup>, Dayna S Early<sup>3</sup>, Krish Rangunath<sup>2</sup>, John T Anderson<sup>13</sup>, Pradeep Bhandari<sup>14</sup>, Martin Goetz<sup>15</sup>, Ralf Kiesslich<sup>16</sup>, Emmanuel Coron<sup>17</sup>, Laurence B Lovat<sup>1</sup>, Rehan Haidry<sup>1,12</sup>

### Supplementary material

**Table 1s:** Outcomes in the combination therapy subgroups

	<b>Hemospray + adrenaline injection (n = 36)</b>	<b>Hemospray + clips (n=12)</b>	<b>Hemospray + adrenaline + clips (n =31)</b>	<b>Hemospray + adrenaline + thermal (n = 13)</b>	<b>Hemospray + adrenaline + thermal + clips</b>
<b>Haemostasis</b>	32/36 (89%)	11/12 (92%)	28/31 (90%)	10/13 (77%)	7/7 (100%)
<b>Median Rockall score (IQR)</b>	7 (6-8)	7 (6-8)	7 (6-8)	8 (6-8)	7 (6-8)
<b>Median Rockall score (IQR)</b>	13 (10-14)	15 (12-16)	11 (8-14)	13 (11-14)	12 (12-15)
<b>Re-bleeding</b>	4/30 (13%)	2/9 (22%)	4/26 (15%)	1/10 (10%)	1/6 (17%)
<b>7- day mortality</b>	2/33 (6%)	3/10 (30%)	2/28 (7%)	1/10 (10%)	1/6 (17%)
<b>30-day mortality</b>	5/33 (15%)	0	4/28 (14%)	0	0