

The 2020 release of the ExoMol database: molecular line lists for exoplanet and other hot atmospheres

Jonathan Tennyson^{1*}, Sergei N. Yurchenko¹, Ahmed F. Al-Refaie¹, Victoria H. J. Clark¹, Katy L. Chubb^{1,2}, Eamon K. Conway^{1,9}, Akhil Dewan¹, Maire N. Gorman^{1,3}, Christian Hill^{1,4}, A. E. Lynas-Gray^{1,7,8}, Thomas Mellor¹, Laura K. McKemmish^{1,5}, Alec Owens¹, Oleg L. Polyansky¹, Mikhail Semenov¹, Wilfrid Somogyi¹, Giovanna Tinetti¹, Apoorva Upadhyay¹, Ingo Waldmann¹, Yixin Wang^{1,6}, Samuel Wright¹, Olga P. Yurchenko¹

¹Department of Physics and Astronomy, University College London, London, WC1E 6BT, UK

²Present address: SRON Netherlands Institute for Space Research, Sorbonnelaan 2, 3584 CA, Utrecht, Netherlands

³ Present address: Department of Physics, Aberystwyth University, Ceredigion, UK, SY23 3BZ, UK

⁴Present address: Nuclear Data Section, International Atomic Energy Agency, Vienna A-1400, Austria

⁵ Present address: School of Chemistry, University of New South Wales, 2052 Sydney, Australia

⁶ Present address: Nankai University, 94 Weijin Road, Tianjin, China

⁷ Department of Physics, University of Oxford, England

⁸ Department of Physics and Astronomy, University of the Western Cape, South Africa

⁹ Atomic and Molecular Physics Division, Center for Astrophysics | Harvard & Smithsonian, Cambridge, MA. USA.

Abstract

The ExoMol database (www.exomol.com) provides molecular data for spectroscopic studies of hot atmospheres. While the data is intended for studies of exoplanets and other astronomical bodies, the dataset is widely applicable. The basic form of the database is extensive line lists; these are supplemented with partition functions, state lifetimes, cooling functions, Landé g-factors, temperature-dependent cross sections, opacities, pressure broadening parameters, k -coefficients and dipoles. This paper presents the latest release of the database which has been expanded to consider 80 molecules and 190 isotopologues totaling over 700 billion transitions. While the spectroscopic data is

concentrated at infrared and visible wavelengths, ultraviolet transitions are being increasingly considered in response to requests from observers. The core of the database comes from the ExoMol project which primarily uses theoretical methods, albeit usually fine-tuned to reproduce laboratory spectra, to generate very extensive line lists for studies of hot bodies. The data has recently been supplemented by line lists deriving from direct laboratory observations, albeit usually with the use of *ab initio* transition intensities. A major push in the new release is towards accurate characterisation of transition frequencies for use in high resolution studies of exoplanets and other bodies.

Keywords: infrared, visible, Einstein A coefficients, transition frequencies, partition functions, cooling functions, lifetimes, cross sections, k coefficients, Landé g -factors, pressure broadening

1. Introduction

The ExoMol project was started in 2011 [1] with the purpose of providing molecular line lists for studies of exoplanets and other (hot) atmospheres. Besides data demands for exoplanets [2], other hot astronomical bodies with significant molecular content in their atmospheres include cool stars [3] and brown dwarfs [3, 4]. On Earth, similar spectroscopic data are required to study flames [5, 6], discharge plasmas [7], explosions [8] and the gases emitted from smoke stacks [9]. In addition, bodies in non-local thermodynamic equilibrium (non-LTE) such as comets and masers give observable emissions from highly excited states [10–12].

ExoMol data are proving very popular among the exoplanet atmospheric modellers owing to their extensive coverage of the molecular species and their completeness as function of both frequency and, very importantly, temperature. ExoMol data have been incorporated into many radiative transfer and retrieval codes including Tau-REx [13–15], Phoenix [16], NEMESIS [17], CHIMERA [18], HELIOS [19] and HELIOS-r2 [20], ATMO [21–23], ARCiS [24], ARTES [25], HyDRA [26], GENESIS [27], petitRADTRANS [28], PLATON [29], VSTAR [30, 31], BART [32] and Pyrat Bay [33]. In addition the ExoMol database has been used to support a variety of other studies. Examples including the analysis of gaseous ammonia in Jupiter [34], tentative detection of H₂S in Uranus [35], detection of CaO in meteors [36] and SiO in B[e] Supergiants [37], Late-type stars [38] or circumstellar environment [39]. Detections in the atmospheres of exoplanets are discussed below. More diverse applications include modeling laser analysis of Solar System objects [40, 41], isotope abundance quantification in stars [42], search for molecules highly sensitive to proton-to-electron mass ratio variation [43, 44], models of molecular steering [45, 46], design of THz lasers [47], design of dark matter detection schemes [48] and the design of novel propulsion systems [49].

A number of diatomic molecules for which ExoMol provides extensive hot line

lists have been detected in sunspots. These include SiO [50], SiH [51], SH [52], BeH [53], VO [54], as well as MgH, MgO, TiO, C₂ and CaH which have all long been observed in sunspots [55]. In addition the spectrum of water is well-known in sunspots [56–60]; studies of water spectra in sunspots should be aided by the new POKAZATEL water line list [61] which is designed for studies at temperatures above 3000 K such as those encountered in solar umbra and penumbra.

ExoMol line lists have been published as a series in the journal Monthly Notices of the Royal Astronomical Society (see below) and a first data release in 2016 [62], henceforth ExoMol2016, provided full documentation for the ExoMol database which can be found at www.exomol.com. The basic form of the database is extensive line lists; these are supplemented with partition functions, state lifetimes, cooling functions, Landé g-factors, temperature-dependent cross sections, opacities, pressure broadening parameters, *k*-coefficients and dipoles. The ExoMol website also offers an extensive bibliography database on research literature on molecules relevant for ExoMol applications. So far the bibliography database contains 6709 sources.

Since ExoMol started there have been major changes in exoplanet science which have driven further expansion and development of the ExoMol database. The first is the discovery of rocky exoplanets orbiting so near to their host star that their surfaces are likely to be molten. The atmospheres of these planets are thought to contain a variety of species not considered previously [63–66]. Secondly higher temperature planets have been observed than anyone anticipated, the poster child for hot exoplanets is Kelt-9b which is thought to reach temperatures of 4000 K [67], hotter than most of the stars in our Galaxy! This has led to the need to construct line list for key species over an extended temperature range. Third is the development of Doppler-shift high resolution spectroscopy of exoplanets [68, 69] which has proved a powerful tool for detecting molecules but fails in the absence of highly accurate molecular line lists [70]. These develop-

ments have led us to both expand the range of molecules included in the database and to begin a systematic attempt to improve the accuracy of the line positions for the line lists contained in the database. Progress on both of these objectives is described below as well work on extending the coverage of the database into the ultraviolet.

Of course ExoMol is not the only database providing spectroscopic data for atmospheric studies. For the Earth’s atmosphere, databases HITRAN [71–73] and GEISA [74, 75] provide comprehensive and validated datasets for approximately 50 key molecules. However, these databases are designed for studies at room temperature and below and do not contain the necessary data for adequately calculating radiative transport in hot bodies. The HITEMP database was constructed to extend HITRAN to higher temperatures. The latest, 2010, release [76] only contained data on 5 molecules (OH, NO, CO, H₂O and CO₂) and improved hot line lists are available for all these molecules; an update to HITEMP is currently in progress [77–79]. Other relevant databases include TheoReTS [80], which contains hot line lists for 8 polyatomic molecules, and Kurucz’s compilation of data, which is very complete for atomic sources but contains data only on about ten diatomic molecules [81] all of which are covered by ExoMol, usually to higher accuracy. The MoLLIST data base of Bernath and co-workers [82] contains empirically derived line lists for about 20 diatomic species. These line lists have recently been incorporated into the ExoMol database [83], see Table 2 below. Databases of hot molecular spectra for other specialised applications include those for combustion [6, 84] and studies of laser-induced plasmas [85]. ExoMol line lists together with data from other sources will serve as the main data source for exoplanet stands planned for the upcoming space missions Ariel and JWST.

The methodology [65, 86–91] and software [92, 93] developed and used by the ExoMol project has been extensively documented elsewhere. Here we will only

consider those aspects which impinge on the new ExoMol2020 release.

2. Database coverage

The ExoMol project aims at complete coverage of the spectroscopic properties of molecules which are deemed to be important in hot astrophysical environments. Coverage concerns (a) the molecular species considered, including isotopologues; (b) the spectroscopic and thermodynamic properties considered; (c) the frequency range considered and (d) the upper temperature range for which the data are reasonably complete. Both the required temperature and frequency range completeness are to some extent a judgement on what is required for astronomical and other studies.

Unlike ExoMol2016, the ExoMol2020 database essentially provides a complete dataset for modelling hot atmospheres. While new species are still being added, commonly at the request of users, the database now contains molecular data for 80 molecules and covers the key molecules thought to be important for exoplanetary studies. In many cases there is more than one line list for a given isotopologue. For this reason it is our practice to name each line list, including (unnamed) ones we have imported from other sources. In the case of multiple line lists the ExoMol website provides a recommended line list for the given species. This recommendation is for studies of hot atmospheres; other available line lists may be more suitable for room temperature studies.

For ease of understanding we have divided the description of line lists provided into three tables. Tables 1 and 2 summarise the molecules for which data are provided by ExoMol and give the characteristics of the line list in each case. Table 1 details line lists which have been formally published as part of the ExoMol project while Table 2 shows line lists imported from other sources which have been recast in the ExoMol format (see Wang *et al* [83] for example); these line

lists are fully integrated into the database. Table 3 lists the main extra line lists available through the ExoMol website; this table is not comprehensive but the other line lists available on the ExoMol website and not listed in Table 1 – 3 are largely of historic interest only.

Table 1: Datasets created by the ExoMol project and included in the ExoMol database.

Paper	Molecule	N_{iso}	T_{max}	N_{elec}	N_{lines}^a	DSName	Reference
I	BeH	1	2000	1	16 400	Yadin	Yadin et al. [94]
I	MgH	3	2000	1	10 354	Yadin	✓ Yadin et al. [94]
I	CaH	1	2000	1	15 278	Yadin	✓ Yadin et al. [94]
II	SiO	5	9000	1	254 675	EJBT	✓ Barton et al. [95]
III	HCN/HNC	1 ^a	4000	1	399 000 000	Harris	✓ Barber et al. [96]
IV	CH ₄	1	2000	1	34 153 806 005	YT34to10	✓ Yurchenko and Tennyson [97], Yurchenko et al. [98]
V	NaCl	2	3000	1	702 271	Barton	✓ Barton et al. [99]
V	KCl	4	3000	1	1 326 765	Barton	✓ Barton et al. [99]
VI	PN	2	5000	1	142 512	YYLT	✓ Yorke et al. [100]
VII	PH ₃	1	1500	1	16 803 703 395	SAITY	✓ Sousa-Silva et al. [101]
VIII	H ₂ CO	1	1500	1	10 000 000 000	AYTY	✓ Al-Refaie et al. [102]
IX	AlO	4	8000	3	4 945 580	ATP	✓ Patrascu et al. [103]
X	NaH	2	7000	2	79 898	Rivlin	✓ Rivlin et al. [104]
XI	HNO ₃	1	500	1	6 722 136 109	AlJS	✓ Pavlyuchko et al. [105]
XII	CS	8	3000	1	548 312	JnK	✓ Paulose et al. [106]
XIII	CaO	1	5000	5	21 279 299	VBATHY	✓ Yurchenko et al. [107]
XIV	SO ₂	1	2000	1	1 300 000 000	ExoAmes	✓ Underwood et al. [108]
XV	H ₂ O ₂	1	1250	1	20 000 000 000	APTY	✓ Al-Refaie et al. [109]
XVI	H ₂ S	1	2000	1	115 530 3730	AYT2	✓ Azzam et al. [110]
XVII	SO ₃	1	800	1	21 000 000 000	UYT2	✓ Underwood et al. [111]
XVIII	VO	1	2000	13	277 131 624	VOMYT	✓ McKemmish et al. [112]
XIX	H ₂ ^{17,18} O	2	3000	1	519 461 789	HotWat78	✓ Polyansky et al. [113]
XX	H ₃ ⁺	1 ^b	3000	1	11 500 000 000	MiZATEP	✓ Mizus et al. [114]
XXI	NO	6	5000	2	2 281 042	NOName	✓ Wong et al. [115]
XXII	SiH ₄	1	1200	1	62 690 449 078	OY2T	✓ Owens et al. [116]
XXIII	PO	1	5000	1	2 096 289	POPS	✓ Prajapat et al. [117]
XXIII	PS	1	5000	3	30 394 544	POPS	✓ Prajapat et al. [117]
XXIV	SiH	4	5000	3	1 724 841	SIGHTLY	✓ Yurchenko et al. [118]
XXV	SiS	12	5000	1	91 715	UCTY	✓ Upadhyay et al. [119]
XXVI	NS	6	5000	1	3 479 067	SNaSH	✓ Yurchenko et al. [120]
XXVI	HS	6	5000	1	219 463	SNaSH	Yurchenko et al. [120]
XXVII	C ₂ H ₄	1	700	1	60 000 000 000	MaYTY	✓ Mant et al. [121]
XXVIII	AlH	3	5000	3	40 000	AlHambra	✓ Yurchenko et al. [122]
XXIX	CH ₃ Cl	2	1200	1	166 279 593 333	OYT	✓ Owens et al. [123]
XXX	H ₂ ¹⁶ O	1 ^c	5000	1	5 745 071 340	POKAZATEL	✓ Polyansky et al. [61]
XXXI	C ₂	3	5000	8	6 080 920	8states	✓ Yurchenko et al. [124]
XXXII	MgO	5	5000	5	72 833 173	LiTY	✓ Li et al. [125]
XXXIII	TiO	5	5000	13	59 000 000	Toto	✓ McKemmish et al. [126]
XXXIV	PH	1	4000	2	65 055	LaTY	✓ Langleben et al. [127]
XXXV	NH ₃	1 ^d	1500	1	16 900 000	CoYuTe	✓ Coles et al. [128]
XXXVI	SH	2	3000	2	572 145	GYT	✓ Gorman et al. [129]
XXXVII	HCCH	1	2000	1	4 347 381 911	aCeTY	✓ Chubb et al. [130]
XXXVIII	SiO ₂	1	3000	1	32 951 275 437	OYT3	✓ Owens et al. [131]
XXXIX	CO ₂	1	3000	1	7 996 570 390	UCL-4000	✓ Yurchenko et al. [132]
XL	H ₃ O ⁺	1	1500	1	2 089 331 073	eXeL	✓ Yurchenko et al. [133]

Paper Number in series published in Mon. Not. R. Astron. Soc.

N_{iso} Number of isotopologues considered;

T_{max} Maximum temperature for which the line list is complete;

N_{elec} Number of electronic states considered;

N_{lines} Number of lines: value is for the main isotope.

✓ indicates line list recommended for studies of hot atmospheres.

^a The Larner line list for H¹³CN/HN¹³C due to Harris et al. [134] is recommended.

^b The ST line list for H₂D⁺ due to Sochi and Tennyson [135] is recommended.

^c The VTT line list for HDO due to Voronin et al. [136] is recommended.

^d There is a room temperature ¹⁵NH₃ line list due to Yurchenko [137].

Table 2: Datasets not created as part of the ExoMol project but included in the ExoMol database.

Molecule	N_{iso}	T_{max}	N_{elec}	N_{lines}	DSName	Reference	Methodology
H ₂	1	10000	1	4712	RACPPK	✓ Roueff et al. [138]	Ab initio
CH	1	5000	4	52201	MoLLIST	✓ Masseron et al. [139]	Empirical
NH	1	5000	1	12150	MoLLIST	✓ Brooke et al. [140, 141], Fernando et al. [142]	Empirical
OH	1	5000	2	54276	MoLLIST	✓ Brooke et al. [143]	Empirical
AlCl	2	5000	1	20245	MoLLIST	✓ Yousefi and Bernath [144]	Empirical
AlF	1	5000	1	40490	MoLLIST	✓ Yousefi and Bernath [144]	Empirical
OH ⁺	1	5000	2	12044	MoLLIST	✓ Hodges and Bernath [145], Hodges et al. [146]	Empirical
CaF	1	5000	6	14817	MoLLIST	✓ Hou and Bernath [147]	Empirical
MgF	1	5000	3	8136	MoLLIST	✓ Hou and Bernath [148]	Empirical
KF	1	5000	2	10572	MoLLIST	✓ Frohman et al. [149]	Empirical
NaF	1	5000	1	7884	MoLLIST	✓ Frohman et al. [149]	Empirical
LiCl	1	5000	4	26260	MoLLIST	✓ Bittner and Bernath [150]	Empirical
LiF	1	5000	2	10621	MoLLIST	✓ Bittner and Bernath [150]	Empirical
MgH	2	5000	1	14179	MoLLIST	✓ GharibNezhad et al. [151]	Empirical
TiH	1	5000	3	181080	MoLLIST	✓ Burrows et al. [152]	Empirical
CrH	1	5000	2	13824	MoLLIST	✓ Chowdhury et al. [153]	Empirical
FeH	1	5000	2	93040	MoLLIST	✓ Wende et al. [154]	Empirical
HF	2	5000	1	7956	Coxon-Hajig	✓ Coxon and Hajigeorgiou [155]	Empirical
HCl	4	5000	1	2588	HITRAN	✓ Li et al. [156]	Empirical
CP	1	5000	2	28752	MoLLIST	✓ Ram et al. [157]	Empirical
CN	1	5000	3	195120	MoLLIST	✓ Brooke et al. [158]	Empirical
C ₂	1	5000	2	47 570	MoLLIST	✓ Brooke et al. [159]	Empirical
CaH	2	5000	1	6000	MoLLIST	✓ Li et al. [160], Shayesteh et al. [161]	Empirical
N ₂	1	10000	4 ^a	7 182 000	WCCRMT	✓ Western et al. [162]	Empirical
SiO	1	5000	3	6 67 814	Kurucz-SiO	Kurucz [81]	Empirical
ScH	1	5000	6	1 152 827	LYT	✓ Lodi et al. [163]	Ab initio
LiH	1	12000	1	18 982	CLT	✓ Coppola et al. [164]	Ab initio
LiH ⁺	1	12000	1	332	CLT	✓ Coppola et al. [164]	Ab initio
CO	9	9000	1	752 976	Li2015	✓ Li et al. [165]	Empirical
HeH ⁺	4	9000	1	1430	ADSJAAM	✓ Amaral et al. [166]	Ab initio
HD ⁺	1	9000	1	10285	ADSJAAM	✓ Amaral et al. [166]	Ab initio
HD	1	9000	1	5939	ADSJAAM	✓ Amaral et al. [166]	Ab initio
CH ₃ F	1	300	1	139 188 215	OYKYT	✓ Owens et al. [167]	Ab initio
AsH ₃	1	300	1	3 600 000	CYT18	✓ Coles et al. [168]	Ab initio
P ₂ H ₂ ^b	2	300	1	10 667 208 951	OY-Trans	✓ Owens and Yurchenko [169]	Ab initio
P ₂ H ₂ ^b	2	300	1	11 020 092 365	OY-Cis	✓ Owens and Yurchenko [169]	Ab initio
PF ₃	1	300	1	68 000 000 000	MCYTY	✓ Mant et al. [170]	Ab initio
CH ₃	1	1500	1	2 058 655 166	AYYJ	✓ Adam et al. [171]	Ab initio
BeH	3	5000	2	592308	Darby-Lewis	✓ Darby-Lewis et al. [172]	ExoMol
CO ₂	13	4000	1	298 323 789	Ames-2016	Huang et al. [173, 174]	ExoMol-like
SiH ₂	1	2000	1	254 061 207	CATS	✓ Clark et al. [175]	ExoMol
YO	1	5000	6	3520133	SSYT	✓ Yurchenko et al. [176]	Ab initio

N_{iso} Number of isotopologues considered;

T_{max} Maximum temperature for which the line list is complete;

N_{elec} Number of electronic states considered;

N_{lines} Number of lines: value is for the main isotope.

✓ indicates line list recommended for studies of hot atmospheres.

^a The WCCRMT line list considers triplet states only.

^b There are separate line lists for cis and trans P₂H₂.

Table 3: Supplementary datasets available from the website.

Molecule	N_{iso}	T_{max}	N_{elec}	N_{lines}	DSName	Reference	Methodology
H_3^+	2 ^a	4000	1	3 070 571	NMT	Neale et al. [177]	ExoMol
H_2O	2 ^b	3000	1	505 806 202	BT2	Barber et al. [178]	ExoMol
NH_3	2 ^c	1500	1	1 138 323 351	BYTe	Yurchenko et al. [179]	ExoMol
HeH^+	4	10000	1	1 431	Engel	Engel et al. [180]	Ab initio
HD^+	1	12000	1	10 119	CLT	Coppola et al. [164]	Ab initio
CO_2	13	300	1	161944	Zak	Zak et al. [181, 182, 183]	ExoMol
CO_2	1	4000	1	628 324 454	CSSD-4000	Tashkun and Perevalov [184]	Empirical
H_2O	1	300	1		WAT_UV296 [185]	ExoMol	

N_{iso} Number of isotopologues considered;

T_{max} Maximum temperature for which the line list is complete;

N_{elec} Number of electronic states considered;

N_{lines} Number of lines; value is for the main isotope.

3. Individual line lists

An overview of the line lists in the ExoMol database is given in Fig. 1.

One general issue is that Medvedev and co-workers [165, 186] identified a numerical problem with the intensities of high overtone transitions computed with the standard compilation of the diatomic vibration-rotation program LEVEL [187]. Our diatomic line lists computed with LEVEL or DUO [188] have been adjusted to remove transitions affected by this issue. Medvedev *et al.* [189] recently identified similar issues with triatomic systems but tests suggest that in practice our triatomic line lists seem largely unaffected by the problem.

Below we consider some of the line lists presented in the ExoMol database and listed in Tables 1 and 2. We restrict our discussion to issues not covered in the ExoMol2016 release or the original publications. We start by considering the 42 molecules for which line lists have been created as part of the ExoMol series as listed in Table 1.

AlH	AICl	AlO	BeH	C ₂	CaO	CaF	CaH	CH	CN	CO	CP
CrH	CS	FeH	FeO	H ₂	H ₃ ⁺	H ₃ O ⁺	HCl	HF	LiCl	LiF	LiH
LiH ⁺	KCl	KF	MgF	MgH	MgO	N ₂	NaCl	NaF	NaH	NaF	NiH
NH	NS	NO	OH	O ₂	PH	PN	PO	PS	ScH	SiC	SiH
SiS	SH	SiO	SiS	SO	TiO	TiH	TiF	VO	VN	YO	ZnS
C ₃	CaOH	CO ₂	CH ₂	H ₃ ⁺	H ₂ D ⁺	HCN	HNC	H ₂ O	HDO	H ₂ S	KOH
NaOH	N ₂ O	OCS	PO ₂	SiH ₂	SiO ₂	SO ₂	AsH ₃	CH ₃	C ₂ H ₂	HOOH	H ₂ CO
NH ₃	SiH ₃	PH ₃	P ₂ H ₂	PF ₃	SO ₃	CH ₄	CH ₃ D	CH ₃ Cl	CH ₃ F	SiH ₄	HNO ₃
C ₂ H ₄	C ₂ H ₆	C ₃ H ₈									

Figure 1: Molecular line lists: red indicates line lists in progress, blue corresponds to the line lists suggested for molecules specific for hot rocky exoplanets and green indicates line lists which contain data applicable for high resolution.

3.1. Diatomics

3.1.1. AlH, paper XVIII

There is a new AlHambra line list for AlH [122]. A MARVEL (measured active rotational-vibrational energy levels [190]) project was performed as part of this project meaning that many transitions are predicted with experimental accuracy. The AlHambra line list has been updated to give the uncertainty in the energy in the states file which allows users to determine the uncertainty in a given transition wavenumber.

3.1.2. AlO, paper IX

No change. The APTY line list was recently used to make a detection of AlO in an exoplanet atmosphere, the ultra-hot Jupiter exoplanet WASP-33 b [191] and hot Jupiter exoplanet WASP-43 b [192].

3.1.3. BeH, paper I

The BeH “Yadin” line list was one of the first constructed by ExoMol [94]; however, the line list only included transitions within the X ² Σ^+ ground electronic

state. Recently Darby-Lewis et al. [172] constructed line lists for BeH, BeD and BeT which consider both the ground and first excited ($A\ ^2\Pi$) states. At the same time Darby-Lewis *et al.* performed a MARVEL analysis and used their empirical levels in the fit. The Darby-Lewis line list is therefore more accurate and more extensive than the Yadin one which it replaces in the ExoMol database and is therefore recommended. Yadin is still available on the website though renamed to Yadin-BeH in order to avoid conflict with the “Yadin” line lists for MgH and CaH, which remain to be recommended for the IR region.

3.1.4. C_2 , paper XXXI

There are two new line lists for C_2 in the ExoMol2020 release: ExoMol 8state [124] and the empirical MoLLIST [159]. The MoLLIST line list only considers the much-observed Swan band while 8state covers the 8 band systems which interconnect the 8 lowest electronic states of C_2 . As 8state uses empirical MARVEL energy levels [193] where available it should be as accurate as MoLLIST for the Swan band. Use of 8state is therefore recommended. Since MoLLIST line lists are recommended for other molecules and in order to avoid conflicts with the ‘recommended’ flag for 8states, the C_2 MoLLIST line list is now referenced to as MoLLIST-C2 on ExoMol.

An update of the C_2 MARVEL data has just been completed [194]; this has been used to improve the 8state energies and hence transition wavenumbers. This latest version includes uncertainties in the states file.

3.1.5. CaH , Paper I

The “Yadin” CaH line list only considers transitions within the $X\ ^2\Sigma^+$ ground electronic state [94]. MoLLIST provides a rovibronic line list for the $X\ ^2\Sigma^+ - X\ ^2\Sigma^+$, $A\ ^2\Pi - X\ ^2\Sigma^+$, $B\ ^2\Sigma^+ - X\ ^2\Sigma^+$ and $E\ ^2\Pi - X\ ^2\Sigma^+$ systems due to Li et al. [160], Shayesteh et al. [161], Ram et al. [195]. Both line lists are included in the

ExoMol database; work is in progress on creating a single unified line list which will extend the range of rovibronic transitions considered. In the meantime both line lists are recommended.

3.1.6. CaO, Paper XIII

No change.

3.1.7. CS, Paper XII

No change. We note that a line list for CS has also recently been supplied by Hou and Wei [196] and Xing et al. [197] have extended consideration to rovibronic transitions for the lowest, A $^1\Pi$ – X $^1\Sigma^+$, allowed electronic band.

3.1.8. KCl, Paper V

No change

3.1.9. MgH, Paper I

The ExoMol “Yadin” line list for MgH only considers transitions within the X $^2\Sigma^+$ ground electronic state [94]. MoLLIST provides a rovibronic line list containing A $^2\Pi$ – X $^2\Sigma^+$ and B' $^2\Sigma^+$ – X $^2\Sigma^+$ transitions due to GharibNezhad et al. [151]. Both line lists are included in the ExoMol database; work is in progress on creating a single unified line list which will extend the range of rovibronic transitions considered. In the meantime both line lists are recommended.

3.1.10. MgO, Paper XXXII

There is a new LiTY line list for MgO [125].

3.1.11. NaCl, Paper V

No change.

3.1.12. NaH, Paper X

No change.

3.1.13. NO, XXI

There is a new NOName line list for NO [115]. This line list was constructed using a combination of standard ExoMol and empirical methodologies, and also included a MARVEL project. Its transition frequencies should therefore be close to experimental accuracy. NOName has largely been adopted for the new release of HITEMP [77].

NOName only covers transitions between levels which lie within the X $^2\Pi$ ground electronic state of NO. A new line list which includes ultraviolet rovibronic transitions of NO is currently being constructed.

3.1.14. PH, Paper XXXIV

There is a new LaTY line list for PH [127].

3.1.15. PN, Paper VI

No change. An extended line list covering visible and ultra violet (UV), accompanied by a MARVEL project is currently in progress.

3.1.16. PO, Paper XXIII

There is a new POPS line list for PO [117].

3.1.17. PS, Paper XXIII

There is a new POPS line list for PS [117].

3.1.18. SH, Paper XXXVI

There are two new ExoMol line lists for SH: SNaSH [120] which only covers transitions within the X $^2\Pi$ ground electronic state of SH and the newer GYT line list of Gorman et al. [129]. GYT considers both transitions within X $^2\Pi$ and vibronic transitions in the A $^2\Sigma^+ - X ^2\Pi$ band system. As GYT also improves the accuracy of the X state transitions, it is recommended for all applications.

The SNaSH SH line list is also renamed to SNaSH-SH in order to avoid a conflict and retained only for completeness.

3.1.19. SiO, Paper II

The ExoMol SiO line lists only consider transitions within the X ${}^1\Sigma^+$ ground electronic state [95]; these line lists have been widely used including a recent determination of isotopologue ratios in Arcturus [198]. For this release of ExoMol, an empirical and less accurate SiO line list by Kurucz [81] covering the X–X, A–X and E–X electronic bands was added. Given the importance of SiO for lava planets [65], construction of a comprehensive (accurate and complete) rovibronic line list for SiO covering both IR and ultraviolet would be useful.

3.1.20. SiH, Paper XXIV

There is a new SiGHTLY line list for SiH [118].

3.1.21. SiS, Paper XXV

There is a new UCTY line list for SiS [119].

3.1.22. SN, Paper XXVI

There is a new SNaSH line list for SN [120].

3.1.23. TiO, Paper XXXIII

There is a new Toto line list for TiO [126]. The line list for the main isotopologue (${}^{48}\text{Ti}{}^{16}\text{O}$) used empirical energies from an associated MARVEL study [199] while the energy levels of other isotopologues were improved using the procedure of Polyansky et al. [113]. The Toto line list has been updated to give MARVEL uncertainties, where available, in the States file. For other states the uncertainties were estimated as follows: 1 cm^{-1} for all levels in vibronic states that contain levels determined by MARVEL, 10 cm^{-1} for levels in those electronic states who

excitation energy (T_e/T_0) is known from experiment and 100 cm^{-1} for all other levels.

TiO is a particularly important species in the atmosphere of cool stars and has been detected in exoplanets [200, 201]. The Toto line list is significantly better at reproducing stellar spectra than the line lists due to Plez [202] and Schwenke [203] which it supersedes. Furthermore a recent study of Ti isotope abundances in two M-dwarf stars demonstrated the accuracy of the line lists for several isotopologues of TiO [42]. However, TiO remains a challenging system to treat using *ab initio* electronic structure methods (see McKemmish et al. [204]) and further work is required before there is a definitive line list for TiO.

3.1.24. VO, Paper XVIII

There is a new VOMYT line list for VO [112] which has been used to tentatively identify VO in the atmosphere of exoplanets [205, 206]. We note that VO spectra display particularly large splittings due to hyperfine effects as well as transitions which are only allowed due to these effects [207]. VOMYT does not include hyperfine effects in its spectroscopic model meaning that the line list unsuitable for high resolution studies. We plan to address this problem by both performing a hyperfine resolved MARVEL study and extending DUO to allow for hyperfine effects.

3.2. Triatomics

3.2.1. CO₂, Paper XXXIX

Line lists for hot CO₂ have been constructed by NASA Ames [173, 173] using methodologies similar to those employed by ExoMol. The CDSD-4000 hot line list due to Tashkun and Perevalov [184] and the room temperature line lists due to Zak et al. [181, 182, 183] are also available on the ExoMol website. The CDSD-4000 line list is based on the use of an effective Hamiltonian which tends to miss contributions from unobserved hot bands.

A new ExoMol line list UCL-4000 for CO₂ has been produced [132] using an accurate *ab initio* dipole moment surface (DMS) by Polyansky et al. [208] and empirical potential energy surface (PES) Ames-2016 by Huang et al. [209]; where possible computed energies have been replaced by empirical one derived from HITRAN. The UCL-4000 line list covers the wavenumber range 0–20000 cm⁻¹ and should be applicable for temperatures up to 4000 K [132]. It is recommended for use in high temperature applications.

3.2.2. HCN/HNC, Paper III

The combined HCN/HNC ExoMol line list of Barber *et al* [96] made extensive use of empirical energy levels due to Mellau [210, 211]. This has enabled the line list to be successfully used in high-resolution Doppler spectroscopy studies of exoplanets [212–214].

3.2.3. H₂O, Papers XIX and XXX

Water is one of the most widely studied molecules and its spectrum has been detected in a variety of exoplanets, in many cases as the only clearly identifiable molecule [206, 215]. Good water line lists have been available for some time, in particular the Ames line list of Partridge and Schwenke [216] and the BT2 line list due to Barber et al. [178]. While BT2 was significantly more complete than Ames, the Ames line list was more accurate, particularly at infrared wavelengths. A new line list, known as POKAZATEL [61] has been generated by the ExoMol project. POKAZATEL is complete in the sense that it contains transitions between all bound rotation-vibration states in the molecule and thus is reliable for temperatures above 3000 K where the earlier line lists are not valid. POKAZATEL is also intrinsically more accurate than the Ames line list and, indeed, as many of its energy levels have been replaced by empirical energy levels generated using MARVEL [217], key transition frequencies are actually reproduced to

experimental accuracy. POKAZATEL should therefore be used in preference to the earlier line lists.

It has become apparent that the rotation-vibration spectrum of water is responsible for weak but observable near-UV absorption in the Earth's atmosphere [218]. The POKAZATEL line list, which used the *ab initio* LTP2001S DMS of Lodi et al. [219], appears to underestimate the strength of water absorption in the blue and near-UV. As a response to this Conway and co-workers have recently developed a more accurate global dipole moment surface [220]. Through a significant number of comparisons against high quality experimental and theoretical sources of spectroscopic data [221, 222], their results suggest there may be advantages of using their line lists in particular regions, primarily at short wavelengths. For this reason, a new room temperature H_2^{16}O line list called WAT_UV296 [185] has been computed. This line list is available on the website and should be used for room temperature studies of spectra with $\lambda < 0.5 \mu\text{m}$. A high temperature companion to this line list will form the basis of the updated HITEMP database [79].

Polyansky et al. [113] provided the HotWat78 line lists for H_2^{17}O and H_2^{18}O . The energy levels of this line list were improved using MARVEL energies [223, 224]. Furthermore, Polyansky et al. [113] developed a method which gives excellent isotopologue energy levels for states only observed for H_2^{16}O . Finally the VTT line list due to Voronin et al. [136] remains the recommended one for HDO; a new, improved HDO line list is currently under construction [225].

A new room temperature line list CKYKKY for H_2O containing electric quadrupole moments was computed using an *ab initio* quadrupole moment surface [226] and accurate empirical PES by Mizus et al. [227] with the variational program TROVE [228]. The energies were replaced by the H_2O IUPAC MARVEL values [217] or HITRAN values [229], if available.

3.2.4. H_2S , Paper XVI

There is a new ExoMol AYT2 line list for hydrogen sulphide due to Azzam et al. [110]. Since completion of this line list a MARVEL project has been performed for H_2S [230]; the AYT2 line list is being updated to use these improved energy levels.

3.2.5. H_3^+

The new ExoMol MiZATeP line list of Mizus et al. [114] replaces that of Neale et al. [177] (NMT). The new line list uses empirical energy levels from the MARVEL study of Furtenbacher et al. [231]. Astronomy makes wide use of H_3^+ line lists and is reliant on *ab initio* line intensities since no absolute line intensities have been measured in the laboratory, see Petrignani et al. [232]. This astronomical work on H_3^+ has recently been reviewed by Miller et al. [233]. The ST H_2D^+ line list due to Sochi and Tennyson [135] plus the newly generated D_3^+ and D_2H^+ line lists will be MARVELised and released in the near future.

3.2.6. SO_2 , Paper XVII

The ExoAmes line list for SO_2 [193] is unchanged since the ExoMol2016 release. However, a MARVEL set of energy levels for SO_2 are now available [234] and will be used to update ExoAmes in the near future.

3.3. Tetratomics

3.3.1. $HCCH$, Paper XXXVII

The new aCeTY line list for acetylene due to Chubb et al. [130] has recently been released. This line list already incorporated the MARVEL energies of Chubb et al. [235] and gives uncertainties in the energy levels as part of the States file.

3.3.2. H_2CO , Paper VIII

The AYTY formaldehyde line list [102] is in the process of being updated with empirical energy levels produced by a recent MARVEL study [236]; this will make the line list suitable for high resolution studies.

3.3.3. H_2O_2 , Paper XV

There is a new ExoMol APTY line list for hydrogen peroxide [237]. We note because of difficulties with assigning spectra, even at room temperature HITRAN is very incomplete for H_2O_2 . APTY should give complete coverage at infrared wavelengths. Recently APTY was used to suggest the importance of H_2O_2 as a greenhouse gas on oxidised Early Mars [238].

3.3.4. H_3O^+ , Paper XL

A line list for the hydronium ion, H_3O^+ , has recently been constructed [133] in response to a laboratory study by Bourgalais et al. [239] which suggested both that hydronium is likely to be a dominant molecular ion in gaseous exoplanets and that it should be detectable by upcoming space missions.

3.3.5. NH_3 , Paper XXXV

A new ExoMol line list for ammonia called CoYuTe [128] has recently been completed. CoYuTe, which also uses MARVEL energy levels [188], is both more accurate and more extensive than the BYTe line list [179] it replaces. In particular, CoYuTe was found to provide a good model of the Jovian optical absorption spectrum due to ammonia [34] although with a shift in positions of the main bands. More laboratory work or analysis of existing laboratory work on the visible spectrum of ammonia will be required to resolve this issue. An illustration of the ammonia absorption cross sections computed using CoYuTe is given in Fig. 2.




Figure 2: Absorption cross sections generated using the ExoMol line list CoYuTe for spectrum of NH_3 at $T = 300 \text{ K}$ and 2000 K , with the Doppler profile on a wavenumber grid of 1 cm^{-1} . The cross sections were computed using the ExoCross program, which is developed to work with two file line list structure adopted by ExoMol.

3.3.6. PH_3 , Paper VII

No change. Sousa-Silva et al. [240] use the SALTY PH_3 line list to assess the detectability of a possible PH_3 biosignature. A MARVEL project on the PH_3 experimental data is under way, which will improve the quality of the line positions in SALTY.

3.3.7. SO_3 , Paper XVII

There is a new UYT2 line list for SO_3 [111].

3.4. Pentatomics

3.4.1. CH_3Cl , Paper XXIX

There are new OYT line lists for methyl chloride [123] covering both major isotopes of chlorine.

3.4.2. CH₄, Paper IV

The ExoMol YT10to10 line list [97] has been extended to higher temperatures [98] to give the YT34to10 line list. However, at present the TheoReTS line list of Rey et al. [241] is the most accurate methane line list available and is recommended for detailed studies. This line list forms the basis of the very recent update to HITEMP [78].

The importance of methane and the huge number of lines required to accurately represent the opacity of hot methane [31] has led to this system being the primary focus of studies aimed at compacting these lines into a more manageable form [98, 242, 243]. This issue is discussed in section 5.3.2.

3.4.3. HNO₃, Paper XI

No change. We note that, like H₂O₂, because of difficulties with assigning spectra even at room temperature, HITRAN is very incomplete for nitric acid. The AlJS line list gives complete coverage at infrared wavelengths.

3.4.4. SiH₄, Paper XXII

There is a new OY2T line list for silane [116].

3.5. Larger molecules

3.5.1. C₂H₄, Paper XXVII

There is a new MaYTY line list for ethylene [121].

3.6. MoLLIST species

A number of diatomic species have line lists provided by the MoLLIST website [82]. These have been incorporated into the ExoMol database, which now includes empirically-derived line lists for the following species [83]: CH [139]; NH [140–142]; OH [143]; AlCl [144]; AlF [144]; OH⁺ [145, 146]; CaF [147]; MgF [148]; KF and NaF [149]; LiCl and LiF [150]; MgH [151]; TiH [152]; CrH [153]; FeH [154]; CP [157]; CN [158]; and CaH [160, 161]. See Table 2 for further details.

3.7. HITRAN

In general the line lists contained in the HITRAN database [229] do not provide the temperature coverage needed for astrophysical applications. However, for a few diatomic species very extensive line lists have been constructed. The line lists presented by Li et al. [156] for HCl and Li et al. [165] for CO are both valid over an extended temperature range and have been included in the ExoMol database.

The HF line list from Li et al. [244] and Coxon and Hajigeorgiou [155] is also adopted directly from HITRAN.

3.8. N₂

The empirical WCCRMT nitrogen molecule line list of Western et al. [162] is somewhat unusual in that it only considers transitions between excited, triplet states of the molecule. However, N₂ is an important molecule with very limited spectral signature at long wavelengths and WCCRMT may prove useful in hot or irradiated, usually non-LTE environments. For the partition function of N₂ the ground electronic state data from TIPS [245] were used.

3.9. HD, HD⁺, HeH⁺

One and two electron diatomics are important primordial species and may well play a role elsewhere. Recently Amaral et al. [166] created new line lists using high level *ab initio* procedures. These line lists have been included in the database with the name ADSJAAM. In particular the ADSJAAM line lists for HeH⁺ and HD supersedes those due to Engel et al. [180] and Coppola et al. [164], respectively.

3.10. H₂

A new external line list RACPPK by Roueff et al. [138] for the ground electronic state of H₂ was converted to the ExoMol format. The line list contains

a combination of electric quadrupole and magnetic dipole transitions computed from first principles.

3.11. Other *ab initio* line lists

A number of other *ab initio* line lists are also available. In general *ab initio* line lists are only accurate for systems with very few electrons. The CLT line lists for LiH and LiH⁺ due to Coppola et al. [164] fall into the few-electron category and should be reliable. The other *ab initio* line lists provided, namely those for ScH [163], CH₃F [167], AsH₃ [168], P₂H₂ [169], PF₃ [170], CH₃ [171] and YO [176] must be regarded as intrinsically less accurate than other line lists provided by the database. In the case of YO, work is in progress aimed at producing an empirical line list based on available experimental data. P₂H₂ occurs as two distinct isomers, the cis and trans forms. Separate line lists are provided for cis-P₂H₂ and trans-P₂H₂.

3.12. SiH₂

A new line list for SiH₂ [175] has been recently constructed in its ground electronic state using the standard ExoMol technology [93]. The set of experimental data or their quality were, however, very limited which may affect the quality of the hot bands, which basically has to rely on the quality of the *ab initio* PES.

3.13. NiH

Empirical line lists for three isotopologues of NiH (⁵⁸NiH, ⁶⁰NiH, ⁶²NiH) known as HRV. These line lists were constructed by Vallon et al. [246] and Harker et al. [247]. However, at present these line lists are not available in ExoMol format or accessible through the API.

3.14. Partition functions

The temperature dependent partition functions for most of the ExoMol line lists are computed using the corresponding energy levels as collected in the States files as

$$Q(T) = \sum_i g_{\text{ns}}^{(i)} J_i(J_i + 1) \exp\left(-\frac{c_2 \tilde{E}_i}{T}\right), \quad (1)$$

where $g_{\text{ns}}^{(i)}$ is the state dependent nuclear statistical weight, J_i is the total angular momentum of the state i , c_2 is the second radiation constant (K cm) and \tilde{E}_i is the corresponding energy term value (cm^{-1}). ExoMol uses the HITRAN convention [248], where the entire, integer $g_i = g_{\text{ns}}^{(i)} J_i(J_i + 1)$ factors are explicitly included.

Some of the line lists in the ExoMol database do not provide sufficiently complete sets states for a proper evaluation of the molecular partition functions. For example, most of the MoLLIST line lists are constructed from or based on the measured data only and therefore can be severely incomplete. In these cases the partition functions are either taken from external sources, such as the TIPS database [245], generated using the empirical expansions, such as by Irwin [249], Sauval and Tatum [250], Irwin [251], Barklem and Collet [252] or extrapolated using simplified models [253]. The external partition functions often include the $g_{\text{ns}}^{(i)}$ factors in the astrophysical convention (see, e.g. Pavlenko et al. [198]) and therefore have to be transformed to the HITRAN convention.

We strongly recommend that users use these partition functions rather than attempting to compute their own using the ExoMol States files. These States files are not constructed with a view to delivering reliable partition functions and in a number of cases use of them has been found to lead to problems.

3.15. VUV sections

A section is provided based on measured ultraviolet cross sections for key species. For many molecules their ultraviolet spectrum is a mixture of line and

quasi-continuum absorption which cannot be represented as a line list. The new VUV section will provide temperature-dependent absorption cross sections. Recent results suggest the VUV absorption by H₂O enhances the production of OH, which plays an important molecule in atmospheric chemistry of exoplanetary atmospheres [254]. Currently, the VUV cross sections are provided for H₂O, H₂, CO₂, SO₂, NH₃, H₂CO and C₂H₄ measured by Fateev's lab at the Danish Technical University (DTU) [255] and for CO₂ by Venot et al. [256]. The temperature and wavelength coverage is illustrated in Table 4.

The cross sections are given in the common two-column ASCII format separated by spaces, where the first column contains the wavelength in nm and the second column contains the absorption cross sections in cm²/molecule using Fortran format: (F10.3,1x,E13.6). The VUV cross section file names have the following structure

'<ISOTOPOLYGE>_<DATASET>_<RANGE>_T<TEMP>K_P<PRESSURE>bar_<STEP>.nm', where ISOTOPOLYGE is the isotopologue name, DATASET is the name of the line list, RANGE is the wavelength range in nm, TEMP is the temperature in K, PRESSURE is the pressure in bar, STEP is the wavelength step in nm. For example, the States file of the VUV line list for CO₂ the filename:

12C-1202__Venot-2018__116.90-230.00__T0800K__P0bar__0.03.nm.

Table 4: VUV absorption cross sections on ExoMol, grid spacing 0.01–0.015 nm, presented in natural abundance.

Molecule	Temperature (K)	Range (nm)
DTU data		
H ₂ O	423	110–230
H ₂ O	573	110–230
H ₂ O	1630	182–237
H ₂ O	1773	182–237
CO ₂	1160	109–324
SO ₂	423	110–230
NH ₃	289	113–201
NH ₃	296	113–201
H ₂ CO	303	110–230
H ₂ CO	353	110–230
H ₂ CO	423	110–230
H ₂ CO	573	110–230
C ₂ H ₄	562	113–201
Venot et al. [256]		
CO ₂	170–800 K	115–220

4. Data Provided

Table 5 provides a summary of different types of data provided. The website provides two routes to accessing these data. Users can search by molecule which will show all the types of data available for each isotopologues. Alternatively it is possible to search by data type in which case a list of molecules for which data of the specified type is available. The following section lists the data types and their file name extensions for the various data and metadata provided by ExoMol. A

more technical specification of the data structures and how to access data using the application program interface (API) is provided in Section 5.

Table 5: Summary of data provided by the ExoMol Database

Data type
Line lists
Absorption cross sections
VUV absorption cross sections
Pressure broadening coefficients
Temperature dependent super-lines (histograms)
Partition functions
Cooling functions
Specific heat - heat capacity
Examples of ExoCross input files
Temperature and pressure dependent opacities
Spectroscopic Models

4.1. Data Structure

Table 6: Specification of the ExoMol file types. (Contents in brackets are optional.)

File extension	N_{files}	File DSname	Contents
.all	1	Master	Single file defining contents of the ExoMol database..
.def	N_{tot}	Definition	Defines contents of other files for each isotopologue.
.states	N_{tot}	States	Energy levels, quantum numbers, lifetimes, (Landé g -factors, Uncertainties).
.trans	^a	Transitions	Einstein A coefficients, (wavenumber).
.broad	N_{mol}	Broadening	Parameters for pressure-dependent line profiles.
.cross	^b	Cross sections	Temperature or temperature and pressure-dependent cross sections.
.kcoef	^c	k -coefficients	Temperature and pressure-dependent k -coefficients.
.pf	N_{tot}	Partition function	Temperature-dependent partition function, (cooling function).
.dipoles	N_{tot}	Dipoles	Transition dipoles including phases.
.super	^d	Super-lines	Temperature dependent super-lines (histograms) on a wavenumber grid.
.nm	^e	VUV cross sections	Temperature and pressure dependent VUV cross-sections (wavelength, nm).
.fits, .h5, .kta	^f	Opacities	Temperature and pressure dependent opacities for radiative-transfer applications.
.overview	N_{mol}	Overview	Overview of datasets available.
.readme	N_{iso}	Readme	Specifies data formats.
.model	N_{iso}	Model	Model specification.

N_{files} total number of possible files;

N_{mol} Number of molecules in the database;

N_{tot} is the sum of N_{iso} for the N_{mol} molecules in the database;

N_{iso} Number of isotopologues considered for the given molecule.

^a There are N_{tot} sets of .trans files but for molecules with large numbers of transitions the .trans files are subdivided into wavenumber regions.

^b There are N_{cross} sets of .cross files for isotopologue.

^c There are N_{kcoef} sets of .kcoef files for each isotopologue.

^d There are N_T sets of T -dependent super-lines. ^e There are N_{VUV} sets of VUV cross sections. ^f Set of opacity files in the format native to specific radiative-transfer programs.

The general ExoMol approach is molecule-by-molecule: a comprehensive line list is created for a particular molecule and made available in the database. The line list for each isotopologue is stored as a separate data structure which can be accessed directly or via the application program interface (API) described in the following sections. Table 6 specifies the file types that can be available for each isotopologue. The .states and .trans files are the heart of the ExoMol data structure [257] and define what has become known as the “ExoMol format”. These files are available for all isotopologues; other files may not be. ExoMol format files can be used and provided by the effective Hamiltonian code PGOPHER [258].

A manual providing technical specifications of the database is included as

supplementary data to this article and is included on the website where it will be updated as the data base evolves. Only the relatively few changes that have been implemented since the 2016 release are discussed below.

4.2. Super-lines

Super-lines [80, 98] represent a novel, compact way of storing the opacity data. Super-lines are constructed as temperature-dependent intensity histograms by summing all absorption coefficients within a wavenumber bin centred around a grid point $\tilde{\nu}_k$. For each $\tilde{\nu}_k$ the total absorption intensity $I_k(T)$ is computed as a sum of absorption line intensities I_{if} from all $i \rightarrow f$ transitions falling into the wavenumber bin $[\tilde{\nu}_k - \Delta\tilde{\nu}_k/2 \dots \tilde{\nu}_k + \Delta\tilde{\nu}_k/2]$ at the given temperature T . Each grid point $\tilde{\nu}_k$ is then treated as a line position of an artificial transition (super-line) with an effective absorption intensity $I_k(T)$. The number of data points in the super-lines can be drastically reduced without significant loss of accuracy when computing pressure-dependent cross sections and can be combined with standard, pressure, temperature and frequency dependent line profiles. The super-lines cannot be associated with any specific upper/lower states and therefore the line broadening parameters used cannot depend on quantum numbers.

The super-line list files have the format of cross sections represented by two columns, wavenumbers (cm^{-1}) and super-line absorption intensity ($\text{cm}/\text{molecule}$). The Fortran format is `(F12.6,1x,ES14.8)`. The super-lines are computed on a grid of temperatures from 100 K to T_{\max} . For the resolution an adaptive grid of $R = 1\,000\,000$ is used. For the technical details of super-line construction and numerical tests see Yurchenko et al. [98].

Super-lines are currently available for H_2O , NH_3 , CH_3Cl , C_2H_2 and C_2H_4 .

4.3. Specific heat

The specific heat at constant pressure C_p is computed on a grid of temperatures of 1 K from the molecule partition function as given by [259]

$$C_p(T) = R \left[\frac{Q''}{Q} - \left(\frac{Q'}{Q} \right)^2 \right] + \frac{5R}{2}, \quad (2)$$

where the second term is the translational contribution,

$$Q'(T) = T \frac{dQ}{dT},$$

$$Q''(T) = T^2 \frac{d^2Q}{dT^2} + 2Q' \quad (3)$$

and R is the gas constant. The specific heat values are given in units of J mol⁻¹ K⁻¹ and are currently provided for H₂O (taken from Furtenbacher et al. [259]) and CH₄ (computed using the YT10to10 ro-vibrational energies [97]) only (see Fig. 3).

4.4. Line shifts using the diet format

For high resolution applications it will be important to take into account the pressure dependent line shifts. To this end a line shift diet has been introduced using the same data structure used for pressure dependent line broadening diet [260]. An example for H₂O is given in Table 7.




Figure 3: Specific heat at constant pressure of H_2O as computed by Furtenbacher et al. [259] and CH_4 generated the YT10to10 ro-vibrational energies [97].

Table 7: File `1H-1602_a0.shift` A pressure line `.shift` file for H_2O : portion of the file (upper part); field specification (lower part).

	Code	Shift	J
	a0	0.0001	0
	a0	0.0001	1
	a0	0.0001	2
	a0	0.0001	3
	a0	0.0001	4

Field	Fortran Format	C Format	Description
code	A2	%2s	Code identifying quantum number set following J''^*
γ_{ref}	F15.6	%15.6f	Line shift at reference temperature and pressure in cm^{-1}
J''	I7/F7.1	%7d	Lower J -quantum number

*Code definition: a0 = none

4.5. New broadening parameters

The pressure broadening parameters are provided for 15 molecules and are listed in Table 8 using the pressure-broadening diet [260].

The `.broad` file has a hierarchical structure; each record starts with four compulsory columns: a label defining the broadening scheme ('a0', 'a1', ...), values of γ_{ref} , n and J'' . The compulsory fields are followed by additional quantum numbers when a more detailed specification of the quantum assignments is available. The most basic scheme 'a0' represents broadening with J'' dependence only. The additional basic scheme 'a1' is used for the case with the (J', J'') dependence. Any other schemes (e.g. 'a2', 'a3', 'a4', 'b1', 'b2', 'c1', 'c2') are the molecule specific and should be described as part of the line list specifications. The broadening data format is illustrated in Table 9.

The requirement for enhanced line broadening parameters was identified by Fortney et al. [261] in their study on the need for laboratory data requirements for studies of exoplanetary atmospheres. Recognising this, our plan is significantly enhance the treatment pressure effects in future releases of the ExoMol database.

Table 8: Pressure broadening parameters in ExoMol

Molecule	Broadener	Scheme
CS	air, self	a0
HCl	H ₂ , He, self, air, CO ₂	a0
HF	H ₂ , He	a0
CO	H ₂ , He	a0
NO	air	a0
H ₂ O	H ₂ , He, self, air	a0,a1
CO ₂	air, self	a0
SO ₂	H ₂ , He	a0, a5
HCN	He, He	a0
OCS	He, He	a0
PH ₃	He, He	a0, c1
NH ₃	He, He	a0
H ₂ CO	He, He	a0
C ₂ H ₂	He, He	a0
CH ₄	He, He, CO ₂	a0, a1

Table 9: Example of the three basic broadening schemes in the ExoMol diet, ‘a0’, ‘a1’, ‘a2’.

Label	γ	n	J''	J'	K
a0	0.0860	0.096	0		
a0	0.0850	0.093		1	
...					
a1	0.0860	0.096	0	1	
a1	0.0850	0.093		1	2
...					

4.6. *ExoMolOP: Opacities*

Recently Chubb et al. [262] computed opacity cross sections and k -tables for all molecules available from the ExoMol database plus some atomic data from NIST. These data are formatted for use in various retrieval codes including Tau-REx [13–15], ARCiS [24], petitRADTRANS [28], and NEMESIS [17]. Data were calculated on temperature-pressure grids suitable for characterising a variety of exoplanet and stellar atmospheres. Broadening parameters were taken from the literature where available, with broadening parameters used for a known molecule with a similar dipole moment, where they are not available. Tables of cross sections and k -coefficients are provided on ExoMol as part of the line list webpage. The format of the data and the resolution (grid spacing) is application dependent, for example the opacities for ARCiS, petitRADTRANS, and NEMESIS are given as k -tables at the resolving power of $R = 1000$ while Tau-REx works with cross-sections with $R = 15\,000$.

The opacity cross sections and k -tables will also be made available via the virtual atomic and molecular data centre (VAMDC) portal [263, 264]. We note that ExoMol data has also been extensively used to construct the EXOPLINES molecular absorption cross-Section database for brown dwarf and giant exoplanet atmospheres [265].

4.7. *Post-processing*

ExoMol provides post-processing capabilities through the program ExoCross [266]. ExoCross has many functions such as generating pressure and temperature dependent cross sections, partition functions, specific heat, state-resolved radiative lifetimes, non-LTE spectra, electric dipole, electric quadrupole and magnetic dipole spectra. ExoCross can read data in both ExoMol and HITRAN [267] formats and output it in these formats as well as SPECTRA (<http://spectra.iao.ru/>) and Phoenix formats [16]. Should data be needed in say

HITRAN format, it is strongly recommended that the data is downloaded to a local computer in the much more compact ExoMol format and then processed using ExoCross. Examples of ExoCross input files are provided on the website.

There are also a number of Python utilities available on the ExoMol website. These are now largely redundant as ExoCross provides all functionality required to work with ExoMol line lists. However, we mention them for completeness: Utility `extract_trans.py` reads the `.trans` in `.bz2` format without requiring it to be uncompressed. The `ExoMol_to_HITRAN.py` script converts ExoMol format to HITRAN format; this should be used with caution as ExoMol format is significantly more compact than HITRAN format and the line list files are large. The program `exomol2gf.py` can be used to generate oscillator strengths.

4.8. New web services

New web services include:

- A molecule and line list search.
- A graphical illustration of a line list represented by absorption spectra at two or three temperatures (e.g. 300 K, 2000 K and 5000 K) computed using a Doppler line profile on a grid of 1 cm^{-1} .
- The ExoMol bibliography in the BibTeX format is stored at and version controlled by GitHub at <https://github.com/ExoMol/bib>.

5. ExoMol data formats and API

5.1. Format of the `states` and `trans` files

The formats of the States and Transitions files are specified in Table 10 and 11. Tables 12 and 13 show typical examples of the States (`.states`) and Transitions (`.trans`) files. A significant new feature of the 2020 update is the uncertainty field

(cm^{-1}) in the States file appearing as column 5 after the J values; this currently an optional feature but will become a compulsory column for the ExoMol States file. The uncertainty values typically come from three different sources: (i) the uncertainties of the MARVEL energies ($\sim 0.000001\text{--}0.1 \text{ cm}^{-1}$), (ii) uncertainties obtained as the fitting observed – calculated error of energies as part of the refinement to experimental data (energies or line positions) participated in the refinement ($\sim 0.001\text{--}1 \text{ cm}^{-1}$) and (iii) roughly estimated uncertainties for all other states that cannot be verified against existing experiment ($\sim 0.1\text{--}10 \text{ cm}^{-1}$). The uncertainty in the States file are given using the same format as the energy term values, i.e. with six decimal places after the decimal point, see Table 10.

To date the new standard of the States file has been applied to the only a number of key/ recent line lists namely H₂O (POKAZATEL) [61], AlH (AlHambra) [122], C₂ 8states [124, 194], HCCH (aCeTY) [130], CO₂ (UCL-4000) [132], H₃O⁺ (eXeL) [133] and TiO (Toto) [126]. A rolling programme is in place for updating the other files so the uncertainty is uniformly available for all sources.

Table 10: Updated specification of the ExoMol States file.

Field	Fortran Format	C Format	Description
i	I12	%12d	State ID
E	F12.6	%12.6f	State energy in cm^{-1}
g_{tot}	I6	%6d	State degeneracy
J	I7/F7.1	%7d/%7.1f	J -quantum number (integer/half-integer)
(ΔE)	F12.6	%12.6f	Energy uncertainty in cm^{-1} (currently optional)
(τ)	ES12.4	%12.4e	Lifetime in s (optional)
(g)	F10.6	%10.6f	Landé g -factor (optional)
(Extra)	-	-	Extra quantum numbers, any format (optional)

Table 11: Specification of the Transitions file.

Field	Fortran Format	C Format	Description
i	I12	%12d	Upper state ID
f	I12	%12d	Lower state ID
A	ES10.4	%10.4e	Einstein A coefficient in s^{-1}
$\tilde{\nu}_{fi}$	E15.6	%15.6e	Transition wavenumber in cm^{-1} (optional).

Table 12: An example of an extract from the final States file for UCL-4000 of CO₂ [132].

<i>i</i>	\tilde{E}	<i>g_{tot}</i>	<i>J</i>	unc.	Γ	<i>e/f</i>	<i>n</i> ₁	<i>n</i> ₂ ^{lin}	<i>l</i> ₂	<i>n</i> ₃
1	0.000000	1	0	0.000500	A1	e	0	0	0	0
2	1285.408200	1	0	0.000500	A1	e	0	2	0	0
3	1388.184200	1	0	0.005000	A1	e	1	0	0	0
4	2548.366700	1	0	0.000500	A1	e	1	2	0	0
5	2671.142957	1	0	0.005000	A1	e	2	0	0	0
6	2797.136000	1	0	0.005000	A1	e	1	2	0	0
7	3792.681898	1	0	0.005000	A1	e	1	4	0	0
8	3942.541358	1	0	0.005000	A1	e	3	0	0	0
9	4064.274256	1	0	0.005000	A1	e	3	0	0	0
10	4225.096148	1	0	0.005000	A1	e	1	4	0	0
11	4673.325200	1	0	0.000500	A1	e	0	0	0	2
12	5022.349428	1	0	0.005000	A1	e	1	6	0	0
13	5197.252900	1	0	0.005000	A1	e	3	2	0	0
14	5329.645446	1	0	0.005000	A1	e	4	0	0	0
15	5475.553054	1	0	0.000500	A1	e	3	2	0	0
16	5667.644584	1	0	0.005000	A1	e	2	4	0	0
17	5915.212302	1	0	0.000500	A1	e	0	2	0	2

i: State counting number.

\tilde{E} : State energy in cm⁻¹.

g_{tot}: Total state degeneracy.

J: Total angular momentum.

unc.: Uncertainty in cm⁻¹.

Γ : Total symmetry index in C_{2v}(M).

e/f: Kronig rotationless parity.

*n*₁: Normal mode stretching symmetry (*A*₁) quantum number.

*n*₂^{lin} : Normal mode linear molecule bending (*A*₁) quantum number.

*l*₂: Normal mode vibrational angular momentum quantum number.

*n*₃: Normal mode stretching asymmetric (*B*₁) quantum number.

Table 13: Extract from the Transitions file for CaO.

f	i	A_{fi}	$\tilde{\nu}_{fi}$
10571	10884	9.5518E-06	120.241863
21053	21375	1.9515E-05	120.242886
8726	9672	1.8658E-04	120.243522
11655	11950	5.0065E-06	120.243733
93209	93967	5.7055E-03	120.244192
2228	3175	7.3226E-07	120.244564
46727	46432	1.0599E-04	120.244658
44436	44774	1.4626E-04	120.245583
29037	28723	1.8052E-04	120.245669
4458	4805	1.0431E-08	120.246396
69313	68434	5.0531E-06	120.248178
22640	22985	1.1281E-07	120.248891
57027	56721	7.1064E-06	120.250180

f : Upper state counting number; i : Lower state counting number; A_{fi} : Einstein-A coefficient in s^{-1} ; $\tilde{\nu}_{fi}$: transition wavenumber in cm^{-1} .

5.2. Formats for other data types

Table 14 shows the format of the `.broad` files containing the line broadening parameters. Table 15 shows the format of the cross section `.cross` and super-lines `.super` files. Tables 16 and 17 show the format of the partition function `.pf` and specific heat `.cp` files.

Table 14: Specification of the mandatory part of the pressure broadening parameters file.

Field	Fortran Format	C Format	Description
code	A2	%2s	Code identifying quantum number set following J''
γ_{ref}	F6.4	%6.4f	Lorentzian half-width at reference temperature and pressure in $\text{cm}^{-1}/\text{bar}$
n	F6.3	%6.3f	Temperature exponent
J''	I7/F7.1	%7d/%7.1f	Lower J -quantum number integer/half-integer

Fortran format, J integer: (A2,1x,F6.4,1x,F6.3,1x,I7)

or J half-integer: (A2,1x,F6.4,1x,F6.3,1x,F7.1)

Table 15: Specification of the `.cross` cross section and `.stick` file format

Field	Fortran Format	C Format	Description
$\tilde{\nu}_i$	F12.6	%12.6f	Central bin wavenumber, cm^{-1}
σ_i	ES14.8	%14.8e	Absorption cross section, $\text{cm}^2 \text{ molec}^{-1}$
α_i	ES14.8	%14.8e	Absorption coefficient, cm molec^{-1}
Fortran format: (F12.6,1x,ES14.8)			

Table 16: Specification of the `.pf` partition function file.

Field	Fortran Format	C Format	Description
T	F8.1	%8.1d	Temperature in K
$Q(T)$	F15.4	%15.4d	Partition function (dimensionless).
$W(T)$	ES12.4	%12.4e	Cooling function in $\text{ergs s}^{-1} \text{ molecule}^{-1}$ (if available).
Fortran format: (F8.1,1x,F15.4,1x,ES12.4) or (F8.1,1x,F15.4)			

Table 17: Specification of the .cp specific heat file.

Field	Fortran Format	C Format	Description
T	F8.1	%8.1d	Temperature in K
$C_p(T)$	F15.4	%15.4d	Specific heat function (dimensionless).
Fortran format: (F8.1,1x,F15.4)			

5.3. API

5.3.1. Searching for data through the API

The molecules and isotopologues available in ExoMol are listed in a master file located at: www.exomol.com/exomol.all. Given a molecule or list of isotopologues, ExoMol can be searched for recommended datasets using the API which can be queried using the HTTP GET request method described below. The structure of a JSON file is illustrated in Fig. 4.

```

{<ISOTOPOLOGUE>:
  {'<DATATYPE>':
    {'<DATASET>':
      {'description': <DATASET-DESCRIPTION>,
       'files': [{<FILE-DESCRIPTION>,
                  'url': <URL>,
                  'size': <SIZE>
                },
                ...
              ]
      },
      {'<DATASET>':
        ...
      },
      ...
    },
    {'<DATATYPE>':
      ...
    },
    ...
  },
  {'<ISOTOPOLOGUE>':
    {
      ...
    }
  }
}

```

Figure 4: A JSON API structure used for search queries using the `HTTP GET` request in ExoMol.

To search for all data sets (and their files) related to a single molecule, use the query:

`http://exomol.com/api?molecule=<MOLECULE>`

For example, the URL `http://exomol.com/api?molecule=NH3` returns a JSON data structure describing all files related to the ammonia molecule. Within this, the list of files belonging to the BYTe-15 line list for $^{15}\text{N}^1\text{H}_3$ is accessible by traversing the JSON object with:

`json_result['(15N)(1H)3']['linelist']['BYTe-15']['files']`

and the URL for the States file belonging to this line list is at:

```
jsonresult['(15N)(1H)3']['linelist']['BYTe-15']['files'][1]['url'].
```

Note that molecules are identified by simple text strings (no subscript or superscript symbols) but that special characters must be URL-encoded: for example, the HeH^+ cation is identified by `molecule=HeH%2B`. Most software will provide this “percent-encoding” automatically.

Searches can be further refined by isotopologue by setting the query keyword `isotopologues` to a comma-delimited sequence; for example:

```
http://exomol.com/api/?isotopologues=\(12C\)\(32S\),\(12C\)\(34S\)
```

Restricting the search by datatype is also supported:

```
http://exomol.com/api/?molecule=H2O&datatype=linelist
```

returns the details of linelists for the isotopologues of water, omitting partition functions, cooling functions, opacities, etc. Valid values for the `datatype` parameter are: `linelist`, `energylevels`, `opacity`, `super` (“Super” line lists), `Cp` (heat capacity), `broadening_coefficients`, `coolingfunction`, `partitionfunction`.

5.3.2. Accessing the data via the API

There are a number of ways of accessing the data. First, all data sets are available on the ExoMol website (www.exomol.com) and can be downloaded manually. Many ExoMol line lists contain in excess of 10 billion transitions. In this case the file of transitions are generally split into chunks in frequency. These files are compressed using `.bz2` format.

However, if multiple datasets are required manual downloads are inefficient. The systematic structure of the ExoMol database filesystem allows for automated downloads from within software or using utilities such as `wget` and `curl`. As described in detail in the ExoMol2016 release, the API structures the ExoMol data resource files as URLs of the form:

```
http://exomol.com/db/<MOLECULE>/<ISOTOPOLYGE>/<DATASET>/<FILENAME>
```

where <FILENAME> is structured '<ISOTOPPOLOGUE>_<DATASET>.<DATATYPE>'.

For example, the States file of the YYLT line list for $^{31}\text{P}^{15}\text{N}$ is obtainable at the URL http://exomol.com/db/PN/31P-15N/YYLT/31P-15N__YYLT.states.bz2.

Furthermore, each dataset has a *manifest* that lists the data files it is comprised of and their sizes (in bytes). The manifest is a single text file located at the above URL with the filename '<ISOTOPPOLOGUE>_<DATASET>.manifest'. For example, the contents of http://exomol.com/db/PN/31P-15N/YYLT/31P-15N__YYLT.manifest are:

```
http://exomol.com/db/PN/31P-15N/YYLT/31P-15N__YYLT.states.bz2 142161
http://exomol.com/db/PN/31P-15N/YYLT/31P-15N__YYLT.trans.bz2 1803083
http://exomol.com/db/PN/31P-15N/YYLT/31P-15N__YYLT(pf 175000
http://exomol.com/db/PN/31P-15N/YYLT/31P-15N__YYLT.def 4761
http://exomol.com/db/PN/31P-15N/YYLT/31P-15N__YYLT.manifest 323
```

6. Moving to higher spectral resolution

Table 18: Molecules of importance for the ExoMol project with published MARVEL datasets.

Molecule	N_{iso}	N_{elec}	N_{levels}	Reference(s)
H ₂ O	7	1	18 486	Tennyson et al. [217, 223, 224, 268]
update	1	1	19 200	Tóbiás et al. [269], Furtenbacher et al. [270]
H ₃ ⁺	3	1	652	Furtenbacher et al. [231, 271]
NH ₃	1	1	4951	Al Derzi et al. [188]
update	1	1	4936	Furtenbacher et al. [272]
C ₂	1	14	5699	Furtenbacher et al. [193]
update	1	20	7087	McKemmish et al. [194]
TiO	1	9	10 564	McKemmish et al. [126, 199]
HCCH	1	1	11 213	Chubb et al. [235]
SO ₂	3	1	15 130	Tóbiás et al. [234]
H ₂ S	1	1	11 213	Chubb et al. [230]
ZrO	1	10	8088	McKemmish et al. [273]
O ₂	1	6	4279	Furtenbacher et al. [274]
NH	1	4	1058	Chubb et al. [235]
CaOH	1	5	1954	Wang et al. [275]
H ₂ CO	1	1	4841	Al-Derzi et al. [236]
NO	1	1	4106	Wong et al. [115]
AlH	2	2	331	Yurchenko et al. [122]
BeH	3	2	1264	Darby-Lewis et al. [172]
CN	1	10	7779	Syme and McKemmish [276]

N_{iso} Number of isotopologues considered;

N_{elec} Number of electronic states considered;

N_{levels} Number of energy levels extracted: value is for the main isotopologue.

Transit spectroscopy of exoplanets has thus far been performed at rather low resolution; however, very precise spectroscopic data are required for high

resolution Doppler spectroscopy [277–279]. This has proved to be an issue for important species [70]. Indeed the ExoMol datasets described above were generally constructed with greater emphasis on completeness than obtaining very precise transition frequencies [280]. In practice, however, the use of empirical energy levels in the States file means that some transition frequencies are indeed reproduced with high (experimental) accuracy; however, with the ExoMol2016 data structure it was not possible to tell how accurately a particular transition was predicted.

Although some work has been done on using ExoMol data to provide the input for high resolution studies [214], there is a clear need to adapt the database to provide the laboratory data needed for these studies. For this reason we have updated the data structure to allow the uncertainty in a particular transition frequency to be determined. This is done via uncertainties in the energy levels which are now specified (optionally) in the States file, see Table 10.

To improve the accuracy of the predicted spectra, the ExoMol States files are being systematically updated using empirically-determined energy levels. The main means of doing this is via the MARVEL (measured active rotational-vibrational energy levels) procedure [190, 269, 281]. MARVEL inverts available high resolution spectra for a given isotopologue to give a list of empirical energy levels with associated uncertainties. For levels determined by MARVEL these uncertainties are now given in the States file. Otherwise the (usually much larger) uncertainty arising from the calculation used to generate the line list is given. By combining the uncertainty of upper and lower states (ΔE_u , ΔE_l) using the standard formula

$$\Delta E_\nu = \frac{1}{\sqrt{2}} \sqrt{(\Delta E_u)^2 + (\Delta E_l)^2} \quad (4)$$

gives the uncertainty in the transition wavenumber, ΔE_ν .

To help improve the accuracy of key line lists we have been running MAR-

VEL projects on relevant molecules. Table 18 lists the astronomically important molecules for which MARVEL studies have been completed. We note that a number of these studies [199, 230, 235, 273, 282] have been performed as part of the so-called ORBYTS schools outreach project, see Sousa-Silva et al. [283] for a discussion of this. We are in the process of working through all the molecules in the ExoMol database running MARVEL projects for those isotopologues for which there are enough high accuracy laboratory data available to justify this activity.

7. Other Future development

The ExoMol project already maintains extensive molecule by molecule bibliography files. These are stored on LaTeX’s BibTex format and are freely accessible at <https://github.com/ExoMol/bib/tree/master/exomol>. Our aim is to make referencing to the original data as easy and convenient as possible, including automatic generation of the list of references to cite in the appropriate format such as BibTex, Endnote etc. Such services have been developed and offered by HITRAN [284].

We are in the process of moving the database to a more powerful platform. After the move we plan to offer the ability to compute cross sections for a given species, temperature and pressure on the fly. This will be done using super-lines which greatly reduce the computing time. We will also facilitate the computation of k -tables for a given atmospheric model.

Finally the move should allow much greater integration with the code Tau-REx [13, 14]. Tau-REx is an open source retrieval code for exoplanetary atmospheres which has just undergone a major upgrade [15]. Integration with the Tau-REx-III upgrade via a graphical interface is currently in progress.

8. Conclusions

The ExoMol database presented here is a molecule-by-molecule set of comprehensive line lists for modelling spectra and other properties of hot gases. The choice of molecules is dictated by the need to model the atmospheres of exoplanets and other hot astronomical objects, but the spectroscopic data have much wider applications than this. We are still in the process of adding molecules to the database and are receptive to suggestions of other key species to include. In addition we are working on improving the accuracy of the line positions, particularly for strong lines, enhancing the treatment of pressure broadening, and extending the range of wavelengths covered into the ultra-violet for molecules where this is considered important. In addition, we also plan to expand the database to consider temperature dependent photodissociation.

Acknowledgments

We thank the many scientists who have contributed directly or indirectly to the ExoMol project including Ala'a Azzam, Emma Barton, Bob Barber, Phillip Coles, Lorenzo Lodi, Barry Mant, Andrei Patrascu, Anatoly Pavlyuchko, Clara Sousa-Silva, Tom Rivlin, Daniel Underwood, Andrei Yachmenev and Peter Bernath. This work was supported by the European Research Council (ERC) under the Advanced Investigator ExoMol Project 267219 and the ExoAI project 758892, and STFC through grants ST/H008586/1 and ST/K00333X/1.

References

- [1] Tennyson, J., Yurchenko, S.N.. ExoMol: molecular line lists for exoplanet and other atmospheres. *Mon Not R Astron Soc* 2012;425:21–33. doi:[10.1111/j.1365-2966.2012.21440.x](https://doi.org/10.1111/j.1365-2966.2012.21440.x).
- [2] Tinetti, G., Encrenaz, T., Coustenis, A.. Spectroscopy of planetary atmospheres in our galaxy. *Astron Astrophys Rev* 2013;21:1–65. doi:[10.1007/s00159-013-0063-6](https://doi.org/10.1007/s00159-013-0063-6).

- [3] Bernath, P.F.. Molecular astronomy of cool stars and sub-stellar objects. *Int Rev Phys Chem* 2009;28:681–709. doi:10.1080/01442350903292442.
- [4] Allard, F., Hauschildt, P.H., Alexander, D.R., Starrfield, S.. Model atmosphere of very low mass stars and brown dwarfs. *Ann Rev Astron Astrophys* 1997;35:137–177.
- [5] Bordbar, M.H., Wecel, G., Hyppanen, T.. A line by line based weighted sum of gray gases model for inhomogeneous CO₂-H₂O mixture in oxy-fired combustion. *Cumbust Flame* 2014;161:2435–2445. doi:10.1016/j.combustflame.2014.03.013.
- [6] Consalvi, J.L., Liu, F.. A database of narrow-band parameters for fuels commonly encountered in fire applications. *Fire Safety J* 2015;78:202 – 218. doi:10.1016/j.firesaf.2015.10.002.
- [7] Chu, Y.N., Wang, H.M., Li, J.Q., Cheng, P., Cao, D.Z.. Emission observation for electronically excited state SO(c ¹ Σ^-) in gas phase. *Chem Phys Lett* 2002;366:147–152. doi:10.1016/S0009-2614(02)01525-7.
- [8] Carney, J.R., Lightstone, J.M., Piecuch, S., Koch, J.D.. Water temperature and concentration measurements within the expanding blast wave of a high explosive. *Measure Sci Tech* 2011;22:045601. doi:10.1088/0957-0233/22/4/045601.
- [9] Evseev, V., Fateev, A., Clausen, S.. High-resolution transmission measurements of CO₂ at high temperatures for industrial applications. *J Quant Spectrosc Radiat Transf* 2012;113:2222–2233. doi:10.1016/j.jqsrt.2012.07.015.
- [10] Dello Russo, N., DiSanti, M.A., Magee-Sauer, K., Gibb, E.L., Mumma, M.J., Barber, R.J., et al. Accurate rotational temperature retrievals from hot-band line near 2.9 μ m. *Icarus* 2004;168:186–200.
- [11] Dello Russo, N., Bonev, B.P., DiSanti, M.A., Gibb, E.L., Mumma, M.J., Magee-Sauer, K., et al. Water production rates, rotational temperatures and spin temperatures in comets C/1999 H1 (Lee), C/1999 S4 and C/2001 A2. *Astrophys J* 2005;621:537–544.
- [12] Gray, M.D., Baudry, A., Richards, A.M.S., Humphreys, E.M.L., Sobolev, A.M., Yates, J.A.. The physics of water masers observable with ALMA and SOFIA: model predictions for evolved stars. *Mon Not R Astron Soc* 2016;456:374–404. doi:10.1093/mnras/stv2437.
- [13] Waldmann, I.P., Tinetti, G., Barton, E.J., Yurchenko, S.N., Tennyson, J.. Tau-REx I: a next generation retrieval code for exoplanetary atmospheres. *Astrophys J* 2015;802:107. doi:10.1088/0004-637X/802/2/107.
- [14] Waldmann, I.P., Rocchetto, M., Tinetti, G., Barton, E.J., Yurchenko, S.N., Tennyson, J.. τ -REx II: Retrieval of Emission Spectra. *Astrophys J* 2015;813:13.
- [15] Al-Refaie, A.F., Changeat, Q., Waldmann, I.P., Tinetti, G.. TauREx

- III: A fast, dynamic and extendable framework for retrievals. arXiv e-prints 2019;:arXiv:1912.07759 [arXiv:1912.07759](https://arxiv.org/abs/1912.07759).
- [16] Jack, D., Hauschildt, P.H., Baron, E.. Time-dependent radiative transfer with PHOENIX. *Astron Astrophys* 2009;502:1043–1049.
 - [17] Irwin, P.G.J., Teanby, N.A., de Kok, R., Fletcher, L.N., Howett, C.J.A., Tsang, C.C.C., et al. The NEMESIS planetary atmosphere radiative transfer and retrieval tool. *J Quant Spectrosc Radiat Transf* 2008;109:1136 – 1150. doi:[10.1016/j.jqsrt.2007.11.006](https://doi.org/10.1016/j.jqsrt.2007.11.006).
 - [18] Line, M.R., Wolf, A.S., Zhang, X., Knutson, H., Kammer, J.A., Ellison, E., et al. A Systematic Retrieval Analysis of Secondary Eclipse Spectra. I. A Comparison of Atmospheric Retrieval Techniques. *Astrophys J* 2013;775:137.
 - [19] Malik, M., Grosheintz, L., Mendonça, J.M., Grimm, S.L., Lavie, B., Kitzmann, D., et al. HELIOS: An Open-Source, GPU-Accelerated Radiative Transfer Code For Self-Consistent Exoplanetary Atmospheres. *Astrophys J* 2017;153:56. doi:[10.3847/1538-3881/153/2/56](https://doi.org/10.3847/1538-3881/153/2/56).
 - [20] Kitzmann, D., Heng, K., Oreshenko, M., Grimm, S.L., Apai, D., Bowler, B.P., et al. Helios-r2: A New Bayesian, Open-source Retrieval Model for Brown Dwarfs and Exoplanet Atmospheres. *Astrophys J* 2020;890:174. doi:[{10.3847/1538-4357/ab6d71}](https://doi.org/10.3847/1538-4357/ab6d71).
 - [21] Drummond, B., Tremblin, P., Baraffe, I., Amundsen, D.S., Mayne, N.J., Venot, O., et al. The effects of consistent chemical kinetics calculations on the pressure-temperature profiles and emission spectra of hot Jupiters. *Astron Astrophys* 2016;594:A69. doi:[{10.1051/0004-6361/20162.8799}](https://doi.org/10.1051/0004-6361/20162.8799).
 - [22] Goyal, J.M., Mayne, N., Sing, D.K., Drummond, B., Tremblin, P., Amundsen, D.S., et al. A library of ATMO forward model transmission spectra for hot Jupiter exoplanets. *Mon Not R Astron Soc* 2018;473:5158–5185. doi:[{10.1093/mnras/stx3015}](https://doi.org/10.1093/mnras/stx3015).
 - [23] Phillips, M.W., Tremblin, P., Baraffe, I., Chabrier, G., Allard, N.F., Spiegelman, F., et al. A new set of atmosphere and evolution models for cool T-Y brown dwarfs and giant exoplanets. *Astron Astrophys* 2020;637:A38. doi:[{10.1051/0004-6361/201937381}](https://doi.org/10.1051/0004-6361/201937381).
 - [24] Min, M., Ormel, C.W., Chubb, K.L., Helling, C., Kawashima, Y.. The ARCiS framework for exoplanet atmospheres: Modelling philosophy and retrieval. *Astron Astrophys* 2020;(In preparation).
 - [25] Stolk, T., Min, M., Stam, D.M., Molliere, P., Dominik, C., Waters, L.B.F.M.. Polarized scattered light from self-luminous exoplanets Three-dimensional scattering radiative transfer with ARTES. *Astron Astrophys* 2017;607:A42. doi:[{10.1051/0004-6361/201730780}](https://doi.org/10.1051/0004-6361/201730780).

- [26] Gandhi, S., Madhusudhan, N.. Retrieval of Exoplanet Emission Spectra with HyDRA. *Mon Not R Astron Soc* 2018;474:271–288. doi:[10.1093/mnras/stx2748](https://doi.org/10.1093/mnras/stx2748).
- [27] Gandhi, S., Madhusudhan, N.. GENESIS: New Self-Consistent Models of Exoplanetary Spectra. *Mon Not R Astron Soc* 2017;472:2334–2355. doi:[10.1093/mnras/stx1601](https://doi.org/10.1093/mnras/stx1601).
- [28] Molliére, P., Wardenier, J.P., van Boekel, R., Henning, T., Molaverdikhani, K., Snellen, I.A.G.. petitRADTRANS - A Python radiative transfer package for exoplanet characterization and retrieval. *Astron Astrophys* 2019;627:A67. doi:[10.1051/0004-6361/201935470](https://doi.org/10.1051/0004-6361/201935470).
- [29] Zhang, M., Chachan, Y., Kempton, E.M.R., Knutson, H.A.. Forward modeling and retrievals with PLATON, a fast open-source tool. *Publ Astron Soc Pac* 2019;131(997):034501. doi:[10.1088/1538-3873/aaf5ad](https://doi.org/10.1088/1538-3873/aaf5ad).
- [30] Bailey, J., Kedziora-Chudczer, L.. Modelling the spectra of planets, brown dwarfs and stars using VSTAR. *Mon Not R Astron Soc* 2012;419:1913–1929. doi:[10.1111/j.1365-2966.2011.19845.x](https://doi.org/10.1111/j.1365-2966.2011.19845.x).
- [31] Yurchenko, S.N., Tennyson, J., Bailey, J., Hollis, M.D.J., Tinetti, G.. Spectrum of hot methane in astronomical objects using a comprehensive computed line list. *Proc Nat Acad Sci* 2014;111:9379–9383. doi:[10.1073/pnas.1324219111](https://doi.org/10.1073/pnas.1324219111).
- [32] BART: Bayesian Atmospheric Radiative Transfer code. *Astrophys J Suppl* 2020;.
- [33] Cubillos, P.E., *et al.*, . Yet Another Open-source Radiative-Transfer Code for Exoplanet Modeling 2020;.
- [34] Irwin, P.G.J., Bowles, N., Braude, A.S., Garland, R., Calcutt, S., Coles, P.A., *et al.* Analysis of gaseous ammonia (NH_3) absorption in the visible spectrum of Jupiter - Update. *Icarus* 2019;321:572–582. doi:[10.1016/j.icarus.2018.12.008](https://doi.org/10.1016/j.icarus.2018.12.008).
- [35] Irwin, P.G.J., Toledo, D., Garland, R., Teanby, N.A., Fletcher, L.N., Orton, G.S., *et al.* Probable detection of hydrogen sulphide (H_2S) in Neptune’s atmosphere. *Icarus* 2019;321:550–563. doi:[10.1016/j.icarus.2018.12.014](https://doi.org/10.1016/j.icarus.2018.12.014).
- [36] Berezhnoy, A.A., Borovicka, J., Santos, J., Rivas-Silva, J.F., Sandoval, L., Stolyarov, A.V., *et al.* The CaO orange system in meteor spectra. *Planet Space Sci* 2018;151:27–32. doi:[10.1016/j.pss.2017.10.007](https://doi.org/10.1016/j.pss.2017.10.007).
- [37] Kraus, M., Oksala, M.E., Cidale, L.S., Arias, M.L., Torres, A.F., Borges Fernandes, M.. Discovery of SiO band emission from galactic B[e] supergiants. *Astrophys J Lett* 2015;800:L20. doi:[10.1088/2041-8205/800/2/L20](https://doi.org/10.1088/2041-8205/800/2/L20).
- [38] Pavlenko, Y.V.. SiO Bands in the Infrared Spectra of Late-Type Stars. *Kinemat Phys Celest Bodies* 2019;.

- [39] Evans, A., Pavlenko, Y.V., Banerjee, D.P.K., Munari, U., Gehrz, R.D., Woodward, C.E., et al. Gas phase SiO in the circumstellar environment of the recurrent nova T Coronae Borealis. *Mon Not R Astron Soc* 2019;486:3498–3505. doi:10.1093/mnras/stz1071.
- [40] Hughes, G.B., Lubin, P., Meinholt, P., O'Neill, H., Brashears, T., Zhang, Q., et al. Stand-off molecular composition analysis. In: Taylor, E W and Cardimona, D A, , editor. Nanophotonics and macrophotronics for space environments IX; vol. 9616 of *Proceedings of SPIE*. 2015, p. 961603. doi:{10.1117/12.2186795}.
- [41] Hughes, G.B., Lubin, P., Cohen, A., Madajian, J., Kulkarni, N., Zhang, Q., et al. Remote laser evaporative molecular absorption spectroscopy. In: Hughes, G B, , editor. Planetary Defense and Space Environment Applications; vol. 9981 of *Proc. SPIE*. SPIE; 2016,doi:10.1117/12.2242730.
- [42] Pavlenko, Y.V., Yurchenko, S.N., McKemmish, L.K., Tennyson, J.. Analysis of the TiO isotopologues in stellar optical spectra. *Astron Astrophys* 2020;.
- [43] Owens, A., Yurchenko, S.N., Špirko, V.. Anomalous phosphine sensitivity coefficients as probes for a possible variation of the proton-to-electron mass ratio. *Mon Not R Astron Soc* 2018;473:4986–4992. doi:10.1093/mnras/stx2696.
- [44] Syme, A.M., Mousley, A., Cunningham, M., McKemmish, L.K.. Diatomic rovibronic transitions as potential probes for proton-to-electron mass ratio across cosmological time. *Australian J of Chem* 2020;doi:10.1071/CH19448.
- [45] Owens, A., Zak, E.J., Chubb, K.L., Yurchenko, S.N., Tennyson, J., Yachmenev, A.. Simulating electric-field interactions with polar molecules using spectroscopic databases. *Sci Rep* 2017;45068:7. doi:10.1038/srep45068.
- [46] Owens, A., Yachmenev, A.. Richmol: A general variational approach for rovibrational molecular dynamics in external electric fields. *J Chem Phys* 2018;148(12):124102.
- [47] Buchanan, Z., Martin-Drumel, M.A., Eliet, S., Turut, J., Mouret, G., Hindle, F., et al. Building a Database for QCL Pumped Far-IR Lasers. In: 74th International Symposium on Molecular Spectroscopy. 2019,doi:10.15278/isms.2019.MJ10.
- [48] Essig, R., Pérez-Ríos, J., Ramani, H., Slone, O.. Direct detection of nuclear scattering of sub-gev dark matter using molecular excitations. *Phys Rev Research* 2019;1:033105. doi:10.1103/PhysRevResearch.1.033105.
- [49] Markov, N.. A quantum propulsion method. In: Georgiev, P and Soares, C G, , editor. Sustainable development and innovations in marine technologies; vol. 3 of *Proceedings in Marine Technology and Ocean Engineering*. Maritime Assoc Mediterranean; Tech Univ Varna. ISBN 978-0-367-81008-5; 978-0-367-40951-7; 2020, p. 247–251. 18th In-

ternational Congress of the Maritime-Association-of-the-Mediterreanean (IMAM), Varna, BULGARIA, SEP 09-11, 2019.

- [50] Joshi, G.C., Punetha, L.M., Pande, M.C.. (A–X) system of SiO in sunspots. *Solar Phys* 1979;62:77–82. doi:10.1007/BF00150133.
- [51] Sriramachandran, P., Ambika, C., Shanmugavel, R.. The rotational temperatures of Silicon hydride isotopomer lines in sunspot umbrae. *New Astron* 2012;17(3):281–284. doi:10.1016/j.newast.2011.07.008.
- [52] Joshi, G.C., Pande, M.C.. On the presence of SH in the sunspot spectrum. *Sol Phys* 1979;62:69–75. doi:10.1007/BF00150132.
- [53] Shanmugavel, R., Bagare, S.P., Rajamanickam, N., Balachandra Kumar, K.. Identification of beryllium hydride isotopomer lines in sunspot umbral spectra. *Serb Astron J* 2008;176:51–58. doi:10.2298/SAJ0876051S.
- [54] Sriramachandran, P., Bagare, S.P., Rajamanickam, N., Balachandrakumar, K.. Presence of lao, sco and vo molecular lines in sunspot umbral spectra. *Sol Phys* 2008;252:267–281. doi:10.1007/s11207-008-9261-1.
- [55] Sotirovski, P.. The Molecular Spectrum of Sunspot Umbrae. *Astron Astrophys* 1971;14:319.
- [56] Polyansky, O.L., Zobov, N.F., Viti, S., Tennyson, J., Bernath, P.F., Wallace, L.. Water in the sun: line assignments based on variational calculations. *Science* 1997;277:346–349.
- [57] Polyansky, O.L., Zobov, N.F., Viti, S., Tennyson, J., Bernath, P.F., Wallace, L.. High temperature rotational transitions of water in sunspot and laboratory spectra. *J Mol Spectrosc* 1997;186:422–447.
- [58] Zobov, N.F., Polyansky, O.L., Tennyson, J., Shirin, S.V., Nassar, R., Hirao, T., et al. Using laboratory spectroscopy to identify lines in the K and L-band spectrum of water in a sunspot. *Astrophys J* 2000;530:994–998.
- [59] Tereszchuk, K., Bernath, P.F., Zobov, N.F., Shirin, S.V., Polyansky, O.L., Libeskind, N.I., et al. Laboratory spectroscopy of hot water near 2-microns and sunspot spectroscopy in the H-band region. *Astrophys J* 2002;577:496–500.
- [60] Sonnabend, G., Wirtz, D., Schieder, R., Bernath, P.. High-resolution infrared measurements of H₂O and SiO in sunspots. *Solar Phys* 2006;233:205–213. doi:10.1007/s11207-006-2488-9.
- [61] Polyansky, O.L., Kyuberis, A.A., Zobov, N.F., Tennyson, J., Yurchenko, S.N., Lodi, L.. ExoMol molecular line lists XXX: a complete high-accuracy line list for water. *Mon Not R Astron Soc* 2018;480:2597–2608. doi:10.1093/mnras/sty1877.

- [62] Tennyson, J., Yurchenko, S.N., Al-Refaie, A.F., Barton, E.J., Chubb, K.L., Coles, P.A., et al. The ExoMol database: molecular line lists for exoplanet and other hot atmospheres. *J Mol Spectrosc* 2016;327:73–94. doi:10.1016/j.jms.2016.05.002.
- [63] Ito, Y., Ikoma, M., Kawahara, H., Nagahara, H., Kawashima, Y., Nakamoto, T.. Theoretical emission spectra of atmospheres of hot rocky super-earths. *Astrophys J* 2015;801:144. doi:10.1088/0004-637X/801/2/144.
- [64] Fegley Jr., B., Jacobson, N.S., Williams, K.B., Plane, J.M.C., Schaefer, L., Lodders, K.. Solubility of Rock in Steam Atmospheres of Planets. *Astrophys J* 2016;824:103. doi:10.3847/0004-637X/824/2/103.
- [65] Tennyson, J., Yurchenko, S.N.. Laboratory spectra of hot molecules: data needs for hot super-earth exoplanets. *Mol Astrophys* 2017;8:1–18. doi:10.1016/j.molap.2017.05.002.
- [66] Herbort, O., Woitke, P., Helling, C., Zerkle, A.. The atmospheres of rocky exoplanets I. Outgassing of common rock and the stability of liquid water. *Astron Astrophys* 2020;636:A71. doi:{10.1051/0004-6361/201936614}.
- [67] Gaudi, B.S., Stassun, K.G., Collins, K.A., Beatty, T.G., Zhou, G., Latham, D.W., et al. A giant planet undergoing extreme-ultraviolet irradiation by its hot massive-star host. *Nature* 2017;546:514+. doi:10.1038/nature22392.
- [68] Snellen, I.. High-dispersion spectroscopy of extrasolar planets: from CO in hot jupiters to O₂ in exo-earths. *Phil Trans Royal Soc London A* 2014;372:20130075. doi:10.1098/rsta.2013.0075.
- [69] Birkby, J.L.. Spectroscopic direct detection of exoplanets. *Handbook of Exoplanets* 2018;:1485–1508.
- [70] Hoeijmakers, H.J., de Kok, R.J., Snellen, I.A.G., Brogi, M., Birkby, J.L., Schwarz, H.. A search for TiO in the optical high-resolution transmission spectrum of HD 209458b: Hindrance due to inaccuracies in the line database. *Astron Astrophys* 2015;575:A20. doi:{10.1051/0004-6361/201424794}.
- [71] Rothman, L.S., Jacquemart, D., Barbe, A., Benner, D.C., Birk, M., Brown, L.R., et al. The *HITRAN* 2004 molecular spectroscopic database. *J Quant Spectrosc Radiat Transf* 2005;96:139–204.
- [72] Rothman, L.S., Gordon, I.E., Barbe, A., Benner, D.C., Bernath, P.F., Birk, M., et al. The *HITRAN* 2008 molecular spectroscopic database. *J Quant Spectrosc Radiat Transf* 2009;110:533–572.
- [73] Down, M.J., Hill, C., Yurchenko, S.N., Tennyson, J., Brown, L.R., Kleiner, I.. Re-analysis of ammonia spectra: Updating the HITRAN ¹⁴NH₃ database. *J Quant Spectrosc*

- Radiat Transf 2013;130:260–272. doi:10.1016/j.jqsrt.2013.05.027.
- [74] Jacquinet-Husson, N., Crepeau, L., Armante, R., Boutammine, C., Chédin, A., Scott, N.A., et al. The 2009 edition of the GEISA spectroscopic database. J Quant Spectrosc Radiat Transf 2011;112:2395–2445.
- [75] Jacquinet-Husson, N., Armante, R., Scott, N.A., Chédin, A., Crépeau, L., Boutammine, C., et al. The 2015 edition of the GEISA spectroscopic database. J Mol Spectrosc 2016;327:31–72. doi:10.1016/j.jms.2016.06.007.
- [76] Rothman, L.S., Gordon, I.E., Barber, R.J., Dothe, H., Gamache, R.R., Goldman, A., et al. HITRAN, the High-Temperature Molecular Spectroscopic Database. J Quant Spectrosc Radiat Transf 2010;111:2139–2150.
- [77] Hargreaves, R.J., Gordon, I.E., Rothman, L.S., Tashkun, S.A., Perevalov, V.I., Lukashevskaya, A.A., et al. Spectroscopic line parameters of NO, NO₂, and N₂O for the HITRAN database. J Quant Spectrosc Radiat Transf 2019;232:35 – 53. doi:10.1016/j.jqsrt.2019.04.040.
- [78] Hargreaves, R.J., Gordon, I.E., Rey, M., Nikitin, A.V., Tyuterev, V.G., Kochanov, R.V., et al. An accurate, extensive, and practical line list of methane for the HITRAN database. Astrophys J Suppl 2020;247:55.
- [79] Conway, E.K., Gordon, I.E., Hargreaves, R.J., Gamache, R.R., Polyansky, O.L., Tennyson, J.. An accurate high temperature water line list for the HITRAN database. J Quant Spectrosc Radiat Transf 2020;In Preparation.
- [80] Rey, M., Nikitin, A.V., Babikov, Y.L., Tyuterev, V.G.. TheoReTS – an information system for theoretical spectra based on variational predictions from molecular potential energy and dipole moment surfaces. J Mol Spectrosc 2016;327:138–158. doi:10.1016/j.jms.2016.04.006.
- [81] Kurucz, R.L.. Including all the lines. Can J Phys 2011;89:417–428.
- [82] Bernath, P.F.. MOLIST: Molecular Line Lists, Intensities and Spectra. J Quant Spectrosc Radiat Transf 2020;240:106687. doi:10.1016/j.jqsrt.2019.106687.
- [83] Wang, Y., Tennyson, J., Yurchenko, S.N.. Empirical line lists in the ExoMol database. Atoms 2020;8:7. doi:10.3390/atoms8010007.
- [84] Rivière, P., Soufiani, A.. Updated band model parameters for H₂O, CO₂, CH₄ and CO radiation at high temperature. Intern J Heat Mass Transf 2012;55:3349 – 3358. doi:10.1016/j.ijheatmasstransfer.2012.03.019.
- [85] Parigger, C.G., Woods, A.C., Surmick, D.M., Gautam, G., Witte, M.J., Hornkohl, J.. Computation of diatomic molecular spectra for selected transitions of aluminum monoxide,

- cyanide, diatomic carbon, and titanium monoxide. *Spectra Chimica Acta B* 2015;107:132–138. doi:10.1016/j.sab.2015.02.018.
- [86] Tennyson, J.. Accurate variational calculations for line lists to model the vibration rotation spectra of hot astrophysical atmospheres. *WIREs Comput Mol Sci* 2012;2:698–715. doi:10.1002/wcms.94.
- [87] Tennyson, J.. Vibration-rotation transition dipoles from first principles. *J Mol Spectrosc* 2014;298:1–6. doi:10.1016/j.jms.2014.01.012.
- [88] Tennyson, J., Hulme, K., Naim, O.K., Yurchenko, S.N.. Radiative lifetimes and cooling functions for astrophysically important molecules. *J Phys B: At Mol Opt Phys* 2016;49:044002. doi:10.1088/0953-4075/49/4/044002.
- [89] Tennyson, J., Lodi, L., McKemmish, L.K., Yurchenko, S.N.. The ab initio calculation of spectra of open shell diatomic molecules. *J Phys B: At Mol Opt Phys* 2016;49:102001.
- [90] Tennyson, J.. Perspective: Accurate ro-vibrational calculations on small molecules. *J Chem Phys* 2016;145:120901. doi:10.1063/1.4962907.
- [91] Semenov, M., Yurchenko, S.N., Tennyson, J.. Predicted Landé g-factors for open shell diatomic molecules. *J Mol Spectrosc* 2017;330:57–62. doi:10.1016/j.jms.2016.11.004.
- [92] Yurchenko, S.N., Lodi, L., Tennyson, J., Stolyarov, A.V.. Duo: a general program for calculating spectra of diatomic molecules. *Comput Phys Commun* 2016;202:262–275. doi:10.1016/j.cpc.2015.12.021.
- [93] Tennyson, J., Yurchenko, S.N.. The ExoMol project: Software for computing molecular line lists. *Intern J Quantum Chem* 2017;117:92–103. doi:10.1002/qua.25190.
- [94] Yadin, B., Vaness, T., Conti, P., Hill, C., Yurchenko, S.N., Tennyson, J.. ExoMol Molecular linelists: I The rovibrational spectrum of BeH, MgH and CaH the $X^2\Sigma^+$ state. *Mon Not R Astron Soc* 2012;425:34–43.
- [95] Barton, E.J., Yurchenko, S.N., Tennyson, J.. ExoMol Molecular linelists – II. The ro-vibrational spectrum of SiO. *Mon Not R Astron Soc* 2013;434:1469–1475.
- [96] Barber, R.J., Strange, J.K., Hill, C., Polyansky, O.L., Mellau, G.C., Yurchenko, S.N., et al. ExoMol line lists – III. An improved hot rotation-vibration line list for HCN and HNC. *Mon Not R Astron Soc* 2014;437:1828–1835. doi:10.1093/mnras/stt2011.
- [97] Yurchenko, S.N., Tennyson, J.. ExoMol line lists IV: The rotation-vibration spectrum of methane up to 1500 K. *Mon Not R Astron Soc* 2014;440:1649–1661.
- [98] Yurchenko, S.N., Amundsen, D.S., Tennyson, J., Waldmann, I.P.. A hybrid line list for CH₄ and hot methane continuum . *Astron Astrophys* 2017;605:A95. doi:10.1051/0004-6361/201731026.

- [99] Barton, E.J., Chiu, C., Golpayegani, S., Yurchenko, S.N., Tennyson, J., Frohman, D.J., et al. ExoMol Molecular linelists – V. The ro-vibrational spectra of NaCl and KCl. *Mon Not R Astron Soc* 2014;442:1821–1829. doi:[10.1093/mnras/stu944](https://doi.org/10.1093/mnras/stu944).
- [100] Yorke, L., Yurchenko, S.N., Lodi, L., Tennyson, J.. ExoMol line lists VI: A high temperature line list for Phosphorus Nitride. *Mon Not R Astron Soc* 2014;445:1383–1391. doi:[10.1093/mnras/stu1854](https://doi.org/10.1093/mnras/stu1854).
- [101] Sousa-Silva, C., Al-Refaie, A.F., Tennyson, J., Yurchenko, S.N.. ExoMol line lists - VII. the rotation-vibration spectrum of phosphine up to 1500 K. *Mon Not R Astron Soc* 2015;446:2337–2347. doi:[10.1093/mnras/stu2246](https://doi.org/10.1093/mnras/stu2246).
- [102] Al-Refaie, A.F., Yurchenko, S.N., Yachmenev, A., Tennyson, J.. ExoMol line lists - VIII: A variationally computed line list for hot formaldehyde. *Mon Not R Astron Soc* 2015;448:1704–1714. doi:[10.1093/mnras/stv091](https://doi.org/10.1093/mnras/stv091).
- [103] Patrascu, A.T., Tennyson, J., Yurchenko, S.N.. ExoMol molecular linelists: VII: The spectrum of AlO. *Mon Not R Astron Soc* 2015;449:3613–3619. doi:[10.1093/mnras/stv507](https://doi.org/10.1093/mnras/stv507).
- [104] Rivlin, T., Lodi, L., Yurchenko, S.N., Tennyson, J., Le Roy, R.J.. ExoMol line lists X: The spectrum of sodium hydride. *Mon Not R Astron Soc* 2015;451:5153–5157. doi:[10.1093/mnras/stv979](https://doi.org/10.1093/mnras/stv979).
- [105] Pavlyuchko, A.I., Yurchenko, S.N., Tennyson, J.. ExoMol line lists XI: A Hot Line List for nitric acid. *Mon Not R Astron Soc* 2015;452:1702–1706. doi:[10.1093/mnras/stv1376](https://doi.org/10.1093/mnras/stv1376).
- [106] Paulose, G., Barton, E.J., Yurchenko, S.N., Tennyson, J.. ExoMol Molecular linelists – XII. Line lists for eight isotopologues of CS. *Mon Not R Astron Soc* 2015;454:1931–1939. doi:[10.1093/mnras/stv1543](https://doi.org/10.1093/mnras/stv1543).
- [107] Yurchenko, S.N., Blissett, A., Asari, U., Vasilios, M., Hill, C., Tennyson, J.. ExoMol Molecular linelists – XIII. The spectrum of CaO. *Mon Not R Astron Soc* 2016;456:4524–4532. doi:[10.1093/mnras/stv2858](https://doi.org/10.1093/mnras/stv2858).
- [108] Underwood, D.S., Tennyson, J., Yurchenko, S.N., Huang, X., Schwenke, D.W., Lee, T.J., et al. ExoMol line lists XIV: A line list for hot SO₂. *Mon Not R Astron Soc* 2016;459:3890–3899. doi:[10.1093/mnras/stw849](https://doi.org/10.1093/mnras/stw849).
- [109] Al-Refaie, A.F., Polyansky, O.L., Ovsyannikov, R.I., Tennyson, J., Yurchenko, S.N.. ExoMol line lists XV: A hot line-list for hydrogen peroxide. *Mon Not R Astron Soc* 2016;461:1012–1022. doi:[10.1093/mnras/stw1295](https://doi.org/10.1093/mnras/stw1295).
- [110] Azzam, A.A.A., Yurchenko, S.N., Tennyson, J., Naumenko, O.V.. ExoMol line lists XVI: A Hot Line List for H₂S. *Mon Not R Astron Soc* 2016;460:4063–4074. doi:[10.1093/mnras/stw1133](https://doi.org/10.1093/mnras/stw1133).

- [111] Underwood, D.S., Tennyson, J., Yurchenko, S.N., Clausen, S., Fateev, A.. ExoMol line lists XVII: A line list for hot SO₃. *Mon Not R Astron Soc* 2016;462:4300–4313. doi:10.1093/mnras/stw1828.
- [112] McKemmish, L.K., Yurchenko, S.N., Tennyson, J.. ExoMol Molecular linelists – XVIII. The spectrum of Vanadium Oxide. *Mon Not R Astron Soc* 2016;463:771–793. doi:10.1093/mnras/stw1969.
- [113] Polyansky, O.L., Kyuberis, A.A., Lodi, L., Tennyson, J., Ovsyannikov, R.I., Zobov, N.. ExoMol molecular line lists XIX: high accuracy computed line lists for H₂¹⁷O and H₂¹⁸O. *Mon Not R Astron Soc* 2017;466:1363–1371. doi:10.1093/mnras/stw3125.
- [114] Mizus, I.I., Alijah, A., Zobov, N.F., Kyuberis, A.A., Yurchenko, S.N., Tennyson, J., et al. ExoMol molecular line lists XX: a comprehensive line list for H₃⁺. *Mon Not R Astron Soc* 2017;468:1717–1725. doi:10.1093/mnras/stx502.
- [115] Wong, A., Yurchenko, S.N., Bernath, P., Mueller, H.S.P., McConkey, S., Tennyson, J.. ExoMol Line List XXI: Nitric Oxide (NO). *Mon Not R Astron Soc* 2017;470:882–897. doi:10.1093/mnras/stx1211.
- [116] Owens, A., Yurchenko, S.N., Yachmenev, A., Thiel, W., Tennyson, J.. ExoMol molecular line lists XXII. The rotation-vibration spectrum of silane up to 1200 K. *Mon Not R Astron Soc* 2017;471:5025–5032. doi:10.1093/mnras/stx1952.
- [117] Prajapat, L., Jagoda, P., Lodi, L., Gorman, M.N., Yurchenko, S.N., Tennyson, J.. ExoMol molecular line lists XXIII. Spectra of PO and PS. *Mon Not R Astron Soc* 2017;472:3648–3658. doi:10.1093/mnras/stx2229.
- [118] Yurchenko, S.N., Sinden, F., Lodi, L., Hill, C., Gorman, M.N., Tennyson, J.. ExoMol Molecular linelists – XXIV: A new hot line list for silicon monohydride, SiH. *Mon Not R Astron Soc* 2018;473:5324–5333. doi:10.1093/mnras/stx2738.
- [119] Upadhyay, A., Conway, E.K., Tennyson, J., Yurchenko, S.N.. ExoMol Molecular linelists – XXV: A hot line list for silicon sulphide, SiS. *Mon Not R Astron Soc* 2018;477:1520–1527. doi:10.1093/mnras/sty998.
- [120] Yurchenko, S.N., Bond, W., Gorman, M.N., Lodi, L., McKemmish, L.K., Nunn, W., et al. ExoMol Molecular linelists – XXVI: spectra of SH and NS. *Mon Not R Astron Soc* 2018;478:270–282. doi:10.1093/mnras/sty939.
- [121] Mant, B.P., Yachmenev, A., Tennyson, J., Yurchenko, S.N.. ExoMol molecular line lists - XXVII: spectra of C₂H₄. *Mon Not R Astron Soc* 2018;478:3220 – 3232. doi:10.1093/mnras/sty1239.
- [122] Yurchenko, S.N., Williams, H., Leyland, P.C., Lodi, L., Tennyson, J.. ExoMol line lists

- XXVIII: The rovibronic spectrum of AlH. *Mon Not R Astron Soc* 2018;479:1401–1411. doi:10.1093/mnras/sty1524.
- [123] Owens, A., Yachmenev, A., Tennyson, J., Thiel, W., Yurchenko, S.N.. ExoMol Molecular line lists XXIX: The rotation-vibration spectrum of methyl chloride up to 1200 K. *Mon Not R Astron Soc* 2018;479:3002–3010. doi:10.1093/mnras/sty1542.
- [124] Yurchenko, S.N., Szabo, I., Pyatenko, E., Tennyson, J.. ExoMol Molecular line lists XXXI: The spectrum of C₂. *Mon Not R Astron Soc* 2018;480:3397–3411. doi:10.1093/mnras/sty2050.
- [125] Li, H.Y., Tennyson, J., Yurchenko, S.N.. ExoMol molecular line lists XXXII: the rovibronic spectrum of MgO. *Mon Not R Astron Soc* 2019;486:2351–2365. doi:10.1093/mnras/stz912.
- [126] McKemmish, L.K., Masseron, T., Hoeijmakers, J., Pérez-Mesa, V.V., Grimm, S.L., Yurchenko, S.N., et al. ExoMol Molecular line lists – XXXIII. The spectrum of Titanium Oxide. *Mon Not R Astron Soc* 2019;488:2836–2854.
- [127] Langleben, J., , Yurchenko, S.N., Tennyson, J.. ExoMol line list XXXIV: A Rovibrational Line List for Phosphinidene (PH) in its X ³ Σ ⁻ and a ¹ Δ Electronic States. *Mon Not R Astron Soc* 2019;488:2332. doi:10.1093/mnras/stz1856-2342.
- [128] Coles, P.A., , Yurchenko, S.N., Tennyson, J.. ExoMol molecular line lists XXXV: a rotation-vibration line list for hot ammonia. *Mon Not R Astron Soc* 2019;490:4638–4647. doi:10.1093/mnras/stz2778.
- [129] Gorman, M., Yurchenko, S.N., Tennyson, J.. ExoMol Molecular linelists – XXXVI. X ² Π – X ² Π and A ² Σ ⁺ – X ² Π transitions of SH. *Mon Not R Astron Soc* 2019;490:1652–1665. doi:10.1093/mnras/stz2517/5565070.
- [130] Chubb, K.L., Tennyson, J., Yurchenko, S.N.. ExoMol Molecular linelists – XXXVII: spectra of acetylene. *Mon Not R Astron Soc* 2020;493:1531–1545. doi:10.1093/mnras/staa229.
- [131] Owens, A., Conway, E.K., Tennyson, J., Yurchenko, S.N.. ExoMol Molecular linelists – XXXVIII: High-temperature molecular line list of silicon dioxide (SiO₂). *Mon Not R Astron Soc* 2020;495:1927–1933. doi:10.1093/mnras/staa1287.
- [132] Yurchenko, S.N., Mellor, T.M., Freedman, R.S., Tennyson, J.. ExoMol molecular line lists XXXIX: Ro-vibrational molecular line list for CO₂. *Mon Not R Astron Soc* 2020;doi:10.1093/mnras/staa1874.
- [133] Yurchenko, S.N., Tennyson, J., Miller, S., Melnikov, V.V., O'Donoghue, J., Moore, L.. ExoMol molecular line lists XL: Ro-vibrational molecular line list for the hydronium

- ion (H_3O^+). Mon Not R Astron Soc 2020;.
- [134] Harris, G.J., Larner, F.C., Tennyson, J., Kaminsky, B.M., Pavlenko, Y.V., Jones, H.R.A.. A $\text{H}^{13}\text{CN}/\text{HN}^{13}\text{C}$ linelist, model atmospheres and synthetic spectra for carbon stars. Mon Not R Astron Soc 2008;390:143–148.
- [135] Sochi, T., Tennyson, J.. A computed line list for the H_2D^+ molecular ion. Mon Not R Astron Soc 2010;405:2345–2350.
- [136] Voronin, B.A., Tennyson, J., Tolchenov, R.N., Lugovskoy, A.A., Yurchenko, S.N.. A high accuracy computed line list for the HDO molecule. Mon Not R Astron Soc 2010;402:492–496.
- [137] Yurchenko, S.N.. A theoretical room-temperature line list for $^{15}\text{NH}_3$. J Quant Spectrosc Radiat Transf 2015;152:28–36. doi:10.1016/j.jqsrt.2014.10.023.
- [138] Roueff, E., Abgrall, H., Czachorowski, P., Pachucki, K., Puchalski, M., Komasa, J.. The full infrared spectrum of molecular hydrogen. Astron Astrophys 2019;630:A58. doi:10.1051/0004-6361/201936249.
- [139] Masseron, T., Plez, B., Van Eck, S., Colin, R., Daoutidis, I., Godefroid, M., et al. CH in stellar atmospheres: an extensive linelist. Astron Astrophys 2014;571:A47. doi:10.1051/0004-6361/201423956.
- [140] Brooke, J.S.A., Bernath, P.F., Western, C.M., van Hemert, M.C., Groenenboom, G.C.. Line strengths of rovibrational and rotational transitions within the $X\ ^3\Sigma^-$ ground state of NH. J Chem Phys 2014;141:054310. doi:10.1063/1.4891468.
- [141] Brooke, J.S.A., Bernath, P.F., Western, C.M.. Note: Improved line strengths of rovibrational and rotational transitions within the $X\ ^3\Sigma^-$ ground state of NH. J Chem Phys 2015;143:026101. doi:10.1063/1.4923422.
- [142] Fernando, A.M., Bernath, P.F., Hodges, J.N., Masseron, T.. A new linelist for the $A\ ^3\Pi - X\ ^3\Sigma^-$ transition of the NH free radical. J Quant Spectrosc Radiat Transf 2018;217:29–34. doi:10.1016/j.jqsrt.2018.05.021.
- [143] Brooke, J.S.A., Bernath, P.F., Western, C.M., Sneden, C., Afsar, M., Li, G., et al. Line strengths of rovibrational and rotational transitions in the $X\ ^2\Pi$ ground state of OH. J Quant Spectrosc Radiat Transf 2016;138:142–157. doi:10.1016/j.jqsrt.2015.07.021.
- [144] Yousefi, M., Bernath, P.F.. Line Lists for AlF and AlCl in the $X^1\Sigma^+$ Ground State. Astrophys J Suppl 2018;237:8. doi:10.3847/1538-4365/aacc6a.
- [145] Hodges, J.N., Bernath, P.F.. Fourier Transform Spectroscopy of the $A\ ^3\Pi - X\ ^3\Sigma^-$ Transition of OH^+ . Astrophys J 2017;840:81. doi:10.3847/1538-4357/aa6bf5.
- [146] Hodges, J.N., Bittner, D.M., Bernath, P.F.. Improved Ultraviolet and Infrared Oscillator

- Strengths for OH⁺. *Astrophys J* 2018;855. doi:10.3847/1538-4357/aaa7ef.
- [147] Hou, S., Bernath, P.F.. Line list for the ground state of CaF. *J Quant Spectrosc Radiat Transf* 2018;210:44 – 51. doi:10.1016/j.jqsrt.2018.02.011.
- [148] Hou, S., Bernath, P.F.. Line list for the MgF ground state. *J Quant Spectrosc Radiat Transf* 2017;203:511 – 516.
- [149] Frohman, D.J., Bernath, P.F., Brooke, J.S.A.. Molecular line lists: The ro-vibrational spectra of NaF and KF. *J Quant Spectrosc Radiat Transf* 2016;169:104–110. doi:10.1016/j.jqsrt.2015.10.004.
- [150] Bittner, D.M., Bernath, P.F.. Line Lists for LiF and LiCl in the X ¹ Σ^+ Ground State. *Astrophys J Suppl* 2018;235:8. doi:10.3847/1538-4365/aa9846.
- [151] GharibNezhad, E., Shayesteh, A., Bernath, P.F.. Einstein A coefficients for rovibronic lines of the A ² Π → X ² Σ^+ and B' ² Σ^+ → X ² Σ^+ transitions of MgH. *Mon Not R Astron Soc* 2013;432:2043–2047. doi:10.1093/mnras/stt510.
- [152] Burrows, A., Dulick, M., Bauschlicher, C.W., Bernath, P.F., Ram, R.S., Sharp, C.M., et al. Spectroscopic constants, abundances, and opacities of the tih molecule. *Astrophys J* 2005;624:988–1002. doi:10.1086/429366.
- [153] Chowdhury, P.K., Merer, A.J., Rixon, S.J., Bernath, P.F., Ram, R.S.. Low-N lines of the A ⁶ Σ^+ - X ⁶ Σ^+ (1,0) band of CrH. *Phys Chem Chem Phys* 2006;8:822–826. doi:10.1039/b514188e.
- [154] Wende, S., Reiners, A., Seifahrt, A., Bernath, P.F.. Crires spectroscopy and empirical line-by-line identification of feh molecular absorption in an m dwarf. *Astron Astrophys* 2010;523:A58. doi:10.1051/0004-6361/201015220.
- [155] Coxon, J.A., Hajigeorgiou, P.G.. Improved direct potential fit analyses for the ground electronic states of the hydrogen halides: HF/DF/TF, HCl/DCl/TCl, HBr/DBr/TBr and HI/DI/TI. *J Quant Spectrosc Radiat Transf* 2015;151:133 – 154. doi:10.1016/j.jqsrt.2014.08.028.
- [156] Li, G., Gordon, I.E., Bernath, P.F., Rothman, L.S.. Direct fit of experimental ro-vibrational intensities to the dipole moment function: Application to HCl . *J Quant Spectrosc Radiat Transf* 2011;112:1543 – 1550. doi:10.1016/j.jqsrt.2011.03.014.
- [157] Ram, R.S., Brooke, J.S.A., Western, C.M., Bernath, P.F.. Einstein A-values and oscillator strengths of the A ² Π – X ² Σ^+ system of CP. *J Quant Spectrosc Radiat Transf* 2014;138:107–115. doi:10.1016/j.jqsrt.2014.01.030.
- [158] Brooke, J.S.A., Ram, R.S., Western, C.M., Li, G., Schwenke, D.W., Bernath, P.F.. Einstein A coefficients and oscillator strengths for the A ² Π – X ² Σ^+ (red) and B ² Σ^+ –

- X $^2\Sigma^+$ violet systems and rovibrational transitions in the X $^2\Sigma^+$ state of CN. *Astrophys J Suppl* 2014;210:23. doi:10.1088/0067-0049/210/2/23.
- [159] Brooke, J.S.A., Bernath, P.F., Schmidt, T.W., Bacskay, G.B.. Line strengths and updated molecular constants for the C₂ Swan system. *J Quant Spectrosc Radiat Transf* 2013;124:11 – 20. doi:10.1016/j.jqsrt.2013.02.025.
- [160] Li, G., Harrison, J.J., Ram, R.S., Western, C.M., Bernath, P.F.. Einstein A coefficients and absolute line intensities for the E $^2\Pi$ - X $^2\Sigma^+$ transition of CaH. *J Quant Spectrosc Radiat Transf* 2012;113:67–74. doi:10.1016/j.jqsrt.2011.09.010.
- [161] Shayesteh, A., Ram, R.S., Bernath, P.F.. Fourier transform emission spectra of the A $^2\Pi \rightarrow$ X $^2\Sigma^+$ and B $^2\Sigma^+ \rightarrow$ X $^2\Sigma^+$ band systems of CaH. *J Mol Spectrosc* 2013;288:46–51. doi:10.1016/j.jms.2013.04.009.
- [162] Western, C.M., Carter-Blatchford, L., Crozet, P., Ross, A.J., Morville, J., Tokaryk, D.W.. The spectrum of N₂ from 4,500 to 15,700 cm⁻¹ revisited with PGOPHER. *J Quant Spectrosc Radiat Transf* 2018;219:127–141. doi:10.1016/j.jqsrt.2018.07.017.
- [163] Lodi, L., Yurchenko, S.N., Tennyson, J.. The calculated rovibronic spectrum of scandium hydride, ScH. *Mol Phys* 2015;113:1559–1575. doi:10.1080/00268976.2015.1029996.
- [164] Coppola, C.M., Lodi, L., Tennyson, J.. Radiative cooling functions for primordial molecules. *Mon Not R Astron Soc* 2011;415:487–493.
- [165] Li, G., Gordon, I.E., Rothman, L.S., Tan, Y., Hu, S.M., Kassi, S., et al. Rovibrational line lists for nine isotopologues of the CO molecule in the X $^1\Sigma^+$ ground electronic state. *Astrophys J Suppl* 2015;216:15. doi:10.1088/0067-0049/216/1/15.
- [166] Amaral, P.H.R., Diniz, L.G., Jones, K.A., Stanke, M., Alijah, A., Adamowicz, L., et al. Benchmark Rovibrational Linelists and Einstein A-coefficients for the Primordial Molecules and Isotopologues. *Astrophys J* 2019;878(2):95. URL: <https://doi.org/10.3847/2F1538-4357%2Fab1f65>. doi:10.3847/1538-4357/ab1f65.
- [167] Owens, A., Yachmenev, A., Küpper, J., Yurchenko, S.N., Thiel, W.. The rotation-vibration spectrum of methyl fluoride from first principles. *Phys Chem Chem Phys* 2018;21:3496–3505. doi:10.1039/C8CP01721B.
- [168] Coles, P.A., Yurchenko, S.N., Kovacich, R.P., Hobby, J., Tennyson, J.. A variationally computed room temperature line list for AsH₃. *Phys Chem Chem Phys* 2019;21:3264–3277. doi:10.1039/C8CP07110A.
- [169] Owens, A., Yurchenko, S.N.. Theoretical rotation-vibration spectroscopy of cis- and trans-diphosphene (P₂H₂) and the deuterated species P₂HD. *J Chem Phys* 2019;150(19):194308. doi:10.1063/1.5092767.

- [170] Mant, B.P., Chubb, K.L., Yachmenev, A., Tennyson, J., Yurchenko, S.N.. The Infrared Spectrum of PF₃ and the Analysis of Rotational Energy Clustering Effect. Mol Phys 2019;118:e1581951. doi:10.1080/00268976.2019.1581951.
- [171] Adam, A.Y., Yachmenev, A., Yurchenko, S.N., Jensen, P.. A Variationally Computed IR Line List for the Methyl Radical CH₃. J Phys Chem A 2019;123:22. doi:10.1021/acs.jpca.9b02919.
- [172] Darby-Lewis, D., Tennyson, J., Lawson, K.D., Yurchenko, S.N., Stamp, M.F., Shaw, A., et al. Synthetic spectra of BeH, BeD and BeT for emission modelling in JET plasmas. J Phys B: At Mol Opt Phys 2018;51:185701. doi:10.1088/1361-6455/aad6d0.
- [173] Huang, X., Freedman, R.S., Tashkun, S.A., Schwenke, D.W., Lee, T.J.. Semi-empirical ¹²C¹⁶O₂ IR line lists for simulations up to 1500 K and 20,000 cm⁻¹. J Quant Spectrosc Radiat Transf 2013;130:134–146. doi:10.1016/j.jqsrt.2013.05.018.
- [174] Huang, X., Schwenke, D.W., Freedman, R.S., Lee, T.J.. Ames-2016 line lists for 13 isotopologues of CO₂ : Updates, consistency, and remaining issues. J Quant Spectrosc Radiat Transf 2017;203:224 – 241. doi:10.1016/j.jqsrt.2017.04.026.
- [175] Clark, V.H.J., Owens, A., Tennyson, J., Yurchenko, S.N.. The high-temperature rotation-vibration spectrum and rotational clustering of silylene (SiH₂). J Quant Spectrosc Radiat Transf 2020;246:106929. doi:10.1016/j.jqsrt.2020.106929.
- [176] Yurchenko, S.N., Smirnov, A.N., Solomonik, V.G., Tennyson, J.. Spectroscopy of YO from first principles. Phys Chem Chem Phys 2019;21:22794–22810. doi:10.1039/C9CP03208H.
- [177] Neale, L., Miller, S., Tennyson, J.. Spectroscopic properties of the H₃⁺ molecule: a new calculated linelist. Astrophys J 1996;464:516–520.
- [178] Barber, R.J., Tennyson, J., Harris, G.J., Tolchenov, R.N.. A high accuracy computed water line list. Mon Not R Astron Soc 2006;368:1087–1094.
- [179] Yurchenko, S.N., Barber, R.J., Tennyson, J.. A variationally computed hot line list for NH₃. Mon Not R Astron Soc 2011;413:1828–1834. doi:10.1111/j.1365-2966.2011.18261.x.
- [180] Engel, E.A., Doss, N., Harris, G.J., Tennyson, J.. An opacity for HeH⁺ and its effect on models of cool, metal poor stars. Mon Not R Astron Soc 2005;357:471–477.
- [181] Zak, E.J., Tennyson, J., Polyansky, O.L., Lodi, L., Tashkun, S.A., Perevalov, V.I.. A room temperature CO₂ line list with *ab initio* computed intensities. J Quant Spectrosc Radiat Transf 2016;177:31–42. doi:10.1016/j.jqsrt.2015.12.022.
- [182] Zak, E.J., Tennyson, J., Polyansky, O.L., Lodi, L., Zobov, N.F., Tashkun, S.A., et al.

- Room temperature line lists for CO₂ symmetric isotopologues with *ab initio* computed intensities. J Quant Spectrosc Radiat Transf 2017;189:267 – 280. doi:10.1016/j.jqsrt.2016.11.022.
- [183] Zak, E.J., Tennyson, J., Polyansky, O.L., Lodi, L., Zobov, N.F., Tashkun, S.A., et al. Room temperature line lists for CO₂ asymmetric isotopologues with *ab initio* computed intensities. J Quant Spectrosc Radiat Transf 2017;203:265–281. doi:10.1016/j.jqsrt.2017.01.037.
 - [184] Tashkun, S.A., Perevalov, V.I.. CDSD-4000: High-resolution, high-temperature carbon dioxide spectroscopic databank. J Quant Spectrosc Radiat Transf 2011;112:1403–1410. doi:10.1016/j.jqsrt.2011.03.005.
 - [185] Conway, E.K., Gordon, I.E., Tennyson, J., Polyansky, O.L., Yurchenko, S.N., Chance, K.. A semi-empirical potential energy surface and line list for H₂¹⁶O extending into the near-ultraviolet. Atmos Chem Phys 2020;.
 - [186] Medvedev, E.S., Meshkov, V.V., Stolyarov, A.V., Gordon, I.E.. Peculiarities of high-overtone transition probabilities in carbon monoxide. J Chem Phys 2015;143:154301. doi:10.1063/1.4933136.
 - [187] Le Roy, R.J.. Level: A computer program for solving the radial schrödinger equation for bound and quasibound levels. J Quant Spectrosc Radiat Transf 2017;186:167 – 178. doi:10.1016/j.jqsrt.2016.05.028.
 - [188] Al Derzi, A.R., Furtenbacher, T., Yurchenko, S.N., Tennyson, J., Császár, A.G.. MARVEL analysis of the measured high-resolution spectra of ¹⁴NH₃. J Quant Spectrosc Radiat Transf 2015;161:117–130. doi:10.1016/j.jqsrt.2015.03.034.
 - [189] Medvedev, E.S., Ushakov, V.G., Conway, E.K., Upadhyay, A., Gordon, I.E., Tennyson, J.. Empirical normal intensity distribution law for overtone vibrational spectra of triatomic molecules. J Quant Spectrosc Radiat Transf 2020;252:107084. doi:10.1016/j.jqsrt.2020.107084.
 - [190] Furtenbacher, T., Császár, A.G., Tennyson, J.. MARVEL: measured active rotational-vibrational energy levels. J Mol Spectrosc 2007;245:115–125. doi:10.1016/j.jms.2007.07.005.
 - [191] von Essen, C., Mallonn, M., Welbanks, L., Madhusudhan, N., Pinhas, A., Bouy, H., et al. An optical transmission spectrum of the ultra-hot Jupiter WASP-33 b First indication of aluminum oxide in an exoplanet. Astron Astrophys 2019;622:A71. doi:10.1051/0004-6361/201833837.
 - [192] Chubb, K.L., Min, M., Kawashima, Y., Helling, C., Waldmann, I.. Aluminium oxide

- in the atmosphere of hot jupiter wasp-43b. 2020. [arXiv:2004.13679](#).
- [193] Furtenbacher, T., Szabó, I., Császár, A.G., Bernath, P.F., Yurchenko, S.N., Tennyson, J.. Experimental energy levels and partition function of the $^{12}\text{C}_2$ molecule. *Astrophys J Suppl* 2016;224:44. doi:[10.3847/0067-0049/224/2/44](#).
- [194] McKemmish, L.K., Syme, A.M., Borsovszky, J., Yurchenko, S.N., Tennyson, J., Furtenbacher, T., et al. Incorporating new experiments in diatomic spectral databases: An update to the $^{12}\text{C}_2$ MARVEL database and ExoMok line list of $^{12}\text{C}_2$. *Mon Not R Astron Soc* 2020;
- [195] Ram, R.S., Tereszchuk, K., Gordon, I.E., Walker, K.A., Bernath, P.F.. Fourier transform emission spectroscopy of the $E\ ^2\Pi - X\ ^2\Sigma^+$ transition of CaH and CaD. *J Mol Spectrosc* 2011;266:86–91. doi:[10.1016/j.jms.2011.03.009](#).
- [196] Hou, S., Wei, Z.. Line Lists for the $X\ ^1\Sigma^+$ State of CS. *Astrophys J Suppl* 2020;246:14. doi:[10.3847/1538-4365/ab61ef](#).
- [197] Xing, W., Shi, D., Sun, J.. Rovibrational transition properties of the $X\ ^1\Sigma^+$ and $A\ ^1\Pi$ states of carbon monosulfide. *Mol Phys* 2020;0:1–13. doi:[10.1080/00268976.2020.1759831](#).
- [198] Pavlenko, Y.V., Yurchenko, S.N., Tennyson, J.. Analysis of first overtone bands of isotopologues of CO and SiO in stellar spectra. *Astron Astrophys* 2020;633:A52.
- [199] McKemmish, L.K., Masseron, T., Sheppard, S., Sandeman, E., Schofield, Z., Furtenbacher, T., et al. MARVEL analysis of the measured high-resolution spectra of $^{48}\text{Ti}^{16}\text{O}$. *Astrophys J Suppl* 2017;228:15. doi:[10.3847/1538-4365/228/2/15](#).
- [200] Sedaghati, E., Boffin, H.M.J., MacDonald, R.J., Gandhi, S., Madhusudhan, N., Gibson, N.P., et al. Detection of titanium oxide in the atmosphere of a hot Jupiter. *Nature* 2017;549:238+. doi:[10.1038/nature23651](#).
- [201] Nugroho, S.K., Kawahara, H., Masuda, K., Hirano, T., Kotani, T., Tajitsu, A.. High-resolution Spectroscopic Detection of TiO and a Stratosphere in the Day-side of WASP-33b. *Astrophys J* 2017;154:221. doi:[10.3847/1538-3881/aa9433](#).
- [202] Plez, B.. A new TiO line list. *Astron Astrophys* 1998;337:495–500.
- [203] Schwenke, D.W.. Opacity of TiO from a coupled electronic state calculation parametrized by ab initio and experimental data. *Faraday Discuss* 1998;109:321–334. doi:[10.1039/a800070k](#).
- [204] McKemmish, L.K., Yurchenko, S.N., Tennyson, J.. Ab initio calculations to support accurate modelling of the rovibronic spectroscopy calculations of vanadium monoxide (VO). *Mol Phys* 2016;114:3232–3248. doi:[10.1080/00268976.2016.1225994](#).

- [205] Evans, T.M., Sing, D.K., Wakeford, H.R., Nikolov, N., Ballester, G.E., Drummond, B., et al. Detection of H₂O and evidence for tio/vo in an ultra-hot exoplanet atmosphere. *Astrophys J* 2016;822:L4. doi:10.3847/2041-8205/822/1/14.
- [206] Tsiaras, A., Waldmann, I.P., Zingales, T., Rocchetto, M., Morello, G., Damiano, M., et al. A population study of gaseous exoplanets. *Astron J* 2018;155:156. doi:10.3847/1538-3881/aaaf75.
- [207] Cheung, A.S.C., Hansen, R.C., Merer, A.J.. Laser spectroscopy of vo- analysis of the rotational and hyperfine structure of the c4-sigma-x4-sigma-(o, o) band. *J Mol Spectrosc* 1982;91:165–208. doi:10.1016/0022-2852(82)90039-X.
- [208] Polyansky, O.L., Bielska, K., Ghysels, M., Lodi, L., Zobov, N.F., Hodges, J.T., et al. High accuracy CO₂ line intensities determined from theory and experiment. *Phys Rev Lett* 2015;114:243001. doi:10.1103/PhysRevLett.114.243001.
- [209] Huang, X., Schwenke, D.W., Freedman, R.S., Lee, T.J.. Ames-2016 line lists for 13 isotopologues of co₂: Updates, consistency, and remaining issues. *J Quant Spectrosc Radiat Transf* 2017;203:224–241. doi:10.1016/j.jqsrt.2017.04.026.
- [210] Mellau, G.C.. Complete experimental rovibrational eigenenergies of HCN up to 6880 cm⁻¹ above the ground state. *J Chem Phys* 2011;134:234303.
- [211] Mellau, G.C.. Highly excited rovibrational states of HNC. *J Mol Spectrosc* 2011;269:77–85.
- [212] Hawker, G.A., Madhusudhan, N., Cabot, S.H.C., Gandhi, S.. Evidence for Multiple Molecular Species in the Hot Jupiter HD 209458b. *Astrophys J Lett* 2018;863:L11. doi:{10.3847/2041-8213/aac49d}.
- [213] Cabot, S.H.C., Madhusudhan, N., Hawker, G.A., Gandhi, S.. On the robustness of analysis techniques for molecular detections using high-resolution exoplanet spectroscopy. *Mon Not R Astron Soc* 2019;482:4422–4436. doi:10.1093/mnras/sty2994.
- [214] Gandhi, S., Brogi, M., Yurchenko, S.N., Tennyson, J., Coles, P.A., Webb, R.K., et al. Molecular Cross Sections for High Resolution Spectroscopy of super Earths, warm Neptunes and Hot Jupiters. *Mon Not R Astron Soc* 2020;495:224–237. doi:10.1093/mnras/staa981.
- [215] Tsiaras, A., Waldmann, I.P., Tinetti, G., Tennyson, J., Yurchenko, S.N.. Water vapour in the atmosphere of the habitable-zone eight-Earth-mass planet K2 18b. *Nature Astron* 2019;3:1086–1091. doi:10.1038/s41550-019-0878-9.
- [216] Partridge, H., Schwenke, D.W.. The determination of an accurate isotope dependent potential energy surface for water from extensive ab initio calculations and experimental

- data. J Chem Phys 1997;106:4618–4639. doi:10.1063/1.473987.
- [217] Tennyson, J., Bernath, P.F., Brown, L.R., Campargue, A., Carleer, M.R., Császár, A.G., et al. IUPAC critical evaluation of the rotational-vibrational spectra of water vapor. Part III. Energy levels and transition wavenumbers for H₂¹⁶O. J Quant Spectrosc Radiat Transf 2013;117:29–80. doi:10.1016/j.jqsrt.2012.10.002.
- [218] Lampel, J., Pöhler, D., Polyansky, O.L., Kyuberis, A.A., Zobov, N.F., Tennyson, J., et al. Detection of water vapour absorption around 363 nm in measured atmospheric absorption spectra and its effect on DOAS evaluations. Atmos Chem Phys 2017;17:1271–1295. doi:10.5194/acp-2016-388.
- [219] Lodi, L., Tennyson, J., Polyansky, O.L.. A global, high accuracy *ab initio* dipole moment surface for the electronic ground state of the water molecule. J Chem Phys 2011;135:034113. doi:10.1063/1.3604934.
- [220] Conway, E.K., Kyuberis, A.A., Polyansky, O.L., Tennyson, J., Zobov, N.. A highly accurate *ab initio* dipole moment surface for the ground electronic state of water vapour for spectra extending into the ultraviolet. J Chem Phys 2018;149:084307. doi:10.1063/1.5043545.
- [221] Conway, E.K., Gordon, I.E., Kyuberis, A.A., Polyansky, O.L., Tennyson, J., Zobov, N.F.. Accurate line lists for H₂¹⁶O, H₂¹⁸O and H₂¹⁷O with extensive comparisons to theoretical and experimental sources including the HITRAN2016 database. J Quant Spectrosc Radiat Transf 2020;241:106711. doi:10.1016/j.jqsrt.2019.106711.
- [222] Conway, E.K., Gordon, I.E., Polyansky, O.L., Tennyson, J.. Use of the complete basis set limit for computing highly accurate *ab initio* dipole moments. J Chem Phys 2020;152:024105. doi:10.1063/1.5135931.
- [223] Tennyson, J., Bernath, P.F., Brown, L.R., Campargue, A., Carleer, M.R., Császár, A.G., et al. IUPAC critical Evaluation of the Rotational-Vibrational Spectra of Water Vapor. Part I. Energy Levels and Transition Wavenumbers for H₂¹⁷O and H₂¹⁸O. J Quant Spectrosc Radiat Transf 2009;110:573–596. doi:10.1016/j.jqsrt.2009.02.014.
- [224] Tennyson, J., Bernath, P.F., Brown, L.R., Campargue, A., Carleer, M.R., Császár, A.G., et al. IUPAC critical Evaluation of the Rotational-Vibrational Spectra of Water Vapor. Part II. Energy Levels and Transition Wavenumbers for HD¹⁶O, HD¹⁷O, and HD¹⁸O. J Quant Spectrosc Radiat Transf 2010;111:2160–2184. doi:10.1016/j.jqsrt.2010.06.012.
- [225] Zobov, N.F., Koshelev, M.A., Makaro, D.S., Boyarkin, O.V., Tyuterev, V.G., Tennyson, J., et al. Global study of the HDO spectrum - analysis of experiment and high

- temperature line list. *J Quant Spectrosc Radiat Transf* 2020;In Preparation.
- [226] Campargue, A., Kassi, S., Yachmenev, A., Kyuberis, A.A., Küpper, J., Yurchenko, S.N.. Observation of electric-quadrupole infrared transitions in water vapor. *Phys Rev Research* 2020;2:023091. URL: <https://link.aps.org/doi/10.1103/PhysRevResearch.2.023091>. doi:10.1103/PhysRevResearch.2.023091.
- [227] Mizus, I.I., Kyuberis, A.A., Zobov, N.F., Makhnev, V.Y., Polyansky, O.L., Tennyson, J.. High accuracy water potential energy surface for the calculation of infrared spectra. *Phil Trans Royal Soc London A* 2018;376:20170149. doi:10.1098/rsta.2017.0149.
- [228] Yurchenko, S.N., Thiel, W., Jensen, P.. Theoretical ROVibrational Energies (TROVE): A robust numerical approach to the calculation of rovibrational energies for polyatomic molecules. *J Mol Spectrosc* 2007;245:126–140. doi:10.1016/j.jms.2007.07.009.
- [229] Gordon, I.E., Rothman, L.S., Hill, C., Kochanov, R.V., Tan, Y., Bernath, P.F., et al. The *HITRAN* 2016 molecular spectroscopic database. *J Quant Spectrosc Radiat Transf* 2017;203:3–69. doi:10.1016/j.jqsrt.2017.06.038.
- [230] Chubb, K.L., Naumenko, O.V., Keely, S., Bartolotto, S., MacDonald, S., Mukhtar, M., et al. MARVEL analysis of the measured high-resolution rovibrational spectra of H₂S. *J Quant Spectrosc Radiat Transf* 2018;218:178 – 186. doi:10.1016/j.jqsrt.2018.07.012.
- [231] Furtenbacher, T., Szidarovszky, T., Fábri, C., Császár, A.G.. MARVEL analysis of the rotational-vibrational states of the molecular ions H₂D⁺ and D₂H⁺. *Phys Chem Chem Phys* 2013;15:10181–10193. doi:10.1039/c3cp44610g.
- [232] Petrignani, A., Berg, M., Wolf, A., Mizus, I.I., Polyansky, O.L., Tennyson, J., et al. Visible intensities of the triatomic hydrogen ion from experiment and theory. *J Chem Phys* 2014;141:241104. doi:10.1063/1.4904440.
- [233] Miller, S., Geballe, T.R., Stallard, T., Tennyson, J.. Thirty years of H₃⁺ astronomy. *Rev Mod Phys* 2020;.
- [234] Tóbiás, R., Furtenbacher, T., Császár, A.G., Naumenko, O.V., Tennyson, J., Flaud, J.M., et al. Critical Evaluation of Measured Rotational-Vibrational Transitions of Four Sulphur Isotopologues of S¹⁶O₂. *J Quant Spectrosc Radiat Transf* 2018;208:152–163. doi:10.1016/j.jqsrt.2018.01.006.
- [235] Chubb, K.L., Joseph, M., Franklin, J., Choudhury, N., Furtenbacher, T., Császár, A.G., et al. MARVEL analysis of the measured high-resolution spectra of C₂H₂. *J Quant Spectrosc Radiat Transf* 2018;204:42–55. doi:10.1016/j.jqsrt.2017.08.018.
- [236] Al-Derzi, A.R., Furtenbacher, T., Yurchenko, S.N., Tennyson, J., Császár, A.G.. MARVEL analysis of the measured high-resolution spectra of formaldehyde. *J Quant Spectrosc Radiat Transf* 2018;204:42–55. doi:10.1016/j.jqsrt.2017.08.018.

Spectrosc Radiat Transf 2020;To be submitted.

- [237] Al-Refaie, A.F., Ovsyannikov, R.I., Polyansky, O.L., Yurchenko, S.N., Tennyson, J.. A variationally calculated room temperature line-list for H₂O₂. *J Mol Spectrosc* 2015;318:84–90. doi:10.1016/j.jms.2015.10.004.
- [238] Ito, Y., Hashimoto, G.L., Takahashi, Y.O., Ishiwatari, M., Kuramoto, K.. H₂O₂-induced Greenhouse Warming on Oxidized Early Mars. *Astrophys J* 2020;893:168. doi:10.3847/1538-4357/ab7db4.
- [239] Bourgalais, J., Carrasco, N., Changeat, Q., Venot, O., Javanović, L., Pernot, P., et al. Ions in the thermosphere of exoplanets: Observational constraints revealed by innovative laboratory experiments. *Astrophys J* 2020;895:77. doi:10.3847/1538-4357/ab8e2d.
- [240] Sousa-Silva, C., Seager, S., Ranjan, S., Petkowski, J.J., Zhan, Z., Hu, R., et al. Phosphine as a biosignature gas in exoplanet atmospheres. *Astrobiology* 2020;20:235–268. doi:10.1089/ast.2018.1954.
- [241] Rey, M., Nikitin, A.V., Tyuterev, V.G.. Theoretical hot methane line list up T=2000 K for astrophysical applications. *Astrophys J* 2014;789:2. doi:10.1088/0004-637X/789/1/2.
- [242] Hargreaves, R.J., Bernath, P.F., Bailey, J., Dulick, M.. Empirical line lists and absorption cross sections for methane at high temperatures. *Astrophys J* 2015;813:12.
- [243] Rey, M., Nikitin, A.V., Tyuterev, V.G.. Accurate theoretical methane line lists in the infrared up to 3000 K and quasi-continuum absorption/emission modeling for astrophysical applications. *Astrophys J* 2017;847:105.
- [244] Li, G., Gordon, I.E., Hajigeorgiou, P.G., Coxon, J.A., Rothman, L.S.. Reference spectroscopic data for hydrogen halides, Part II: The line lists. *J Quant Spectrosc Radiat Transf* 2013;130:284–295. doi:10.1016/j.jqsrt.2013.07.019.
- [245] R. R. Gamache et al., . Total Internal Partition Sums for 167 isotopologues of 53 molecules important in planetary atmospheres: application to HITRAN2016 and beyond. *J Quant Spectrosc Radiat Transf* 2017;203:70–87. doi:10.1016/j.jqsrt.2017.03.045.
- [246] Vallon, R., Richard, C., Crozet, P., Wannous, G., Ross, A.. Laboratory measurements of nih by fourier transform dispersed fluorescence. *Astrophys J* 2009;696:172–175. doi:10.1088/0004-637X/696/1/172.
- [247] Harker, H., Richard, C., Tourasse, G., Crozet, P., Ross, A.J.. Zeeman spectroscopy of NiH: Lande factors of three Omega=3/2 excited electronic states. *J Mol Spectrosc* 2013;292:28–34. doi:{10.1016/j.jms.2013.09.005}.
- [248] Šimečková, M., Jacquemart, D., Rothman, L.S., Gamache, R.R., Goldman, A.. Einstein A coefficients and statistical weights for molecular absorption transitions in the

- HITRAN database. *J Quant Spectrosc Radiat Transf* 2006;98:130–155.
- [249] Irwin, A.W.. Polynomial partition function approximations of 344 atomic and molecular species. *Astrophys J Suppl* 1981;45:621–633. doi:10.1086/190730.
 - [250] Sauval, A.J., Tatum, J.B.. A set of partition functions and equilibrium constants for 300 diatomic molecules of astrophysical interest. *Astrophys J Suppl* 1984;56:193–209.
 - [251] Irwin, A.W.. The partition functions of JANAF polyatomic molecules that significantly affect the stellar atmospheric equation of state. *Astron Astrophys Suppl* 1988;74:145–160.
 - [252] Barklem, P.S., Collet, R.. Partition functions and equilibrium constants for diatomic molecules and atoms of astrophysical interest. *Astron Astrophys* 2016;588:A96. doi:10.1051/0004-6361/201526961.
 - [253] Sousa-Silva, C., Hesketh, N., Yurchenko, S.N., Hill, C., Tennyson, J.. High Temperature partition functions and thermodynamic data for ammonia and phosphine. *J Quant Spectrosc Radiat Transf* 2014;142:66–74. doi:10.1016/j.jqsrt2014.03.012.
 - [254] Ranjan, S., Schwieterman, E.W., Harman, C., Fateev, A., Sousa-Silva, C., Seager, S., et al. Photochemistry of Anoxic Abiotic Habitable Planet Atmospheres: Impact of New H₂O Cross-Sections. 2020. arXiv:2004.04185.
 - [255] Fateev, A., Clausen, S., Wang, Y., Yurchenko, S.N., Tennyson, J.. Experimental vuv cross sections in the exomol database. *J Quant Spectrosc Radiat Transf* 2020;In preparation.
 - [256] Venot, O., Bénilan, Y., Fray, N., Gazeau, M.C., Lefèvre, F., Es-sebbar, E., et al. VUV-absorption cross section of carbon dioxide from 150 to 800 K and applications to warm exoplanetary atmospheres. *Astron Astrophys* 2018;609:A34. doi:10.1051/0004-6361/201731295.
 - [257] Tennyson, J., Hill, C., Yurchenko, S.N.. Data structures for ExoMol: Molecular line lists for exoplanet and other atmospheres. In: 6th international conference on atomic and molecular data and their applications ICAMDATA-2012; vol. 1545 of *AIP Conference Proceedings*. AIP, New York; 2013, p. 186–195. doi:10.1063/1.4815853.
 - [258] Western, C.M.. PGOPHER: A program for simulating rotational, vibrational and electronic spectra . *J Quant Spectrosc Radiat Transf* 2017;186:221–242. doi:10.1016/j.jqsrt.2016.04.010.
 - [259] Furtenbacher, T., Szidarovszky, T., Hruby, J., Kyuberis, A.A., Zobov, N.F., Polyansky, O.L., et al. Definitive high-temperature ideal-gas thermochemical functions of the H₂¹⁶O molecule. *J Phys Chem Ref Data* 2016;45:043104. doi:10.1063/1.4967723.
 - [260] Barton, E.J., Hill, C., Czurylo, M., Li, H.Y., Hyslop, A., Yurchenko, S.N., et al. The

- ExoMol diet of line-by-line pressure-broadening parameters. *J Quant Spectrosc Radiat Transf* 2017;203:490–495. doi:10.1016/j.jqsrt.2017.01.028.
- [261] Fortney, J.J., Robinson, T.D., Domagal-Goldman, S., Del Genio, A.D., Gordon, I.E., Gharib-Nezhad, E., et al. The need for laboratory measurements and ab initio studies to aid understanding of exoplanetary atmospheres. 2019. ArXiv:1905.07064.
 - [262] Chubb, K.L., Rocchetto, M., Yurchenko, S.N., Min, M., Waldmann, I., Barstow, J.K., et al. The ExoMolOP Database: Cross-sections and K-tables for Molecules of Interest in High-Temperature Exoplanet Atmospheres. *Astron Astrophys* 2020;.
 - [263] Dubernet, M.L., Boudon, V., Culhane, J.L., Dimitrijevic, M.S., Fazliev, A.Z., Joblin, C., et al. Virtual Atomic and Molecular Data Centre. *J Quant Spectrosc Radiat Transf* 2010;111:2151–2159.
 - [264] Dubernet, M.L., Antony, B.K., Ba1, Y.A., Babikov, Y.L., Bartschat, K., Boudon, V., et al. The Virtual Atomic and Molecular Data Centre (VAMDC) Consortium for Astrophysics. *J Phys B: At Mol Opt Phys* 2016;49:074003. doi:10.1088/0953-4075/49/7/074003.
 - [265] Gharib-Nezhad, E., Iyer, A.R., Line, M.R., Freedman, R.S., Marley, M.S., Batalha, N.E.. EXOPLINES: Molecular Absorption Cross-Section Database for Brown Dwarf and Giant Exoplanet Atmospheres. *Astrophys J Suppl* 2020;.
 - [266] Yurchenko, S.N., Al-Refaie, A.F., Tennyson, J.. ExoCross: A general program for generating spectra from molecular line lists. *Astron Astrophys* 2018;614:A131. doi:10.1051/0004-6361/201732531.
 - [267] Rothman, L.S., Gordon, I.E., Babikov, Y., Barbe, A., Benner, D.C., Bernath, P.F., et al. The HITRAN 2012 molecular spectroscopic database. *J Quant Spectrosc Radiat Transf* 2013;130:4 – 50. doi:10.1016/j.jqsrt.2013.07.002.
 - [268] Tennyson, J., Bernath, P.F., Brown, L.R., Campargue, A., Császár, A.G., Daumont, L., et al. IUPAC critical evaluation of the rotational-vibrational spectra of water vapor. Part IV. Energy levels and transition wavenumbers for $D_2^{16}O$, $D_2^{17}O$ and $D_2^{18}O$. *J Quant Spectrosc Radiat Transf* 2014;142:93–108. doi:10.1016/j.jqsrt.2014.03.019.
 - [269] Tóbiás, R., Furtenbacher, T., Tennyson, J., Császár, A.G.. Accurate empirical rovibrational energies and transitions of $H_2^{16}O$. *Phys Chem Chem Phys* 2019;21:3473–3495. doi:10.1039/c8cp05169k.
 - [270] Furtenbacher, T., Tóbiás, R., Tennyson, J., Polyansky, O.L., Császár, A.G.. W2020: A Database of Validated Rovibrational Experimental Transitions and Empirical Energy Levels of $H_2^{16}O$. *J Phys Chem Ref Data* 2020;49:033101.

- [271] Furtenbacher, T., Szidarovszky, T., Mátyus Edit Fábi, C., Császár, A.G.. Analysis of the Rotational-Vibrational States of the Molecular Ion H₃⁺. *J Chem Theory Comput* 2013;9:5471–5478. doi:10.1021/ct4004355.
- [272] Furtenbacher, T., Coles, P.A., Tennyson, J., Yurchenko, S.N., Yu, S., Drouin, B., et al. Empirical rovibational energy of ammonia up to 7500 cm⁻¹. *J Quant Spectrosc Radiat Transf* 2020;251:107027. doi:10.1016/j.jqsrt.2020.107027.
- [273] McKemmish, L.K., Borsovszky, J., Goodhew, K.L., Sheppard, S., Bennett, A.F.V., Martin, A.D.J., et al. MARVEL analysis of the measured high-resolution spectra of ⁹⁰Zr¹⁶O. *Astrophys J* 2018;867:33. doi:10.3847/1538-4357/aadd19.
- [274] Furtenbacher, T., Horváth, M., Koller, D., Sólyom, P., Balogh, A., Balogh, I., et al. MARVEL analysis of the measured high-resolution rovibronic spectra and definitive ideal-gas thermochemistry of the ¹⁶O₂ molecule. *J Phys Chem Ref Data* 2019;48:023101. doi:10.1063/1.5083135.
- [275] Wang, Y., Owens, A., Tennyson, J., Yurchenko, S.N.. MARVEL analysis of the measured high-resolution rovibronic spectra of the calcium monohydroxide radical (CaOH). *Astrophys J Suppl* 2020;248:9. doi:10.3847/1538-4365/ab85cb.
- [276] Syme, A.M., McKemmish, L.K.. Experimental energy levels of ¹²C¹⁴N through Marvel analysis. *Mon Not R Astron Soc* 2020;
- [277] de Kok, R.J., Brogi, M., Snellen, I.A.G., Birkby, J., Albrecht, S., de Mooij, E.J.W.. Detection of carbon monoxide in the high-resolution day-side spectrum of the exoplanet HD 189733b. *Astron Astrophys* 2013;554:A82. doi:10.1051/0004-6361/201321381.
- [278] Birkby, J.L., de Kok, R.J., Brogi, M., de Mooij, E.J.W., Schwarz, H., Albrecht, S., et al. Detection of water absorption in the day side atmosphere of HD 189733 b using ground-based high-resolution spectroscopy at 3.2 μm. *Mon Not R Astron Soc* 2013;436:L35–L39. doi:10.1093/mnrasl/slt107.
- [279] Brogi, M., de Kok, R.J., Birkby, J.L., Schwarz, H., Snellen, I.A.G.. Carbon monoxide and water vapor in the atmosphere of the non-transiting exoplanet HD 179949 b. *Astron Astrophys* 2014;565:A124. doi:10.1051/0004-6361/201423537.
- [280] Tennyson, J.. The ExoMol project: Molecular opacity calculations at University College London. In: Mendoza, C., Turck-Chiéze, S., Colgan, J., editors. *Workshop on Astrophysical Opacities*; vol. 515 of *Astron. Soc. Pac. Conf. Ser.* Astronomical Society of the Pacific; 2018, p. 137–144.
- [281] Furtenbacher, T., Császár, A.G.. MARVEL: measured active rotational-vibrational energy levels. II. Algorithmic improvements. *J Quant Spectrosc Radiat Transf* 2012;113:929–

935.

- [282] Darby-Lewis, D., Shah, H., Joshi, D., Khan, F., Kauwo, M., Sethi, N., et al. MARVEL analysis of the measured high-resolution spectra of NH. *J Mol Spectrosc* 2019;362:69–76. doi:10.1016/j.jms.2019.06.002.
- [283] Sousa-Silva, C., McKemmish, L.K., Chubb, K.L., Baker, J., Barton, E.J., Gorman, M.N., et al. Original Research By Young Twinkle Students (ORBYTS): When can students start performing original research? *Phys Educ* 2018;53:015020. doi:10.1088/1361-6552/aa8f2a.
- [284] Skinner, F.M., Gordon, I.E., Hill, C., Hargreaves, R.J., Lockhart, K.E., Rothman, L.S.. Referencing Sources of Molecular Spectroscopic Data in the Era of Data Science: Application to the HITRAN and AMBDAS Databases. *Atoms* 2020;8:16. doi:10.3390/atoms8020016.