

Highly linked tournaments

Alexey Pokrovskiy

Methods for Discrete Structures, Freie Universität, Berlin, Germany

Abstract

A (possibly directed) graph is k -linked if for any two disjoint sets of vertices $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ there are vertex disjoint paths P_1, \dots, P_k such that P_i goes from x_i to y_i . A theorem of Bollobás and Thomason says that every $22k$ -connected (undirected) graph is k -linked. It is desirable to obtain analogues for directed graphs as well. Although Thomassen showed that the Bollobás-Thomason Theorem does not hold for general directed graphs, he proved an analogue of the theorem for tournaments—there is a function $f(k)$ such that every strongly $f(k)$ -connected tournament is k -linked. The bound on $f(k)$ was reduced to $O(k \log k)$ by Kühn, Lapinskas, Osthus, and Patel, who also conjectured that a linear bound should hold. We prove this conjecture, by showing that every strongly $452k$ -connected tournament is k -linked.

Keywords: Connectivity of tournaments, linkedness, linkage structures.

1. Introduction

A graph is connected if there is a path between any two vertices. A graph is k -connected if it remains connected after the removal of any set of $(k-1)$ -vertices. This could be seen as a notion of how robust the graph is. For example, if the graph represents a communication network, then the connectedness measures how many nodes need to fail before communication becomes impossible.

Similar notions make sense for directed graphs, except in that context we usually want a *directed path* between every pair of vertices. If this holds, we say that the directed graph is *strongly connected*. A directed graph is strongly k -connected if it remains strongly connected after the removal of any set of $(k-1)$ -vertices. In this paper, when dealing with connectedness of directed graphs we will always mean strong connectedness.

Connectedness is a fundamental notion in graph theory, and there are countless theorems which involve it. Perhaps the most important of these is Menger's Theorem, which provides an alternative characterization of k -connectedness. Menger's Theorem says that a graph is k -connected if, and only if, there are k internally vertex-disjoint paths between any pair of vertices. Menger's Theorem has the following simple corollary:

Corollary 1.1. *If G is k -connected then for any two disjoint sets of vertices $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ there are vertex-disjoint paths P_1, \dots, P_k such that P_i goes from x_i to $y_{\sigma(i)}$ for some permutation σ of $[k]$.*

This corollary is proved by constructing a new graph H from G by adding two vertices x and y such that x is joined to $\{x_1, \dots, x_k\}$ and y is joined to $\{y_1, \dots, y_k\}$. It is easy to see that H is k -connected, and so, by Menger's Theorem, has k vertex-disjoint $x-y$ paths. Removing the vertices x and y produces the required paths P_1, \dots, P_k . It is not hard to see that the converse of Corollary 1.1 holds for graphs on at least $2k$ vertices as well.

Notice that in Corollary 1.1, we had no control over where the path P_i starting at x_i ends up—it could end at any of the vertices y_1, \dots, y_k . In practice we might want to have control over

Email address: alja123@gmail.com (Alexey Pokrovskiy)

this. This leads to the notion of k -linkedness. A graph is k -linked if for any two disjoint sets of vertices $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ there are vertex disjoint paths P_1, \dots, P_k such that P_i goes from x_i to y_i .

Linkedness is a stronger notion than connectedness. A natural question is whether a k -connected graph must also be ℓ -linked for some ℓ (which may be smaller than k). Larman and Mani [8], and Jung [4] were the first to show that this is indeed the case—they showed that there is a function $f(k)$ such that every $f(k)$ -connected graph is k -linked. This result uses a theorem of Mader [9] about the existence of large topological complete minors in graphs with many edges. The first bounds on $f(k)$ were exponential in k , but Bollobás and Thomason showed that a linear bound on the connectedness suffices [3].

Theorem 1.2 (Bollobás and Thomason). *Every $22k$ -connected graph is k -linked.*

The constant 22 has since been reduced to 10 by Thomas and Wollan [11].

Much of the above discussion holds true for directed graphs as well (when talking about *strong* k -connectedness and *directed* paths). Menger’s Theorem remains true, as does Corollary 1.1. A directed graph is k -linked if for two disjoint sets of vertices $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ there are vertex disjoint directed paths P_1, \dots, P_k such that P_i goes from x_i to y_i . Somewhat surprisingly there is no function $f(k)$ such that every strongly $f(k)$ -connected directed graph is k -linked. Indeed Thomassen constructed directed graphs of arbitrarily high connectedness which are not even 2-linked [13]. Thus, there is a real difference between the directed and undirected cases. For tournaments however the situation is better (A tournament is a directed graph which has exactly one directed edge between any two vertices). There Thomassen showed that there is a constant C , such that every Ck -connected tournament is k -linked [12]. Kühn, Lapinskas, Osthus, and Patel improved the bound on the connectivity to $10^4 k \log k$.

Theorem 1.3 (Kühn, Lapinskas, Osthus, and Patel, [6]). *All strongly $10^4 k \log k$ -connected tournaments are k -linked.*

This theorem is proved using a beautiful construction utilizing the asymptotically optimal sorting networks of Ajtai, Komlós, and Szemerédi [1]. The proof is based on building a small sorting network inside the tournament, which is combined with the directed version of Corollary 1.1 in order to reorder the endpoints of the paths so that P_i goes from x_i to y_i . We refer to [6] for details.

Since sorting networks on k inputs require size at least $k \log k$, it is unlikely that this approach can give a $o(k \log k)$ bound in Theorem 1.3. Nevertheless, Kühn, Lapinskas, Osthus, and Patel conjectured that a linear bound should be possible.

Conjecture 1.4 (Kühn, Lapinskas, Osthus, and Patel, [6]). *There is a constant C such that every strongly Ck -connected tournament is k -linked.*

There has also been some work for small k . Bang-Jensen showed that every 5-connected tournament is 2-linked [2]. Here the value “5” is optimal.

The main result of this paper is a proof of Conjecture 1.4.

Theorem 1.5. *Every strongly $452k$ -connected tournament is k -linked.*

The above theorem is proved using the method of “linkage structures in tournaments” recently introduced in [6] and [7]. Informally, a linkage structure L in a tournament T , is a small subset of $V(T)$ with the property that for many pairs of vertices x, y outside L , there is a path from x to y most of whose vertices are contained in L . Such structures can be found in highly connected tournaments, and they have various applications such as finding Hamiltonian cycles [6, 10] or partitioning tournaments into highly connected subgraphs [5, 7]. Linkage structures were introduced in the same paper where Conjecture 1.4 was made. However in the past they were constructed using Theorem 1.3 to first show that a tournament is highly linked. In our paper the perspective is different—the linkage structures are built using only connectedness, and then linkedness follows as a corollary of the presence of the linkage structures.

In the next section we prove the above theorem. In Section 3, we make some concluding remarks and pose an open problem.

2. Proof of Theorem 1.5

A directed path P is a sequence of vertices v_1, v_2, \dots, v_k in a directed graph such that $v_i v_{i+1}$ is an edge for all $i = 1, \dots, k-1$. The vertex v_1 is called the *start* of P , and v_k the *end* of P . The *length* of P is the number of edges it has which is $|P| - 1$. The vertices v_2, \dots, v_{k-1} are the *internal vertices* of P . Two paths are said to be internally disjoint if their internal vertices are distinct.

A tournament T is transitive if for any three vertices $x, y, z \in V(T)$, if xy and yz are both edges, then xz is also an edge. It is easy to see that a tournament is transitive exactly when it has an ordering (v_1, v_2, \dots, v_k) of $V(T)$ such that the edges of T are $\{v_i v_j : i < j\}$. We say that v_1 is the *tail* of T , and v_k is the *head* of T .

The *out-neighbourhood* of a vertex v in a directed graph, denoted $N^+(v)$, is the set of vertices u for which vu is an edge. Similarly, the *in-neighbourhood*, denoted $N^-(v)$, is the set of vertices u for which uv is an edge. The *out-degree* of v is $d^+(v) = |N^+(v)|$, and the *in-degree* of v is $d^-(v) = |N^-(v)|$. A useful fact is that every tournament, T , has a vertex of out-degree at least $(|T| - 1)/2$, and a vertex of in-degree at least $(|T| - 1)/2$. To see this, observe that since T has $\binom{|T|}{2}$ edges, its average in and out-degrees are both $(|T| - 1)/2$.

We will need the following lemma which says that in any tournament, we can find two large sets such that there is a linkage between them.

Lemma 2.1. *Let n and m be two integers with $m \leq n/11$. Every tournament T on n vertices, contains two disjoint sets of vertices $\{x_1, \dots, x_m\}$ and $\{y_1, \dots, y_m\}$ such that for any permutation σ of $[m]$, there are vertex-disjoint paths P_1, \dots, P_m such that P_i goes from x_i to $y_{\sigma(i)}$.*

Proof. Let x_1, \dots, x_m be a set of m vertices in T of largest out-degrees i.e. any set such that any vertex u outside it satisfies $d^+(u) \leq d^+(x_i)$ for all i . Let y_1, \dots, y_m be a set of m vertices in T of largest in-degrees. Since $m \leq n/11$, we can choose $\{x_1, \dots, x_m\}$ and $\{y_1, \dots, y_m\}$ to be disjoint. Indeed if $I = \{x_1, \dots, x_m\} \cap \{y_1, \dots, y_m\}$ is not empty then using the properties of the x_i s and y_i s we have $d^+(z) = d^+(x_i)$ for any $x_i \in I$. Therefore for any vertices $z_1, \dots, z_{m-|I|}$ not in $\{x_1, \dots, x_m\} \cup \{y_1, \dots, y_m\}$, the set $\{x_1, \dots, x_m\} \setminus I + z_1 + \dots + z_{m-|I|}$ is a set of largest out-degree vertices which is disjoint from $\{y_1, \dots, y_m\}$.

Recall that every tournament T has a vertex of out-degree at least $(|T| - 1)/2$. This means that $d^+(x_i) \geq (n - m)/2$ for each $i = 1, \dots, m$ (since otherwise, there would be a vertex in $T \setminus \{x_1, \dots, x_m\} + x_i$ of out-degree larger than x_i , contradicting the choice of x_i). Similarly, we obtain $d^-(y_i) \geq (n - m)/2$ for each $i = 1, \dots, m$.

For each i and $j \leq m$, let $X_{i,j} = N^+(x_i) \setminus (N^-(y_j) + y_j) + x_i$, $Y_{i,j} = N^-(y_j) \setminus (N^+(x_i) + x_i) + y_j$, $I_{i,j} = N^+(x_i) \cap N^-(y_j)$, and $M_{i,j}$ a maximum matching of edges directed from $X_{i,j}$ to $Y_{i,j}$.

Notice that we have

$$|X_{i,j} \setminus V(M_{i,j})| = |N^+(x_i) + x_i - y_i| - |I_{i,j}| - e(M_{i,j}) \geq \frac{1}{2}(n - m) - |I_{i,j}| - e(M_{i,j}).$$

Similarly, we obtain $|Y_{i,j} \setminus V(M_{i,j})| \geq \frac{1}{2}(n - m) - |I_{i,j}| - e(M_{i,j})$.

Since M is maximal, all the edges between $X_{i,j} \setminus V(M_{i,j})$ and $Y_{i,j} \setminus V(M_{i,j})$ go from $Y_{i,j}$ to $X_{i,j}$. Therefore, if $\frac{1}{2}(n - m) - |I_{i,j}| - e(M_{i,j}) \geq m$ holds, then the lemma follows by choosing x'_1, \dots, x'_m to be any m vertices in $Y_{i,j} \setminus V(M_{i,j})$, and y'_1, \dots, y'_m to be any m vertices in $X_{i,j} \setminus V(M_{i,j})$. This ensures that we can always choose length 1 paths P_1, \dots, P_m as in the lemma.

Therefore, we can suppose that $\frac{1}{2}(n - m) - |I_{i,j}| - e(M_{i,j}) < m$. Combining this with $m \leq n/11$ we obtain that $|I_{i,j}| + e(M_{i,j}) > 4m$ for every i and j .

Notice that for all i and j , there are exactly $|I_{i,j}| + e(M_{i,j}) \geq 4m + 1$ internally vertex disjoint paths of length ≤ 3 between x_i and y_j . This allows us to construct vertex disjoint paths P_1, \dots, P_m each of length ≤ 3 , such that P_i goes from x_i to $y_{\sigma(i)}$ (where σ is an arbitrary permutation of $[m]$). Indeed, assuming we have constructed the paths P_1, \dots, P_k , then we have $|V(P_1) \cup \dots \cup V(P_k) \cup \{x_{k+2}, \dots, x_m\} \cup \{y_{\sigma(k+2)}, \dots, y_{\sigma(m)}\}| \leq 4m$, and so one of the $4m + 1$ internally vertex disjoint paths between x_{k+1} and $y_{\sigma(k+1)}$ must be disjoint from $V(P_1) \cup \dots \cup V(P_k) \cup \{x_{k+2}, \dots, x_m\} \cup \{y_{\sigma(k+2)}, \dots, y_{\sigma(m)}\}$. We let P_{k+1} be this path, and then repeat this process until we have the required m paths. \square

A set of vertices S in-dominates another set B , if for every $b \in B \setminus S$, there is some $s \in S$ such that bs is an edge. Notice that by this definition, a set in-dominates itself. An *in-dominating set* in a tournament T is any set S which in-dominates $V(T)$. Notice that by repeatedly pulling out vertices of largest in-degree and their in-neighbourhoods from T , we can find an in-dominating set of order at most $\lceil \log_2 |T| \rceil$. For our purposes we will study sets which are constructed by pulling out some fixed number of vertices by this process.

Definition 2.2. We say that a sequence (v_1, v_2, \dots, v_k) of vertices of a tournament T is a *partial greedy in-dominating set* if v_1 is a maximum in-degree vertex in T , and for each i , v_i is a maximum in-degree vertex in the subtournament of T on $N^+(v_1) \cap N^+(v_2) \cap \dots \cap N^+(v_{i-1})$.

Partial greedy out-dominating sets are defined similarly, by letting v_i be a maximum out-degree vertex in $N^-(v_1) \cap N^-(v_2) \cap \dots \cap N^-(v_{i-1})$ at each step.

Notice that every partial greedy in-dominating set is a transitive tournament with head v_k and tail v_1 .

For small k , partial greedy in-dominating sets do not necessarily dominate all the vertices in a tournament. A crucial property of partial greedy in-dominating sets is that the vertices they do not dominate have large out-degree. The following is a version of a lemma appearing in [6].

Lemma 2.3. Let (v_1, v_2, \dots, v_k) be a partial greedy in-dominating set in a tournament T . Let E be the set of vertices which are not in-dominated by A . Then every $u \in E$ satisfies $d^+(u) \geq 2^{k-1}|E|$.

Proof. The proof is by induction on k . The initial case is when $k = 1$. In this case we have $E = N^+(v_1)$ where v_1 is a maximum in-degree vertex in T . For any $u \in E$, we must have $d^-(u) \leq d^-(v_1) = |T \setminus E - v_1| = |T| - |E| - 1$. Therefore we have $d^+(u) = |T \setminus N^-(u) - u| = |T| - d^-(u) - 1 \geq |E|$ as required.

Now suppose that the lemma holds for $k = k_0$. Let (v_1, \dots, v_{k_0+1}) be a partial greedy in-dominating set in T , and let $E_0 = N^+(v_1) \cap \dots \cap N^+(v_{k_0})$. By induction we have $d^+(u) \geq 2^{k_0-1}|E_0|$ for every $u \in E_0$. By definition v_{k_0+1} is a maximum in-degree vertex in E_0 . Let $E = E_0 \cap N^+(v_{k_0+1})$ be the set of vertices not in-dominated by (v_1, \dots, v_{k_0+1}) . Since v_{k_0+1} is a maximum in-degree vertex in E_0 , we have $|N^-(v_{k_0+1}) \cap E_0| \geq (|E_0| - 1)/2$ which implies $|E| = |E_0| - |(N^-(v_{k_0+1}) + v_{k_0+1}) \cap E_0| \leq |E_0|/2$. Combining this with the inductive hypothesis, we obtain $d^+(u) \geq 2^{k_0-1}|E_0| \geq 2^{k_0}|E|$, completing the proof. \square

We are now ready to prove the main result of this paper.

Proof of Theorem 1.5. Let T be a strongly $452k$ -connected tournament. Notice that this means that all vertices in T have in-degree and out-degree at least $452k$.

Let x_1, \dots, x_k and y_1, \dots, y_k be vertices in T as in the definition of k -linkedness. We will construct vertex disjoint paths from x_i to y_i . Let $T' = T \setminus \{x_1, \dots, x_k, y_1, \dots, y_k\}$.

Let D_1^- be a partial greedy in-dominating set in T' on 2 vertices. Then, for all $i = 2, \dots, 55k$, let D_i^- be a partial greedy in-dominating set on 2 vertices in $T' \setminus (D_1^- \cup \dots \cup D_{i-1}^-)$.

Similarly, let D_1^+ be a partial greedy out-dominating set on 2 vertices in $T' \setminus (D_1^- \cup \dots \cup D_{55k}^-)$. Then, for all $i = 2, \dots, 55k$, let D_i^+ be a partial greedy out-dominating set on 2 vertices in $T' \setminus (D_1^+ \cup \dots \cup D_{i-1}^+ \cup D_1^- \cup \dots \cup D_{55k}^-)$.

Let $X = D_1^+ \cup \dots \cup D_{55k}^+ \cup D_1^- \cup \dots \cup D_{55k}^- \cup \{x_1, \dots, x_k, y_1, \dots, y_k\}$. For each i , let E_i^- be the set of vertices in $T \setminus X$ which are not in-dominated by D_i^- , and E_i^+ the set of vertices in $T \setminus X$ which are not out-dominated by D_i^+ . By Lemma 2.3, we have $d^+(v) \geq 2|E_i^-|$ for every $v \in E_i^-$, and also $d^-(v) \geq 2|E_i^+|$ for every $v \in E_i^+$.

Let T^- be the set of heads of D_1^-, \dots, D_{55k}^- , and T^+ the set of tails of D_1^+, \dots, D_{55k}^+ . Apply Lemma 2.1 to T^- in order to find two subsets X^- and Y^- of order $5k$ of $V(T^-)$, such that for any bijection $f : X^- \rightarrow Y^-$, there is a set of $5k$ vertex-disjoint paths in T^- with each path joining x to $f(x)$ for some $x \in X^-$. Apply Lemma 2.1 to T^+ in order to find two subsets X^+ and Y^+ of order $5k$ of $V(T^+)$, such that for any bijection $f : X^+ \rightarrow Y^+$, there is a set of $5k$ vertex-disjoint paths in T^+ with each path joining x to $f(x)$ for some $x \in X^+$. Reorder $(D_1^-, \dots, D_{55k}^-)$ so that X^- is the set of heads of D_1^-, \dots, D_{55k}^- . Reorder $(D_1^+, \dots, D_{55k}^+)$ so that Y^+ is the set of tails

of D_1^+, \dots, D_{5k}^+ . Notice that since each partial greedy dominating set is on 2 vertices, we have $|X| \leq 222k$. By Menger's Theorem, since T is $452k$ -connected, there is a set of vertex-disjoint paths Q_1, \dots, Q_{5k} in $(T \setminus X) \cup Y^- \cup X^+$ such that each path Q_i starts in Y^- and ends in X^+ .

Recall that all out-degrees in T are at least $452k$ and $|X| \leq 222k$. Therefore, for each $i = 1, \dots, k$ we can choose an out-neighbour x'_i of x_i which is not in X . Similarly for each i we can choose an in-neighbour y'_i of y_i which is not in X . In addition we can ensure that $x'_1, \dots, x'_k, y'_1, \dots, y'_k$ are all distinct. Let $X' = X \cup \{x'_1, \dots, x'_k, y'_1, \dots, y'_k\}$.

Notice that each vertex $v \in E_i^-$ satisfies $d^+(v) \geq 2|E_i^-|$ and $d^+(v) \geq 452k \geq 2|X'| + 4k$. Averaging these, we get $d^+(v) \geq |E_i^-| + |X'| + 2k$ and so v has at least $2k$ out-neighbours outside of $E_i^- \cup X'$. Similarly each $v \in E_i^+$ has at least $2k$ in-neighbours outside of $E_i^+ \cup X'$. Therefore, for each i , we choose x''_i to be either equal to x'_i if $x'_i \notin E_i^-$ or we choose x''_i to be an out-neighbour of x'_i in $T \setminus (E_i^- \cup X')$. Similarly, for each i , we choose y''_i to be either equal to y'_i if $y'_i \notin E_i^+$ or we choose y''_i to be an in-neighbour of y'_i in $T \setminus (E_i^+ \cup X')$. We can also choose the vertices $x''_1, \dots, x''_k, y''_1, \dots, y''_k$ so that they are all distinct (since when $x'' \neq x'$ and $y'' \neq y'$ are always at least $2k$ choices for x''_i and y''_i respectively).

For each $i = 1, \dots, k$, let Q_i^- be a path from x''_i to the head of D_i^- whose internal vertices are all in D_i^- . The facts that D_i^- is transitive and $x''_i \notin E_i^-$ ensure that we can do this. Similarly, for each i let Q_i^+ be a path from the tail of D_i^+ to y''_i whose internal vertices are all in D_i^+ .

Notice that at least k of the paths Q_1, \dots, Q_{5k} are disjoint from $\{x'_1, \dots, x'_k, y'_1, \dots, y'_k, x''_1, \dots, x''_k, y''_1, \dots, y''_k\}$. Let Q'_1, \dots, Q'_k be some choice of such paths.

Since Q_i^- ends in X^- and Q'_i starts in Y^- , Lemma 2.1 implies that we can choose disjoint paths P_1^-, \dots, P_k^- in T^- such that P_i^- is from the end of Q_i^- to the start of Q'_i . Similarly we can choose disjoint paths P_1^+, \dots, P_k^+ in T^+ such that P_i^+ is from the end of Q'_i to the start of Q_i^+ .

Now for each i we join x_i to x'_i to Q_i^- to P_i^- to Q'_i to P_i^+ to Q_i^+ to y'_i to y_i in order to obtain the required vertex-disjoint paths from the x_i s to the y_i s. \square

3. Concluding remarks

It would be interesting to reduce the constant 452 in Theorem 1.5. It is not hard to find minor improvements to our proof in Section 2 which improve this constant by a little bit.

The optimal dependence of the connectedness on k in Theorem 1.5 may be difficult to determine. For $k = 2$, Bang-Jensen showed that every 5-connected tournament is 2-linked [2], and that the value "5" here is best possible. It is not clear how to extend Bang-Jensen's construction of 4-connected tournaments which are not 2-linked to higher k . However, for general k , it is easy to construct tournaments which are roughly $2k$ -connected, but not k -linked.

Proposition 3.1. *For all $n \geq 6k$, there is a $(2k-1)$ -connected tournament T on n vertices which is not k -linked.*

Proof. Let $n = 2k - 2 + a + b$ for some $a, b \geq 2k$. The vertices of T consist of three disjoint sets S , A , and B of with $|S| = 2k - 2$, $|A| = a$, and $|B| = b$. The edges between S , A , and B are directed from S to A , from A to B , and from B to S i.e. the digraph between these sets is a blow up of a cyclicly oriented triangle. The edges within S , A , and B are oriented arbitrarily.

To see that T is $(2k-1)$ -connected, notice that any subtournament of T containing a vertex from each of S , A , and B is strongly connected. From this it is easy to see that the set S is the unique minimum separating set in T .

To see that T is not k -linked consider two sets $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ with $S = \{x_1, \dots, x_{k-1}, y_1, \dots, y_{k-1}\}$, $x_k \in B$, and $y_k \in A$. Now if there were disjoint paths P_1, \dots, P_k with P_i going from x_i to y_i , then we would have $S \subseteq V(P_1) \cup \dots \cup V(P_{k-1})$. But then there could not be a path P_k from x_k to y_k avoiding P_1, \dots, P_{k-1} since all the edges between A and B are directed towards B . \square

While the construction above is certainly not optimal (at least for $k = 2$), the author believes that it might be essentially correct for tournaments with no small degree vertices.

Conjecture 3.2. *For every k , there is a $d = d(k)$ such that the following holds. Every strongly $2k$ -connected tournament with $\delta^-(T), \delta^+(T) \geq d$ is k -linked.*

In the above conjecture $\delta^-(T)$ and $\delta^+(T)$ are the minimum in-degree and out-degree in T respectively.

One motivation for the above conjecture is a related result of Thomas and Wollan [11] about undirected graphs which says that every $2k$ -connected graph which is sufficiently dense is k -linked. Specifically, they showed that every $2k$ -connected graph with at least $5kn$ edges is k -linked [11]. Thus Conjecture 3.2 could be seen as asking for an analogue of this result for tournaments but with a minimum degree condition instead of an average degree condition.

Acknowledgement

The author would like to thank Codruț Grosu for suggesting a simplification in the proof of Lemma 2.1, as well as a careful reviewer for helpful comments.

This research was supported by the Methods for Discrete Structures, Berlin graduate school (GRK 1408).

References

- [1] M. Ajtai, J. Komlós, and E. Szemerédi. Sorting in $c \log n$ parallel steps. *Combinatorica*, 3:1–19, 1983.
- [2] J. Bang-Jensen. On the 2-linkage problem for semicomplete digraphs. *Ann. Discrete Math.*, 41:23–38, 1989.
- [3] B. Bollobás and A. Thomason. Highly linked graphs. *Combinatorica*, 16:313–320, 1996.
- [4] H. A. Jung. Verallgemeinerung des n -fachen zusammenhangs für Graphen. *Math. Ann.*, 187:95–103, 1970.
- [5] D. Kühn, J. Kim, and D. Osthus. Bipartitions of highly connected tournaments. *arXiv:1411.1533*, 2014.
- [6] D. Kühn, J. Lapinskas, D. Osthus, and V. Patel. Proof of a conjecture of Thomassen on Hamilton cycles in highly connected tournaments. *Proc. London Math. Soc.*, to appear, 2014.
- [7] D. Kühn, D. Osthus, and T. Townsend. Proof of a tournament partition conjecture and an application to 1-factors with prescribed cycle lengths. *Combinatorica*, to appear, 2014.
- [8] D. G. Larman and P. Mani. On the existence of certain configurations within graphs and the 1-skeletons of polytopes. *Proc. London Math. Soc.*, 20:144–160, 1974.
- [9] W. Mader. Homomorphieeigenschaften und mittlere Kantendichte von Graphen. *Math. Ann.*, 174:265–268, 1967.
- [10] A. Pokrovskiy. Edge disjoint Hamiltonian cycles in highly connected tournaments. *Preprint*, 2014.
- [11] R. Thomas and P. Wollan. An improved extremal function for graph linkages. *European J. Combin.*, 26:309–324, 2005.
- [12] C. Thomassen. Connectivity in tournaments. In *Graph Theory and Combinatorics, a volume in honour of Paul Erdős (B. Bollobás, ed.)*, pages 305–313. Academic Press, London, 1984.
- [13] C. Thomassen. Note on highly connected non-2-linked digraphs. *Combinatorica*, 11:393–395, 1991.