

RH: Distemper vaccination of African wild dogs

MODIFIED LIVE DISTEMPER VACCINES CARRY LOW MORTALITY RISK FOR
CAPTIVE AFRICAN WILD DOGS, *LYCAON PICTUS*

5

Rosie Woodroffe, BA, DPhil

From the Institute of Zoology, Regent's Park, London NW1 4RY, UK

10 E-mail rosie.woodroffe@ioz.ac.uk

15

20

25

30 Abstract: Recently, canine distemper virus (CDV) has been linked to population
declines in the endangered African wild dog (*Lycaon pictus*). As CDV appears able to
persist in wildlife, threats to free-ranging wild dogs cannot be eliminated by vaccinating
domestic dogs. Conservation managers may therefore consider CDV vaccination of wild
dogs in highly threatened populations. For use in field conservation, the ideal CDV
35 vaccine would be safe, immunogenic, and readily available in Africa. The CDV vaccine
type most commonly used for domestic dogs (modified live vaccine) is available in
Africa, and apparently immunogenic in wild dogs, but has been linked to fatal vaccine-
induced distemper in captive wild dogs. However, alternatives are either ineffective
(inactivated vaccine) or difficult to obtain in Africa (recombinant vaccine). Data from a
40 questionnaire survey of zoo vaccination practices were therefore combined with studbook
tracing to assess the safety of modified live CDV vaccine in captive African wild dogs.
Among 135 wild dog pups given modified live CDV vaccine for the first time, there was
a single, unconfirmed, case of potential vaccine-induced distemper. Pups given modified
live vaccine survived better than those given inactivated vaccine or no vaccine. Although
45 studbook tracing revealed higher overall pup survival at zoos which responded to the
questionnaire than at zoos which did not, tracing of all pups born during a 20-year period
which lived long enough to be vaccinated (n=698 pups in 155 litters) revealed no
mortality events consistent with vaccine-induced distemper. Modified live CDV vaccine
thus appears to carry low mortality risks for African wild dog pups in captivity, and may
50 warrant trials in free-ranging populations.

INTRODUCTION

55 The threat to wildlife populations from infectious disease is increasingly recognised⁴⁰,
but this recognition seldom prompts intervention in the wild.^{52,63} In mammals, the
pathogens which threaten wildlife populations are often shared with domestic animals,
yet vaccines which are safe and effective in domestic animals may be ineffective or even
harmful in wildlife.^{8,54,63} Hence, concerns about vaccine safety have impeded the
60 development of field vaccination programmes for several wildlife species.^{30,6}

The African wild dog, *Lycaon pictus*, is an endangered species which is threatened by
infectious disease. Rabies has been considered the greatest disease threat, since it was
confirmed in the Serengeti wild dog population just before that population went extinct,
and has also thwarted reintroduction attempts in Namibia and South Africa.^{18,27,65,49,23}
65 Canine distemper virus (CDV) was previously thought to be a less serious threat than
rabies, because seropositive wild dogs were reported from many healthy populations,
suggesting widespread nonlethal exposure.^{13,41,31,5,2} Moreover, prior to 2016, the only
major CDV outbreak confirmed in wild dogs had occurred in captivity, where packs were
held in close proximity; the only two confirmed outbreaks in wild populations were
70 restricted to single packs.^{1,20,57} However, in 2016-7 CDV killed ≥ 20 packs in Kenya's
Ewaso ecosystem, three packs in South Africa, and one pack in Tanzania.^{15,21,32,38} There
is evidence, therefore, that CDV can represent a serious threat to free-ranging wild dog
populations. This pattern of highly variable mortality, with widespread nonlethal
exposure and occasional mass mortality, is also a feature of CDV epidemiology in some
75 other carnivore species.^{47,37,56,10}

Strategic plans for wild dog conservation call for mitigation of disease threats.^{24,26,25}

For rabies, mass vaccination of domestic dogs is likely to reduce these threats, as wild dogs acquire rabies mainly from domestic dogs and, fortuitously, ongoing strategies to end human rabies deaths entail domestic dog vaccination on the very large scales needed to protect wild dogs.^{42,64,66,9,45,67} In contrast, domestic dog vaccination may contribute little to protecting wild dog populations from CDV. The virus appears not to persist in domestic dog populations at the local scale, with several studies suggesting persistence in wildlife.^{42,19,10} Wild dog exposure to CDV is not associated with domestic dog contact, and mass CDV vaccination of domestic dogs in the Serengeti ecosystem failed to prevent new wildlife infections.^{61,41,66} Where there is a recognised need to protect free-ranging wild dogs from CDV, conservation managers may therefore need to consider vaccination of wild dogs themselves.

For use in the wild, the ideal CDV vaccine would be safe, immunogenic, and readily available in African wild dog range states. Unfortunately, it is not clear that any of the three types of CDV vaccine (modified live, inactivated, and recombinant) meet all three criteria. Modified live CDV vaccine is widely used in domestic dogs, including in Africa; however it has been reported to cause fatal vaccine-induced distemper in African wild dogs, as well as some other carnivore species.^{8,35,58,16,22} Inactivated vaccine carries no such risk of vaccine-induced distemper; however it has not prompted strong immune responses and failed to prevent mass mortality at a wild dog captive breeding facility in Tanzania.^{36,62,57} Recombinant CDV vaccine has been shown to trigger immune responses in captive wild dogs, but supply issues can limit access to this vaccine type, and

restrictions on the import of genetically modified organisms to many African countries could hinder its use in wild populations.^{7,29,11}

100 For North American zoos, husbandry guidelines recommend the use of recombinant CDV vaccines, to avoid any risk of vaccine-induced distemper.⁴ In Europe, however, recombinant CDV vaccines are not available and modified live vaccine is widely used. Limited studies suggest that modified live CDV vaccine is immunogenic in captive wild dogs.^{59,51} A recent study of captive tigers (*Panthera tigris*) found that
105 modified live CDV vaccines were markedly more immunogenic than recombinant vaccines.⁴⁸

The immunogenicity and widespread availability of modified live CDV vaccines mean that this vaccine type would be the best tool for field conservation, if the risks of vaccine-induced distemper were acceptably low. Although several cases of vaccine-
110 induced distemper have been reported in African wild dogs, cases when modified live distemper vaccine caused no ill-effects are seldom reported, and so the risk of vaccine-induced distemper is uncertain. Moreover, risks may have changed over time in response to changing conditions. A widely-used vaccine strain (the Rockborn strain, adapted to canine kidney cell culture) which was controversially linked to vaccine-induced
115 distemper was withdrawn during the 1990s, potentially reducing the risk in subsequent years.³³ Additionally, there is some evidence that the immunosuppressive effect of parvovirus infection may increase risks of vaccine-induced distemper, and so the extent of parvovirus control may influence the safety of modified live CDV vaccine.²⁸ Finally, the use of multivalent preparations including modified live vaccines against both

120 distemper and parvovirus has been tentatively linked to the risk of vaccine-induced
distemper.⁵³

This study therefore evaluated the mortality risks associated with administering
modified live CDV vaccine to African wild dogs in captivity, by combining studbook
data with data from a questionnaire survey of zoos, conducted in the years before
125 recombinant CDV vaccines became available.

MATERIALS AND METHODS

Published reports

130 Published accounts were used to characterise the mortality patterns associated with
vaccine-induced distemper, including timing of clinical signs and death relative to
vaccination, and mortality rate within affected litters.^{35,58,16}

Questionnaire survey

135 Questionnaire data were gathered in 2000, by sending paper forms to veterinary staff
responsible for the care of wild dogs held in captivity. Questionnaires were sent to 55
collections worldwide, listed on the International Species Information System (ISIS) as
holding wild dogs within their collections at the time, and to two unlisted private
collections known to hold wild dogs. The questionnaire asked for data on current and past
140 policy on vaccinating wild dogs (with reasons for any changes), and whether any wild
dogs had ever become sick or died following the administration of vaccines. In addition,
data were requested on the vaccination histories (vaccination dates, vaccine brand names

and, where possible, batch numbers) of any wild dogs held. Data were not requested on incidents of wild-type distemper. To maximise responses, the questionnaire was worded
145 to reassure participants that information from individual collections would be held in confidence. Questionnaires were sent a second time to collections that had not responded 4 months after the initial mailing.

Where zoos provided individual vaccination records, an independent estimate of mortality post-vaccination was obtained by using studbooks (which cover the European
150 and North American captive populations) to trace the survival of vaccinated animals.^{46,60} As questionnaires were sent worldwide, not just to the regions covered by the studbooks, the fates of some animals mentioned in the questionnaire responses could not be traced in this way. A Cox proportional hazards model (including litter identity as a random effect, fitted using package *coxme* in the statistical package *R*) was used to compare the survival
155 of pups which were, and were not, given modified live CDV vaccine.^{55,44} The Cox proportional hazards model is a type of survival analysis, which allows assessment of associations between independent variables (e.g., vaccination) and the time taken for an event to occur (e.g., death).¹² A hazard ratio less than one indicates a lower probability of death (i.e., higher survival). The inclusion of a random effect (here, litter identity)
160 accounts for the statistical non-independence of pups from the same litter, as they are genetically related, exposed to the same environmental conditions, and likely to receive vaccine from the same batch.

Studbook analyses

165 Questionnaire surveys will give inaccurate results if respondents represent a biased
sample of the population. To assess whether the zoos which returned their questionnaires
were representative, studbook data were used to estimate pup survival across the
European and North American captive populations during the period covered by most
questionnaire responses (1980-2000). Pup survival was compared between zoos which
170 did, and did not, respond to the questionnaire survey using a Cox proportional hazards
model including litter identity as a random effect, fitted using package *coxme* in the
statistical package *R*.^{55,44}

175 RESULTS

Published reports

All of the published reports of vaccine-induced distemper involved pups given their
first vaccinations (Table 1). All pups had been given modified live CDV vaccine in a
polyvalent formulation which also included parvovirus vaccine; however, some
180 formulations included modified live parvovirus vaccine while others contained
inactivated parvovirus vaccine (Table 1).

Clinical signs were observed 14.2 days post-vaccination on average (SD 5.6 days),
with death occurring 17.8 days (SD 6.2 days) post-vaccination (Table 1). Across the four
affected litters, 20 of 21 pups died (Table 1; 95.2% mortality, exact binomial 95%
185 confidence interval (CI) 76.2-99.9%), with deaths occurring across a period of 2-11 days
within each litter (Table 1).

Questionnaire responses

190 Questionnaires were returned by 36 collections (a 63% response rate), of which 26
reported having vaccinated wild dogs against CDV. Of 13 zoos which gave details of
vaccines used, 12 had used modified live CDV vaccines, in either monovalent (Fervac
[United Vaccines, Fitchburg WI 53593, USA], Galaxy-D [Schering Plough, Omaha, NE
68138, USA], Fromm-D [Solvay, Mendota Heights, MN 55120, USA]), or multivalent
195 (Candur [Behringwerke, Marburg 35041 Germany], Canvac [Dyntec, Terezin 41155,
Czech Republic], Duramune [Elanco, Fort Dodge, IA 50501, USA], Kavak [Fort Dodge,
Overland Park, KS 66210, USA], Nobivac [Intervet, Millsboro, DE 19966, USA],
Protech [Arthur Webster Pty, Sydney, Australia], Vanguard [Pfizer, Ramsgate CT13
9ND, UK]) preparations. Two captive facilities had used both inactivated and modified
200 live CDV vaccines, and one had used inactivated vaccine exclusively (CDV-ISCOM,
Rotterdam 3720, Netherlands).

One hundred and thirty-five wild dogs were reported to have received modified live
CDV vaccine for the first time at known age, although the vaccine brand was only
reported for 37 of these animals. These first vaccinations were given at a mean age of 61
205 days (SD 43 days, range 40-217 days), with 90% of animals having received their first
vaccination by the age of 67 days (Figure 1A).

Of 26 zoos which reported having vaccinated wild dogs against CDV, 25 reported no
observations of ill-effects post-vaccination. One zoo reported a case of a single wild dog
pup which died, aged 82 days, 10 days after its second dose of modified-live CDV
210 vaccine. Neither the brand nor type of vaccine (monovalent/polyvalent) was reported.

The *post mortem* report provided a diagnosis of canine distemper based on clinical symptoms, necropsy, and histopathology, and CDV was confirmed by a fluorescent antibody test. However, failure to isolate the virus in cell culture led the virologist consulted to suggest that this was likely to have been a wild-type virus, since “...*vaccine strains of CDV are highly cell-adapted and usually will propagate in cell cultures...*” (Unpublished necropsy report). This animal had seven littermates; one died aged 43 days, on the day of first vaccination (and so could not have died of vaccine-induced distemper), and another died aged 57 days, following bite-wounding. Five other littermates survived to adulthood. This incident therefore entailed lower post-vaccination mortality than the published cases shown in Table 1 (28.6% *vs* 95.2% mortality; Fisher exact $P = 0.001$). Nevertheless, if the two animals which died post-vaccination are assumed to have contracted vaccine-induced distemper, it would suggest a maximum of two deaths among 135 vaccinated animals, giving a vaccine-associated mortality estimate of 1.5% (CI 0.2-5.3%). If this incident did not represent vaccine-induced distemper, the estimate would be 0% (CI 0-2.7%).

Studbook tracing revealed that 118 pups given a modified live CDV vaccine in Europe or North America (aged 44 days on average) experienced higher survival over the subsequent 180 days than 17 pups (aged 44 days) given either no CDV vaccine ($n=14$) or inactivated CDV vaccine ($n=3$; Cox proportional hazards model including litter identity as a random effect, effect of modified live vaccine: hazard ratio = 0.142, $P = 0.012$, Figure 1B). The survival of pups reported to have been given modified live distemper vaccine in a monovalent formulation ($n=15$) could not be compared with that of pups given polyvalent formulations containing parvovirus vaccine which was either inactivated

(n=6) or modified live (n=15), because none of these 36 animals died in the 180 days
235 following vaccination.

Studbook analyses

Studbooks recorded mortality at all ages: of 1,459 pups born in the years 1980-2000,
40% died in the first five days of life, and 42% survived more than six months (Figure
240 1C).

The minimum age for vaccine-induced distemper (45 days) was estimated by adding
the minimum reported age at first vaccination (40 days; Figure 1A) to a conservative
estimate of the time from vaccination to death, based on published studies (5 days,
calculated as the mean [17.8 days] minus 2SD [12.4 days]; Table 1). The likely maximum
245 age for vaccine-induced distemper (99 days) was calculated by adding the age by which
90% of pups had received their first vaccination (67 days; Figure 1A) to the maximum
reported time from vaccination to death (32 days; Table 1). Among zoos which were sent
questionnaires, pup survival at ages 45-99 days was consistently higher at zoos which
responded to the questionnaire than at those which did not respond (Cox proportional
250 hazards model including litter identity as a random effect, effect of zoo response: hazard
ratio = 0.322, P = 0.045, Figure 1D).

As survival was consistently higher at zoos which responded to the questionnaire, an
alternative estimate of potentially vaccine-induced mortality was derived, independent of
the questionnaire results. Litters were identified as potential cases of vaccine-induced
255 distemper if more than one pup died at age 45-99 days, since published accounts reported
mortality of over 75% in affected litters (Table 1). To help interpret mortality patterns,

data were also collated on deaths aged 20-44 days and 100-175 days in these litters (Table 2).

Among 155 litters with members surviving to 45 days (totalling 698 pups), 18 litters met this criterion, with 53 pups dying aged 45-99 days (Table 2). If all of these litters had received modified live CDV vaccine, and all of these deaths had reflected vaccine-induced distemper, the vaccine-associated mortality would be 7.6% (CI 5.7-9.8%). However, these 18 incidents represented marked lower mortality within affected litters than the published cases (42.4% vs 95.2%, Fisher exact $P < 0.001$). The only case of 100% mortality aged 45-99 days involved a litter from a zoo which had reported that it did not routinely vaccinate wild dog pups against CDV (Table 2). Moreover, in 10 of the 18 litters, the reported deaths were part of a series which extended before or after the period when pups were 45-99 days old (Table 2). One additional litter (which experienced no mortality aged 45-99 days, and is therefore not shown in Table 2) experienced 100% mortality (of three pups) aged 100-175 days, but this litter was confirmed to have received inactivated CDV vaccine, not modified live vaccine. With no litters showing the brief episodes of high mortality reported in the published cases, at the age when vaccine-induced distemper would be expected, the incidence of apparent vaccine-associated mortality in pups aged ≥ 45 days appears to be 0/698 (0%, CI 0-0.53%). However, it is not known how many of the 698 pups had in fact received modified live CDV vaccine.

DISCUSSION

280

The results presented here suggest that modified live distemper vaccine carries a low mortality risk for captive African wild dogs. Zoos reported only one potential case of vaccine-induced distemper among 135 pups given modified live distemper vaccine, and this case was not confirmed, differing in several ways from published accounts. Although
285 the zoos which responded to the questionnaire represented a biased sample of wild dog litters with relatively low mortality, an independent evaluation of studbook records revealed no patterns of mortality similar to those described in published accounts of vaccine-induced distemper. Indeed, studbook tracing revealed that wild dog pups which received a modified live distemper vaccine experienced significantly lower mortality than
290 those which did not.

The difference in pup mortality recorded at zoos which did, and did not, respond to the questionnaire represents a form of non-response bias, a well-known source of bias in postal surveys.⁵⁰ Non-response bias was evaluated because zoo staff might have been reluctant to report cases of vaccine-induced distemper, and such reluctance is one
295 potential explanation for the higher pup mortality observed at non-responding zoos (Figure 1D). However, this explanation is not consistent with analyses of studbook data, which did not reveal mortality patterns similar to published cases of vaccine-induced distemper, in either responding or non-responding zoos. An alternative explanation is that non-responding zoos may not have routinely vaccinated wild dog pups against CDV, a
300 practice which was also associated with lower pup survival (Figure 1B). Reassuringly, the questionnaire survey and the studbook analysis both suggested consistently low risks of vaccine-induced distemper.

Wild dog pups given either no CDV vaccine or inactivated vaccine experienced lower survival than those given modified-live CDV vaccine (Figure 1B). Potentially, this pattern might reflect mortality from wild-type CDV, which has been reported from some captive facilities, including one where wild dogs had been given inactivated CDV vaccine.^{3,39,57} However, this pattern might also have reflected the protection against other canine pathogens afforded by the multivalent vaccines that most zoos used. Additionally, routine pup vaccination could indicate greater attention to veterinary care, improving survival. Irrespective of the mechanism behind it, this finding showed that modified live CDV vaccine was not associated with elevated pup mortality.

This study revealed a single case of confirmed distemper in a vaccinated wild dog pup, suspected by the virologist to reflect infection with wild-type virus rather than a vaccine strain. This observation is consistent with a study of domestic dogs, in which molecular analyses revealed that 10 puppies with suspected vaccine-induced distemper were instead infected with wild-type CDV.³⁴ This finding may also cast doubt on the published cases of vaccine-induced distemper, none of which could rule out the possibility of wild-type CDV infection.^{35,58,16}

It is surprising that four cases of presumed vaccine-induced distemper were published during the 1980s (Table 1), when similar mortality events were not recorded in studbooks during 1980-2000 (Table 2). Three of the published cases involved litters born outside the captive populations monitored by the studbooks, and the fourth case (reported from an unspecified zoo on an unspecified date) could not be identified within studbooks. The literature on CDV vaccines' reversion to virulence suggests several possible explanations for these potential cases of vaccine-induced distemper. Vaccine strain variation is one

explanation; however, the four published cases of presumed vaccine-induced distemper involved at least two different vaccine strains, including the Onderstepoort strain which is widely considered safe (Table 1).^{14,33} Likewise, immunosuppression caused by coinfection with parvovirus (or even modified live parvovirus vaccine) has been proposed as a cause of vaccine-induced distemper,^{17,28} but only two of the four reported cases involved modified live parvovirus vaccine (Table 1). Alternatively, host genetic susceptibility has been linked to vaccine-induced distemper in domestic dogs, and it is possible that the careful genetic management of captive populations (which developed from the 1980s onwards, and is the purpose of maintaining studbooks) might have reduced the risk of vaccine-induced distemper.^{17,43}

Evidence concerning the safety of vaccines in captive animals is, of course, only one component of the information needed to evaluate vaccination as a conservation tool for free-ranging animals. Evidence would also be needed of vaccine safety and effectiveness in the wild, where animals are more likely to be nutritionally stressed, as well as immunologically challenged by other pathogens. The apparently low mortality risk associated with modified live CDV vaccines in captive African wild dogs, combined with the apparently high immunogenicity of such vaccines, and their widespread availability in Africa, suggest that conservation managers may wish to consider a field trial to evaluate them further.

345

Acknowledgments: The author wishes to thank the staff from the various captive facilities who provided (often detailed) data to this study.

LITERATURE CITED

350

1. Alexander KA, Kat PW, Munson LA, Kalake A, Appel MJG. Canine distemper-related mortality among wild dogs (*Lycaon pictus*) in Chobe National Park, Botswana. *J Zoo Wildl Med.* 1996;27(3):426-427.

2. Alexander KA, McNutt JW, Briggs MB, Standers PE, Funston P, Hemson G, Keet D, van Vuuren M. Multi-host pathogens and carnivore management in southern Africa. *Comp Immunol Microbiol Infect Dis.* 2010;33(3):249-265.

3. Appel MJG, Yates RA, Foley GL, Bernstein JJ, Santinelli S, Spelman LH, Miller LD, Arp LH, Anderson M, Barr M, Pearcekelling S, Summers BA. Canine distemper epizootic in lions, tigers, and leopards in North America. *J Vet Diagn Invest.* 1994;6(3):277-288.

4. Association of Zoos and Aquariums Canid TAG. Large Canid (Canidae) Care Manual. Silver Spring, MD: Association of Zoos and Aquariums; 2012.

5. Berentsen AR, Dunbar MR, Becker MS, M'Soka J, Droge E, Sakuya NM, Matandiko W, McRobb R, Hanlon CA. Rabies, canine distemper, and canine parvovirus exposure in large carnivore communities from two Zambian ecosystems. *Vector Borne Zoonotic Dis.* 2013;13(9):643-649.

365

6. Bingham J, Foggin CM, Gerber H, Hill FW, Kappeler A, King AA, Perry BD, Wandeler AI. Pathogenicity of SAD rabies vaccine given orally in chacma baboons (*Papio ursinus*). *Vet Rec.* 1992;131(3):55-56.
- 370 7. Birhanu FM. Genetically modified organisms in Africa: regulating a threat or an opportunity? In: Bodiguel L, Cardwell M, (eds.). *The regulation of genetically modified organisms: comparative approaches.* Oxford: Oxford University Press; 2010. p. 227-253.
8. Carpenter JW, Appel MJG, Erickson RC, Novilla MN. Fatal vaccine-induced canine distemper virus infection in black-footed ferrets. *J Am Vet Med Assoc.* 1976;169(9):961-964.
- 375 9. Cleaveland S, Hampson K. Rabies elimination research: juxtaposing optimism, pragmatism and realism. *Proc R Soc Lond B Biol Sci.* 2017;284(1869):9.
10. Clifford DA, Mazet JK, Dubovi EJ, Garcelon DK, Coonan TJ, Conrad PA, Munson L. Pathogen exposure in endangered island fox (*Urocyon littoralis*) populations: implications for conservation management. *Biol Conserv.* 2006;131(2):230-243.
- 380 11. Connolly M, Thomas P, Woodroffe R, Raphael BL. Comparison of oral and intramuscular recombinant canine distemper vaccination in African wild dogs (*Lycaon pictus*). *J Zoo Wildl Med.* 2013;44(4):882-888.
- 385 12. Cox DR. Regression models and life tables. *J R Stat Soc B.* 1972;34(2):187-220.

13. Creel S, Creel NM, Munson L, Sanderlin D, Appel MJG. Serosurvey for selected viral diseases and demography of African wild dogs in Tanzania. *J Wildl Dis.* 1997;33(4):823-832.
- 390 14. Deem SL, Spelman LH, Yates RA, Montali RJ. Canine distemper in terrestrial carnivores: A review. *J Zoo Wildl Med.* 2000;31(4):441-451.
15. Du Plessis C. Canine distemper virus inoculations at HIP.
<https://wildlifeact.com/blog/canine-distemper-virus-inoculations-hip/>; 2016.
16. Durchfeld B, Baumgartner W, Herbst W, Brahm R. Vaccine-associated canine
395 distemper infection in a litter of African hunting dogs (*Lycaon pictus*). *Zentralbl Veterinarmed B.* 1990;37(1-10):203-212.
17. Fairley RA, Knesl O, Pesavento PA, Elias BC. Post-vaccinal distemper encephalitis in two Border Collie cross littermates. *N Z Vet J.* 2015;63(2):117-120.
18. Gascoyne SC, King AA, Laurenson MK, Borner M, Schildger B, Barrat J.
400 Aspects of rabies infection and control in the conservation of the African wild dog (*Lycaon pictus*) in the Serengeti region, Tanzania. *Onderstepoort J Vet Res.* 1993;60(4):415-420.
19. Gilbert M. Understanding and managing canine distemper virus as a disease threat to Amur tigers. PhD thesis, University of Glasgow; 2016.

- 405 20. Goller KV, Fyumagwa RD, Nikolin V, East ML, Kilewo M, Speck S, Muller T, Matzke M, Wibbelt G. Fatal canine distemper infection in a pack of African wild dogs in the Serengeti ecosystem, Tanzania. *Vet Microbiol.* 2010;146(3-4):245-252.
21. Grumeti Fund. Wild dog report.
<https://www.grumetifund.org/blog/updates/wild-dog-report/>; 2018.
- 410 22. Henke SE. Effects of modified live-virus canine distemper vaccines in gray foxes. *J Wildl Rehabil.* 1997;20(2):3-7.
23. Hofmeyr M, Bingham J, Lane EP, Ide A, Nel L. Rabies in African wild dogs (*Lycaon pictus*) in the Madikwe Game Reserve, South Africa. *Vet Rec.* 2000;146:50-52.
24. International Union for the Conservation of Nature/Species Survival
415 Commission. Regional conservation strategy for the cheetah and wild dog in eastern Africa. <http://www.cheetahandwilddog.org/WP/staging/9849/wp-content/uploads/2017/06/Regional-Conservation-Strategy-Eastern-Africa.pdf>; 2008.
25. International Union for the Conservation of Nature/Species Survival
Commission. Regional conservation strategy for the cheetah and wild dog in southern
420 Africa. http://www.cheetahandwilddog.org/WP/staging/9849/wp-content/uploads/2017/06/Regional-Conservation-Strategy-English_2016.pdf; 2016.
26. International Union for the Conservation of Nature/Species Survival
Commission. Regional conservation strategy for the cheetah and wild dog in West,

- Central and North Africa. http://www.cheetahandwilddog.org/WP/staging/9849/wp-content/uploads/2017/06/regional-conservation-strategy_nwc-africa-1.pdf; 2012.
- 425
27. Kat PW, Alexander KA, Smith JS, Munson L. Rabies and African wild dogs in Kenya. *Proc R Soc Lond B Biol Sci.* 1995;262(1364):229-233.
28. Krakowka S, Olsen RG, Axthelm MK, Rice J, Winters K. Canine parvovirus infection potentiates canine distemper encephalitis attributable to modified live-virus vaccine. *J Am Vet Med Assoc.* 1982;180:137-139.
- 430
29. Lau E. Merial: PureVax for ferrets coming back this week. <https://news.vin.com/vinnews.aspx?articleId=22494>; 2012.
30. Laurenson K, Shiferaw F, Sillero-Zubiri C. Disease, domestic dogs and the Ethiopian wolf: the current situation. In: Sillero-Zubiri C, Macdonald DW, (eds.). *The Ethiopian wolf: Status survey and conservation action plan.* Gland: IUCN; 1997. p. 32-42.
- 435
31. Laurenson K, van Heerden J, Stander P, van Vuuren MJ. Seroepidemiological survey of sympatric domestic and wild dogs (*Lycaon pictus*) in Tsumkwe District, north-eastern Namibia. *Onderstepoort J Vet Res.* 1997;64(4):313-316.
- 440
32. Loots AK, Mokgokong PS, Mitchell E, Venter EH, Kotze A, Dalton DL. Phylogenetic analysis of canine distemper virus in South African wildlife. *PLoS One.* 2018;13(7):e0199993.

33. Martella V, Blixenkron-Moller M, Elia G, Lucente MS, Cirone F, Decaro N, Nielsen L, Banyai K, Carmichael LE, Buonavoglia C. Lights and shades on an historical
445 vaccine canine distemper virus, the Rockborn strain. *Vaccine*. 2011;29(6):1222-1227.
34. Martella V, Elia G, Lucente MS, Decaro N, Lorusso E, Banyai K, Blixenkron-Moller M, Lan NT, Yamaguchi R, Cirone F, Carmichael LE, Buonavoglia C. Genotyping canine distemper virus (CDV) by a hemi-nested multiplex PCR provides a rapid approach for investigation of CDV outbreaks. *Vet Microbiol*. 2007;122(1-2):32-42.
- 450 35. McCormick AE. Canine distemper in African hunting dogs (*Lycaon pictus*) – possibly vaccine induced. *J Zoo Wildl Med*. 1983;14(2):66-71.
36. Montali RJ, Bartz CR, Teare JA, Allen JT, Appel MJG, Bush M. Clinical trials with canine distemper vaccines in exotic carnivores. *J Am Vet Med Assoc*. 1983;183(11):1163-1167.
- 455 37. Munson L, Terio KA, Kock R, Mlengeya T, Roelke ME, Dubovi E, Summers B, Sinclair ARE, Packer C. Climate extremes promote fatal co-infections during canine distemper epidemics in African lions. *PLoS One*. 2008;3(6):e2545.
38. Mutinda M, Cook A, Kinya S. Canine distemper outbreak in wild and domestic carnivores in Laikipia ecosystem of Kenya. Nairobi: Kenya Wildlife Service;
460 2017.
39. Nagao Y, Nishio Y, Shiomoda H, Tamaru S, Shimojima M, Goto M, Une Y, Sato A, Ikebe Y, Maeda K. An outbreak of canine distemper virus in tigers (*Panthera*

tigris): possible transmission from wild animals to zoo animals. J Vet Med Sci. 2012;74(6):699-705.

- 465 40. Pedersen AB, Jones KE, Nunn CL, Altizer S. Infectious diseases and extinction risk in wild mammals. *Conserv Biol.* 2007;21(5):1269-1279.
41. Prager KC, Mazet JAK, Munson L, Dubovi EJ, Szykman Gunther M, Davies-Mostert H, McNutt JW, Rasmussen GSA, Terio K, Cleaveland S, Mills MGL, Woodroffe R. The effect of protected areas on pathogen exposure in endangered African wild dog
470 (*Lycaon pictus*) populations. *Biol Conserv.* 2012;150(1):15-22.
42. Prager KC, Mazet JK, Dubovi EJ, Frank LG, Munson L, Rupprecht CE, Wagner A, Woodroffe R. Rabies and canine distemper virus in wild and domestic carnivores in northern Kenya: Are domestic dogs the reservoir? *EcoHealth.* 2013;9(4):483.
- 475 43. Princée FPG. *Exploring Studbooks for Wildlife Management and Conservation.* Springer; 2016.
44. R Core Team. *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing <http://www.R-project.org>; 2015.
45. Republic of Kenya. Strategic plan for the elimination of human rabies in
480 Kenya 2014-2030. <http://zdukenya.org/wp-content/uploads/2012/09/National-Rabies-Elimination-Strategy.pdf>; 2012.

46. Rhodes S, Quick M, Long S. North American regional studbook – African wild dog (*Lycaon pictus*). Chicago: Association of Zoos and Aquariums/Chicago Zoological Society; 2007.
- 485 47. Roelke-Parker ME, Munson L, Packer C, Kock R, Cleaveland S, Carpenter M, O'Brien SJ, Popischil A, Hofmann-Lehmann R, Lutz H, Mwamengele GLM, Mgasas MN, Machange GA, Summers B, Appel MJG. A canine distemper virus epidemic in Serengeti lions (*Panthera leo*). *Nature*. 1996;379(6564):441-445.
- 490 48. Sadler RA, Ramsay E, McAloose D, Rush R, Wilkes RP. Evaluation of two canine distemper virus vaccines in captive tigers (*Panthera tigris*). *J Zoo Wildl Med*. 2016;47(2):558-563.
- 495 49. Scheepers JL, Venzke KAE. Attempts to reintroduce African wild dogs *Lycaon pictus* into Etosha National Park, Namibia. *S Afr J Wildl Res*. 1995;25(4):138-140.
- 495 50. Sheikh K, Mattingly S. Investigating non-response bias in mail surveys. *Epidemiol Community Health*. 1981;35(4):293-296.
51. Spencer J, Burroughs R. Antibody response to canine distemper vaccine in African wild dogs. *J Wildl Dis*. 1992;28(3):443-444.
- 500 52. Stringer AP, Linklater W. Everything in moderation: principles of parasite control for wildlife conservation. *BioScience*. 2014;64(10):932-937.

53. Sutherland Smith MR, Rideout BA, Mikolon AB, Appel MJG, Morris PJ, Shima AL, Janssen DJ. Vaccine-induced canine distemper in European mink, *Mustela lutreola*. *J Zoo Wildl Med*. 1997;28(3):312-318.

54. Swenson J, Orr K, Bradley GA. Hemorrhagic and necrotizing hepatitis
505 associated with administration of a modified live canine adenovirus-2 vaccine in a maned
wolf (*Chrysocyon brachyurus*). *J Zoo Wildl Med*. 2012;43(2):375-383.

55. Therneau T. Mixed effects Cox models. [http://cran.r-
project.org/web/packages/coxme/coxme.pdf](http://cran.r-project.org/web/packages/coxme/coxme.pdf); 2014.

56. Timm SF, Munson L, Summers BA, Terio KA, Dubovi EJ, Rupprecht CE,
510 Kapil S, Garcelon DK. A suspected Canine Distemper epidemic as the cause of a
catastrophic decline in Santa Catalina island foxes (*Urocyon littoralis catalinae*). *J Wildl
Dis*. 2009;45(2):333-343.

57. van de Bildt MWG, Kuiken T, Visee AM, Lema S, Fitzjohn TR, Osterhaus
ADME. Distemper outbreak and its effect on African wild dog conservation. *Emerg
515 Infect Dis*. 2002;8(2):211-213.

58. van Heerden J, Bainbridge N, Burroughs REJ, Kriek NPJ. Distemper-like
disease and encephalitozoonosis in wild dogs (*Lycaon pictus*). *J Wildl Dis*.
1989;25(1):70-75.

59. van Heerden J, Bingham J, van Vuuren M, Burroughs REJ, Stylianides E.
520 Clinical and serological response of wild dogs (*Lycaon pictus*) to vaccination against

canine distemper, canine parvovirus infection and rabies. *J S Afr Vet Assoc.* 2002;73(1):8-12.

60. Verberkmoes W, Verberkmoes H. European regional studbook – African wild dog (*Lycaon pictus*). Kerkrade: GaiaPark, Kerkrade Zoo; 2007.

525 61. Viana M, Cleaveland S, Matthiopoulos J, Halliday J, Packer C, Craft ME, Hampson K, Czupryna A, Dobson AP, Dubovi E, Eblate E, Fyumagwa RD, Hoare R, Hopcraft JGC, Horton DL, Kaare MT, Kanellos T, Lankester F, Mentzel C, Mlengeya T, Mzimhiri I, Takahashi E, Willett B, Haydon DT, Lembo T. Dynamics of a morbillivirus at the domestic–wildlife interface: Canine distemper virus in domestic dogs and lions. *Proc Natl Acad Sci USA.* 2015;112(5):1464-1469.

530

62. Visee AM, Fitzjohn T, Fitzjohn L, Lema S. The Mkomazi Project African wild dog Report 1998. Netherlands: George Adamson Wildlife Preservation Trust; 1998.

63. Woodroffe R. Managing disease risks to wild mammals. *Anim Conserv.* 1999;2(3):185-193.

535 64. Woodroffe R, Donnelly CA. Risk of contact between endangered African wild dogs *Lycaon pictus* and domestic dogs: opportunities for pathogen transmission. *J Appl Ecol.* 2011;48(6):1345-1354.

65. Woodroffe R, Ginsberg JR, Macdonald DW. The African wild dog: Status survey and conservation action plan. Gland: IUCN; 1997.

- 540 66. Woodroffe R, Prager KC, Munson L, Conrad PA, Dubovi EJ, Mazet JA.
Contact with domestic dogs increases pathogen exposure in endangered African wild
dogs (*Lycaon pictus*). PLoS One. 2012;7(1):e30099.
67. World Health Organisation/Food and Agriculture Organisation/Office
International des Epizooties. Zero by 30: The global strategic plan to end human deaths
545 from dog-mediated rabies by 2030.
<https://apps.who.int/iris/bitstream/handle/10665/272756/9789241513838-eng.pdf?ua=1;>
2018.

550 Table 1. Characteristics of presumed vaccine-induced distemper in African wild dogs as reported in the literature. All published cases involved pups being given their first dose of modified-live distemper vaccine, in a polyvalent formulation including canine parvovirus (CPV) vaccine. “NR” indicates data information which was not reported.

Litter	Individual	Year	Vaccine type	Age at vaccination (days)	Days to clinical signs	Days to death	Ref
A	A1	<1989	Candur SHLP,	57	18	22	16
A	A2	<1989	Behringwerke,	57	24	30	16
A	A3	<1989	Marburg 35041.	57	25	31	16
A	A4	<1989	Strain: Rockborn CPV: inactivated	57	26	32	16
B	B1	1981		150	10	12	35
B	B2	1981	Paramune 5, Dellen	150	10	13	35
B	B3	1981	Laboratories, Omaha,	150	10	15	35
B	B4	1981	NE 68134, USA.	150	10	17	35
B	B5	1981	Strain: unknown	150	10	19	35
B	B6	1981	CPV: inactivated	150	10	20	35
B	B7	1981		150	10	survived	35
C	C1	<1987	Quantum 4, Pitman-	56	13	15	58
C	C2	<1987	Moore, Washington	56	13	15	58
C	C3	<1987	Crossing NJ 08560,	56	13	15	58
C	C4	<1987	USA.	56	13	16	58
C	C5	<1987	Strain: Onderstepoort	56	13	16	58
C	C6	<1987	CPV: live	56	13	15	58
D	D1	<1987	Quantum 6, Pitman-	77	NR	13	58
D	D2	<1987	Moore, Washington	77	NR	13	58
D	D3	<1987	Crossing NJ 08560, USA.	77	NR	13	58
D	D4	<1987	Strain: Onderstepoort CPV: live	77	NR	14	58
mean \pm SD				91.5 \pm 43.1	14.2 \pm 5.6	17.8 \pm 6.2	

Table 2. Captive African wild dog litters identified through studbook tracing to have experienced ≥ 2 deaths aged 45-99 days (when the risk of vaccine-induced distemper was estimated to be highest). Deaths before and after this age are also shown; “na” (not applicable) indicates that no deaths were recorded within the stated time window. CDV vaccination status is reported as “unknown” if zoos were not sent questionnaires, or did not return them, as “yes”/ “no” if zoos provided vaccination records for these individuals, and “probably”/ “probably not” if zoos only indicated their typical vaccination practices.

560

Zoo	Litter	Pups alive at 45 days	Mortality 45-99 days	Days of age at death			CDV vaccinated?
				20-44	45-99	100-175	
A	E284	5	40%	na	65,65	na	unknown
A	E354	4	50%	na	79,87	105,114	unknown
B	E286	8	25%	na	77,84	na	unknown
B	E287	10	30%	29	56,56,72	na	unknown
B	E288	5	80%	39	48,65,65,77	na	unknown
B	E289	8	50%	35	45,54,88,91	na	unknown
C	N51	8	25%	na	56,56	na ^a	unknown
D	E166	8	63%	na	79,81,82,82,92	na	unknown
E	E197	9	22%	na	50,50	na	unknown
F	E257	3	67%	35	53,70	na	unknown
F	E260	10	20%	36	65,68	127,155	unknown
G	E92	8	100%	39	49,53,53,56,58,63,66,66	na	probably not
G	E95	8	25%	29,41	53,55	na	no
H	E233	6	50%	na	85,86,91	na	probably not
I	E390	9	33%	na	62,74,80	na	probably
J	N6	7	28%	43	57,82	na	yes - live
K	N46	6	50%	na	50,54,55	na	unknown
L	E266	3	67%	33,40	62,71	na	unknown
Total		125	42.4%				

^athree littermates transferred and lost to monitoring at 101 days of age

565

Figure 1 Vaccination of captive African wild dogs against Canine Distemper Virus

(CDV). Panel A shows age at first vaccination for 135 pups given modified live CDV vaccine; the dotted line indicates the age by which 90% of pups had been vaccinated.

Panel B compares the survival of pups first given modified live CDV vaccine at known age in North America and Europe, with those given either no vaccine or an inactivated

570 vaccine. Panel C shows the survival, from birth to 180 days, of 1,459 pups born in North America and Europe in 1980-2000, with shading showing the age (45-99 days) when the risk of vaccine-induced distemper was estimated to be highest.

Panel D compares the survival, from 45-99 days, of pups born at zoos which did, and did not, respond to the

575 questionnaire about vaccination practices.

