

References

1. State of the Nation, stroke statistics [Internet]. Stroke association. 2017 [cited 15 November 2019]. Available from: https://www.stroke.org.uk/sites/default/files/state_of_the_nation_2017_final_1.pdf
2. Koohi N, Vickers DA, Lakshmanan R, et al. Hearing Characteristics of Stroke Patients: Prevalence and Characteristics of Hearing Impairment and Auditory Processing Disorders in Stroke Patients. *Journal of the American Academy of Audiology*. (2017) 28: 491-505.
3. Landi F, Onder G, Cesari M, et al. Functional decline in frail community-dwelling stroke patients. *European journal of neurology : the official journal of the European Federation of Neurological Societies*. (2006) 13: 17-23.
4. Savva GM, Stephan BC; Alzheimer's Society Vascular Dementia Systematic Review Group. Epidemiological studies of the effect of stroke on incident dementia: a systematic review. *Stroke* 2010;41:e41–e46
5. Livingston G, Sommerlad A, Orgeta V, et al. Dementia prevention, intervention, and care. *Lancet*. (2017) 390: 2673-2734.
6. Koohi, N., Vickers, D. A., Utoomprurkporn, N., Werring, D. J., & Bamiou, D. E. (2019). A Hearing Screening Protocol for Stroke Patients: An Exploratory Study. *Frontiers in neurology*, 10, 842. doi:10.3389/fneur.2019.00842
7. Rey B, Frischknecht R, Maeder P, et al. Patterns of recovery following focal hemispheric lesions: relationship between lasting deficit and damage to specialized networks. *Restorative neurology and neuroscience*. (2007) 25: 285-294.
8. American Speech-Language-Hearing Association 1997. Guidelines for Audiological Screening. Rockville MD: Author.
9. Yueh B, Collins MP, Souza PE, et al. Long-term effectiveness of screening for hearing loss: the screening for auditory impairment--which hearing assessment test (SAI-WHAT) randomized trial. *J Am Geriatr Soc*. (2010) 58: 427-434.