

Table 1. Baseline characteristics and group differences

	All
	646
Age, mean (SD)	55.39 (13)
Sex	
Female	432 (66.9)
Comorbidities	
Hypertension, n (%)	236 (36.5)
Hypercholesterolemia, n (%)	54 (8.4)
Diabetes Mellitus, n (%)	23 (3.6)
Heart disorder, n (%)	57 (8.8)
Smoker, n (%)	276 (42.7)
Severity grade	
Hunt and Hess Score	
• 1, n (%)	155 (24.1)
• 2, n (%)	218 (33.9)
• 3, n (%)	100 (15.5)
• 4, n (%)	99 (15.4)
• 5, n (%)	72 (11.2)
WFNS Score	
• 1, n (%)	179 (27.8)
• 2, n (%)	207 (32.1)
• 3, n (%)	77 (12)
• 4, n (%)	100 (15.5)
• 5, n (%)	81 (12.6)
Radiological characteristics	
Diameter Blood, mean (SD)	10.4 (4.6)
Aneurysm location	
• MCA	148 (22.9)
• ICA	75 (11.6)
• ACA/ACOM	249 (38.5)
• PCOM	80 (12.4)
• Posterior circulation	94 (14.6)
Multiple aneurysms	167 (25.9)
Fisher Grade	
• 1	15 (2.3)
• 2	2 (0.3)
• 3	621 (96.1)
• 4	8 (1.2)
BNI Grade	
• 1	20 (3.1)
• 2	77 (11.9)
• 3	190 (29.4)

• 4	266 (41.2)
• 5	93 (14.4)
Treatment characteristics	
Hydrocephalus	432 (66.9)
EVD, n (%)	378 (58.6)
Clipping, n (%)	339 (52.6)
Coiling, n (%)	302 (46.8)
Complications	
Rebleeding	55 (8.5)
CVS	268 (41.5)
DCI	148 (22.9)
Infarct	152 (23.5)
Outcome at 1 year (n=504)	
• GOS 1, n (%)	91 (18.1)
• GOS 2, n (%)	4 (0.8)
• GOS 3, n (%)	51 (10.1)
• GOS 4, n (%)	103 (20.4)
• GOS 5, n (%)	255 (50.6)

Data are presented in number with percentages in brackets unless stated otherwise

ACA = Anterior cerebral artery; ACOM = anterior communicating artery; BNI = Barrow Neurological Institute; CVS = cerebral vasospasm; DCI = delayed cerebral ischemia; EVD = External ventricular drainage; GOS = Glasgow Outcome Scale; ICA = internal carotid artery; MCA = middle cerebral artery; PCOM = posterior communicating artery; SD = standard deviation; WFNS = World Federation of Neurologic Surgeons