

Table 3. Incidence of DCI and functional outcome according to Neidert et al.¹

DCI			
	N=145/621 (%)	OR (95%CI)	Overall p-value
BNI Grade (only Fisher 3)¹			p=.05
1	n/a	n/a	
2	13/ <u>72</u> (18.1)	0.67 (0.31-1.44)	
3	34/ <u>190</u> (17.9)	0.66 (0.36-1.21)	
4	75/ <u>266</u> (28.2)	1.2 (0.7-2.05)	
5	23/ <u>93</u> (24.7)	Reference group	
Unfavorable outcome at 1 year			
	N=143/481 (%)	OR (95%CI)	Overall p-value
BNI Grade (only Fisher 3)¹			p<0.001
1	n/a	n/a	
2	7/ <u>53</u> (13.21)	0.14 (0.06-0.35)	
3	31/ <u>146</u> (21.23)	0.25 (0.14-0.45)	
4	65/ <u>205</u> (31.71)	0.43 (0.25-0.73)	
5	40/ <u>77</u> (51.95)	Reference group	

BNI = Barrow Neurological Institute; CI = Confidence Interval; DCI = delayed cerebral ischemia; n = number; n/a = not applicable; OR = Odds Ratio

1. Neidert MC, Maldaner N, Stienen MN, et al. The Barrow Neurological Institute Grading Scale as a Predictor for Delayed Cerebral Ischemia and Outcome After Aneurysmal Subarachnoid Hemorrhage: Data From a Nationwide Patient Registry (Swiss SOS). *Neurosurgery* 2018 doi: 10.1093/neuros/nyx609 [published Online First: 2018/01/20]