

Table 4. Multivariable model for creation of a risk score for prediction of unfavorable outcome

	OR	95% CI	P value
Age (binary)	<u>2.18</u>	<u>1.3</u> - <u>3.66</u>	.003
<u>Sex</u>	<u>0.76</u>	<u>0.46</u> - <u>1.28</u>	<u>0.31</u>
<u>Smoker</u>	<u>1.42</u>	<u>0.86</u> - <u>2.36</u>	<u>0.17</u>
BNI			.07
- 1 (reference group)			
- 2	<u>0.63</u>	<u>0.11</u> - <u>3.53</u>	
- 3	<u>0.55</u>	<u>0.11</u> - <u>2.71</u>	
- 4	<u>0.95</u>	<u>0.2</u> - <u>4.53</u>	
- 5	<u>1.63</u>	<u>0.32</u> - <u>8.3</u>	
WFNS			<.001
- 1 (reference group)			
- 2	<u>0.95</u>	<u>0.44</u> - <u>2.08</u>	
- 3	<u>3.34</u>	<u>1.44</u> - <u>7.75</u>	
- 4	<u>4.2</u>	<u>1.8</u> - <u>9.77</u>	
- 5	<u>8.12</u>	<u>3.35</u> - <u>19.65</u>	
Clipping	<u>2.56</u>	<u>1.56</u> - <u>4.22</u>	<.001
Hydrocephalus	<u>3.66</u>	<u>1.71</u> - <u>7.85</u>	.001
Rebleeding	<u>4.94</u>	<u>2.36</u> - <u>10.33</u>	<.001

BNI = Barrow Neurological Institute; CI = Confidence Interval; OR = Odds Ratio; WFNS = World Federation of Neurologic Surgeons