

MISS ZHENGZHENG HUANG (Orcid ID : 0000-0002-0196-5109)

DR MARTIN SCHULTZ (Orcid ID : 0000-0003-3458-1811)

Article type : Original Article

Cardiorespiratory fitness, fatness and the acute blood pressure response to exercise in adolescence

Zhengzheng Huang¹, Chloe Park,² Nish Chaturvedi,^{2,3} Laura D. Howe,⁴ James E. Sharman,¹ Alun D. Hughes,^{2,3} & Martin G. Schultz.¹

1) Menzies Institute for Medical Research, University of Tasmania, Hobart, Australia

2) Department of Population Science and Experimental Medicine, Institute of Cardiovascular Science, University College London, London, UK.

3) MRC Unit for Lifelong Health and Ageing at UCL, London, UK

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/SMS.13976](https://doi.org/10.1111/SMS.13976)

This article is protected by copyright. All rights reserved

4) MRC Integrative Epidemiology Unit at the University of Bristol, Bristol Medical School, University of
Bristol, Bristol, UK

Short title: Fitness, fatness and exercise blood pressure

Total word count: 2788

Tables: 2

Figures: 1

Supplemental files: 3

Corresponding author:

Dr Martin G. Schultz

Menzies Institute for Medical Research, College of Health and Medicine,

University of Tasmania, Hobart, 7000, Australia

Telephone: +61 (0) 3 6226 4264

Fax: +61 (0) 3 6226 7704

Email: Martin.Schultz@utas.edu.au

Abstract

Objective. Exaggerated exercise blood pressure (BP) is associated with cardiovascular risk factors in adolescence. Cardiorespiratory fitness and adiposity (fatness) are independent contributors to cardiovascular risk, but their interrelated associations with exercise BP are unknown. This study aimed to determine the relationships between fitness, fatness and the acute BP response to exercise in a large birth cohort of adolescents. **Methods.** 2292 adolescents from the Avon Longitudinal Study of Parents and Children (aged 17.8 ± 0.4 years, 38.5% male) completed a submaximal exercise step-test that allowed fitness ($VO_{2\text{ max}}$) to be determined from workload and heart rate using a validated equation. Exercise BP was measured immediately on test cessation and fatness calculated as the ratio of total fat mass to total body mass measured by DXA. **Results.** Post-exercise systolic BP decreased stepwise with tertile of fitness (146 (18); 142 (17); 141 (16) mmHg) but increased with tertile of fatness (138 (15); 142 (16); 149 (18) mmHg). In separate models, fitness and fatness were associated with post-exercise systolic BP adjusted for sex, age,

height, smoking and socioeconomic status (standardized β : -1.80, 95%CI: -2.64, -0.95 mmHg/SD and 4.31, 95%CI: 3.49, 5.13 mmHg/SD). However, when fitness and fatness were included in the same model, only fatness remained associated with exercise BP (4.65, 95%CI: 3.69, 5.61 mmHg/SD). **Conclusion.** Both fitness and fatness are associated with the acute BP response to exercise in adolescence. The fitness-exercise BP association was not independent of fatness, implying the cardiovascular protective effects of cardiorespiratory fitness may only be realised with more-favourable body composition.

Key words. ALSPAC; Exercise; Blood pressure; Cardiorespiratory fitness; Body Composition; Adolescent

Introduction

An exaggerated blood pressure (BP) response to exercise has been shown to be a risk factor for cardiovascular disease morbidity and mortality.^{1, 2} Evidence of the prognostic importance of exaggerated exercise BP has mostly been drawn from studies involving middle-to-older age and clinical populations. However, several studies also indicate that associations between exercise BP and markers of cardiovascular disease (including raised left-ventricular (LV) mass, increased aortic stiffness and impaired vascular function) may be present even in adolescence.^{3, 4} This highlights a need to understand the determinants of exercise BP in youth.

Poor cardiorespiratory fitness and raised adiposity (fatness) are important contributors to an adverse cardiovascular risk profile in adolescence,⁵⁻⁷ and are related to future cardiometabolic disease outcomes.⁸⁻¹⁰ Cardiorespiratory fitness and fatness are also amongst several factors known to influence the acute BP response to exercise. Indeed, current evidence suggests that higher cardiorespiratory fitness is associated with improved exercise BP in adults.^{11, 12} Moreover, our recent analysis from the Avon Longitudinal Study of Parents and Children (ALSPAC) showed that cardiorespiratory fitness also modifies the association between exercise BP and cardiac structure in 15-17 year old adolescents.¹³ Similarly, improved body composition (a decrease in body fatness) following exercise training has been shown to attenuate exercise BP in sedentary older people.¹⁴ Despite this, no study has directly assessed the individual and combined contributions of cardiorespiratory fitness and fatness to exercise BP in young people. Better understanding of the factors that contribute to elevated exercise BP in youth is of clinical importance since it may aid early detection of cardiovascular disease risk. Therefore, the aim of our study was to understand the

82 separate and joint associations between cardiorespiratory fitness, fatness and the acute BP response to
83 exercise in a large group of adolescents.

84 Methodology

85 **Participants.** Data were drawn from the Avon Longitudinal Study of Parents and Children (ALSPAC), a
86 large ongoing population-based birth cohort study in the United Kingdom. Information on the study cohort
87 has been reported previously,^{15,16} and details of all data available is fully searchable via an online data
88 dictionary (<http://www.bristol.ac.uk/alspac/researchers/our-data/>). The total sample size for analyses using
89 any data collected after age seven is 15,454 pregnancies, that resulted in 15,589 fetuses, 14,901 who were
90 alive at 1 year of age. The current study involved a complete-case cross-sectional analysis of data collected
91 during the 17-year-old sweep of clinics, which were conducted over a two-year period (2008-2010) at the
92 ALSPAC research clinic. Of the 5215 participants who took part in these clinics, a further 2923 individuals
93 were excluded due to missing or incomplete exercise test data allowing valid estimation of
94 cardiorespiratory fitness (VO_2 max), missing post-exercise BP or dual-energy x-ray absorptiometry (DXA)
95 body composition. A further 83 participants were excluded because systolic BP did not increase from pre-
96 exercise to post-exercise. This left a total of 2209 individuals for this analysis (see supplementary figure 1
97 for participant flow). As this was a complete case analysis, a further 716 were excluded from regression
98 analyses because of either missing height or socioeconomic status (SES). Ethical approval for the study
99 was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees.

100 **Assessment of cardiorespiratory fitness.** Participants undertook a modified Tecumseh Step Test during
101 the mornings of a clinic visit that consisted of stepping onto and off a step (20 cm high) with alternating
102 feet at a rate of 24 steps/minute for 3 minutes. The tempo of steps was kept constant using a metronome.
103 The test was terminated upon completion or on request of the participant. HR was recorded prior to,
104 immediately following and after 3 minutes of recovery from the test. Using this test data, it was possible to
105 calculate an estimate of cardiorespiratory fitness (VO_2 max) using modifications of the equations developed
106 by Astrand and Van Dobeln (equations 1 and 2).^{17, 18}

107 **Equation 1.**

108
$$(Male) VO_2 max = 1.29 \sqrt{\frac{Load [kg m min^{-1}]}{Exercise heart rate [bpm] - 60}} e^{-0.0088}$$

109 **Equation 2.**

$$(Female) VO_2 max = 1.18 \sqrt{\frac{Load [kg m min^{-1}]}{Exercise heart rate [bpm] - 60}} e^{-0.0090}$$

Load was calculated as the product of force and distance, where force was considered equivalent to individual body weight (kg) and distance the product of step rate (steps/min) and step height (m). Exercise HR was considered the HR recorded immediately on cessation of the exercise test. Separate equations were utilised for males and females. Pragmatic physiological limits for estimated VO₂ max were set according to the normal range for males and females <29 years of age, (26.5 - 60.5 ml.kg.min⁻¹ males, 23.7 - 54.5 ml.kg.min⁻¹ females outlined with the American College of Sports Medicine guidelines.¹⁹ When the equations produced VO₂ max values outside these limits, the result was deemed invalid and individuals were excluded from the analysis (as indicated in Supplementary figure 1).

Assessment of fatness. A DXA scanner (Lunar Prodigy, GE Medical Systems, Madison, WI, USA) was used to objectively assess body composition. Total fat mass, lean mass and bone mass (kg) were measured and summed to represent total mass. The primary measure of ‘fatness’ was fat proportion of total mass, calculated as fat mass divided by total mass and expressed as a percentage. Other measures of body composition or stature were also assessed, including lean proportion of total mass (calculated as the percentage of lean mass in total mass), central adiposity (trunk fat) as a proportion of total fat mass, height (measured using a Harpenden Stadiometer to the nearest 0.1cm) and weight (recorded to the nearest 0.1 kg using Tanita TBF 305 scales). Body mass index (BMI) was calculated as weight (kg)/ height squared (m²).

Blood Pressure. BP was measured using a validated automatic monitor (Omron 705 IT, Omron Electronic Components Europe BV) with an appropriate size cuff. Pre-exercise BP was measured prior to the step test, while the participant was standing with both arms down at sides. The participant was asked to wear the BP cuff continuously throughout the exercise test. Post- and recovery-exercise BP were measured and recorded immediately upon completion of the test (within the first 30 seconds) and after three minutes of recovery respectively. Delta exercise BP was defined as post- minus pre-exercise BP. Using the same device as the exercise BP measures, a ‘clinic BP’ was measured in triplicate after a minimum of 5 minutes seated rest on the same day, but prior to, the exercise test. The average of the last two readings was recorded as the clinic BP.

Blood biochemistry and socioeconomic status. Non-fasting blood was taken and biochemistry analysis including glucose, total cholesterol, triglycerides, glucose, high-density lipoprotein (HDL) and low-density lipoprotein (LDL) performed following locally established procedures outlined previously.²⁰ SES was determined based on the occupation of the mother’s partner (usually the father) and classified into eight classes (1, higher managerial and professional through to 8, never worked and long-term unemployed).²¹

Statistical analysis. Data were analysed using Stata (version 16.0, Texas, 77845, USA, Stata Corp LLC). Normality of variables was assessed using the Shapiro-Wilk test and by visualization of distributions. The distribution of categorical variables was analyzed with frequencies and contingency table arrays. Cardiorespiratory fitness and fatness were standardized to age and sex for regression analyses and group comparisons. This was achieved via creation of cardiorespiratory fitness and fatness z scores by first dividing the study population into age (rounded to the nearest 1 year, ranging from 16-19 years) and sex (male and female) strata groups. Multivariable linear regression analyses were performed to determine the association between cardiorespiratory fitness or fatness and exercise systolic BP, followed by mutual adjustments to examine whether each variable (fitness and fatness) were still associated with exercise BP after adjustment for the other. Outcome variables included pre-, post-, recovery-, delta exercise systolic BP, with fitness or fatness (z scores) input as the primary independent variable in each model. Age (years), sex, height (cm), socioeconomic status (SES) and smoking status were included in regression models as covariates. Results were presented as β coefficients (95% confidence interval, CI) per 1 unit increment in standardized cardiorespiratory fitness or fatness. Assumptions for linear regression were assessed by inspection of residuals and variance inflation factors (VIF) >5 or a tolerance level <0.10 interpreted as indicating collinearity. Sex-pooled models were established for each BP outcome since there was no sex-cardiorespiratory fitness or -fatness interaction on any of the exercise BP outcomes. Statistical significance was defined as $P < 0.05$. Data are presented as means (95% confidence intervals) or mean (standard deviation) unless otherwise indicated.

Results

Participant characteristics by cardiorespiratory fitness and fatness. Participant characteristics are outlined by cardiorespiratory fitness and fatness levels in table 1. The proportion of males, lean proportion of total mass and estimated VO_2 max was higher with greater cardiorespiratory fitness, whereas total cholesterol, LDL cholesterol level, clinic systolic and diastolic BP and HR, height, weight, BMI, total fat mass, total lean mass, fat proportion of total mass (fatness), delta exercise HR were lower. Total cholesterol, triglycerides, LDL cholesterol level, clinic systolic and diastolic BP, weight, BMI, total fat mass, total lean mass, fat proportion of total mass (fatness) and delta exercise HR were higher with greater fatness. The proportion of males was lowest in the middle fatness level. Clinic HR, height, lean proportion of total mass and estimated VO_2 max were lower with greater fatness level. Those with a higher level of cardiorespiratory fitness had higher paternal SES compared to those with lower cardiorespiratory fitness levels. Those with a higher level of fatness had lower paternal SES compared to those with lower fatness levels.

173 **Exercise BP by cardiorespiratory fitness and fatness.** Pre-exercise systolic and diastolic, post-exercise
174 systolic and diastolic, recovery-exercise systolic and diastolic BPs were lower with higher level of
175 cardiorespiratory fitness. Delta diastolic BP was similar in the moderate and high level of cardiorespiratory
176 fitness but slightly lower in those with a low level of cardiorespiratory fitness. Pre-exercise systolic and
177 diastolic, post-exercise systolic and diastolic, recovery-exercise systolic and delta exercise systolic BPs
178 were higher in those with higher level of fatness. Recovery-exercise diastolic and delta diastolic BP was
179 similar between low and moderate level of cardiorespiratory fitness groups, but slightly higher with a high
180 level of fatness (see supplementary table 1 for data and statistical significance).

181

182 **The association of cardiorespiratory fitness and fatness with exercise BP.** Cardiorespiratory fitness was
183 associated with pre-, post-, recovery-exercise systolic BP (model 1, table 2). After adjustment for sex, age,
184 height, smoking status and socioeconomic status, cardiorespiratory fitness remained associated with pre-,
185 post- and recovery-exercise systolic BP (model 2, table 2). When fatness was added into the model (model
186 3, table 2), the association between cardiorespiratory fitness and pre- and post-exercise systolic BP was
187 attenuated, with only recovery-exercise systolic BP remaining associated with cardiorespiratory fitness.
188 The association between cardiorespiratory fitness and delta-exercise systolic BP become stronger (model 3,
189 table 2).

190 Fatness was associated with pre-, post-, recovery-, delta exercise systolic BP (model 4, table 2). These
191 associations remained after adjustment for sex, age, height, smoking status and socioeconomic status
192 (model 5, table 2). With the further addition of cardiorespiratory fitness to the model (model 6, table 2),
193 fatness remained independently associated with pre-, post-, recovery-, delta exercise systolic BP.
194 Substituting fatness for leanness led to the results as presented in supplementary table 2. When lean
195 proportion of total mass was added into the model 3 (supplementary table 2), the association between
196 cardiorespiratory fitness and pre- and post-exercise systolic BP was attenuated, with only recovery-
197 exercise, delta-exercise systolic BP remaining associated with cardiorespiratory fitness (model 3,
198 supplementary table 2). All other associations between leanness and all exercise BPs were as per those of
199 fatness. Substituting fatness for central adiposity led to the results as presented in Supplementary table x,
200 which were broadly like those of fatness.

201 Although not statistically significant, pre-, post-, and recovery-exercise systolic BP was 3-8 mmHg higher
202 in those with high fatness compared to those with low fatness at any level of cardiorespiratory fitness
203 (Figure 1 a,b,c). Delta-exercise systolic BP was higher in those with high fatness compared to those with

low fatness at low and moderate levels of cardiorespiratory fitness, but similar in those with a high level of cardiorespiratory fitness (Figure 1 d).

206

207

Discussion

In this cohort of adolescents, cardiorespiratory fitness was negatively associated with exercise systolic BP, whilst DXA measured fatness was found to be positively associated with exercise systolic BP. The association between exercise systolic BP and cardiorespiratory fitness was not independent of fatness, whereas fatness remained strongly associated with exercise BP after adjustment for cardiorespiratory fitness. Moreover, for any level of cardiorespiratory fitness, exercise systolic BP was higher in those with greater fatness. This suggests that fatness may be more closely related to exercise BP than cardiorespiratory fitness in adolescence.

It is well known that low cardiorespiratory fitness increases risk of chronic disease and mortality across the lifespan,²² satisfying criteria to be considered a fundamental cardiovascular disease risk factor.²³ The importance of attaining and improving cardiorespiratory fitness in youth is also clear. For example, results of the large 'Coronary Artery Risk Development in Young Adults (CARDIA)' study indicated that each additional minute of exercise completed during a treadmill exercise test (when aged 18-30 years) was associated with a 15% risk reduction in all-cause mortality, independent of traditional cardiovascular risk factors.²⁴ Several other studies also show clear associations between adolescent cardiorespiratory fitness and general cardiovascular health.^{6, 10, 25} Cardiorespiratory fitness is also known to influence the acute BP response to exercise. Indeed, Cardiorespiratory fitness typically shares a negative linear association with exercise BP, such that higher fitness appears to mitigate an abnormal rise in exercise systolic BP,^{11, 12} therefore being associated with a more favourable exercise BP response and improved cardiovascular health. Whilst we found such a linear relationship between cardiorespiratory fitness and exercise BP in our general population sample of adolescents, one other analysis in young adults indicated that both low and high fit individuals may exhibit elevated exercise BP, a relationship that is more 'U-shaped'.²⁶ We also recently found cardiorespiratory fitness to modify the association between exercise BP and cardiac structure in adolescence, such that those with both low and high cardiorespiratory fitness levels share a similar exercise BP-cardiac structure relationship. Such a dependency on cardiorespiratory fitness would imply that elevated exercise BP should not always be considered the result of pathological change, but also physiologically induced adaptation. Indeed, those with high levels of cardiorespiratory fitness (such as well-trained athletes) may have elevated exercise BP, but do not share the same underlying cardiovascular dysfunction as low fit individuals.²⁷ Thus, the differing aetiology of the cardiorespiratory fitness-exercise

236 BP relationship implies there is unlikely to be a single or direct pathway between cardiorespiratory fitness
237 and exercise BP.

238 In the current study the cardiorespiratory fitness-exercise BP association was not independent of fatness.
239 On the one hand this could be interpreted to suggest that fatness is more strongly associated with exercise
240 BP than is cardiorespiratory fitness. Some studies have indicated that exercise training that improves
241 fatness also induces small-to-moderate reductions in exercise systolic BP.¹⁴ Whilst our study did not
242 involve any exercise training intervention, exercise systolic BP was step-wise higher with each level of
243 fatness. Moreover, for any level of cardiorespiratory fitness, exercise BP was higher in those with greater
244 fatness. This finding is consistent with a study of apparently healthy young men and women, in which
245 individuals with a higher waist circumference had elevated exercise systolic BP, irrespective of their
246 cardiorespiratory fitness level.²⁸ This could suggest that the benefits of greater cardiorespiratory fitness on
247 exercise SBP may not be fully realised without improvements in adiposity.

248 Beyond cardiorespiratory fitness, high fatness is an important cardiovascular disease risk factor. A large
249 study of 6th grade students in the United States indicated that fatness was associated with many
250 cardiovascular disease risk markers (including BP, cholesterol, glucose and insulin) independently of
251 fitness.²⁹ Similarly, percent body fat has been shown to have a greater influence on overall CVD risk in
252 adolescence than cardiorespiratory fitness measured as aerobic power.³⁰ Obesity is a known correlate of
253 metabolic induced vascular dysfunction (including raised arterial stiffness),³¹ that could give rise to a
254 'pathological' elevation in exercise BP when the exercise-induced demand for blood flow is not offset by
255 normal vascular compliance. Indeed, steeper trajectories of central adiposity through adolescence and
256 young adulthood are associated with greater large-artery (carotid) stiffness at mid-life, largely independent
257 of other cardiovascular risk factors including cardiorespiratory fitness.³² We have also previously found
258 associations between exercise systolic BP and aortic stiffness in the ALSPAC cohort.⁴ This cohort at the
259 age of 17 is, however, generally healthy and largely free of any clinical cardiovascular disease. Whilst this
260 allows determination of the associations between cardiorespiratory fitness, fatness and exercise BP not
261 subject to reverse causality by disease, we cannot attribute causality in a cross-sectional study. Further
262 studies are required to fully understand the mechanisms by which fatness contributes to elevations in
263 exercise BP.

264 **Limitations.** The study sample is relatively homogenous with respect to ethnicity, being of predominately
265 white European decent. Thus, our results may not be applicable to other population groups.
266 Cardiorespiratory fitness was estimated using an equation with parameters derived from a sub-maximal
267 exercise test. In the absence of an objective cardiorespiratory fitness test (such as the gold-standard VO₂

max test), we are unable to fully understand the relationship between cardiorespiratory fitness and exercise BP.

Perspectives. Cardiorespiratory fitness is negatively associated with exercise BP, whilst DXA measured fatness is positively associated with exercise systolic BP. The association between exercise systolic BP and cardiorespiratory fitness was not independent of fatness, thus the cardiorespiratory fitness-exercise BP association is accounted for by fatness. This implies that the cardiovascular protective effects of cardiorespiratory fitness on exercise BP may only be fully realised with more favourable body composition. While further longitudinal studies are needed to describe the future cardiovascular disease risk associated with raised exercise BP in youth, interventions targeted at improving both cardiorespiratory fitness and fatness could have both an independent and additive effect on reducing exercise BP and preventing cardiovascular disease risk.

279

280

281

282

283

Acknowledgments

We are extremely grateful to all the families who took part in the study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviews, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists, and nurses.

287

Funding

The UK Medical Research Council and Wellcome Trust (217065/Z/19), together with the University of Bristol, provide core support for the ALSPAC study. A comprehensive list of grants funding the ALSPAC study is available on the ALSPAC website: <http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf>. ADH received support from the Wellcome Trust (086676/7/08/Z) and the British Heart Foundation (PG/06/145 & CS/15/6/31468) in relation to the work presented in this analysis. MGS is supported by a National Heart Foundation of Australia Future Leader Research Fellowship (reference 102553). ADH and NC work in a unit that receives support from the UK Medical Research Council (Programme Code MC_UU_12019/1). LDH is supported by a Career Development Award from the UK

296 Medical Research Council (MR/M020894/1) and works in a unit funded by the UK Medical Research
297 Council (MC_UU_00011/6) and the University of Bristol. This publication is the work of the authors and
298 the corresponding author will serve as guarantor for the contents of this paper.

299 Data Availability Statement: The data that support the findings of this study are available from the
300 corresponding author upon reasonable request. Moreover, the ALSPAC study investigators encourage and
301 facilitate data sharing upon formal request.

302

303

304

305

306

307

Figure Legends

308 **Figure 1.** Exercise systolic BP by cardiorespiratory fitness and fatness level: a: pre-exercise systolic BP, b:
309 post-exercise systolic BP, c: recovery-exercise systolic BP, d: delta-exercise systolic BP. Square: high
310 fatness level, triangle: low fatness level. Differences in exercise BP between high and low fatness assessed
311 by between group t-test at each fitness tertile. Values are presented as mean and error bars represent
312 standard error.

313

314

315

316

317

318

319

320

321

322

323

324

325

326

References.

- 327 1. Schultz MG, Otahal P, Picone DS, et al. Clinical Relevance of Exaggerated Exercise Blood
328 Pressure. *J Am Coll Cardiol*. 2015;66:1843-1845.
- 329 2. Schultz MG, Otahal P, Cleland VJ, et al. Exercise-induced hypertension, cardiovascular
330 events, and mortality in patients undergoing exercise stress testing: a systematic review and meta-
331 analysis. *Am J Hypertens*. 2013;26:357-66.
- 332 3. Lambiase MJ, Dorn J, Thurston RC, et al. Flow-mediated dilation and exercise blood
333 pressure in healthy adolescents. *J Sci Med Sport*. 2014;17:425-429.
- 334 4. Schultz MG, Park C, Fraser A, et al. Submaximal exercise blood pressure and
335 cardiovascular structure in adolescence. *Int J Cardiol*. 2019;275:152-157.
- 336 5. Eisenmann JC, Welk GJ, Ihmels M, et al. Fatness, fitness, and cardiovascular disease risk
337 factors in children and adolescents. *Med Sci Sports Exerc*. 2007;39:1251-1256.
- 338 6. Ortega FB, Ruiz JR, Castillo MJ, et al. Physical fitness in childhood and adolescence: a
339 powerful marker of health. *Int J Obes*. 2008;32:1-11.
- 340 7. Steele RM, Brage S, Corder K, et al. Physical activity, cardiorespiratory fitness, and the
341 metabolic syndrome in youth. *J Appl Physiol* 2008;105:342-351.

- 342 8. Mintjens S, Menting MD, Daams JG, et al. Cardiorespiratory fitness in childhood and
343 adolescence affects future cardiovascular risk factors: a systematic review of longitudinal studies.
344 *Sports Med.* 2018;48:2577-2605.
- 345 9. Ruiz JR, Cavero-Redondo I, Ortega FB, et al. Cardiorespiratory fitness cut points to avoid
346 cardiovascular disease risk in children and adolescents; what level of fitness should raise a red flag?
347 A systematic review and meta-analysis. *British Journal of Sports Medicine.* 2016;50:1451-1458.
- 348 10. García-Hermoso A, Ramírez-Vélez R, García-Alonso Y, et al. Association of
349 Cardiorespiratory Fitness Levels During Youth With Health Risk Later in Life: A Systematic
350 Review and Meta-analysis. *JAMA pediatrics.* 2020;174:952-960.
- 351 11. Kokkinos P, Pittaras A, Narayan P, et al. Exercise capacity and blood pressure associations
352 with left ventricular mass in prehypertensive individuals. *Hypertension.* 2007;49:55-61.
- 353 12. Kokkinos PF, Andreas PE, Coutoulakis E, et al. Determinants of exercise blood pressure
354 response in normotensive and hypertensive women: role of cardiorespiratory fitness. *J Cardiopulm*
355 *Rehabil Prev.* 2002;22:178-183.
- 356 13. Huang Z, Fonseca R, Sharman JE, et al. The influence of fitness on exercise blood pressure
357 and its association with cardiac structure in adolescence. *Scand J Med Sci Sports.* 2020;30:1033-
358 1039.
- 359 14. Barone BB, Wang N-Y, Bacher AC, et al. Decreased exercise blood pressure in older
360 adults after exercise training: contributions of increased fitness and decreased fatness. *Br J Sports*
361 *Med.* 2009;43:52-56.
- 362 15. Boyd A, Golding J, Macleod J, et al. Cohort Profile: the 'children of the 90s'--the index
363 offspring of the Avon Longitudinal Study of Parents and Children. *Int J Epidemiol.* 2013;42:111-
364 27.
- 365 16. Fraser A, Macdonald-Wallis C, Tilling K, et al. Cohort profile: the Avon Longitudinal
366 Study of Parents and Children: ALSPAC mothers cohort. *International journal of epidemiology.*
367 2013;42:97-110.

- 368 17. Hughes AD and Chaturvedi N. Estimation of maximal oxygen consumption and heart rate
369 recovery using the Tecumseh sub-maximal step test and their relationship to cardiovascular risk
370 factors. *Artery Res.* 2017;18:29-35.
- 371 18. Von Döbeln W, Åstrand I and Bergström A. An analysis of age and other factors related to
372 maximal oxygen uptake. *J Appl Physiol.* 1967;22:934-938.
- 373 19. *American College of Sports Medicine: Guidelines for exercise testing and prescription, 9th*
374 *edition.* Ninth Edition ed: Lippincott, Williams & Wilkins; 2013.
- 375 20. Futema M, Cooper JA, Charakida M, et al. Screening for familial hypercholesterolaemia in
376 childhood: Avon Longitudinal Study of Parents and Children (ALSPAC). *Atherosclerosis.*
377 2017;260:47-55.
- 378 21. Rose D, Pevalin DJ and O'Reilly K. *The National Statistics Socio-economic Classification: origins, development and use.* Palgrave Macmillan Basingstoke; 2005.
- 380 22. Harber MP, Kaminsky LA, Arena R, et al. Impact of Cardiorespiratory Fitness on All-
381 Cause and Disease-Specific Mortality: Advances Since 2009. *Prog Cardiovasc Dis.* 2017;60:11-
382 20.
- 383 23. Ross R, Blair SN, Arena R, et al. Importance of Assessing Cardiorespiratory Fitness in
384 Clinical Practice: A Case for Fitness as a Clinical Vital Sign: A Scientific Statement From the
385 American Heart Association. *Circulation.* 2016;134:e653-e699.
- 386 24. Shah RV, Murthy VL, Colangelo LA, et al. Association of Fitness in Young Adulthood
387 With Survival and Cardiovascular Risk: The Coronary Artery Risk Development in Young Adults
388 (CARDIA) Study. *JAMA internal medicine.* 2016;176:87-95.
- 389 25. Ortega FB, Labayen I, Ruiz JR, et al. Improvements in Fitness Reduce the Risk of
390 Becoming Overweight across Puberty. *Medicine & Science in Sports & Exercise.* 2011;43:1891-
391 1897.

- 392 26. Prasad VK, Drenowatz C, Hand GA, et al. Association between cardiorespiratory fitness
393 and submaximal systolic blood pressure among young adult men: a reversed J-curve pattern
394 relationship. *J Hypertens*. 2015;33:2239-44.
- 395 27. Currie KD, Sless RT, Notarius CF, et al. Absence of resting cardiovascular dysfunction in
396 middle-aged endurance-trained athletes with exaggerated exercise blood pressure responses. *J*
397 *Hypertens*. 2017;35:1586-1593.
- 398 28. Huot M, Arsenault BJ, Gaudreault V, et al. Insulin resistance, low cardiorespiratory fitness,
399 and increased exercise blood pressure: contribution of abdominal obesity. *Hypertension*.
400 2011;58:1036-42.
- 401 29. Jago R, Drews KL, McMurray RG, et al. Fatness, fitness, and cardiometabolic risk factors
402 among sixth-grade youth. *Med Sci Sports Exerc*. 2010;42:1502-10.
- 403 30. Ondrak KS, McMurray RG, Bangdiwala SI, et al. Influence of aerobic power and percent
404 body fat on cardiovascular disease risk in youth. *The Journal of adolescent health : official*
405 *publication of the Society for Adolescent Medicine*. 2007;41:146-52.
- 406 31. Seifalian AM, Filippatos TD, Joshi J, et al. Obesity and arterial compliance alterations.
407 *Curr Vasc Pharmacol*. 2010;8:155-168.
- 408 32. Ferreira I, van de Laar RJ, Prins MH, et al. Carotid stiffness in young adults: a life-course
409 analysis of its early determinants: the Amsterdam Growth and Health Longitudinal Study.
410 *Hypertension*. 2012;59:54-61.

Table 1. Participant characteristics by cardiorespiratory fitness and fatness level

	Cardiorespiratory fitness			Fatness		
	Low	Moderate	High	Low	Moderate	High
Age, years	17.8 (0.4)	17.8 (0.4)	17.8 (0.4)	17.8 (0.4)	17.8 (0.4)	17.8 (0.4)
Male, percentage	180 (38.0)	186 (37.2)	213 (40.8)	197 (39.0)	187 (35.8)	195 (41.7)
Total cholesterol, mmol/l	3.9 (0.7)	3.8 (0.7)	3.7 (0.7)	3.7 (0.7)	3.8 (0.7)	4.0 (0.7)
^Triglycerides, mmol/l	0.9 (0.6-1.1)	0.8 (0.6-1.0)	0.8 (0.6-0.9)	0.8 (0.6-0.9)	0.8 (0.6-1.0)	1.0 (0.7-1.2)
^Glucose, mmol/l	5.0 (0.4)	5.0 (0.4)	5.0 (0.7)	5.0 (0.3)	5.0 (0.4)	5.1 (0.7)
^HDL, mmol/l	1.2 (0.3)	1.3 (0.3)	1.3 (0.3)	1.3 (0.3)	1.3 (0.3)	1.2 (0.3)
^LDL, mmol/l	2.3 (0.6)	2.2 (0.6)	2.0 (0.6)	2.0 (0.6)	2.2 (0.6)	2.3 (0.7)
Clinic SBP, mmHg	119 (11)	116 (11)	114 (10)	114 (10)	115 (10)	120 (11)
Clinic DBP, mmHg	66 (7)	65 (7)	63 (7)	63 (7)	64 (7)	67 (7)
Clinic HR, bpm	76 (10)	74 (9)	71 (9)	75 (10)	72 (9)	73 (10)
Height, cm	172.2 (9.6)	171.4 (9.1)	170.0 (9.1)	171.5 (9.6)	170.8 (9.3)	171.2 (9.0)
Weight, kg	76.8 (14.5)	66.3 (11.5)	63.3 (10.6)	59.3 (8.1)	66.3 (8.6)	81.1 (13.4)
Body mass index, kg/m ²	26.1 (4.8)	22.7 (3.4)	22.0 (3.1)	20.1 (1.7)	22.7 (2.0)	27.6 (3.7)
Total fat mass, kg	27.3 (11.0)	18.8 (8.6)	16.1 (7.2)	11.9 (4.6)	19.0 (4.8)	31.6 (8.7)
Total lean mass, kg	45.8 (9.7)	44.3 (9.5)	44.1 (9.6)	44.5 (9.7)	44.2 (9.6)	45.5 (9.6)
Fat proportion of total mass (fatness), %	35.3 (10.5)	28.2 (10.5)	25.6 (9.7)	20.7 (8.3)	29.3 (7.8)	39.4 (8.0)
Lean proportion of total mass, %	60.8 (10.3)	67.6 (10.3)	70.2 (9.6)	75.0 (8.2)	66.6 (7.7)	56.8 (7.8)
Estimated VO ₂ max, ml/kg/min	39.6 (4.3)	47.1 (3.6)	53.8 (3.7)	50.0 (5.6)	47.7 (6.4)	43.2 (7.2)
Delta exercise HR, bpm	24 (14)	24 (12)	24 (10)	22 (11)	24 (12)	26 (14)


HDL, high-density lipoprotein; LDL, low-density lipoprotein. Data are presented as mean (SD) or n (%). Triglycerides was presented as median (interquartile range (IQR)). Fitness levels classified as low (first tertile), moderate (second tertile) and high (third tertile) level of age-sex standardised estimated VO₂ max. Fatness were classified as low (first tertile), moderate (second tertile) and high (third tertile) level of age-sex standardised fat proportion of total mass. ^ total n= 1521. Delta exercise HR: post-exercise HR less pre-exercise HR.

Table 2. Cardiorespiratory fitness, fatness and exercise systolic BP.

	Cardiorespiratory fitness			Fatness		
	Model 1	Model 2	Model 3	Model 4	Model 5	Model 6
Outcome	β (95% CI)	β (95% CI)	β (95% CI)	β (95% CI)	β (95% CI)	β (95% CI)
Pre-exercise SBP (n=1576)	-1.79 (-2.44, -1.15) **	-1.84 (-2.45, -1.23) **	-0.53(-1.24, 0.17)	3.01 (2.37, 3.65) **	2.78 (2.18, 3.37) **	2.50 (1.80, 3.20) **
Post-exercise SBP (n=1597)	-1.70 (-2.54, -0.87) **	-1.81 (-2.66, -0.97) **	0.64 (-0.32, 1.61)	4.39 (3.58, 5.21) **	4.31 (3.50, 5.13) **	4.65 (3.69, 5.61) **
Recovery-exercise SBP (n=1556)	-2.59 (-3.23, -1.95) **	-2.62 (-3.24, -2.00) **	-1.23 (-1.94, -0.52) *	3.51 (2.88, 4.14) **	3.32 (2.71, 3.94) **	2.69 (1.98, 3.40) **
Delta exercise SBP (n=1576)	-0.15 (-0.69, 0.72)	-0.04 (-0.74, -0.65)	1.12 (0.31, 1.93) *	1.51 (0.80, 2.22) **	1.66 (0.97, 2.35) **	2.24 (1.43, 3.04) **

Fitness and fatness are age and sex standardized. Data are outcome (β , 95%CI) mmHg per one age-sex standardised unit of fitness or fatness increment of exercise systolic BP.

Model 1 - univariable; **Model 2** – model 1 plus adjustment for age (years), sex, height (cm), smoking status and socioeconomic status (SES); **Model 3** - model 2 plus adjustment for one standardized unit of age-sex standardised fatness; **Model 4** – univariable; **Model 5** – model 4 plus adjustment for age (years), sex, height (cm), smoking status and SES; **Model 6** - model 5 plus adjustment for one standardised unit of age-sex standardised fitness. **P<0.001, *P<0.05.



sms_13976_f1.tif