

Measurements of CP -Violating Asymmetries and Branching Fractions in B Meson Decays to η'/K

- B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ P. Robbe,¹ V. Tisserand,¹
A. Zghiche,¹ A. Palano,² A. Pompili,² J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ I. Ofte,⁴
B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ E. Charles,⁵
C. T. Day,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵
Yu. G. Kolomensky,⁵ J. F. Kral,⁵ G. Kukartsev,⁵ C. LeClerc,⁵ M. E. Levi,⁵ G. Lynch,⁵ L. M. Mir,⁵ P. J. Oddone,⁵
T. J. Orimoto,⁵ M. Pripstein,⁵ N. A. Roe,⁵ A. Romosan,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ A. V. Telnov,⁵ W. A. Wenzel,⁵
T. J. Harrison,⁶ C. M. Hawkes,⁶ D. J. Knowles,⁶ R. C. Penny,⁶ A. T. Watson,⁶ N. K. Watson,⁶ T. Deppermann,⁷ K. Goetzen,⁷
H. Koch,⁷ B. Lewandowski,⁷ M. Pelizaeus,⁷ K. Peters,⁷ H. Schmuecker,⁷ M. Steinke,⁷ N. R. Barlow,⁸ W. Bhimji,⁸
J. T. Boyd,⁸ N. Chevalier,⁸ W. N. Cottingham,⁸ C. Mackay,⁸ F. F. Wilson,⁸ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹
D. Thiessen,⁹ P. Kyberd,¹⁰ A. K. McKemey,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹
E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ E. P. Solodov,¹¹ A. N. Yushkov,¹¹ D. Best,¹²
M. Chao,¹² D. Kirkby,¹² A. J. Lankford,¹² M. Mandelkern,¹² S. McMahon,¹² R. K. Mommens,¹² W. Roethel,¹²
D. P. Stoker,¹² C. Buchanan,¹³ H. K. Hadavand,¹⁴ E. J. Hill,¹⁴ D. B. MacFarlane,¹⁴ H. P. Paar,¹⁴ Sh. Rahatlou,¹⁴
U. Schwanke,¹⁴ V. Sharma,¹⁴ J. W. Berryhill,¹⁵ C. Campagnari,¹⁵ B. Dahmes,¹⁵ N. Kuznetsova,¹⁵ S. L. Levy,¹⁵
O. Long,¹⁵ A. Lu,¹⁵ M. A. Mazur,¹⁵ J. D. Richman,¹⁵ W. Verkerke,¹⁵ J. Beringer,¹⁶ A. M. Eisner,¹⁶ C. A. Heusch,¹⁶
W. S. Lockman,¹⁶ T. Schalk,¹⁶ R. E. Schmitz,¹⁶ B. A. Schumm,¹⁶ A. Seiden,¹⁶ M. Turri,¹⁶ W. Walkowiak,¹⁶
D. C. Williams,¹⁶ M. G. Wilson,¹⁶ J. Albert,¹⁷ E. Chen,¹⁷ M. P. Dorsten,¹⁷ G. P. Dubois-Felsmann,¹⁷ A. Dvoretskii,¹⁷
D. G. Hitlin,¹⁷ I. Narsky,¹⁷ F. C. Porter,¹⁷ A. Ryd,¹⁷ A. Samuel,¹⁷ S. Yang,¹⁷ S. Jayatilleke,¹⁸ G. Mancinelli,¹⁸
B. T. Meadows,¹⁸ M. D. Sokoloff,¹⁸ T. Barillari,¹⁹ F. Blanc,¹⁹ P. Bloom,¹⁹ P. J. Clark,¹⁹ W. T. Ford,¹⁹ U. Nauenberg,¹⁹
A. Olivas,¹⁹ P. Rankin,¹⁹ J. Roy,¹⁹ J. G. Smith,¹⁹ W. C. van Hoek,¹⁹ L. Zhang,¹⁹ J. L. Harton,²⁰ T. Hu,²⁰ A. Soffer,²⁰
W. H. Toki,²⁰ R. J. Wilson,²⁰ J. Zhang,²⁰ D. Altenburg,²¹ T. Brandt,²¹ J. Brose,²¹ T. Colberg,²¹ M. Dickopp,²¹
R. S. Dubitzky,²¹ A. Hauke,²¹ H. M. Lacker,²¹ E. Maly,²¹ R. Müller-Pfefferkorn,²¹ R. Nogowski,²¹ S. Otto,²¹
K. R. Schubert,²¹ R. Schwierz,²¹ B. Spaan,²¹ L. Wilden,²¹ D. Bernard,²² G. R. Bonneau,²² F. Brochard,²²
J. Cohen-Tanugi,²² Ch. Thiebaux,²² G. Vasileiadis,²² M. Verderi,²² A. Khan,²³ D. Lavin,²³ F. Muheim,²³ S. Playfer,²³
J. E. Swain,²³ J. Tinslay,²³ C. Bozzi,²⁴ L. Piemontese,²⁴ A. Sarti,²⁴ E. Treadwell,²⁵ F. Anulli,^{1,*} R. Baldini-Ferroli,²⁶
A. Calcaterra,²⁶ R. de Sangro,²⁶ D. Falciai,²⁶ G. Finocchiaro,²⁶ P. Patteri,²⁶ I. M. Peruzzi,^{26,*} M. Piccolo,²⁶ A. Zallo,²⁶
A. Buzzo,²⁷ R. Contri,²⁷ G. Crosetti,²⁷ M. Lo Vetere,²⁷ M. Macri,²⁷ M. R. Monge,²⁷ S. Passaggio,²⁷ F. C. Pastore,²⁷
C. Patrignani,²⁷ E. Robutti,²⁷ A. Santroni,²⁷ S. Tosi,²⁷ S. Bailey,²⁸ M. Morii,²⁸ M. L. Aspinwall,²⁹ D. A. Bowerman,²⁹
P. D. Dauncey,²⁹ U. Egede,²⁹ I. Eschrich,²⁹ G. W. Morton,²⁹ J. A. Nash,²⁹ P. Sanders,²⁹ G. P. Taylor,²⁹ G. J. Grenier,³⁰
S.-J. Lee,³⁰ U. Mallik,³⁰ J. Cochran,³¹ H. B. Crawley,³¹ J. Lamsa,³¹ W. T. Meyer,³¹ S. Prell,³¹ E. I. Rosenberg,³¹ J. Yi,³¹
M. Davier,³² G. Grosdidier,³² A. Höcker,³² S. Laplace,³² F. Le Diberder,³² V. Lepeltier,³² A. M. Lutz,³² T. C. Petersen,³²
S. Plaszczynski,³² M. H. Schune,³² L. Tantot,³² G. Wormser,³² R. M. Bionta,³³ V. Brigljević,³³ C. H. Cheng,³³
D. J. Lange,³³ D. M. Wright,³³ A. J. Bevan,³⁴ J. R. Fry,³⁴ E. Gabathuler,³⁴ R. Gamet,³⁴ M. Kay,³⁴ D. J. Payne,³⁴
R. J. Sloane,³⁴ C. Touramanis,³⁴ J. J. Back,³⁵ G. Bellodi,³⁵ P. F. Harrison,³⁵ H. W. Shorthouse,³⁵ P. Strother,³⁵ P. B. Vidal,³⁵
G. Cowan,³⁶ H. U. Flaecher,³⁶ S. George,³⁶ M. G. Green,³⁶ A. Kurup,³⁶ C. E. Marker,³⁶ T. R. McMahon,³⁶ S. Ricciardi,³⁶
F. Salvatore,³⁶ G. Vaitsas,³⁶ M. A. Winter,³⁶ D. Brown,³⁷ C. L. Davis,³⁷ J. Allison,³⁸ R. J. Barlow,³⁸ A. C. Forti,³⁸
P. A. Hart,³⁸ F. Jackson,³⁸ G. D. Lafferty,³⁸ A. J. Lyon,³⁸ J. H. Weatherall,³⁸ J. C. Williams,³⁸ A. Farbin,³⁹ A. Jawahery,³⁹
D. Kovalskyi,³⁹ C. K. Lae,³⁹ V. Lillard,³⁹ D. A. Roberts,³⁹ G. Blaylock,⁴⁰ C. Dallapiccola,⁴⁰ K. T. Flood,⁴⁰
S. S. Hertzbach,⁴⁰ R. Kofler,⁴⁰ V. B. Koptchev,⁴⁰ T. B. Moore,⁴⁰ H. Staengle,⁴⁰ S. Willocq,⁴⁰ R. Cowan,⁴¹ G. Sciolla,⁴¹
F. Taylor,⁴¹ R. K. Yamamoto,⁴¹ D. J. J. Mangeol,⁴² M. Milek,⁴² P. M. Patel,⁴² A. Lazzaro,⁴³ F. Palombo,⁴³ J. M. Bauer,⁴⁴
L. Cremaldi,⁴⁴ V. Eschenburg,⁴⁴ R. Godang,⁴⁴ R. Kroeger,⁴⁴ J. Reidy,⁴⁴ D. A. Sanders,⁴⁴ D. J. Summers,⁴⁴ H. W. Zhao,⁴⁴
C. Hast,⁴⁵ P. Taras,⁴⁵ H. Nicholson,⁴⁶ C. Cartaro,⁴⁷ N. Cavallo,^{47,†} G. De Nardo,⁴⁷ F. Fabozzi,^{47,†} C. Gatto,⁴⁷ L. Lista,⁴⁷
P. Paolucci,⁴⁷ D. Piccolo,⁴⁷ C. Sciacca,⁴⁷ M. A. Baak,⁴⁸ G. Raven,⁴⁸ J. M. LoSecco,⁴⁹ T. A. Gabriel,⁵⁰ B. Brau,⁵¹
T. Pulliam,⁵¹ J. Brau,⁵² R. Frey,⁵² M. Iwasaki,⁵² C. T. Potter,⁵² N. B. Sinev,⁵² D. Strom,⁵² E. Torrence,⁵² F. Colecchia,⁵³
A. Dorigo,⁵³ F. Galeazzi,⁵³ M. Margoni,⁵³ M. Morandin,⁵³ M. Posocco,⁵³ M. Rotondo,⁵³ F. Simonetto,⁵³ R. Stroili,⁵³
G. Tiozzo,⁵³ C. Voci,⁵³ M. Benayoun,⁵⁴ H. Briand,⁵⁴ J. Chauveau,⁵⁴ P. David,⁵⁴ Ch. de la Vaissière,⁵⁴ L. Del Buono,⁵⁴
O. Hamon,⁵⁴ Ph. Leruste,⁵⁴ J. Ocariz,⁵⁴ M. Pivk,⁵⁴ L. Roos,⁵⁴ J. Stark,⁵⁴ S. T'Jampens,⁵⁴ P. F. Manfredi,⁵⁵ V. Re,⁵⁵

L. Gladney,⁵⁶ Q. H. Guo,⁵⁶ J. Panetta,⁵⁶ C. Angelini,⁵⁷ G. Batignani,⁵⁷ S. Bettarini,⁵⁷ M. Bondioli,⁵⁷ F. Bucci,⁵⁷ G. Calderini,⁵⁷ M. Carpinelli,⁵⁷ F. Forti,⁵⁷ M. A. Giorgi,⁵⁷ A. Lusiani,⁵⁷ G. Marchiori,⁵⁷ F. Martinez-Vidal,^{57,‡} M. Morganti,⁵⁷ N. Neri,⁵⁷ E. Paoloni,⁵⁷ M. Rama,⁵⁷ G. Rizzo,⁵⁷ F. Sandrelli,⁵⁷ J. Walsh,⁵⁷ M. Haire,⁵⁸ D. Judd,⁵⁸ K. Paick,⁵⁸ D. E. Wagoner,⁵⁸ N. Danielson,⁵⁹ P. Elmer,⁵⁹ C. Lu,⁵⁹ V. Miftakov,⁵⁹ J. Olsen,⁵⁹ A. J. S. Smith,⁵⁹ E. W. Varnes,⁵⁹ F. Bellini,⁶⁰ G. Cavoto,^{59,60} D. del Re,⁶⁰ R. Faccini,^{14,60} F. Ferrarotto,⁶⁰ F. Ferroni,⁶⁰ M. Gaspero,⁶⁰ E. Leonardi,⁶⁰ M. A. Mazzoni,⁶⁰ S. Morganti,⁶⁰ M. Pierini,⁶⁰ G. Piredda,⁶⁰ F. Safai Tehrani,⁶⁰ M. Serra,⁶⁰ C. Voenia,⁶⁰ S. Christ,⁶¹ G. Wagner,⁶¹ R. Waldi,⁶¹ T. Adye,⁶² N. De Groot,⁶² B. Franek,⁶² N. I. Geddes,⁶² G. P. Gopal,⁶² E. O. Olaiya,⁶² S. M. Xella,⁶² R. Aleksan,⁶³ S. Emery,⁶³ A. Gaidot,⁶³ S. F. Ganzhur,⁶³ P.-F. Giraud,⁶³ G. Hamel de Monchenault,⁶³ W. Kozanecki,⁶³ M. Langer,⁶³ G. W. London,⁶³ B. Mayer,⁶³ G. Schott,⁶³ G. Vasseur,⁶³ Ch. Yecho,⁶³ M. Zito,⁶³ M. V. Purohit,⁶⁴ A. W. Weidemann,⁶⁴ F. X. Yumiceva,⁶⁴ D. Aston,⁶⁵ R. Bartoldus,⁶⁵ N. Berger,⁶⁵ A. M. Boyarski,⁶⁵ O. L. Buchmueller,⁶⁵ M. R. Convery,⁶⁵ D. P. Coupal,⁶⁵ D. Dong,⁶⁵ J. Dorfan,⁶⁵ D. Dujmic,⁶⁵ W. Dunwoodie,⁶⁵ R. C. Field,⁶⁵ T. Glanzman,⁶⁵ S. J. Gowdy,⁶⁵ E. Grauges-Pous,⁶⁵ T. Hadig,⁶⁵ V. Halyo,⁶⁵ T. Hryna'ova,⁶⁵ W. R. Innes,⁶⁵ C. P. Jessop,⁶⁵ M. H. Kelsey,⁶⁵ P. Kim,⁶⁵ M. L. Kocian,⁶⁵ U. Langenegger,⁶⁵ D. W. G. S. Leith,⁶⁵ S. Luitz,⁶⁵ V. Luth,⁶⁵ H. L. Lynch,⁶⁵ H. Marsiske,⁶⁵ S. Menke,⁶⁵ R. Messner,⁶⁵ D. R. Muller,⁶⁵ C. P. O'Grady,⁶⁵ V. E. Ozcan,⁶⁵ A. Perazzo,⁶⁵ M. Perl,⁶⁵ S. Petrak,⁶⁵ B. N. Ratcliff,⁶⁵ S. H. Robertson,⁶⁵ A. Roodman,⁶⁵ A. A. Salnikov,⁶⁵ R. H. Schindler,⁶⁵ J. Schwiening,⁶⁵ G. Simi,⁶⁵ A. Snyder,⁶⁵ A. Soha,⁶⁵ J. Stelzer,⁶⁵ D. Su,⁶⁵ M. K. Sullivan,⁶⁵ H. A. Tanaka,⁶⁵ J. Va'vra,⁶⁵ S. R. Wagner,⁶⁵ M. Weaver,⁶⁵ A. J. R. Weinstein,⁶⁵ W. J. Wisniewski,⁶⁵ D. H. Wright,⁶⁵ C. C. Young,⁶⁵ P. R. Burchat,⁶⁶ T. I. Meyer,⁶⁶ C. Roat,⁶⁶ S. Ahmed,⁶⁷ J. A. Ernst,⁶⁷ W. Bugg,⁶⁸ M. Krishnamurthy,⁶⁸ S. M. Spanier,⁶⁸ R. Eckmann,⁶⁹ H. Kim,⁶⁹ J. L. Ritchie,⁶⁹ R. F. Schwitters,⁶⁹ J. M. Izen,⁷⁰ I. Kitayama,⁷⁰ X. C. Lou,⁷⁰ S. Ye,⁷⁰ F. Bianchi,⁷¹ M. Bona,⁷¹ F. Gallo,⁷¹ D. Gamba,⁷¹ C. Borean,⁷² L. Bosisio,⁷² G. Della Ricca,⁷² S. Dittongo,⁷² S. Grancagnolo,⁷² L. Lanceri,⁷² P. Poropat,^{72,§} L. Vitale,⁷² G. Vuagnin,⁷² R. S. Panvini,⁷³ Sw. Banerjee,⁷⁴ C. M. Brown,⁷⁴ D. Fortin,⁷⁴ P. D. Jackson,⁷⁴ R. Kowalewski,⁷⁴ J. M. Roney,⁷⁴ H. R. Band,⁷⁵ S. Dasu,⁷⁵ M. Datta,⁷⁵ A. M. Eichenbaum,⁷⁵ H. Hu,⁷⁵ J. R. Johnson,⁷⁵ R. Liu,⁷⁵ F. Di Lodovico,⁷⁵ A. K. Mohapatra,⁷⁵ Y. Pan,⁷⁵ R. Prepost,⁷⁵ S. J. Sekula,⁷⁵ J. H. von Wimmersperg-Toeller,⁷⁵ J. Wu,⁷⁵ S. L. Wu,⁷⁵ Z. Yu,⁷⁵ and H. Neal⁷⁶

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

³Institute of High Energy Physics, Beijing 100039, China

⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁶University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁸University of Bristol, Bristol BS8 ITL, United Kingdom

⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹²University of California at Irvine, Irvine, California 92697, USA

¹³University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁴University of California at San Diego, La Jolla, California 92093, USA

¹⁵University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁶University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁷California Institute of Technology, Pasadena, California 91125, USA

¹⁸University of Cincinnati, Cincinnati, Ohio 45221, USA

¹⁹University of Colorado, Boulder, Colorado 80309, USA

²⁰Colorado State University, Fort Collins, Colorado 80523, USA

²¹Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

²²Ecole Polytechnique, LLR, F-91128 Palaiseau, France

²³University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

²⁴Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

²⁵Florida A&M University, Tallahassee, Florida 32307, USA

²⁶Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

²⁷Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

²⁸Harvard University, Cambridge, Massachusetts 02138, USA

²⁹Imperial College London, London, SW7 2BW, United Kingdom

³⁰University of Iowa, Iowa City, Iowa 52242, USA

- ³¹Iowa State University, Ames, Iowa 50011-3160, USA
³²Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
³³Lawrence Livermore National Laboratory, Livermore, California 94550, USA
³⁴University of Liverpool, Liverpool L69 3BX, United Kingdom
³⁵Queen Mary, University of London, E1 4NS, United Kingdom
³⁶University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
³⁷University of Louisville, Louisville, Kentucky 40292, USA
³⁸University of Manchester, Manchester M13 9PL, United Kingdom
³⁹University of Maryland, College Park, Maryland 20742, USA
⁴⁰University of Massachusetts, Amherst, Massachusetts 01003, USA
⁴¹Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
⁴²McGill University, Montréal, QC, Canada H3A 2T8
⁴³Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁴University of Mississippi, University, Mississippi 38677, USA
⁴⁵Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7
⁴⁶Mount Holyoke College, South Hadley, Massachusetts 01075, USA
⁴⁷Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
⁴⁸NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
⁴⁹University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁰Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA
⁵¹The Ohio State University, Columbus, Ohio 43210, USA
⁵²University of Oregon, Eugene, Oregon 97403, USA
⁵³Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁴Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
⁵⁵Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
⁵⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁵⁷Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
⁵⁸Prairie View A&M University, Prairie View, Texas 77446, USA
⁵⁹Princeton University, Princeton, New Jersey 08544, USA
⁶⁰Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶¹Universität Rostock, D-18051 Rostock, Germany
⁶²Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
⁶³DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
⁶⁴University of South Carolina, Columbia, South Carolina 29208, USA
⁶⁵Stanford Linear Accelerator Center, Stanford, California 94309, USA
⁶⁶Stanford University, Stanford, California 94305-4060, USA
⁶⁷State University of New York, Albany, New York 12222, USA
⁶⁸University of Tennessee, Knoxville, Tennessee 37996, USA
⁶⁹University of Texas at Austin, Austin, Texas 78712, USA
⁷⁰University of Texas at Dallas, Richardson, Texas 75083, USA
⁷¹Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷²Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷³Vanderbilt University, Nashville, Tennessee 37235, USA
⁷⁴University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷⁵University of Wisconsin, Madison, Wisconsin 53706, USA
⁷⁶Yale University, New Haven, Connecticut 06511, USA

(Received 31 March 2003; published 13 October 2003)

We present measurements of the branching fractions of the decays $B^+ \rightarrow \eta' K^+$ and $B^0 \rightarrow \eta' K^0$. For $B^0 \rightarrow \eta' K_S^0$ we also measure the time-dependent CP -violation parameters $S_{\eta' K_S^0}$ and $C_{\eta' K_S^0}$, and for $B^+ \rightarrow \eta' K^+$ the time-integrated charge asymmetry \mathcal{A}_{ch} . The data sample corresponds to 88.9×10^6 $B\bar{B}$ pairs produced by $e^+ e^-$ annihilation at the Y(4S). The results are $\mathcal{B}(B^+ \rightarrow \eta' K^+) = (76.9 \pm 3.5 \pm 4.4) \times 10^{-6}$, $\mathcal{B}(B^0 \rightarrow \eta' K^0) = (60.6 \pm 5.6 \pm 4.6) \times 10^{-6}$, $S_{\eta' K_S^0} = 0.02 \pm 0.34 \pm 0.03$, $C_{\eta' K_S^0} = 0.10 \pm 0.22 \pm 0.04$, and $\mathcal{A}_{ch} = 0.037 \pm 0.045 \pm 0.011$.

DOI: 10.1103/PhysRevLett.91.161801

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Nonconservation of CP in the neutral B meson system has been clearly established [1,2] in decays to charmonium such as $B^0 \rightarrow J/\psi K_S^0$. The CP effect arises from the interference between mixing and decay involving the CP -violating phase $\beta = \arg(-V_{cd} V_{cb}^* / V_{td} V_{tb}^*)$ of the

Cabibbo-Kobayashi-Maskawa (CKM) mixing matrix, and appears experimentally as an asymmetry in the time evolution of the $B^0 \bar{B}^0$ meson pair. These decays occur via a CKM-favored (though color suppressed) $b \rightarrow c$ tree amplitude.

Here we report results of a similar analysis of the decay $B^0 \rightarrow \eta' K_S^0$, a CKM-suppressed process that is expected to be dominated by penguin $b \rightarrow s$ transitions, while the tree and electroweak contributions are expected to be small [3–5]. The observed branching fraction is 3–10 times larger than initially expected [3], which has motivated a variety of conjectures by way of explanation, including flavor singlet [4] and charm enhanced [6] terms. A recent next-to-leading order QCD factorization calculation [5] suggests that the decay rate is not significantly enhanced by these mechanisms, but is adequately predicted by constructive interference between the penguin diagrams in which the spectator quark is contained in the η' or in the kaon.

The results presented in this Letter are based on data collected in 1999–2002 with the *BABAR* detector [7] at the PEP-II asymmetric $e^+ e^-$ collider [8] located at the Stanford Linear Accelerator Center. An integrated luminosity of 81.9 fb^{-1} , corresponding to $88.9 \times 10^6 B\bar{B}$ pairs, was recorded at the $\Upsilon(4S)$ resonance (center-of-mass energy $\sqrt{s} = 10.58 \text{ GeV}$).

Charged particles from the $e^+ e^-$ interactions are detected, and their momenta measured, by a combination of a vertex tracker (SVT) consisting of five layers of double-sided silicon microstrip detectors, and a 40-layer central drift chamber, both operating in the 1.5 T magnetic field of a superconducting solenoid. Photons and electrons are detected by a CsI(Tl) electromagnetic calorimeter. Charged particle identification (PID) is provided by the average energy loss (dE/dx) in the tracking devices, and by an internally reflecting ring imaging Cherenkov detector (DIRC) covering the central region.

From a $B^0 \bar{B}^0$ meson pair produced in $\Upsilon(4S)$ decay we reconstruct one of the mesons in the final state $f = \eta' K_S^0$, a CP eigenstate with eigenvalue $\eta_f = -1$. For the time evolution measurement, we also identify the flavor (B^0 or \bar{B}^0) and reconstruct the decay vertex of the partner (B_{tag}). The asymmetric beam configuration in the laboratory frame provides a boost of $\beta\gamma = 0.56$ to the $\Upsilon(4S)$, which allows the determination of the proper decay time difference $\Delta t \equiv t_f - t_{\text{tag}}$ from the vertex separation of the two B meson candidates. The distribution of Δt is

$$F(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} \{ 1 \mp \Delta w \pm (1 - 2w) \times [S_f \sin(\Delta m_d \Delta t) - C_f \cos(\Delta m_d \Delta t)] \}. \quad (1)$$

The upper (lower) sign denotes a decay accompanied by a B^0 (\bar{B}^0) tag, τ is the mean B^0 lifetime, Δm_d is the mixing frequency, and the mistag parameters w and Δw are the average and difference, respectively, of the probabilities that a true B^0 (\bar{B}^0) meson is tagged as a \bar{B}^0 (B^0). The tagging algorithm is described in [1], and has a measured analyzing power [efficiency times $(1 - 2w)^2$] of $(28.1 \pm 0.7)\%$.

The parameter C_f measures direct CP violation. If $C_f = 0$, then $S_f = \sin 2\beta_{\text{eff}}$, with β_{eff} equal to β combined with any weak phase difference arising from multiple amplitudes in the decay. Assuming the tree amplitudes are negligible, a deviation from the value found in charmonium channels can be considered an effect of phases coming from new physics [9]. Direct CP violation can also be detected as an asymmetry $\mathcal{A}_{\text{ch}} = (\Gamma^- - \Gamma^+)/(\Gamma^- + \Gamma^+)$ in the rates $\Gamma^\pm = \Gamma(B^\pm \rightarrow \eta' K^\pm)$.

We reconstruct a B meson candidate by combining a K^+ [10] or K_S^0 with an $\eta' \rightarrow \eta \pi^+ \pi^-$ ($\eta' \eta \pi \pi$) or $\eta' \rho \gamma$ ($\eta' \rho \gamma$). The $K_S^0 \rightarrow \pi^+ \pi^-$, $\eta' \rightarrow \gamma \gamma$, and $\rho^0 \rightarrow \pi^+ \pi^-$ candidates are selected with requirements on the relevant invariant masses similar to those of our previous analysis [11]. Distributions from the data and from Monte Carlo (MC) simulations [12] guide the choice of these selection criteria. For those quantities taken subsequently as observables for fitting we retain sidebands adequate to characterize the background as well as the signal. For charged B decays, the K^+ candidate must have an associated DIRC Cherenkov angle between -5σ and $+2\sigma$ of the value expected for a kaon. This requirement rejects 91% of pions.

The B -meson candidate is characterized by the energy substituted mass $m_{\text{ES}} = \sqrt{(\frac{1}{2}s + \mathbf{p}_0 \cdot \mathbf{p}_B)^2/E_0^2 - |\mathbf{p}_B|^2}$ and energy difference $\Delta E = E_B^* - \frac{1}{2}\sqrt{s}$, where the subscripts 0 and B refer to the initial $\Upsilon(4S)$ and the B candidate, respectively, and the asterisk denotes the $\Upsilon(4S)$ rest frame. The resolutions on these quantities measured for signal events are 29 MeV for ΔE and $2.9 \text{ MeV}/c^2$ for m_{ES} . We require $|\Delta E| \leq 0.2 \text{ GeV}$ and $5.2 \leq m_{\text{ES}} \leq 5.29 \text{ GeV}/c^2$.

Backgrounds arise primarily from combinatorics among continuum events. To reject these we make use of the angle θ_T between the thrust axis of the B candidate in the $\Upsilon(4S)$ frame and that of the rest of the charged tracks and neutral clusters in the event. The distribution of $\cos \theta_T$ is sharply peaked near ± 1 for combinations drawn from jetlike $q\bar{q}$ pairs, and nearly uniform for the isotropic B meson decays; we require $|\cos \theta_T| < 0.9$.

We obtain the yields and decay time evolution from extended unbinned maximum likelihood fits, with input observables Δt , ΔE , m_{ES} , $m_{\eta'}$, and a Fisher discriminant \mathcal{F} . The Fisher discriminant [13] combines four variables: the angles with respect to the beam axis of the B momentum and B thrust axis [in the $\Upsilon(4S)$ frame], and the zeroth and second angular moments of the energy flow (excluding the B candidate) about the B thrust axis.

We use MC simulation to estimate backgrounds from other B decays, including final states with and without charm. These contributions are negligible for the $\eta' \eta \pi \pi$ modes. For $\eta' \rho \gamma$ we include in the fit a $B\bar{B}$ component (which we find to be small).

Since we measure the correlations among the observables to be small in the (background-dominated) data samples entering the fit, we take the probability density

function (PDF) for each event to be a product of the PDFs for the separate observables. The efficiencies and mistag rates w for each of four tagging categories are measured with a large sample (B_{flav}) of decays to fully reconstructed flavor eigenstates [1]. The signatures of the four tagging categories are essentially lepton, K^+ from D^* , K^+ , and a flavor-correlated inclusive class. For each event hypothesis j (signal, $B\bar{B}$ background, continuum background) and tagging category k , we define the PDF (to be evaluated with the observable set for event i) as

$$\mathcal{P}_{j,k}^i = \mathcal{P}_j(m_{\text{ES}})\mathcal{P}_j(\Delta E)\mathcal{P}_j(\mathcal{F})\mathcal{P}_j(m_{\eta'})\mathcal{P}_j(\Delta t; \sigma_{\Delta t}, k). \quad (2)$$

The likelihood function for each decay chain is then

$$\mathcal{L} = \prod_k \exp\left(-\sum_j Y_{j,k}\right) \prod_i^{N_k} \left[\sum_j Y_{j,k} \mathcal{P}_{j,k}^i \right], \quad (3)$$

where $Y_{j,k}$ is the yield of events of hypothesis j found by the fitter in category k , and N_k is the number of category k events in the sample.

The signal PDF factor $\mathcal{P}_{\text{sig}}(\Delta t; \sigma_{\Delta t}, k)$ is equal to the convolution of $F(\Delta t; k)$ [Eq. (1)], with the signal resolution function, determined from the B_{flav} sample; $\sigma_{\Delta t}$ is the error on Δt for a given event. We determine the remaining PDFs from simulation for the signal and $B\bar{B}$ background components, and from $(m_{\text{ES}}, \Delta E)$ sideband data for continuum background. Each of the functions $\mathcal{P}_{\text{sig}}(m_{\text{ES}})$, $\mathcal{P}_{\text{sig}}(\Delta E)$, $\mathcal{P}_j(\mathcal{F})$, $\mathcal{P}_{\text{bkg}}(\Delta t; k)$, and the peaking component of $\mathcal{P}_j(m_{\eta'})$ is parametrized as a Gaussian function, with or without a second or third Gaussian or asymmetric width as required to describe the distribution. Slowly varying distributions (combinatorial background under mass or energy peaks) are represented by linear or quadratic dependences, or for m_{ES} , by the function $x\sqrt{1-x^2} \exp[-\xi(1-x^2)]$, with $x \equiv 2m_{\text{ES}}/\sqrt{s}$ and parameter ξ . Large control samples of B decays to charmed final states of similar topology are used to verify the simulated resolutions in ΔE and m_{ES} .

We compute the branching fractions and \mathcal{A}_{ch} from fits made without Δt or flavor tagging. Seven parameters of the background PDF are free in the fit, along with signal

and continuum background yields, for $\eta'_{\rho\gamma} K$ the $B\bar{B}$ background yield, and for charged modes the signal and background \mathcal{A}_{ch} . We compute the branching fractions from the fitted signal yields, reconstruction efficiencies, daughter branching fractions, and the number of produced B mesons, assuming equal production rates of charged and neutral pairs. To determine the reconstruction efficiency, including any yield bias of the likelihood fit, we apply the method to simulated samples constructed to contain the signal and continuum background populations expected for data.

Table I shows for each decay chain the branching fraction we measure, together with the quantities entering into its computation. The purity estimate is given by the ratio of the signal yield to the effective background plus signal, defined as the square of the error on the yield. In Fig. 1 we show projections onto m_{ES} and ΔE of a subset of the data for which the signal likelihood (computed without the variable plotted) exceeds a mode-dependent threshold that optimizes the sensitivity.

For the time evolution we combine the two decay chains in a single fit with 28 free parameters: S_f , C_f , signal fractions (two), $\eta'_{\rho\gamma} K$ $B\bar{B}$ background yield (one), common background \mathcal{F} PDF parameters (three), and separate background Δt , m_{ES} , ΔE , $m_{\eta'}$ PDF parameters (20). The last four columns of Table I give the tagged subsample yields with their purity, along with S_f and C_f . The S_f and C_f values for $B^+ \rightarrow \eta' K^+$ are included as a control; they are consistent with zero, as expected. We show in Fig. 2 the Δt projections and asymmetry of the combined neutral modes for events selected as for Fig. 1.

Most of the systematic errors on yields, which arise from PDF uncertainties (1%–2%, depending on the decay mode), have already been incorporated into the overall statistical error, because their background parameters are free in the fit. We verify that the likelihood of each fit is consistent with the distribution found in simulation.

The uncertainty in our knowledge of the efficiency is found from auxiliary studies to be 0.8% per charged track, 2.5% per photon, and 4% per K_S^0 . Our estimate of the B production systematic error is 1.1%. The estimate of systematic bias from the fitter itself (0%–4%) comes from fits of simulated samples with varying background

TABLE I. Signal yield, purity P , detection efficiency ϵ , daughter branching fraction product that was forced to 100% in our signal mode simulation, measured branching fraction, background ($\mathcal{A}_{\text{ch}}^{qq}$) and signal (\mathcal{A}_{ch}) charge asymmetries, tagged subsample yield Y_{tag} and purity P_{tag} , S_f , and C_f for each decay chain, and the combined result for each mode, with statistical error.

Mode	Yield	P (%)	ϵ (%)	$\prod \mathcal{B}_i$ (%)	\mathcal{B} (10^{-6})	$\mathcal{A}_{\text{ch}}^{qq}$ (%)	\mathcal{A}_{ch} (%)	Y_{tag}	P_{tag} (%)	S_f	C_f
$\eta'_{\rho\pi\pi} K^+$	268 ± 19	78	25	17.4	71 ± 5	0.6 ± 1.6	-0.1 ± 6.8	183	92	0.08 ± 0.20	-0.16 ± 0.15
$\eta'_{\rho\gamma} K^+$	514 ± 31	55	24	29.5	82 ± 5	-0.9 ± 0.5	6.3 ± 5.9	355	63	-0.07 ± 0.16	-0.14 ± 0.11
$\eta' K^+$					76.9 ± 3.5	-0.8 ± 0.4	3.7 ± 4.5			-0.01 ± 0.13	-0.15 ± 0.09
$\eta'_{\rho\pi\pi} K^0$	48 ± 8	75	22	6.0	42 ± 7			31.6	75	0.75 ± 0.51	-0.21 ± 0.35
$\eta'_{\rho\gamma} K^0$	155 ± 17	59	23	10.1	76 ± 8			77.6	61	-0.41 ± 0.42	0.24 ± 0.27
$\eta' K^0$					60.6 ± 5.6					0.02 ± 0.34	0.10 ± 0.22




FIG. 1 (color online). The B candidate m_{ES} and ΔE projections for $B^+ \rightarrow \eta' K^+$ (a),(b) and $B^0 \rightarrow \eta' K_S^0$ (c),(d). The points with errors represent the data, the solid curves represent the full fit functions, and the dashed curves represent the background functions; the shaded histogram represents the $\eta' \pi \pi K$ subset.

populations. Published data [14] provide the B daughter product branching fraction uncertainties (3.4%). Selection efficiency uncertainties are 1% for $\cos\theta_T$ and 0.5% for PID. As can be seen in Table I, the branching fractions we find for $B^0 \rightarrow \eta' K^0$ are rather different (3 standard deviations) as measured with $\eta' \rightarrow \eta \pi \pi$ or $\eta' \rightarrow \rho \gamma$. Having exhausted other explanations, we attribute this difference to a statistical fluctuation, and include both components in the final measurement.

Using several large inclusive kaon and B -decay samples, we find a systematic uncertainty for \mathcal{A}_{ch} of 1.1% due to the dependence of reconstruction efficiency on the charge of the high momentum K^\pm .

We find systematic uncertainties for $S_{\eta' K_S^0}$ and $C_{\eta' K_S^0}$ by varying within their errors the fit parameters controlling the PDF shapes. We use the B_{flav} sample to determine the errors associated with the signal Δt resolutions, tagging efficiencies, and mistag rates, and published measurements [14] for τ_B and Δm_d . All of these sum to 0.013 (0.014) for $S_{\eta' K_S^0}$ ($C_{\eta' K_S^0}$). The contributions from the m_{ES} , ΔE , $m_{\eta'}$, and \mathcal{F} PDFs are 0.025 (0.014), for $S_{\eta' K_S^0}$ ($C_{\eta' K_S^0}$). We take systematic uncertainties due to SVT alignment (0.01), beam spot (0.01), boost, and z scale (negligible) from previous determinations of these effects [1]. We estimate an uncertainty in $C_{\eta' K_S^0}$ of 0.025 from the effect on some tagside B decays of the interference between the CKM-suppressed $b \rightarrow \bar{u}cd$ amplitude with that of the favored $b \rightarrow c\bar{u}d$ [15].

We have reconstructed about 800 events of $B^+ \rightarrow \eta' K^+$ and 200 of $B^0 \rightarrow \eta' K_S^0$ with which we have measured the branching fractions, the time-integrated charge asymmetry \mathcal{A}_{ch} , and the time-dependent asymmetry parameters $S_{\eta' K_S^0}$ and $C_{\eta' K_S^0}$. We find $S_{\eta' K_S^0} = 0.02 \pm 0.34 \pm 0.03$ and $C_{\eta' K_S^0} = 0.10 \pm 0.22 \pm 0.04$. These are in agreement with a previous measurement by the Belle Collaboration [16]. A nonzero value of $C_{\eta' K_S^0}$ would indicate direct CP nonconservation in the $B^0 \rightarrow \eta' K_S^0$ decay. With $C_{\eta' K_S^0} = 0$,




FIG. 2 (color online). Projections onto Δt for $B^0 \rightarrow \eta' K_S^0$ data (points with errors), the fit function (solid line), and background function (dashed line), for (a) B^0 and (b) \bar{B}^0 tagged events, and (c) the asymmetry between B^0 and \bar{B}^0 tags.

and provided the decay is dominated by amplitudes with a single weak phase, $S_{\eta' K_S^0}$ is equal to $\sin 2\beta$. Our result for $S_{\eta' K_S^0}$ is about 2 standard deviations smaller than the value obtained with $B^0 \rightarrow J/\psi K_S^0$ [1,2], and consistent with zero.

The measured branching fractions are $\mathcal{B}(B^+ \rightarrow \eta' K^+) = (76.9 \pm 3.5 \pm 4.4) \times 10^{-6}$ and $\mathcal{B}(B^0 \rightarrow \eta' K^0) = (60.6 \pm 5.6 \pm 4.6) \times 10^{-6}$, and we find $\mathcal{A}_{\text{ch}} = 0.037 \pm 0.045 \pm 0.011$. The null result for \mathcal{A}_{ch} represents a limit on direct CP nonconservation in $B^+ \rightarrow \eta' K^+$; the 90% C.L. limit range is $[-0.04, 0.11]$, and is consistent with predictions [5]. These values supersede our previous measurements [11], and are more than a factor of 2 more precise than previous results [11,17]. The branching fractions depend on $R_{+/0} \equiv \mathcal{B}(Y(4S) \rightarrow B^+ B^-)/\mathcal{B}(Y(4S) \rightarrow B^0 \bar{B}^0)$, which we have assumed to be unity. To compare the decay rates we form their ratio, making use of measurements [18] of $r_{+/0} \equiv R_{+/0} \times \tau(B^+)/\tau(B^0) = 1.14 \pm 0.06$ (our average); we find

$$\frac{\Gamma(B^+ \rightarrow \eta' K^+)}{\Gamma(B^0 \rightarrow \eta' K^0)} = 1.12 \pm 0.13 \pm 0.06 \pm 0.06,$$

where the last error is from $r_{+/0}$.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), IHEP (China), CEA and

CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

*Also with Università di Perugia, Perugia, Italy.

[†]Also with Università della Basilicata, Potenza, Italy.

[‡]Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain.

[§]Deceased.

- [1] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 201802 (2002).
- [2] *Belle* Collaboration, K. Abe *et al.*, Phys. Rev. D **66**, 071102(R) (2002).
- [3] See, for example, G. Kou and A. I. Sanda, Phys. Lett. B **525**, 240 (2002) for a recent perturbative QCD prediction.
- [4] C-W. Chiang and J. L. Rosner, Phys. Rev. D **65**, 074035 (2002), and references therein; M-Z. Yang and Y-D. Yang, Nucl. Phys. **B609**, 469 (2001).
- [5] M. Beneke and M. Neubert, Nucl. Phys. **B651**, 225 (2003).
- [6] E.V. Shuryak and A. Zhitnitsky, Phys. Rev. D **57**, 2001 (1998).

- [7] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [8] PEP-II Conceptual Design Report, SLAC-R-418, 1993.
- [9] D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
- [10] Except as noted explicitly, we use a particle name to denote either member of a charge conjugate pair.
- [11] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 221802 (2001); Phys. Rev. D **65**, 051101 (2002).
- [12] The *BABAR* detector simulation is based on GEANT 4, S. Agostinelli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [13] CLEO Collaboration, D. M. Asner *et al.*, Phys. Rev. D **53**, 1039 (1996).
- [14] Particle Data Group, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).
- [15] O. Long *et al.*, hep-ex/0303030 [Phys. Rev. D (to be published)].
- [16] *Belle* Collaboration, K.-F. Chen *et al.*, Phys. Lett. B **546**, 196 (2002).
- [17] CLEO Collaboration, S. J. Richichi *et al.*, Phys. Rev. Lett. **85**, 520 (2000); *Belle* Collaboration, K. Abe *et al.*, Phys. Lett. B **517**, 3 (2001).
- [18] CLEO Collaboration, J. P. Alexander *et al.*, Phys. Rev. Lett. **86**, 2737 (2001); *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **65**, 032001 (2002); see also R. Kaiser, A.V. Manohar, and T. Mehen, Phys. Rev. Lett. **90**, 142001 (2003); M. B. Voloshin, hep-ph/0301076.