

## Measurements of $CP$ -Violating Asymmetries in $B^0 \rightarrow K_s^0 \pi^0$ Decays

B. Aubert,<sup>1</sup> R. Barate,<sup>1</sup> D. Boutigny,<sup>1</sup> F. Couderc,<sup>1</sup> J.-M. Gaillard,<sup>1</sup> A. Hicheur,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup> A. Palano,<sup>2</sup> A. Pompili,<sup>2</sup> J. C. Chen,<sup>3</sup> N. D. Qi,<sup>3</sup> G. Rong,<sup>3</sup> P. Wang,<sup>3</sup> Y. S. Zhu,<sup>3</sup> G. Eigen,<sup>4</sup> I. Ofte,<sup>4</sup> B. Stugu,<sup>4</sup> G. S. Abrams,<sup>5</sup> A. W. Borgland,<sup>5</sup> A. B. Breon,<sup>5</sup> D. N. Brown,<sup>5</sup> J. Button-Shafer,<sup>5</sup> R. N. Cahn,<sup>5</sup> E. Charles,<sup>5</sup> C. T. Day,<sup>5</sup> M. S. Gill,<sup>5</sup> A. V. Gritsan,<sup>5</sup> Y. Groysman,<sup>5</sup> R. G. Jacobsen,<sup>5</sup> R. W. Kadel,<sup>5</sup> J. Kadyk,<sup>5</sup> L. T. Kerth,<sup>5</sup> Yu. G. Kolomensky,<sup>5</sup> G. Kukartsev,<sup>5</sup> C. LeClerc,<sup>5</sup> G. Lynch,<sup>5</sup> A. M. Merchant,<sup>5</sup> L. M. Mir,<sup>5</sup> P. J. Oddone,<sup>5</sup> T. J. Orimoto,<sup>5</sup> M. Pripstein,<sup>5</sup> N. A. Roe,<sup>5</sup> M. T. Ronan,<sup>5</sup> V. G. Shelkov,<sup>5</sup> A. V. Telnov,<sup>5</sup> W. A. Wenzel,<sup>5</sup> K. Ford,<sup>6</sup> T. J. Harrison,<sup>6</sup> C. M. Hawkes,<sup>6</sup> S. E. Morgan,<sup>6</sup> A. T. Watson,<sup>6</sup> M. Fritsch,<sup>7</sup> K. Goetzen,<sup>7</sup> T. Held,<sup>7</sup> H. Koch,<sup>7</sup> B. Lewandowski,<sup>7</sup> M. Pelizaeus,<sup>7</sup> M. Steinke,<sup>7</sup> J. T. Boyd,<sup>8</sup> N. Chevalier,<sup>8</sup> W. N. Cottingham,<sup>8</sup> M. P. Kelly,<sup>8</sup> T. E. Latham,<sup>8</sup> F. F. Wilson,<sup>8</sup> T. Cuhadar-Donszelmann,<sup>9</sup> C. Hearty,<sup>9</sup> T. S. Mattison,<sup>9</sup> J. A. McKenna,<sup>9</sup> D. Thiessen,<sup>9</sup> P. Kyberd,<sup>10</sup> L. Teodorescu,<sup>10</sup> V. E. Blinov,<sup>11</sup> A. D. Bukin,<sup>11</sup> V. P. Druzhinin,<sup>11</sup> V. B. Golubev,<sup>11</sup> V. N. Ivanchenko,<sup>11</sup> E. A. Kravchenko,<sup>11</sup> A. P. Onuchin,<sup>11</sup> S. I. Serednyakov,<sup>11</sup> Yu. I. Skovpen,<sup>11</sup> E. P. Solodov,<sup>11</sup> A. N. Yushkov,<sup>11</sup> D. Best,<sup>12</sup> M. Bruinsma,<sup>12</sup> M. Chao,<sup>12</sup> I. Eschrich,<sup>12</sup> D. Kirkby,<sup>12</sup> A. J. Lankford,<sup>12</sup> M. Mandelkern,<sup>12</sup> R. K. Mommsen,<sup>12</sup> W. Roethel,<sup>12</sup> D. P. Stoker,<sup>12</sup> C. Buchanan,<sup>13</sup> B. L. Hartfiel,<sup>13</sup> J. W. Gary,<sup>14</sup> B. C. Shen,<sup>14</sup> K. Wang,<sup>14</sup> D. del Re,<sup>15</sup> H. K. Hadavand,<sup>15</sup> E. J. Hill,<sup>15</sup> D. B. MacFarlane,<sup>15</sup> H. P. Paar,<sup>15</sup> Sh. Rahatlou,<sup>15</sup> V. Sharma,<sup>15</sup> J. W. Berryhill,<sup>16</sup> C. Campagnari,<sup>16</sup> B. Dahmes,<sup>16</sup> S. L. Levy,<sup>16</sup> O. Long,<sup>16</sup> A. Lu,<sup>16</sup> M. A. Mazur,<sup>16</sup> J. D. Richman,<sup>16</sup> W. Verkerke,<sup>16</sup> T. W. Beck,<sup>17</sup> A. M. Eisner,<sup>17</sup> C. A. Heusch,<sup>17</sup> W. S. Lockman,<sup>17</sup> T. Schalk,<sup>17</sup> R. E. Schmitz,<sup>17</sup> B. A. Schumm,<sup>17</sup> A. Seiden,<sup>17</sup> P. Spradlin,<sup>17</sup> D. C. Williams,<sup>17</sup> M. G. Wilson,<sup>17</sup> J. Albert,<sup>18</sup> E. Chen,<sup>18</sup> G. P. Dubois-Felsmann,<sup>18</sup> A. Dvoretzki,<sup>18</sup> D. G. Hitlin,<sup>18</sup> I. Narsky,<sup>18</sup> T. Piatenko,<sup>18</sup> F. C. Porter,<sup>18</sup> A. Ryd,<sup>18</sup> A. Samuel,<sup>18</sup> S. Yang,<sup>18</sup> S. Jayatilake,<sup>19</sup> G. Mancinelli,<sup>19</sup> B. T. Meadows,<sup>19</sup> M. D. Sokoloff,<sup>19</sup> T. Abe,<sup>20</sup> F. Blanc,<sup>20</sup> P. Bloom,<sup>20</sup> S. Chen,<sup>20</sup> P. J. Clark,<sup>20</sup> W. T. Ford,<sup>20</sup> U. Nauenberg,<sup>20</sup> A. Olivas,<sup>20</sup> P. Rankin,<sup>20</sup> J. G. Smith,<sup>20</sup> L. Zhang,<sup>20</sup> A. Chen,<sup>21</sup> J. L. Harton,<sup>21</sup> A. Soffer,<sup>21</sup> W. H. Toki,<sup>21</sup> R. J. Wilson,<sup>21</sup> Q. L. Zeng,<sup>21</sup> D. Altenburg,<sup>22</sup> T. Brandt,<sup>22</sup> J. Brose,<sup>22</sup> T. Colberg,<sup>22</sup> M. Dickopp,<sup>22</sup> E. Feltresi,<sup>22</sup> A. Hauke,<sup>22</sup> H. M. Lacker,<sup>22</sup> E. Maly,<sup>22</sup> R. Müller-Pfefferkorn,<sup>22</sup> R. Nogowski,<sup>22</sup> S. Otto,<sup>22</sup> A. Petzold,<sup>22</sup> J. Schubert,<sup>22</sup> K. R. Schubert,<sup>22</sup> R. Schwierz,<sup>22</sup> B. Spaan,<sup>22</sup> J. E. Sundermann,<sup>22</sup> D. Bernard,<sup>23</sup> G. R. Bonneaud,<sup>23</sup> F. Brochard,<sup>23</sup> P. Grenier,<sup>23</sup> S. Schrenk,<sup>23</sup> Ch. Thiebaux,<sup>23</sup> G. Vasileiadis,<sup>23</sup> M. Verderi,<sup>23</sup> D. J. Bard,<sup>24</sup> A. Khan,<sup>24</sup> D. Lavin,<sup>24</sup> F. Muheim,<sup>24</sup> S. Playfer,<sup>24</sup> M. Andreotti,<sup>25</sup> V. Azzolini,<sup>25</sup> D. Bettoni,<sup>25</sup> C. Bozzi,<sup>25</sup> R. Calabrese,<sup>25</sup> G. Cibinetto,<sup>25</sup> E. Luppi,<sup>25</sup> M. Negrini,<sup>25</sup> L. Piemontese,<sup>25</sup> A. Sarti,<sup>25</sup> E. Treadwell,<sup>26</sup> R. Baldini-Ferroli,<sup>27</sup> A. Calcaterra,<sup>27</sup> R. de Sangro,<sup>27</sup> G. Finocchiaro,<sup>27</sup> P. Patteri,<sup>27</sup> M. Piccolo,<sup>27</sup> A. Zallo,<sup>27</sup> A. Buzzo,<sup>28</sup> R. Capra,<sup>28</sup> R. Contri,<sup>28</sup> G. Crosetti,<sup>28</sup> M. Lo Vetere,<sup>28</sup> M. Macri,<sup>28</sup> M. R. Monge,<sup>28</sup> S. Passaggio,<sup>28</sup> C. Patrignani,<sup>28</sup> E. Robutti,<sup>28</sup> A. Santroni,<sup>28</sup> S. Tosi,<sup>28</sup> S. Bailey,<sup>29</sup> G. Brandenburg,<sup>29</sup> M. Morii,<sup>29</sup> E. Won,<sup>29</sup> R. S. Dubitzky,<sup>30</sup> U. Langenegger,<sup>30</sup> W. Bhimji,<sup>31</sup> D. A. Bowerman,<sup>31</sup> P. D. Dauncey,<sup>31</sup> U. Egede,<sup>31</sup> J. R. Gaillard,<sup>31</sup> G. W. Morton,<sup>31</sup> J. A. Nash,<sup>31</sup> G. P. Taylor,<sup>31</sup> G. J. Grenier,<sup>32</sup> U. Mallik,<sup>32</sup> J. Cochran,<sup>33</sup> H. B. Crawley,<sup>33</sup> J. Lamsa,<sup>33</sup> W. T. Meyer,<sup>33</sup> S. Prell,<sup>33</sup> E. I. Rosenberg,<sup>33</sup> J. Yi,<sup>33</sup> M. Davier,<sup>34</sup> G. Grosdidier,<sup>34</sup> A. Höcker,<sup>34</sup> S. Laplace,<sup>34</sup> F. Le Diberder,<sup>34</sup> V. Lepeltier,<sup>34</sup> A. M. Lutz,<sup>34</sup> T. C. Petersen,<sup>34</sup> S. Plaszczynski,<sup>34</sup> M. H. Schune,<sup>34</sup> L. Tantot,<sup>34</sup> G. Wormser,<sup>34</sup> C. H. Cheng,<sup>35</sup> D. J. Lange,<sup>35</sup> M. C. Simani,<sup>35</sup> D. M. Wright,<sup>35</sup> A. J. Bevan,<sup>36</sup> J. P. Coleman,<sup>36</sup> J. R. Fry,<sup>36</sup> E. Gabathuler,<sup>36</sup> R. Gamet,<sup>36</sup> R. J. Parry,<sup>36</sup> D. J. Payne,<sup>36</sup> R. J. Sloane,<sup>36</sup> C. Touramanis,<sup>36</sup> J. J. Back,<sup>37</sup> C. M. Cormack,<sup>37</sup> P. F. Harrison,<sup>37,\*</sup> G. B. Mohanty,<sup>37</sup> C. L. Brown,<sup>38</sup> G. Cowan,<sup>38</sup> R. L. Flack,<sup>38</sup> H. U. Flaecher,<sup>38</sup> M. G. Green,<sup>38</sup> C. E. Marker,<sup>38</sup> T. R. McMahon,<sup>38</sup> S. Ricciardi,<sup>38</sup> F. Salvatore,<sup>38</sup> G. Vaitsas,<sup>38</sup> M. A. Winter,<sup>38</sup> D. Brown,<sup>39</sup> C. L. Davis,<sup>39</sup> J. Allison,<sup>40</sup> N. R. Barlow,<sup>40</sup> R. J. Barlow,<sup>40</sup> P. A. Hart,<sup>40</sup> M. C. Hodgkinson,<sup>40</sup> G. D. Lafferty,<sup>40</sup> A. J. Lyon,<sup>40</sup> J. C. Williams,<sup>40</sup> A. Farbin,<sup>41</sup> W. D. Hulsbergen,<sup>41</sup> A. Jawahery,<sup>41</sup> D. Kovalskyi,<sup>41</sup> C. K. Lae,<sup>41</sup> V. Lillard,<sup>41</sup> D. A. Roberts,<sup>41</sup> G. Blaylock,<sup>42</sup> C. Dallapiccola,<sup>42</sup> K. T. Flood,<sup>42</sup> S. S. Hertzbach,<sup>42</sup> R. Kofler,<sup>42</sup> V. B. Koptchev,<sup>42</sup> T. B. Moore,<sup>42</sup> S. Saremi,<sup>42</sup> H. Staengle,<sup>42</sup> S. Willocq,<sup>42</sup> R. Cowan,<sup>43</sup> G. Sciolla,<sup>43</sup> F. Taylor,<sup>43</sup> R. K. Yamamoto,<sup>43</sup> D. J. J. Mangeol,<sup>44</sup> P. M. Patel,<sup>44</sup> S. H. Robertson,<sup>44</sup> A. Lazzaro,<sup>45</sup> F. Palombo,<sup>45</sup> J. M. Bauer,<sup>46</sup> L. Cremaldi,<sup>46</sup> V. Eschenburg,<sup>46</sup> R. Godang,<sup>46</sup> R. Kroeger,<sup>46</sup> J. Reidy,<sup>46</sup> D. A. Sanders,<sup>46</sup> D. J. Summers,<sup>46</sup> H. W. Zhao,<sup>46</sup> S. Brunet,<sup>47</sup> D. Côté,<sup>47</sup> P. Taras,<sup>47</sup> H. Nicholson,<sup>48</sup> N. Cavallo,<sup>49</sup> F. Fabozzi,<sup>49,†</sup> C. Gatto,<sup>49</sup> L. Lista,<sup>49</sup> D. Monorchio,<sup>49</sup> P. Paolucci,<sup>49</sup> D. Piccolo,<sup>49</sup> C. Sciacca,<sup>49</sup> M. Baak,<sup>50</sup> H. Bulten,<sup>50</sup> G. Raven,<sup>50</sup> L. Wilden,<sup>50</sup> C. P. Jessop,<sup>51</sup> J. M. LoSecco,<sup>51</sup> T. A. Gabriel,<sup>52</sup> T. Allmendinger,<sup>53</sup> B. Brau,<sup>53</sup> K. K. Gan,<sup>53</sup> K. Honscheid,<sup>53</sup> D. Hufnagel,<sup>53</sup> H. Kagan,<sup>53</sup> R. Kass,<sup>53</sup> T. Pulliam,<sup>53</sup> A. M. Rahimi,<sup>53</sup> R. Ter-Antonyan,<sup>53</sup> Q. K. Wong,<sup>53</sup> J. Brau,<sup>54</sup> R. Frey,<sup>54</sup> O. Igonkina,<sup>54</sup> C. T. Potter,<sup>54</sup> N. B. Sinev,<sup>54</sup> D. Strom,<sup>54</sup> E. Torrence,<sup>54</sup> F. Colecchia,<sup>55</sup> A. Dorigo,<sup>55</sup> F. Galeazzi,<sup>55</sup> M. Margoni,<sup>55</sup> M. Morandin,<sup>55</sup> M. Posocco,<sup>55</sup>

M. Rotondo,<sup>55</sup> F. Simonetto,<sup>55</sup> R. Stroili,<sup>55</sup> G. Tiozzo,<sup>55</sup> C. Voci,<sup>55</sup> M. Benayoun,<sup>56</sup> H. Briand,<sup>56</sup> J. Chauveau,<sup>56</sup> P. David,<sup>56</sup> Ch. de la Vaissière,<sup>56</sup> L. Del Buono,<sup>56</sup> O. Hamon,<sup>56</sup> M. J. J. John,<sup>56</sup> Ph. Leruste,<sup>56</sup> J. Ocariz,<sup>56</sup> M. Pivk,<sup>56</sup> L. Roos,<sup>56</sup> S. T'Jampens,<sup>56</sup> G. Therin,<sup>56</sup> P. F. Manfredi,<sup>57</sup> V. Re,<sup>57</sup> P. K. Behera,<sup>58</sup> L. Gladney,<sup>58</sup> Q. H. Guo,<sup>58</sup> J. Panetta,<sup>58</sup> F. Anulli,<sup>27,59</sup> M. Biasini,<sup>59</sup> I. M. Peruzzi,<sup>27,59</sup> M. Pioppi,<sup>59</sup> C. Angelini,<sup>60</sup> G. Batignani,<sup>60</sup> S. Bettarini,<sup>60</sup> M. Bondioli,<sup>60</sup> F. Bucci,<sup>60</sup> G. Calderini,<sup>60</sup> M. Carpinelli,<sup>60</sup> V. Del Gamba,<sup>60</sup> F. Forti,<sup>60</sup> M. A. Giorgi,<sup>60</sup> A. Lusiani,<sup>60</sup> G. Marchiori,<sup>60</sup> F. Martinez-Vidal,<sup>60,‡</sup> M. Morganti,<sup>60</sup> N. Neri,<sup>60</sup> E. Paoloni,<sup>60</sup> M. Rama,<sup>60</sup> G. Rizzo,<sup>60</sup> F. Sandrelli,<sup>60</sup> J. Walsh,<sup>60</sup> M. Haire,<sup>61</sup> D. Judd,<sup>61</sup> K. Paick,<sup>61</sup> D. E. Wagoner,<sup>61</sup> N. Danielson,<sup>62</sup> P. Elmer,<sup>62</sup> C. Lu,<sup>62</sup> V. Miftakov,<sup>62</sup> J. Olsen,<sup>62</sup> A. J. S. Smith,<sup>62</sup> F. Bellini,<sup>63</sup> G. Cavoto,<sup>62,63</sup> R. Faccini,<sup>63</sup> F. Ferrarotto,<sup>63</sup> F. Ferroni,<sup>63</sup> M. Gaspero,<sup>63</sup> L. Li Gioi,<sup>63</sup> M. A. Mazzone,<sup>63</sup> S. Morganti,<sup>63</sup> M. Pierini,<sup>63</sup> G. Piredda,<sup>63</sup> F. Safai Tehrani,<sup>63</sup> C. Voena,<sup>63</sup> S. Christ,<sup>64</sup> G. Wagner,<sup>64</sup> R. Waldi,<sup>64</sup> T. Adye,<sup>65</sup> N. De Groot,<sup>65</sup> B. Franek,<sup>65</sup> N. I. Geddes,<sup>65</sup> G. P. Gopal,<sup>65</sup> E. O. Olaiya,<sup>65</sup> R. Aleksan,<sup>66</sup> S. Emery,<sup>66</sup> A. Gaidot,<sup>66</sup> S. F. Ganzhur,<sup>66</sup> P.-F. Giraud,<sup>66</sup> G. Hamel de Monchenault,<sup>66</sup> W. Kozanecki,<sup>66</sup> M. Langer,<sup>66</sup> M. Legendre,<sup>66</sup> G. W. London,<sup>66</sup> B. Mayer,<sup>66</sup> G. Schott,<sup>66</sup> G. Vasseur,<sup>66</sup> Ch. Yèche,<sup>66</sup> M. Zito,<sup>66</sup> M. V. Purohit,<sup>67</sup> A. W. Weidemann,<sup>67</sup> F. X. Yumiceva,<sup>67</sup> D. Aston,<sup>68</sup> R. Bartoldus,<sup>68</sup> N. Berger,<sup>68</sup> A. M. Boyarski,<sup>68</sup> O. L. Buchmueller,<sup>68</sup> M. R. Convery,<sup>68</sup> M. Cristinziani,<sup>68</sup> G. De Nardo,<sup>68</sup> D. Dong,<sup>68</sup> J. Dorfan,<sup>68</sup> D. Dujmic,<sup>68</sup> W. Dunwoodie,<sup>68</sup> E. E. Elsen,<sup>68</sup> S. Fan,<sup>68</sup> R. C. Field,<sup>68</sup> T. Glanzman,<sup>68</sup> S. J. Gowdy,<sup>68</sup> T. Hadig,<sup>68</sup> V. Halyo,<sup>68</sup> C. Hast,<sup>68</sup> T. Hryn'ova,<sup>68</sup> W. R. Innes,<sup>68</sup> M. H. Kelsey,<sup>68</sup> P. Kim,<sup>68</sup> M. L. Kocian,<sup>68</sup> D. W. G. S. Leith,<sup>68</sup> J. Libby,<sup>68</sup> S. Luitz,<sup>68</sup> V. Luth,<sup>68</sup> H. L. Lynch,<sup>68</sup> H. Marsiske,<sup>68</sup> R. Messner,<sup>68</sup> D. R. Muller,<sup>68</sup> C. P. O'Grady,<sup>68</sup> V. E. Ozcan,<sup>68</sup> A. Perazzo,<sup>68</sup> M. Perl,<sup>68</sup> S. Petrak,<sup>68</sup> B. N. Ratcliff,<sup>68</sup> A. Roodman,<sup>68</sup> A. A. Salnikov,<sup>68</sup> R. H. Schindler,<sup>68</sup> J. Schwiening,<sup>68</sup> G. Simi,<sup>68</sup> A. Snyder,<sup>68</sup> A. Soha,<sup>68</sup> J. Stelzer,<sup>68</sup> D. Su,<sup>68</sup> M. K. Sullivan,<sup>68</sup> J. Va'vra,<sup>68</sup> S. R. Wagner,<sup>68</sup> M. Weaver,<sup>68</sup> A. J. R. Weinstein,<sup>68</sup> W. J. Wisniewski,<sup>68</sup> M. Wittgen,<sup>68</sup> D. H. Wright,<sup>68</sup> A. K. Yarritu,<sup>68</sup> C. C. Young,<sup>68</sup> P. R. Burchat,<sup>69</sup> A. J. Edwards,<sup>69</sup> T. I. Meyer,<sup>69</sup> B. A. Petersen,<sup>69</sup> C. Roat,<sup>69</sup> S. Ahmed,<sup>70</sup> M. S. Alam,<sup>70</sup> J. A. Ernst,<sup>70</sup> M. A. Saeed,<sup>70</sup> M. Saleem,<sup>70</sup> F. R. Wappler,<sup>70</sup> W. Bugg,<sup>71</sup> M. Krishnamurthy,<sup>71</sup> S. M. Spanier,<sup>71</sup> R. Eckmann,<sup>72</sup> H. Kim,<sup>72</sup> J. L. Ritchie,<sup>72</sup> A. Satpathy,<sup>72</sup> R. F. Schwitters,<sup>72</sup> J. M. Izen,<sup>73</sup> I. Kitayama,<sup>73</sup> X. C. Lou,<sup>73</sup> S. Ye,<sup>73</sup> F. Bianchi,<sup>74</sup> M. Bona,<sup>74</sup> F. Gallo,<sup>74</sup> D. Gamba,<sup>74</sup> C. Borean,<sup>75</sup> L. Bosisio,<sup>75</sup> C. Cartaro,<sup>75</sup> F. Cossutti,<sup>75</sup> G. Della Ricca,<sup>75</sup> S. Dittongo,<sup>75</sup> S. Grancagnolo,<sup>75</sup> L. Lanceri,<sup>75</sup> P. Poropat,<sup>75,§</sup> L. Vitale,<sup>75</sup> G. Vuagnin,<sup>75</sup> R. S. Panvini,<sup>76</sup> Sw. Banerjee,<sup>77</sup> C. M. Brown,<sup>77</sup> D. Fortin,<sup>77</sup> P. D. Jackson,<sup>77</sup> R. Kowalewski,<sup>77</sup> J. M. Roney,<sup>77</sup> H. R. Band,<sup>78</sup> S. Dasu,<sup>78</sup> M. Datta,<sup>78</sup> A. M. Eichenbaum,<sup>78</sup> J. J. Hollar,<sup>78</sup> J. R. Johnson,<sup>78</sup> P. E. Kutter,<sup>78</sup> H. Li,<sup>78</sup> R. Liu,<sup>78</sup> F. Di Lodovico,<sup>78</sup> A. Mihalyi,<sup>78</sup> A. K. Mohapatra,<sup>78</sup> Y. Pan,<sup>78</sup> R. Prepost,<sup>78</sup> S. J. Sekula,<sup>78</sup> P. Tan,<sup>78</sup> J. H. von Wimmersperg-Toeller,<sup>78</sup> J. Wu,<sup>78</sup> S. L. Wu,<sup>78</sup> Z. Yu,<sup>78</sup> and H. Neal<sup>79</sup>

(BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

<sup>2</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

<sup>3</sup>Institute of High Energy Physics, Beijing 100039, China

<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

<sup>6</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

<sup>8</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>9</sup>University of British Columbia, Vancouver, BC, Canada V6T 1Z1

<sup>10</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>11</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>12</sup>University of California at Irvine, Irvine, California 92697, USA

<sup>13</sup>University of California at Los Angeles, Los Angeles, California 90024, USA

<sup>14</sup>University of California at Riverside, Riverside, California 92521, USA

<sup>15</sup>University of California at San Diego, La Jolla, California 92093, USA

<sup>16</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA

<sup>17</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

<sup>18</sup>California Institute of Technology, Pasadena, California 91125, USA

<sup>19</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA

<sup>20</sup>University of Colorado, Boulder, Colorado 80309, USA

<sup>21</sup>Colorado State University, Fort Collins, Colorado 80523, USA

<sup>22</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

<sup>23</sup>Ecole Polytechnique, LLR, F-91128 Palaiseau, France

<sup>24</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

- <sup>25</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy  
<sup>26</sup>Florida A&M University, Tallahassee, Florida 32307, USA  
<sup>27</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy  
<sup>28</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy  
<sup>29</sup>Harvard University, Cambridge, Massachusetts 02138, USA  
<sup>30</sup>Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany  
<sup>31</sup>Imperial College London, London, SW7 2AZ, United Kingdom  
<sup>32</sup>University of Iowa, Iowa City, Iowa 52242, USA  
<sup>33</sup>Iowa State University, Ames, Iowa 50011-3160, USA  
<sup>34</sup>Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France  
<sup>35</sup>Lawrence Livermore National Laboratory, Livermore, California 94550, USA  
<sup>36</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom  
<sup>37</sup>Queen Mary, University of London, E1 4NS, United Kingdom  
<sup>38</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom  
<sup>39</sup>University of Louisville, Louisville, Kentucky 40292, USA  
<sup>40</sup>University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>41</sup>University of Maryland, College Park, Maryland 20742, USA  
<sup>42</sup>University of Massachusetts, Amherst, Massachusetts 01003, USA  
<sup>43</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA  
<sup>44</sup>McGill University, Montréal, QC, Canada H3A 2T8  
<sup>45</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy  
<sup>46</sup>University of Mississippi, University, Mississippi 38677, USA  
<sup>47</sup>Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, QC, Canada H3C 3J7  
<sup>48</sup>Mount Holyoke College, South Hadley, Massachusetts 01075, USA  
<sup>49</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy  
<sup>50</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands  
<sup>51</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA  
<sup>52</sup>Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, USA  
<sup>53</sup>Ohio State University, Columbus, Ohio 43210, USA  
<sup>54</sup>University of Oregon, Eugene, Oregon 97403, USA  
<sup>55</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy  
<sup>56</sup>Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France  
<sup>57</sup>Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy  
<sup>58</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA  
<sup>59</sup>Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy  
<sup>60</sup>Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy  
<sup>61</sup>Prairie View A&M University, Prairie View, Texas 77446, USA  
<sup>62</sup>Princeton University, Princeton, New Jersey 08544, USA  
<sup>63</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy  
<sup>64</sup>Universität Rostock, D-18051 Rostock, Germany  
<sup>65</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom  
<sup>66</sup>DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France  
<sup>67</sup>University of South Carolina, Columbia, South Carolina 29208, USA  
<sup>68</sup>Stanford Linear Accelerator Center, Stanford, California 94309, USA  
<sup>69</sup>Stanford University, Stanford, California 94305-4060, USA  
<sup>70</sup>State University of New York, Albany, New York 12222, USA  
<sup>71</sup>University of Tennessee, Knoxville, Tennessee 37996, USA  
<sup>72</sup>University of Texas at Austin, Austin, Texas 78712, USA  
<sup>73</sup>University of Texas at Dallas, Richardson, Texas 75083, USA  
<sup>74</sup>Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy  
<sup>75</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy  
<sup>76</sup>Vanderbilt University, Nashville, Tennessee 37235, USA  
<sup>77</sup>University of Victoria, Victoria, BC, Canada V8W 3P6  
<sup>78</sup>University of Wisconsin, Madison, Wisconsin 53706, USA  
<sup>79</sup>Yale University, New Haven, Connecticut 06511, USA

(Received 27 February 2004; published 23 September 2004)

We present a measurement of the time-dependent  $CP$ -violating (CPV) asymmetries in  $B^0 \rightarrow K_S^0 \pi^0$  decays based on  $124 \times 10^6$   $Y(4S) \rightarrow B\bar{B}$  decays collected with the BABAR detector at the PEP-II asymmetric-energy  $B$  factory at SLAC. In a sample containing  $122 \pm 16$  signal decays, we obtain the magnitudes of the direct CPV asymmetry  $C_{K_S^0 \pi^0} = 0.40_{-0.28}^{+0.27} \pm 0.09$  and of the CPV asymmetry in the

interference between mixing and decay  $S_{K_S^0 \pi^0} = 0.48_{-0.47}^{+0.38} \pm 0.06$  where the first error is statistical and the second systematic.

DOI: 10.1103/PhysRevLett.93.131805

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

The *BABAR* [1] and Belle [2] Collaborations recently reported observation of  $CP$  violation in  $B$  meson decays through measurements of the time-dependent  $CP$ -violating (CPV) asymmetry in  $B^0$  decays into charmium final states. In the framework of the Standard Model (SM), where  $CP$  violation is a consequence of the presence of a complex phase in the Cabibbo-Kobayashi-Maskawa (CKM) quark mixing matrix [3], these measurements determine the parameter  $\sin 2\beta$ , with  $\beta \equiv \arg(-V_{cd}V_{cb}^*/V_{td}V_{tb}^*)$ . The consistency of the observed value of  $\sin 2\beta$  with the Standard Model expectations provides strong evidence that the CKM mechanism is the dominant source of  $CP$  violation in the quark sector. A major goal of the experimental studies of  $B$  decays is to provide additional information to examine the validity of this conclusion and search for evidence of new physics (NP) in possible deviations from the SM. One avenue for the observation of NP is provided by  $CP$  violation studies of decays dominated by penguin loop-level  $b \rightarrow s\bar{q}q$  ( $q = \{d, s\}$ ) transitions [4,5]. While in the SM the time-dependent CPV asymmetries in these decays measure  $\sin 2\beta$ , additional radiative loop contributions from NP processes may alter this expectation. Presently, the  $B$  factory experiments have explored time-dependent CPV asymmetries in three such decays, which in the SM are dominated by the penguin  $b \rightarrow s\bar{s}s$  transition:  $B^0 \rightarrow \eta' K_S^0$  [6,7],  $B^0 \rightarrow K^+ K^- K_S^0$  [6], and  $B^0 \rightarrow \phi K_S^0$  [6,8]. The latter results hint at a possible deviation from the SM, but are inconclusive.

In this Letter we present the first measurement of the time-dependent CPV asymmetries in the decay  $B^0 \rightarrow K_S^0 \pi^0$ , which has a measured branching fraction  $\mathcal{B}(B^0 \rightarrow K_S^0 \pi^0) = (11.9 \pm 1.5) \times 10^{-6}$  [9]. The CKM and color suppression of the tree-level  $b \rightarrow s\bar{u}u$  transition leads to the expectation that this decay is dominated by a top quark mediated  $b \rightarrow s\bar{d}d$  penguin diagram, which carries a weak phase  $\arg(V_{tb}V_{ts}^*)$ . If other contributions, such as the  $b \rightarrow su\bar{u}$  tree amplitude are ignored, the time-dependent CPV asymmetry is governed by  $\sin 2\beta$  [10]. The deviation from  $\sin 2\beta$  due to standard model contributions with a different weak phase is estimated to be at most 0.2 [11].

The results presented here are based on  $124 \times 10^6$   $Y(4S) \rightarrow B\bar{B}$  decays collected in 1999–2003 with the *BABAR* detector at the PEP-II  $e^+e^-$  collider, located at the Stanford Linear Accelerator Center. The *BABAR* detector, which is fully described in [12], provides charged particle tracking through a combination of a five-layer double-sided silicon micro-strip detector (SVT) and a 40-layer central drift chamber (DCH), both operating in a

1.5 T magnetic field in order to provide momentum measurements. Charged kaon and pion identification is achieved through measurements of particle energy loss ( $dE/dx$ ) in the tracking system and Cherenkov cone angle ( $\theta_c$ ) in a detector of internally reflected Cherenkov light (DIRC). A segmented CsI(Tl) electromagnetic calorimeter (EMC) provides photon detection and electron identification. Finally, the instrumented flux return (IFR) of the magnet allows discrimination of muons from pions.

We search for  $B^0 \rightarrow K_S^0 \pi^0$  decays in hadronic events, which are selected based on charged particle multiplicity and event topology [13]. We reconstruct  $K_S^0 \rightarrow \pi^+ \pi^-$  candidates from pairs of oppositely charged tracks. The two-track combinations must form a vertex with  $\pi^+ \pi^-$  invariant mass within  $3.5\sigma$  of the nominal  $K_S^0$  mass [14] and reconstructed proper lifetime greater than 5 times its uncertainty. We form  $\pi^0 \rightarrow \gamma\gamma$  candidates with an invariant mass  $110 < m_{\gamma\gamma} < 160$  MeV from pairs of photon candidates in the EMC that are isolated from any charged tracks, carry a minimum energy of 30 MeV, and possess the expected lateral shower shapes. Finally, we construct  $B^0 \rightarrow K_S^0 \pi^0$  candidates by combining  $K_S^0$  and  $\pi^0$  candidates in the event. For each  $B$  candidate, two nearly independent kinematic variables are computed, namely, the energy-substituted mass  $m_{ES} = \sqrt{(s/2 + \mathbf{p}_B)^2/E_i^2 - p_B^2}$ , and the energy difference  $\Delta E = E_B^* - \sqrt{s}/2$ . Here,  $(E_i, \mathbf{p}_i)$  is the four-vector of the initial  $e^+e^-$  system,  $\sqrt{s} = \sqrt{E_i^2 - p_i^2}$  is the center-of-mass energy,  $\mathbf{p}_B$  is the reconstructed momentum of the  $B^0$  candidate, and  $E_B^*$  is its energy calculated in the  $e^+e^-$  rest frame. For signal decays, the  $m_{ES}$  distribution peaks near the  $B^0$  mass with a resolution of  $\sim 3.1$  MeV/ $c^2$  and the  $\Delta E$  distribution peaks near zero with a resolution of  $\sim 40$  MeV. Both the  $m_{ES}$  and the  $\Delta E$  distribution exhibit a low-side tail from energy leakage out of the EMC. We select candidates within the window  $5.2 < m_{ES} < 5.29$  GeV/ $c^2$  and  $-150 < \Delta E < 150$  MeV, which includes the signal peak and a “sideband” region for background characterization. For the 1.7% of events with more than one candidate, we select the combination with the smallest  $\chi^2 = \sum_{i=\pi^0, K_S^0} (m_i - m'_i)^2 / \sigma_{m_i}^2$ , where  $m_i$  ( $m'_i$ ) is the measured (nominal) mass and  $\sigma_{m_i}$  is the estimated uncertainty on the mass of particle  $i$ .

For each  $B^0 \rightarrow K_S^0 \pi^0$  candidate, we examine the remaining tracks and neutral candidates in the event to determine if the other  $B$  meson,  $B_{tag}$ , decayed as a  $B^0$  or a  $\bar{B}^0$  (flavor tag). Time-dependent CPV asymmetries are determined by reconstructing the distribution of the difference of the proper decay times,  $\Delta t \equiv t_{CP} - t_{tag}$ , where

the  $t_{CP}$  refers to the signal  $B^0$  and  $t_{\text{tag}}$  to the other  $B$ . At the  $Y(4S)$  resonance, the  $\Delta t$  distribution follows

$$\mathcal{P}_{\overline{B}^0}^{B^0}(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} \times \{1 \pm [S_f \sin(\Delta m_d \Delta t) - C_f \cos(\Delta m_d \Delta t)]\}, \quad (1)$$

where the upper (lower) sign corresponds to  $B_{\text{tag}}$  decaying as  $B^0$  ( $\overline{B}^0$ ),  $\tau$  is the  $B^0$  lifetime averaged over the two mass eigenstates,  $\Delta m_d$  is the mixing frequency,  $C_f$  is the magnitude of direct CPV in the decay to final state  $f$ , and  $S$  the magnitude of CPV in the interference between mixing and decay. For the case of pure penguin dominance, we expect  $S_{K_S^0 \pi^0} = \sin 2\beta$ , and  $C_{K_S^0 \pi^0} = 0$ .

We extract the CPV parameters from an unbinned maximum-likelihood fit to kinematic, event shape, flavor tag, and time structure variables, which are sufficiently independent that we can construct the likelihood from the product of one dimensional probability density functions (PDFs). The PDFs for signal events are parameterized from either more copious fully reconstructed  $B$  decays in data or from simulated samples. For background PDFs we select the functional form from data in the sideband regions of the other observables where backgrounds dominate. We include these regions in the fitted sample and simultaneously extract the parameters of the background PDFs along with the CPV measurements.

The sample of  $B^0 \rightarrow K_S^0 \pi^0$  candidates is dominated by random  $K_S^0 \pi^0$  combinations from  $e^+ e^- \rightarrow q\overline{q}$  ( $q = \{u, d, s, c\}$ ) fragmentation. Monte Carlo studies show that contributions from other  $B$  meson decays can be neglected. We exploit topological observables to discriminate the jetlike  $e^+ e^- \rightarrow q\overline{q}$  events from the more uniformly distributed  $B\overline{B}$  events. In the  $Y(4S)$  rest frame, we compute the angle  $\theta_S^*$  between the sphericity axis [15] of the  $B^0$  candidate and that of the remaining particles in the event. While  $|\cos\theta_S^*|$  is highly peaked near one for  $e^+ e^- \rightarrow q\overline{q}$  events, it is nearly uniformly distributed for  $B\overline{B}$ . We require  $|\cos\theta_S^*| < 0.8$ , eliminating 83% of the background. In addition, we include in the fit a Fisher discriminant variable, which is defined as  $\mathcal{F} = 0.53 - 0.60L_0 + 1.27L_2$ , where  $L_j \equiv \sum_i |\mathbf{p}_i^*| |\cos\theta_i^*|^j$ ,  $\mathbf{p}_i^*$  is the momentum of particle  $i$ , and  $\theta_i^*$  is the angle between  $\mathbf{p}_i^*$  and the sphericity axis of the  $B^0$  candidate.

We use a neural network (NN) to determine the flavor of the  $B_{\text{tag}}$  meson from kinematic and particle identification information [16]. Each event is assigned to one of five mutually exclusive tagging categories, designed to combine flavor tags with similar performance and  $\Delta t$  resolution. We parameterize the performance of this algorithm in a data sample ( $B_{\text{flav}}$ ) of fully reconstructed  $B^0 \rightarrow D^{(*)-} \pi^+ / \rho^+ / a_1^+$  decays. The average effective tagging efficiency obtained from this sample is  $Q = \sum_c \epsilon_S^c (1 - 2w^c)^2 = 0.288 \pm 0.005$ , where  $\epsilon_S^c$  and  $w^c$  are the efficiencies and mistag probabilities, respectively, for events

tagged in category  $c \in \{1 \cdots 5\}$ . For the background, the fraction of events ( $\epsilon_B^c$ ) and the asymmetry in the rate of  $B^0$  versus  $\overline{B}^0$  tags in each tagging category are extracted from the fit to the data.

We compute the proper time difference  $\Delta t$  from the known boost of the  $e^+ e^-$  system and the measured  $\Delta z = z_{CP} - z_{\text{tag}}$ , the difference of the reconstructed decay vertex positions of the  $B^0 \rightarrow K_S^0 \pi^0$  and  $B_{\text{tag}}$  candidate along the boost direction ( $z$ ). A description of the inclusive reconstruction of the  $B_{\text{tag}}$  vertex is given in [13]. For the  $B^0 \rightarrow K_S^0 \pi^0$  decay, where no charged particles are present at the decay vertex, we exploit the fact that the flight distance of the  $B$  meson transverse to the beam direction ( $\sim 30 \mu\text{m}$ ) is small compared to the flight length along the beam ( $\sim 260 \mu\text{m}$ ). We then determine the decay point from the intersection of the  $K_S^0$  trajectory with the interaction region by constraining the  $B$  vertex to the interaction point (IP) in the transverse plane. The position and size of the interaction region are determined on a run-by-run basis from the spatial distribution of vertices from two-track events. The uncertainty in the IP position, which follows from the size of the interaction region (about  $200 \mu\text{m}$  horizontal and  $4 \mu\text{m}$  vertical), is combined with the RMS of the transverse  $B$  flight length distribution to assign an uncertainty to the IP constraint.

Simulation studies indicate that the vertexing procedure provides an unbiased estimate of  $z_{CP}$ . The per-event estimate of the  $\Delta t$  error reflects the strong dependence of the  $z_{CP}$  resolution on the  $K_S^0$  flight direction and the number of SVT layers traversed by its decay daughters. For the 37% of events where both tracks include at least one hit in the inner three SVT layers (at radii from 3.2 cm to 5.4 cm), the mean  $\Delta t$  resolution is comparable to that of decays for which the vertex is directly reconstructed from charged particles originating at the  $B$  decay point [13]. If both tracks have hits in the outer two SVT layers (at radii 9.1 cm to 14.4 cm) but one of the tracks has no hits in the inner three layers ( $\sim 27\%$  of the events), the resolution is nearly 2 times worse. The remaining events provide poor  $\Delta t$  measurements. For these events and for events with  $\sigma_{\Delta t} > 2.5$  ps or  $|\Delta t| > 20$  ps, we do not include  $\Delta t$  information in the fit. However, we account for the contribution of these events in the measurement of  $C_{K_S^0 \pi^0}$ .

We obtain the PDF for the time dependence of signal decays from the convolution of Eq. (1) with a resolution function  $\mathcal{R}(\delta t \equiv \Delta t - \Delta t_{\text{true}}, \sigma_{\Delta t})$ . The resolution function is parameterized as the sum of a ‘‘core’’ and a ‘‘tail’’ Gaussian, each with a width and mean proportional to the reconstructed  $\sigma_{\Delta t}$ , and a third Gaussian centered at zero with a fixed width of 8 ps [13]. We have verified in simulation that the parameters of  $\mathcal{R}(\delta t, \sigma_{\Delta t})$  for  $B^0 \rightarrow K_S^0 \pi^0$  decays are similar to those obtained from the  $B_{\text{flav}}$  sample, even though the distributions of  $\sigma_{\Delta t}$  differ

considerably. Therefore, we obtain these parameters from a fit to the  $B_{\text{flav}}$  sample.

To extract the CPV asymmetries, we maximize the logarithm of the likelihood function

$$\begin{aligned} \mathcal{L}(S_f, C_f, N_S, N_B, f_S, f_B, \vec{\alpha}) &= \frac{e^{-(N_S+N_B)}}{(N_S+N_B)!} \times \prod_{i \in \text{w}/\Delta t} [N_S f_S \epsilon_S^c \mathcal{P}_S(\vec{x}_i, \vec{y}_i; S_f, C_f) + N_B f_B \epsilon_B^c \mathcal{P}_B(\vec{x}_i, \vec{y}_i; \vec{\alpha})] \\ &\times \prod_{i \in \text{w}/\Delta t} [N_S(1-f_S) \epsilon_S^c \mathcal{P}'_S(\vec{x}_i; C_f) + N_B(1-f_B) \epsilon_B^c \mathcal{P}'_B(\vec{x}_i; \vec{\alpha})], \end{aligned}$$

where the second (third) factor on the right-hand side is the contribution from events with (without)  $\Delta t$  information. The probabilities  $\mathcal{P}_S$  and  $\mathcal{P}_B$  are products of PDFs for signal ( $S$ ) and background ( $B$ ) hypotheses evaluated for the measurements  $\vec{x}_i = \{m_{\text{ES}}, \Delta E, \mathcal{F}, \text{tag, tagging category}\}$  and  $\vec{y}_i = \{\Delta t, \sigma_{\Delta t}\}$ . Along with the CPV asymmetries  $S_f$  and  $C_f$ , the fit extracts the yields  $N_S$  and  $N_B$ , the fractions of events with  $\Delta t$  information  $f_S$  and  $f_B$ , and the parameters  $\vec{\alpha}$  which describe the background PDFs.

Fitting the data sample of 4179  $B^0 \rightarrow K_S^0 \pi^0$  candidates, we find  $N_S = 122 \pm 16$  signal decays with  $S_{K_S^0 \pi^0} = 0.48_{-0.47}^{+0.38} \pm 0.06$  and  $C_{K_S^0 \pi^0} = 0.40_{-0.28}^{+0.27} \pm 0.09$ , where the uncertainties are statistical and systematic, respectively. The estimated number of signal decays is consistent with our measurement of the branching fraction [17]. The result for  $C_{K_S^0 \pi^0}$  is consistent with a fit that does not employ  $\Delta t$  information. Fixing  $C_{K_S^0 \pi^0} = 0$ , we obtain  $S_{K_S^0 \pi^0} = 0.41_{-0.48}^{+0.41} \pm 0.06$ . The evaluation of the systematic uncertainties is described below.

Figure 1 shows the  $m_{\text{ES}}$  distributions for a signal-enhanced sample. The event selection is based on a likelihood ratio  $R = \mathcal{P}_S / (\mathcal{P}_B + \mathcal{P}_S)$  calculated without the displayed observable. The dashed and solid curves indicate background and signal-plus-background contributions, respectively, as obtained from the fit, but corrected for the selection on  $R$ . Figure 2 shows distri-




FIG. 1. Distribution of  $m_{\text{ES}}$  for events enhanced in signal decays. The dashed and solid curves represent the background and signal-plus-background contributions, respectively.

butions of  $\Delta t$  for  $B^0$ - and  $\bar{B}^0$ -tagged events, and the asymmetry  $\mathcal{A}_{K_S^0 \pi^0}(\Delta t) = [N_{B^0} - N_{\bar{B}^0}] / [N_{B^0} + N_{\bar{B}^0}]$  as a function of  $\Delta t$ , also for a signal-enhanced sample.

To investigate possible biases introduced in the CPV measurements by the IP-constrained vertexing technique, we examine  $B^0 \rightarrow J/\psi K_S^0$  decays in data, where  $J/\psi \rightarrow \mu^+ \mu^-$  and  $J/\psi \rightarrow e^+ e^-$ . In these events we determine  $\Delta t$  in two ways: by fully reconstructing the  $B^0$  decay vertex using the trajectories of charged daughters of the  $J/\psi$  and the  $K_S^0$  mesons, or by neglecting the  $J/\psi$  contribution to the decay vertex and using the IP constraint and the  $K_S^0$  trajectory only. This study shows that within statistical uncertainties, the IP-constrained  $\Delta t$  measurement is unbiased with respect to the more established technique and that the obtained values of  $S_{J/\psi K_S^0}$  and  $C_{J/\psi K_S^0}$  are consistent. A similar study of  $B^\pm \rightarrow K_S^0 \pi^\pm$  events, where the  $\pi^\pm$  contribution to the decay vertex has been replaced by the IP constraint, yields  $S_{K_S^0 \pi^\pm} = 0.13 \pm 0.19$  and  $C_{K_S^0 \pi^\pm} = 0.06 \pm 0.11$ , which is




FIG. 2. Distributions of  $\Delta t$  for events enhanced in signal decays with  $B_{\text{tag}}$  tagged as (a)  $B^0$  or (b)  $\bar{B}^0$ , and (c) the asymmetry  $\mathcal{A}_{K_S^0 \pi^0}(\Delta t)$ . The dashed and solid curves represent the fitted background and signal-plus-background contributions, respectively. The asymmetry projection corresponds to approximately 36 signal and 25 background events.

consistent with the expectation  $S_{K_S^0\pi^\pm} = 0$  and our previous measurement of the charge asymmetry [17]. We also find that the  $B^0$  lifetime measured in  $B^0 \rightarrow K_S^0\pi^0$  decays and in IP-constrained  $B^0 \rightarrow J/\psi K_S^0$  decays agrees with the world average [14].

To quantify possible systematic effects we examine large samples of simulated  $B^0 \rightarrow K_S^0\pi^0$  and  $B^0 \rightarrow J/\psi K_S^0$  decays. We employ the difference in resolution function parameters extracted from these samples to evaluate uncertainties due to the use of the resolution function  $\mathcal{R}$  extracted from the  $B_{\text{flav}}$  sample. We assign a systematic uncertainty of 0.03 on  $S_{K_S^0\pi^0}$  and 0.02 on  $C_{K_S^0\pi^0}$  due to the uncertainty in  $\mathcal{R}$ . We compare fits to a large sample of simulated nominal and IP-constrained  $B^0 \rightarrow J/\psi K_S^0$  events to account for any potential bias due to the vertexing technique. This latter study yields the difference  $\delta S_{J/\psi K_S^0} = 0.04$ , which we assign as the dominant systematic uncertainty on  $S_{K_S^0\pi^0}$ . We include a systematic uncertainty of 0.03 on  $S_{K_S^0\pi^0}$  and 0.01 on  $C_{K_S^0\pi^0}$  to account for a possible misalignment of the SVT. We consider large variations of the IP position and resolution, which we find to have negligible impact. We assign a systematic uncertainty of 0.09 to  $C_{K_S^0\pi^0}$  due to possible asymmetries in the rate of  $B^0$  versus  $\bar{B}^0$  tags in background events. Finally, we include a systematic uncertainty of 0.02 on both  $S_{K_S^0\pi^0}$  and  $C_{K_S^0\pi^0}$  to account for imperfect knowledge of the PDFs used in the fit.

In summary, we have performed a measurement of the time-dependent CPV asymmetries in  $B^0 \rightarrow K_S^0\pi^0$ . These results supersede our previous measurement of  $C_{K_S^0\pi^0}$  [17], which only relied on time-integrated observables, and introduce the first measurement of  $S_{K_S^0\pi^0}$ .

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan

Foundation, Research Corporation, and Alexander von Humboldt Foundation.

\*Now at Department of Physics, University of Warwick, Coventry, United Kingdom

†Also with Università della Basilicata, Potenza, Italy

‡Also with IFIC, Instituto de Física Corpuscular, CSIC-Universidad de Valencia, Valencia, Spain

§Deceased

- [1] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 091801 (2001).
- [2] BELLE Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **87**, 091802 (2001).
- [3] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [4] Y. Grossman and M. P. Worah, Phys. Lett. B **395**, 241 (1997); M. Ciuchini, E. Franco, G. Martinelli, A. Masiero, and L. Silvestrini, Phys. Rev. Lett. **79**, 978 (1997); D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
- [5] Unless explicitly stated, conjugate decay modes are assumed throughout this paper.
- [6] BELLE Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **91**, 261602 (2003).
- [7] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **91**, 161801 (2003).
- [8] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **93**, 071801 (2004).
- [9] Average computed by the Heavy Flavor Averaging Group from results in A. Bornheim *et al.*, Phys. Rev. D **68**, 052002 (2003); B. Aubert *et al.*, Phys. Rev. Lett. **92**, 201802 (2004).
- [10] R. Fleischer, Phys. Lett. B **365**, 399 (1996).
- [11] M. Gronau, Y. Grossman, and J. L. Rosner, Phys. Lett. B **579**, 331 (2004).
- [12] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [13] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **66**, 032003 (2002).
- [14] Particle Data Group Collaboration, K. Hagiwara *et al.*, Phys. Rev. D **66**, 010001 (2002).
- [15] J. D. Bjorken and S. J. Brodsky, Phys. Rev. D **1**, 1416 (1970).
- [16] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **89**, 201802 (2002).
- [17] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **92**, 201802 (2004).