

Differential Production Cross Section of Z Bosons as a Function of Transverse Momentum at $\sqrt{s} = 1.8$ TeV

B. Abbott,⁴⁵ M. Abolins,⁴² V. Abramov,¹⁸ B. S. Acharya,¹¹ I. Adam,⁴⁴ D. L. Adams,⁵⁴ M. Adams,²⁸ S. Ahn,²⁷ V. Akimov,¹⁶ G. A. Alves,² N. Amos,⁴¹ E. W. Anderson,³⁴ M. M. Baarmand,⁴⁷ V. V. Babintsev,¹⁸ L. Babukhadia,²⁰ A. Baden,³⁸ B. Baldin,²⁷ S. Banerjee,¹¹ J. Bantly,⁵¹ E. Barberis,²¹ P. Baringer,³⁵ J. F. Bartlett,²⁷ A. Belyaev,¹⁷ S. B. Beri,⁹ I. Bertram,¹⁹ V. A. Bezzubov,¹⁸ P. C. Bhat,²⁷ V. Bhatnagar,⁹ M. Bhattacharjee,⁴⁷ G. Blazey,²⁹ S. Blessing,²⁵ P. Bloom,²² A. Boehnlein,²⁷ N. I. Bojko,¹⁸ F. Borchering,²⁷ C. Boswell,²⁴ A. Brandt,²⁷ R. Breedon,²² G. Briskin,⁵¹ R. Brock,⁴² A. Bross,²⁷ D. Buchholz,³⁰ V. S. Burtovoi,¹⁸ J. M. Butler,³⁹ W. Carvalho,³ D. Casey,⁴² Z. Casilum,⁴⁷ H. Castilla-Valdez,¹⁴ D. Chakraborty,⁴⁷ K. M. Chan,⁴⁶ S. V. Chekulaev,¹⁸ W. Chen,⁴⁷ D. K. Cho,⁴⁶ S. Choi,¹³ S. Chopra,²⁵ B. C. Choudhary,²⁴ J. H. Christenson,²⁷ M. Chung,²⁸ D. Claes,⁴³ A. R. Clark,²¹ W. G. Cobau,³⁸ J. Cochran,²⁴ L. Coney,³² W. E. Cooper,²⁷ D. Coppage,³⁵ C. Cretsinger,⁴⁶ D. Cullen-Vidal,⁵¹ M. A. C. Cummings,²⁹ D. Cutts,⁵¹ O. I. Dahl,²¹ K. Davis,²⁰ K. De,⁵² K. Del Signore,⁴¹ M. Demarteau,²⁷ D. Denisov,²⁷ S. P. Denisov,¹⁸ H. T. Diehl,²⁷ M. Diesburg,²⁷ G. Di Loreto,⁴² P. Draper,⁵² Y. Ducros,⁸ L. V. Dudko,¹⁷ S. R. Dugad,¹¹ A. Dyshkant,¹⁸ D. Edmunds,⁴² J. Ellison,²⁴ V. D. Elvira,⁴⁷ R. Engelmann,⁴⁷ S. Eno,³⁸ G. Eppley,⁵⁴ P. Ermolov,¹⁷ O. V. Eroshin,¹⁸ J. Estrada,⁴⁶ H. Evans,⁴⁴ V. N. Evdokimov,¹⁸ T. Fahland,²³ M. K. Fatyga,⁴⁶ S. Feher,²⁷ D. Fein,²⁰ T. Ferbel,⁴⁶ H. E. Fisk,²⁷ Y. Fisyak,⁴⁸ E. Flattum,²⁷ G. E. Forden,²⁰ M. Fortner,²⁹ K. C. Frame,⁴² S. Fuess,²⁷ E. Gallas,²⁷ A. N. Galyaev,¹⁸ P. Gartung,²⁴ V. Gavrilov,¹⁶ T. L. Geld,⁴² R. J. Genik II,⁴² K. Genser,²⁷ C. E. Gerber,²⁷ Y. Gershtein,⁵¹ B. Gibbard,⁴⁸ G. Ginther,⁴⁶ B. Gobbi,³⁰ B. Gómez,⁵ G. Gómez,³⁸ P. I. Goncharov,¹⁸ J. L. González Solís,¹⁴ H. Gordon,⁴⁸ L. T. Goss,⁵³ K. Gounder,²⁴ A. Goussiou,⁴⁷ N. Graf,⁴⁸ P. D. Grannis,⁴⁷ D. R. Green,²⁷ J. A. Green,³⁴ H. Greenlee,²⁷ S. Grinstein,¹ P. Grudberg,²¹ S. Grünendahl,²⁷ G. Guglielmo,⁵⁰ J. A. Guida,²⁰ J. M. Guida,⁵¹ A. Gupta,¹¹ S. N. Gurchiev,¹⁸ G. Gutierrez,²⁷ P. Gutierrez,⁵⁰ N. J. Hadley,³⁸ H. Haggerty,²⁷ S. Hagopian,²⁵ V. Hagopian,²⁵ K. S. Hahn,⁴⁶ R. E. Hall,²³ P. Hanlet,⁴⁰ S. Hansen,²⁷ J. M. Hauptman,³⁴ C. Hays,⁴⁴ C. Hebert,³⁵ D. Hedin,²⁹ A. P. Heinson,²⁴ U. Heintz,³⁹ R. Hernández-Montoya,¹⁴ T. Heuring,²⁵ R. Hirosky,²⁸ J. D. Hobbs,⁴⁷ B. Hoeneisen,⁶ J. S. Hoftun,⁵¹ F. Hsieh,⁴¹ Tong Hu,³¹ A. S. Ito,²⁷ S. A. Jerger,⁴² R. Jesik,³¹ T. Joffe-Minor,³⁰ K. Johns,²⁰ M. Johnson,²⁷ A. Jonckheere,²⁷ M. Jones,²⁶ H. Jöstlein,²⁷ S. Y. Jun,³⁰ S. Kahn,⁴⁸ D. Karmanov,¹⁷ D. Karmgard,²⁵ R. Kehoe,³² S. K. Kim,¹³ B. Klima,²⁷ C. Klopfenstein,²² B. Knuteson,²¹ W. Ko,²² J. M. Kohli,⁹ D. Koltick,³³ A. V. Kostritskiy,¹⁸ J. Kotcher,⁴⁸ A. V. Kotwal,⁴⁴ A. V. Kozelov,¹⁸ E. A. Kozlovsky,¹⁸ J. Krane,³⁴ M. R. Krishnaswamy,¹¹ S. Krzywdzinski,²⁷ M. Kubantsev,³⁶ S. Kuleshov,¹⁶ Y. Kulik,⁴⁷ S. Kunori,³⁸ F. Landry,⁴² G. Landsberg,⁵¹ A. Leflat,¹⁷ J. Li,⁵² Q. Z. Li,²⁷ J. G. R. Lima,³ D. Lincoln,²⁷ S. L. Linn,²⁵ J. Linnemann,⁴² R. Lipton,²⁷ J. G. Lu,⁴ A. Lucotte,⁴⁷ L. Lueking,²⁷ A. K. A. Maciel,²⁹ R. J. Madaras,²¹ R. Madden,²⁵ L. Magaña-Mendoza,¹⁴ V. Manankov,¹⁷ S. Mani,²² H. S. Mao,⁴ R. Markeloff,²⁹ T. Marshall,³¹ M. I. Martin,²⁷ R. D. Martin,²⁸ K. M. Mauritz,³⁴ B. May,³⁰ A. A. Mayorov,¹⁸ R. McCarthy,⁴⁷ J. McDonald,²⁵ T. McKibben,²⁸ J. McKinley,⁴² T. McMahon,⁴⁹ H. L. Melanson,²⁷ M. Merkin,¹⁷ K. W. Merritt,²⁷ C. Miao,⁵¹ H. Miettinen,⁵⁴ A. Mincer,⁴⁵ C. S. Mishra,²⁷ N. Mokhov,²⁷ N. K. Mondal,¹¹ H. E. Montgomery,²⁷ M. Mostafa,¹ H. da Motta,² F. Nang,²⁰ M. Narain,³⁹ V. S. Narasimham,¹¹ A. Narayanan,²⁰ H. A. Neal,⁴¹ J. P. Negret,⁵ P. Nemethy,⁴⁵ D. Norman,⁵³ L. Oesch,⁴¹ V. Oguri,³ N. Oshima,²⁷ D. Owen,⁴² P. Padley,⁵⁴ A. Para,²⁷ N. Parashar,⁴⁰ Y. M. Park,¹² R. Partridge,⁵¹ N. Parua,⁷ M. Paterno,⁴⁶ B. Pawlik,¹⁵ J. Perkins,⁵² M. Peters,²⁶ R. Piegaia,¹ H. Piekarczyk,²⁵ Y. Pischalnikov,³³ B. G. Pope,⁴² H. B. Prosper,²⁵ S. Protopopescu,⁴⁸ J. Qian,⁴¹ P. Z. Quintas,²⁷ R. Raja,²⁷ S. Rajagopalan,⁴⁸ O. Ramirez,²⁸ N. W. Reay,³⁶ S. Reucroft,⁴⁰ M. Rijssenbeek,⁴⁷ T. Rockwell,⁴² M. Roco,²⁷ P. Rubinov,³⁰ R. Ruchti,³² J. Rutherford,²⁰ A. Sánchez-Hernández,¹⁴ A. Santoro,² L. Sawyer,³⁷ R. D. Schamberger,⁴⁷ H. Schellman,³⁰ J. Sculli,⁴⁵ E. Shabalina,¹⁷ C. Shaffer,²⁵ H. C. Shankar,¹¹ R. K. Shivpuri,¹⁰ D. Shpakov,⁴⁷ M. Shupe,²⁰ R. A. Sidwell,³⁶ H. Singh,²⁴ J. B. Singh,⁹ V. Sirotenko,²⁹ P. Slattery,⁴⁶ E. Smith,⁵⁰ R. P. Smith,²⁷ R. Snihur,³⁰ G. R. Snow,⁴³ J. Snow,⁴⁹ S. Snyder,⁴⁸ J. Solomon,²⁸ X. F. Song,⁴ M. Sosebee,⁵² N. Sotnikova,¹⁷ M. Souza,² N. R. Stanton,³⁶ G. Steinbrück,⁵⁰ R. W. Stephens,⁵² M. L. Stevenson,²¹ F. Stichelbaut,⁴⁸ D. Stoker,²³ V. Stolin,¹⁶ D. A. Stoyanova,¹⁸ M. Strauss,⁵⁰ K. Streets,⁴⁵ M. Strovink,²¹ A. Sznajder,³ P. Tamburello,³⁸ J. Tarazi,²³ M. Tartaglia,²⁷ T. L. T. Thomas,³⁰ J. Thompson,³⁸ D. Toback,³⁸ T. G. Trippe,²¹ P. M. Tuts,⁴⁴ V. Vaniev,¹⁸ N. Varelas,²⁸ E. W. Varnes,²¹ A. A. Volkov,¹⁸ A. P. Vorobiev,¹⁸ H. D. Wahl,²⁵ J. Warchol,³² G. Watts,⁵¹ M. Wayne,³² H. Weerts,⁴² A. White,⁵² J. T. White,⁵³ J. A. Wightman,³⁴ S. Willis,²⁹ S. J. Wimpenny,²⁴ J. V. D. Wirjawan,⁵³ J. Womersley,²⁷ D. R. Wood,⁴⁰ R. Yamada,²⁷ P. Yamin,⁴⁸ T. Yasuda,²⁷ P. Yepes,⁵⁴ K. Yip,²⁷ C. Yoshikawa,²⁶ S. Youssef,²⁵ J. Yu,²⁷ Y. Yu,¹³ M. Zanabria,⁵ Z. Zhou,³⁴ Z. H. Zhu,⁴⁶ M. Zielinski,⁴⁶ D. Zieminska,³¹ A. Zieminski,³¹ V. Zutshi,⁴⁶ E. G. Zverev,¹⁷ and A. Zylberstein⁸
(D0 Collaboration)

- ¹Universidad de Buenos Aires, Buenos Aires, Argentina
- ²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil
- ³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil
- ⁴Institute of High Energy Physics, Beijing, People's Republic of China
- ⁵Universidad de los Andes, Bogotá, Colombia
- ⁶Universidad San Francisco de Quito, Quito, Ecuador
- ⁷Institut des Sciences Nucléaires, IN2P3-CNRS, Université de Grenoble 1, Grenoble, France
- ⁸DAPNIA/Service de Physique des Particules, CEA, Saclay, France
- ⁹Panjab University, Chandigarh, India
- ¹⁰Delhi University, Delhi, India
- ¹¹Tata Institute of Fundamental Research, Mumbai, India
- ¹²Kyungshung University, Pusan, Korea
- ¹³Seoul National University, Seoul, Korea
- ¹⁴CINVESTAV, Mexico City, Mexico
- ¹⁵Institute of Nuclear Physics, Kraków, Poland
- ¹⁶Institute for Theoretical and Experimental Physics, Moscow, Russia
- ¹⁷Moscow State University, Moscow, Russia
- ¹⁸Institute for High Energy Physics, Protvino, Russia
- ¹⁹Lancaster University, Lancaster, United Kingdom
- ²⁰University of Arizona, Tucson, Arizona 85721
- ²¹Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720
- ²²University of California, Davis, California 95616
- ²³University of California, Irvine, California 92697
- ²⁴University of California, Riverside, California 92521
- ²⁵Florida State University, Tallahassee, Florida 32306
- ²⁶University of Hawaii, Honolulu, Hawaii 96822
- ²⁷Fermi National Accelerator Laboratory, Batavia, Illinois 60510
- ²⁸University of Illinois at Chicago, Chicago, Illinois 60607
- ²⁹Northern Illinois University, DeKalb, Illinois 60115
- ³⁰Northwestern University, Evanston, Illinois 60208
- ³¹Indiana University, Bloomington, Indiana 47405
- ³²University of Notre Dame, Notre Dame, Indiana 46556
- ³³Purdue University, West Lafayette, Indiana 47907
- ³⁴Iowa State University, Ames, Iowa 50011
- ³⁵University of Kansas, Lawrence, Kansas 66045
- ³⁶Kansas State University, Manhattan, Kansas 66506
- ³⁷Louisiana Tech University, Ruston, Louisiana 71272
- ³⁸University of Maryland, College Park, Maryland 20742
- ³⁹Boston University, Boston, Massachusetts 02215
- ⁴⁰Northeastern University, Boston, Massachusetts 02115
- ⁴¹University of Michigan, Ann Arbor, Michigan 48109
- ⁴²Michigan State University, East Lansing, Michigan 48824
- ⁴³University of Nebraska, Lincoln, Nebraska 68588
- ⁴⁴Columbia University, New York, New York 10027
- ⁴⁵New York University, New York, New York 10003
- ⁴⁶University of Rochester, Rochester, New York 14627
- ⁴⁷State University of New York, Stony Brook, New York 11794
- ⁴⁸Brookhaven National Laboratory, Upton, New York 11973
- ⁴⁹Langston University, Langston, Oklahoma 73050
- ⁵⁰University of Oklahoma, Norman, Oklahoma 73019
- ⁵¹Brown University, Providence, Rhode Island 02912
- ⁵²University of Texas, Arlington, Texas 76019
- ⁵³Texas A&M University, College Station, Texas 77843
- ⁵⁴Rice University, Houston, Texas 77005

(Received 11 October 1999)

We present a measurement of the transverse momentum distribution of Z bosons produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV from data collected by the DØ experiment at the Fermilab Tevatron Collider. We find good agreement between our results and current resummation calculations, and also use our data to extract nonperturbative parameters for a particular version of the resummation formalism. The resulting values are significantly more precise than obtained in previous determinations.

PACS numbers: 14.70.Hp, 13.38.Dg, 13.85.Qk

We report a new measurement [1,2] of the inclusive differential cross section of the Z boson in the dielectron channel as a function of transverse momentum ($d\sigma/dp_T$) with statistics and precision greatly improved over previous measurements [3,4]. The measurement provides a sensitive test of quantum chromodynamics (QCD) at large scales, or high Q^2 . At small transverse momentum (p_T), where the cross section is largest, uncertainties in the phenomenology of vector boson production have contributed significantly to the uncertainty in the extraction of the mass of the W boson. Because of its similar production characteristics, and the fact that the decay electrons can be well measured, the Z provides a unique precision tool for evaluating the veracity of the phenomenology of vector boson production.

In the parton model, Z bosons are produced in collisions of $q\bar{q}$ constituents of the proton and antiproton. The fact that observed Z bosons have finite p_T can be attributed to gluon radiation from the colliding partons prior to their annihilation. In standard perturbative QCD (pQCD), the cross section for Z boson production is calculated by expanding a series in powers of the strong coupling constant α_s . This procedure works well when $p_T^2 \sim Q^2$ with $Q = M_Z$. However, when $p_T \ll Q$, correction terms that are proportional to $\alpha_s \ln(Q^2/p_T^2)$ become significant, and the cross section therefore diverges at small p_T . This difficulty is surmounted by reordering the perturbative series through a technique called *resummation* [5–13]. Although this technique extends the applicability of pQCD to lower values of p_T , a more fundamental barrier is encountered when p_T approaches Λ_{QCD} , the scale characterizing QCD processes. In this region, α_s becomes large and the perturbative calculation is no longer valid. Accounting for the nonperturbative contribution requires the introduction of a phenomenological form factor, which contains several parameters that must be tuned to data [8,10,11]. The universality of the resummation approach is an important idea that requires experimental testing. Its implications are far reaching, ranging from impact on the precision determination of M_W , to the production of Higgs bosons and diphotons [14,15].

The resummation can be carried out in impact-parameter (b) space via a Fourier transform, or in transverse momentum space. Both approaches require a nonperturbative function in order to describe the low- p_T region beyond some cutoff value b_{max} or $p_{T\text{lim}}$, and they merge to fixed-order perturbation theory at $p_T \sim Q$. The current state-of-the-art for the b -space formalism resums terms to next-to-next-to-next leading log and includes fixed-order terms to $\mathcal{O}(\alpha_s^2)$ [11]. Similarly, the p_T -space formalism resums terms to next-to-next leading log and includes fixed-order terms to $\mathcal{O}(\alpha_s)$ [13].

In the b -space formalism, the resummed cross section is modified at large b (above b_{max}) by $\exp[-S_{\text{NP}}(b, Q^2)]$.

The form factor $S_{\text{NP}}(b, Q^2)$ has a general renormalization group invariant form, but requires a specific choice of parametrization when making predictions. A possible choice, suggested by Ladinsky and Yuan [11], is

$$S_{\text{NP}}(b, Q^2) = g_1 b^2 + g_2 b^2 \ln\left(\frac{Q^2}{Q_0^2}\right) + g_1 g_3 b \ln(100 x_i x_j), \quad (1)$$

where x_i and x_j are the fractions of incident hadron momenta carried by the colliding partons and g_i are the nonperturbative parameters. An earlier parametrization by Davies, Webber, and Stirling [8] corresponds to the above with $g_3 \equiv 0$. For measurements at the Fermilab Tevatron at $Q^2 = M_Z^2$, the calculation is most sensitive to the value of g_2 and quite insensitive to the value of g_3 .

In the p_T -space formalism, the resummed cross section is modified at low p_T (below $p_{T\text{lim}}$) by multiplying the cross section by $F_{\text{NP}}(p_T)$. In this case, the form of the nonperturbative function is not constrained by renormalization group invariance. The choice suggested by Ellis and Veseli [13] is

$$\tilde{F}_{\text{NP}}(p_T) = 1 - e^{-\tilde{a} p_T^2}, \quad (2)$$

where $\tilde{a}$ is a nonperturbative parameter.

Previously published measurements of the differential cross section for Z boson production have been limited primarily by statistics (candidate samples of a few hundred events). This measurement is based on a sample of 6407 $Z \rightarrow e^+ e^-$ events, corresponding to an integrated luminosity of $\approx 111 \text{ pb}^{-1}$, collected with the DØ detector [16] in 1994–1996. A recent measurement by the CDF Collaboration has a similar number of events [17].

Electrons are detected in the uranium/liquid-argon calorimeter with a fractional energy resolution of $\approx 15\%/\sqrt{E(\text{GeV})}$. The calorimeter has a transverse granularity at the electron shower maximum of $\Delta\eta \times \Delta\phi = 0.05 \times 0.05$, where η is the pseudorapidity and ϕ is the azimuthal angle. The two electron candidates in the event with the highest transverse energy (E_T), both having $E_T > 25 \text{ GeV}$, are used to reconstruct the Z boson candidate. One electron is required to be in the central region, $|\eta_{\text{det}}| < 1.1$, and the second electron may be either in the central or in the forward region, $1.5 < |\eta_{\text{det}}| < 2.5$, where η_{det} refers to the value of η obtained by assuming that the shower originates from the center of the detector. Offline, both electrons are required to be isolated and to satisfy cluster-shape requirements. Additionally, at least one of the electrons is required to have a matching track in the drift chamber system that points to the reconstructed calorimeter cluster.

Both the acceptance and the theory predictions modified by the DØ detector resolution are calculated using a simulation technique originally developed for measuring the mass of the W boson [18], with minor modifications required by changes in selection criteria. The

four-momentum of the Z boson is obtained by generating the mass of the Z according to an energy-dependent Breit-Wigner line shape. The p_T and rapidity of the Z boson are chosen randomly from two-dimensional grids created using the computer program LEGACY [12], which calculates the Z boson cross section for a given p_T , rapidity, and mass of the Z boson. The positions and energies of the electrons are smeared according to the measured resolutions, and corrected for offsets in energy scale caused by the underlying event and recoil particles that overlap the calorimeter towers. Underlying events are modeled using data from random inelastic $p\bar{p}$ collisions of the same luminosity profile as the Z boson sample. The electron energy and angular resolutions are tuned to reproduce the observed width of the mass distribution at the Z -boson resonance and the difference between the reconstructed vertex positions of the electrons.

We determine the shape of the efficiency of the event selection criteria as a function of p_T using $Z \rightarrow e^+e^-$ events generated with HERWIG [19], smeared with the DØ detector resolutions, and overlaid on randomly selected zero bias $p\bar{p}$ collisions. This simulation models the effects of the underlying event and jet activity on the selection of the electrons. The absolute efficiency is obtained from $Z \rightarrow e^+e^-$ data [20]. The values of the efficiency times acceptance range from (26–37)% for p_T below 200 GeV and is 53% for p_T above 200 GeV.

The primary background arises from multiple-jet production from QCD processes in which two jets pass the electron selection criteria. We use several DØ data sets for estimating this background—direct- γ events, dijet events, and dielectron events in which both electrons fail quality criteria—all of which have very similar kinematic characteristics [1]. The level of the multijet background is determined by fitting the ee invariant mass in the range $60 < M_{ee} < 120$ GeV to a linear combination of Monte Carlo $Z \rightarrow e^+e^-$ signal events (using PYTHIA [21]) and background (from direct- γ events). We assign a systematic uncertainty to this measurement by varying the choice of mass window used in the fit, and by changing the background sample among those mentioned above. We estimate the total multijet background level to be $(4.4 \pm 0.9)\%$. The direct- γ sample is used to parametrize the shape of the background distribution as a function of p_T . Backgrounds from other sources, such as $Z \rightarrow \tau^+\tau^-$, $t\bar{t}$, and diboson production, are negligible.

We use the data corrected for background, acceptance, and efficiency, to determine the best value of the nonperturbative parameter g_2 , given the shape of our data. In the fit, we fix g_1 and g_3 to the values obtained in [11] and vary the value of g_2 . We use the CTEQ4M probability distribution function. The prediction is smeared with the known detector resolutions, and the result fitted to our data. The resulting χ^2 distribution as a function of g_2 is well behaved and parabolic, yielding a value of $g_2 = 0.59 \pm 0.06$ GeV², considerably more precise than

previous determinations. For completeness, we also fit the individual values of g_1 and g_3 , with the other two parameters fixed to their published values [11]. We obtain $g_1 = 0.09 \pm 0.03$ GeV² and $g_3 = -1.1 \pm 0.6$ GeV⁻¹. Both results are consistent with the values of Ref. [11].

To determine the true $d\sigma/dp_T$, we correct the measured cross section for effects of detector smearing, using the ratio of generated to resolution-smeared ansatz p_T distributions. We use the calculation from LEGACY as our ansatz function, with the g_2 determined from our fit. The largest smearing correction occurs at low p_T , where smearing causes the largest fractional change in p_T and where the kinematic boundary at $p_T = 0$ produces non-Gaussian smearing. The correction is 18.5% in the first bin, decreasing to about 2% at 5 GeV. For all p_T values above 5 GeV, the correction is $\lesssim 5\%$. Systematic uncertainties arising from the choice of ansatz function are evaluated by varying g_2 within ± 1 standard deviation of the best-fit values. Additional uncertainties are evaluated by varying

TABLE I. Summary of the results of the measurement of the p_T distribution of the Z boson. The range of p_T corresponds to the intervals used for binning the data. The nominal p_T corresponds to the value of p_T used to plot the data and was obtained from theory. The quantity $d\sigma(Z \rightarrow e^+e^-)/dp_T$ corresponds to the differential cross section in each bin of p_T for $Z \rightarrow e^+e^-$ production. The uncertainty on the differential cross section includes both systematic and statistical uncertainties, but does not include overall normalization uncertainty due to the luminosity of $\pm 4.4\%$.

p_T range (GeV)	Nominal p_T value (GeV)	Number of events	$d\sigma(Z \rightarrow e^+e^-)/dp_T$ (pb/GeV)
0–1	0.6	156	6.04 ± 0.53
1–2	1.5	424	16.2 ± 0.96
2–3	2.5	559	20.4 ± 1.1
3–4	3.5	572	19.7 ± 1.1
4–5	4.5	501	16.2 ± 0.92
5–6	5.5	473	15.0 ± 0.87
6–7	6.5	440	14.1 ± 0.84
7–8	7.5	346	11.1 ± 0.73
8–9	8.5	312	10.0 ± 0.69
9–10	9.5	285	9.29 ± 0.67
10–12	11.0	439	7.25 ± 0.54
12–14	13.0	326	5.45 ± 0.44
14–16	15.0	258	4.45 ± 0.39
16–18	17.0	203	3.54 ± 0.33
18–20	19.0	181	3.21 ± 0.31
20–25	22.3	287	2.06 ± 0.18
25–30	27.3	174	1.29 ± 0.13
30–35	32.3	124	0.962 ± 0.11
35–40	37.4	104	0.840 ± 0.10
40–50	44.5	92	0.373 ± 0.045
50–60	54.5	61	0.251 ± 0.036
60–70	64.5	40	0.163 ± 0.027
70–85	76.6	20	0.053 ± 0.012
85–100	91.7	13	0.034 ± 0.009
100–200	135	15	0.0050 ± 0.0013
200–300	228	2	$0.0004^{+0.0004}_{-0.0003}$

the detector resolutions by ± 1 standard deviation from the nominal values. The effect of these variations is negligible relative to the other uncertainties in the measurement.

Table I shows the values of $d\sigma(Z \rightarrow e^+e^-)/dp_T$. The uncertainties on the data points include statistical and systematic contributions. An additional normalization uncertainty of $\pm 4.4\%$ arises from the uncertainty on the integrated luminosity [20] that is not included in any of the plots nor in the table, but must be taken into account in any fits involving an absolute normalization.

Figure 1 shows the differential cross section corrected for the detector resolutions, compared to the fixed-order calculation and the resummation calculation with three different parametrizations of the nonperturbative region, based on published values of the nonperturbative parameters. Also shown are the fractional differences between data and the resummed predictions. The data are normalized to the measured $Z \rightarrow e^+e^-$ cross section (221 pb [20]), and the predictions are normalized to the absolute theoretical calculation. We observe best agreement with the Ladinsky-Yuan parameters for the b -space formalism; however, we expect that fits to high-statistics newer data using the Davies-Weber-Stirling (b -space) or Ellis-Veseli (p_T -space) parametrizations of the nonperturbative functions could describe our results similarly well. We fur-

ther note that the published values for the Ladinsky-Yuan parametrization are due largely to low energy Drell-Yan data. That they do so well in confrontation with high statistics and high- Q^2 data is intriguing. This is especially significant since these are not just shape comparisons, but involve absolute cross sections.

Figure 2 shows the measured differential cross section compared to the fixed-order absolute cross section calculation and the resummation calculation using the Ladinsky-Yuan parametrization. We observe strong disagreement between the data and the fixed-order prediction in the shape for all but the highest values of p_T . We attribute this to the divergence of the next-to-leading-order calculation at $p_T = 0$, and a significant enhancement of the cross section relative to the prediction at moderate values of p_T . This disagreement confirms the presence of contributions from soft gluon emission, which are accounted for in the resummation formalisms.

In summary, we have measured the inclusive differential cross section of the Z boson as a function of its transverse momentum. With the enhanced precision of this measurement over previous results, we can probe nonperturbative, resummation, and fixed-order QCD effects for all values of p_T . We observe good agreement between data and the b -space resummation calculation of the absolute cross section using the published values of the nonperturbative parameters of Ladinsky-Yuan. Using their form for the




FIG. 1. The differential cross section of the Z boson as a function of p_T compared with resummation calculations using three published parametrizations of the nonperturbative region as well as a fixed-order calculation. Also shown are the fractional differences in absolute cross section between data and each of the resummed predictions. As noted in the text, the large disagreement with the Davies-Weber-Stirling parametrization should not be considered as ruling out the two-parameter description.




FIG. 2. A comparison of data to the resummed and fixed-order [$\mathcal{O}(\alpha_s^2)$] calculations. Also shown are the fractional differences in absolute cross sections between data and the resummed and fixed-order calculations. The uncertainties include both statistical and systematic contributions (other than an overall normalization uncertainty from uncertainty in the luminosity).

parametrization of the nonperturbative region, we obtain $g_2 = 0.59 \pm 0.06 \text{ GeV}^2$.

We thank the Fermilab and collaborating institution staffs for contributions to this work and acknowledge support from the Department of Energy and National Science Foundation (U.S.A.), Commissariat à L'Energie Atomique (France), Ministry for Science and Technology and Ministry for Atomic Energy (Russia), CAPES and CNPq (Brazil), Departments of Atomic Energy and Science and Education (India), Colciencias (Colombia), CONACyT (Mexico), Ministry of Education and KOSEF (Korea), and CONICET and UBACyT (Argentina).

-
- [1] DØ Collaboration, B. Abbott et al., Phys. Rev. D **61**, 032004 (2000).
 - [2] D. Casey, Ph.D. thesis, University of Rochester, 1997 (unpublished) [http://www-d0.fnal.gov/publications_talks/thesis/casey/thesis.ps].
 - [3] UA2 Collaboration, P. Bagnaia et al., Z. Phys. C **47**, 523 (1990).
 - [4] CDF Collaboration, F. Abe et al., Phys. Rev. Lett. **67**, 2937 (1991).
 - [5] J. C. Collins and D. E. Soper, Nucl. Phys. **B193**, 381 (1981); **B213**, 545E (1983); J. C. Collins, D. E. Soper, and G. Sterman, *ibid.* **B250**, 199 (1985).
 - [6] C. T. H. Davies and W. J. Stirling, Nucl. Phys. **B244**, 337 (1984).
 - [7] G. Altarelli, R. K. Ellis, M. Greco, and G. Martinelli, Nucl. Phys. **B246**, 12 (1984).
 - [8] C. T. H. Davies, B. R. Weber, and W. J. Stirling, Nucl. Phys. **B256**, 413 (1985).
 - [9] P. B. Arnold and M. H. Reno, Nucl. Phys. **B319**, 37 (1989); **B330**, 284E (1990).
 - [10] P. B. Arnold and R. P. Kaufman, Nucl. Phys. **B349**, 381 (1991).
 - [11] G. A. Ladinsky and C.-P. Yuan, Phys. Rev. D **50**, 4239 (1994).
 - [12] C. Balazs and C.-P. Yuan, Phys. Rev. D **56**, 5558 (1997).
 - [13] R. K. Ellis and S. Veseli, Nucl. Phys. **B511**, 649 (1998).
 - [14] C. Balazs and C.-P. Yuan, Phys. Rev. D **59**, 114007 (1999).
 - [15] C. Balazs, E. L. Berger, S. Mrenna, and C.-P. Yuan, Phys. Rev. D **57**, 6934 (1999).
 - [16] DØ Collaboration, S. Abachi et al., Nucl. Instrum. Methods Phys. Res., Sect. A **338**, 185 (1994).
 - [17] CDF Collaboration, T. Affolder et al., FERMILAB PUB-99/220-E (to be published).
 - [18] DØ Collaboration, B. Abbott et al., Phys. Rev. D **58**, 092003 (1998).
 - [19] G. Marchesini et al., Comput. Phys. Commun. **67**, 465 (1992).
 - [20] DØ Collaboration, B. Abbott et al., FERMILAB PUB-99/171-E, hep-ex/9906025 (to be published).
 - [21] H. U. Bengtsson and T. Sjostrand, Comput. Phys. Commun. **46**, 43 (1987).