

Annex table 2 Odds ratios (OR) for use of antidepressants during a 2-year follow-up period in participants

living alone compared to participants not living alone. Data: The Health 2000 Study, n = 3471

Predictor: living alone (reference group=not alone	Use of antidepressant medication							
	At baseline			During 2-year follow-up			Incident 2-year use ^a	
	OR (95% CI)	reduction	%	OR (95% CI)	reduction	%	OR (95% CI)	reduction
Adjustment in addition to age and gender:								
1.None	1.61 (1.15-2.25)	0	0	1.41 (1.05-1.89)	0	0	1.19 (0.80-1.76)	0
2.Psychosocial factors ^b	1.43 (1.03-1.99)	30	30	1.29 (0.96-1.73)	29	29	1.14 (0.77-1.70)	26
3.Sociodemographic factors ^c	1.48 (1.04-2.09)	21	21	1.25 (0.93-1.69)	39	39	1.05 (0.70-1.57)	74
4.Health behaviors ^d	1.58 (1.25-2.21)	5	5	1.35 (1.00-1.81)	15	15	1.13 (0.76- 1.68)	32
5.All of the above	1.33 (0.93-1.89)	46	46	1.13 (0.83-1.54)	68	68	0.99 (0.66-1.50)	na ^e

^a New users during the follow-up with no purchases at the baseline (in 2000 or 2001).

^b Low social support at the workplace, low social support in private life, poor job climate, hostile personality.

^c Low educational level, low occupational grade, low income, unemployment, urbanicity, living at rent, housing disadvantages.

^d Regular smoking, alcohol heavy use (>20g women, >40g men), sedentary lifestyle, obesity (BMI >30)