



## OPEN ACCESS

## Optimal management of allergic rhinitis

Glenis K Scadding

## Correspondence to

Dr Glenis K Scadding,  
Department of Allergy and  
Medical Rhinology, Royal  
National Throat Nose and Ear  
Hospital, 330 Gray's Inn Road,  
London WC1Y 8DA, UK;  
[g.scadding@ucl.ac.uk](mailto:g.scadding@ucl.ac.uk)

Received 26 August 2014  
Revised 16 November 2014  
Accepted 21 November 2014  
Published Online First  
2 April 2015

## ABSTRACT

Allergic rhinitis (AR), the most common chronic disease in childhood is often ignored, misdiagnosed and/or mistreated. Undertreated AR impairs quality of life, exacerbates asthma and is a major factor in asthma development. It can involve the nose itself, as well as the organs connected with the nose manifesting a variety of symptoms. Evidence-based guidelines for AR therapy improve disease control. Recently, paediatric AR guidelines have been published by the European Academy of Allergy and Clinical Immunology and are available online, as are a patient care pathway for children with AR and asthma from the Royal College of Paediatrics and Child Health. Management involves diagnosis, followed by avoidance of relevant allergens, with additional pharmacotherapy needed for most sufferers. This ranges, according to severity, from saline sprays, through non-sedating antihistamines, oral or topical, with minimally bioavailable intranasal corticosteroids for moderate/severe disease, possibly plus additional antihistamine or antileukotriene. The concept of rhinitis control is emerging, but there is no universally accepted definition. Where pharmacotherapy fails, allergen-specific immunotherapy, which is uniquely able to alter long-term disease outcomes, should be considered. The subcutaneous form (subcutaneous immunotherapy) in children has been underused because of concerns regarding safety and acceptability of injections. Sublingual immunotherapy is both efficacious and safe for grass pollen allergy. Further studies on other allergens in children are needed. Patient, carer and practitioner education into AR and its treatment are a vital part of management.

## INTRODUCTION

Rhinitis, which can be allergic, infectious or neither of these, is defined as at least two nasal symptoms, including rhinorrhoea, blockage, sneezing and itching. It is a common problem in childhood and adolescence which impacts negatively on physical, social and psychological well-being.

Allergic rhinitis (AR) is an under-recognised inflammatory condition of the nasal mucosa, caused by immunoglobulin E (IgE)-mediated early-phase and late-phase hypersensitivity responses, usually to inhaled allergens, similar to those in allergic asthma.<sup>1–3</sup> Typical allergens include house dust mite, grass and tree pollens, dander from animals such as cat, dog, horse and, occasionally, moulds.<sup>4</sup> Different phenotypes exist; those with obvious symptoms of sneezing and running, who are easily recognised, and others with predominant blockage, where the diagnosis may be missed. Children with AR can present with symptoms related to nasal connections, such as lungs, throat, ears, (table 1) or to quality-of-life impairment, often related to poor

quality sleep and consequent fatigue, poor concentration and school performance.

AR is often part of a systemic inflammatory process associated with other inflammatory conditions, including allergic conjunctivitis (AC), rhinosinusitis and asthma. Asthma shows an increased prevalence in children with both allergic<sup>5</sup> and non-allergic rhinitis.<sup>6</sup> A higher prevalence of asthma is found among those suffering from persistent and more severe rhinitis.<sup>7</sup> Over three-quarters of children with asthma also have AR<sup>8</sup> which is associated with poor asthma control.<sup>9–10</sup> Minimal persistent allergic inflammation of the nasal mucosa<sup>11</sup> synergises with infective inflammation; thus, subjects with AR have more problems with viral colds,<sup>12</sup> and the combination in children of allergic sensitisation, relevant allergen exposure and viral cold gives a high risk for hospital admission for asthma in children.<sup>13</sup> Poor asthma control is found in children with moderate to severe rhinitis, which should be identified and treated.<sup>1–9 10</sup>

AR precedes asthma development in preadolescence, adolescence or adult life and carries a three-fold risk of it persisting into middle age.<sup>14</sup> Bronchial hyper-responsiveness, raised exhaled nitric oxide and reduced lung function have been observed in children with AR.<sup>15</sup>

Pharmacotherapy for AR improves asthma control;<sup>16 17</sup> allergen-specific immunotherapy (SIT) for rhinitis may decrease the progression of rhinitis to asthma.<sup>18</sup>

Rhinitis affects well-being, both physical and psychological,<sup>19 20</sup> with a direct relationship to allergen exposure.<sup>21</sup> Family dynamics can be disturbed.<sup>22</sup> Uncontrolled AR reduces sleep quality<sup>23</sup> impairing concentration, school attendance and performance,<sup>24</sup> including at General Certificate of Secondary Education (GCSE) level.<sup>25</sup> Rhinitis health-related quality of life is reduced, again in direct correlation with allergen exposure.<sup>26</sup>

Sedating antihistamines further reduce learning ability and impinge on examination results.<sup>25 27</sup>

## AETIOLOGY

Environmental factors (tobacco smoke, pollution, infections, diet) acting on a genetic background (family history) contribute to the development of AR which may follow earlier atopic dermatitis<sup>28</sup> but also occurs as the initial manifestation of allergy. Sensitisation may take place via the nose. Local IgE production can occur without evidence of systemic sensitisation.<sup>29</sup>

## MANAGEMENT

Guideline-directed management has been shown to improve disease control.<sup>30</sup> The global Allergic Rhinitis and its Impact on Asthma guideline (ARIA),<sup>1</sup> which includes quality-of-life measures in evaluation



CrossMark

To cite: Scadding GK. *Arch Dis Child* 2015;**100**: 576–582.

**Table 1** Recognising rhinitis in childhood

	Pre-school	School	Adolescent
<b>Classic symptoms and signs of rhinitis</b>	<b>Rhinorrhoea</b> – clear or discoloured discharge, sniffing		
	<b>Pruritus</b> - nose rubbing, the “allergic salute”, “allergic crease”, “paroxysmal sneeze”, may be associated with complaints of an itchy mouth or throat in older children		
	<b>Congestion</b> - mouth breathing, snoring, sleep apnoea, allergic shiners		
<b>Potential atypical presentations</b>	<b>Eustachian tube dysfunction</b> - ear pain on pressure changes (eg flying), reduced hearing, chronic otitis media with effusion		
	<b>Cough</b> – often mislabelled as asthma		
	<b>Poorly controlled asthma</b> – may co-exist with asthma		
	<b>Sleep problems</b> - tired, poor school performance, irritability		
	<b>Prolonged and frequent respiratory tract infections</b>		
	<b>Rhinosinusitis</b> - catarrh, headache, facial pain, halitosis, cough, hyposmia		
	<b>Pollen-food syndrome</b> , particularly with pollen driven allergic rhinitis		

Adapted from Roberts *et al*<sup>33</sup> (with permission). Other rare causes include hormonal rhinitis in teenagers, neurogenic rhinitis and aspirin-sensitive rhinitis.

and treatment, and provides an evidence-based approach to treatment of AR, has recently been updated using GRADE (Grading of Recommendations, Assessment, Development and Evaluations) methodology.<sup>31</sup> The UK has its own guidelines, which are currently being updated.<sup>32</sup> More recent, specifically paediatric guidelines, include those from the European Academy for Allergy and Clinical Immunology (EAACI)<sup>33</sup> and patient pathways from the Royal College of Paediatrics and Child Health.<sup>34</sup> The following is a synopsis of their content.

### Diagnosis

AR is diagnosed by detailed history, including questions about possible asthma, and nasal examination, together with inspection of throat, ears and chest where possible, backed up by specific allergy tests, either skin prick or blood tests, for specific IgE to allergens suggested by the history.

Clinical history: symptom type, duration and frequency and exacerbating factors are the cornerstone for diagnosing and classifying paediatric rhinitis which is characterised by two or more nasal symptoms: itching, sneezing, obstruction and rhinorrhoea. The timing of these in relation to exposure to allergen (ie, specific season or animal) is highly relevant. Upon such exposure, symptoms of AR occur in minutes and last for hours. Late-phase symptoms can include nasal obstruction, hyposmia, postnasal mucous discharge and nasal hyper-reactivity.<sup>35</sup> AC occurs in approximately 50%–70% of patients with AR<sup>6</sup> and is the symptom which best differentiates AR from other forms of rhinitis.

In children, rhinitis may present via associated comorbidities (table 1) depending on the child's age. Nasal obstruction with chronic mouth breathing can sometimes be the only presenting symptom in small children: adenoidal hypertrophy and recurrent viral colds are frequent misdiagnoses. Adenoidal hypertrophy with or without sleep apnoea can, in fact, be associated with AR, and in some studies, a response to AR treatment has been noted with an improvement in sleep or hypoxia.<sup>36</sup>

Adenoidectomy has not been reported as improving AR, although it may have a role in treating chronic paediatric rhinosinusitis.<sup>37</sup> Otitis media with effusion and Eustachian tube dysfunction can be detected in children with AR.<sup>38</sup> In recent years, oral allergy syndrome (OAS), also known as pollen fruit syndrome, has more often been reported by patients suffering with pollen-induced AR, although the paediatric prevalence of this problem is uncertain.<sup>39</sup> The history of oral pruritus and/or local angioedema in response to fresh fruits and vegetables, for example, apples, hazelnuts, carrots, celery and peanuts, in children with seasonal AR can be misdiagnosed as a primary food allergy (FA). In fact, the initial sensitisation is to pollen, with subsequent cross-reactivity to identical molecules in fruits and vegetables. In doubtful cases, molecular allergy diagnosis may help to distinguish OAS from FA.<sup>39</sup>


Pan airway involvement should be considered, both in the history and on examination.

The ARIA classification of AR as mild intermittent, moderate to severe intermittent, mild persistent or moderate to severe persistent may also be made and used as a guide to therapy (figure 1). This classification has been recently validated in children through an epidemiological survey involving 1275 children between 6 and 12 years of age.<sup>40</sup>

Rhinitis has many underlying causes. (table 2). Approximately two-thirds of children and one-third of adult patients with rhinitis will present with AR; the remainder have other forms and some defy classification (idiopathic rhinitis).<sup>1</sup> Recurrent viral colds are more frequent in small children; AR is more common in older ones. Unilateral symptoms, nasal obstruction without other symptoms, mucopurulent discharge, pain or recurrent epistaxis suggest other diagnoses including rarer conditions that can mimic AR.<sup>35</sup> Chronic unremitting rhinitis present from birth should lead to tests for primary ciliary dyskinesia. One-sided persistent nasal blockage, especially with purulent discharge, suggests a foreign body or unilateral choanal atresia. Watery discharge from one side of the

## Review

**Figure 1** The Allergic Rhinitis and its Impact on Asthma guideline classification in untreated patients.<sup>2</sup> The historical classification of allergic rhinitis (AR) as seasonal or perennial in relation to triggering environmental allergens is useful in some geographical regions and is important for pollen immunotherapy and has, therefore, been additionally maintained in the British Society of Allergy and Clinical Immunology UK classification of AR.<sup>1</sup>



nose, especially on bending forward, can represent a cerebrospinal fluid leak. Cystic fibrosis should be sought if nasal polyps are found. Oral contraceptives can cause rhinitis in adolescents.


## EXAMINATION

Allergic children may give the ‘allergic salute’, rubbing their nose upwards, leading to a crease across the nasal bridge, and may also have an extra skin fold or line under their lower

**Table 2** Differential diagnosis of rhinitis in children (level D)

Diagnosis	Pre-school	School	Adolescent
<b>Choanal atresia or stenosis</b>	Obstruction without other features of allergic rhinitis		
<b>Immuno-deficiency</b>	Persisting mucopurulent discharge		
<b>Encephalocele</b>	Unilateral nasal “polyp”		
<b>Adenoidal hypertrophy</b>	Mouth breathing, discoloured nasal secretions, snoring in the absence of other features of allergic rhinitis		
<b>Foreign body</b>	Unilateral discoloured nasal secretions, foul smell		
<b>Rhinosinusitis</b>		Discoloured nasal secretions, headache, facial pain, poor smell, halitosis, cough	
<b>Cystic fibrosis</b>	Bilateral nasal polyps, poor smell, chest symptoms, symptoms of malabsorption, failure to thrive		
<b>Primary ciliary dyskinesia</b>	Persisting mucopurulent discharge without respite between “colds”, bilateral stasis of mucus and secretions at the nasal floor, symptoms from birth		
<b>CSF leakage</b>	Colourless nasal discharge often with a history of trauma		
<b>Coagulopathy</b>	Recurrent epistaxis with minimal trauma		
<b>Septal deviation</b>		Obstruction in the absence of other features of allergic rhinitis	

Adapted from Roberts *et al*<sup>33</sup> (with permission).  
CSF, cerebrospinal fluid.



**Figure 2** Entry to therapy can occur at 1, 2 or 3 year, depending on severity of presenting symptoms. Poor control should lead to a step up, good control to a step down, so that the minimum therapy necessary is used. For seasonal disease, regular therapy should be commenced 2 weeks before the anticipated start of symptoms. \*Oral antihistamines may be better tolerated, while intranasal antihistamines have a more rapid onset of action. \*\*Reconsider diagnosis if not controlled within 1–2 weeks. If <2 years of age and unresponsive to antihistamine within a week, reconsider diagnosis before stepping up therapy. If poorly controlled, consider a short rescue course of a decongestant or low-dose oral prednisolone to gain symptom control; topical ipratropium may be useful for rhinorrhoea (adapted from Roberts *et al* [33] (with permission)).

eyelids (Denny–Morgan lines). Allergic shiners (dark eye shadows beneath the lower eye lid), caused by fluid accumulation in the infraorbital groove, are typical of childhood AR being darker in severe chronic AR.<sup>41</sup> Frequent throat clearing or hoarseness can be another feature. Internal nasal examination, by otoscope or endoscope, is essential, to rule out alternatives such as nasal polyps.<sup>1 33 34</sup>

## INVESTIGATIONS

Confirmation of the diagnosis of AR requires evidence of specific IgE reactivity to airborne allergens relevant to the history via either skin-prick testing or the demonstration of serum-specific IgE. This information is also relevant to environmental control measures and allergen-SIT. Immediate hypersensitivity skin testing provides results within 15 min of performing skin tests, whereas blood tests for specific IgE take days and may be less cost-effective than skin-prick testing, but are useful in patients with dermatographism, severe atopic dermatitis or those unable or unwilling to temporarily stop antihistamine use. All IgE test results must be interpreted in the light of the history since both false negative and false positive (sensitisation without clinical disease) tests occur.<sup>29 42 43</sup> AR is triggered mainly by inhalant allergens, of which house dust mites, grass and tree pollens are the most common in most parts of the world.<sup>5</sup> It should be noted that an isolated positive IgE test with no relevant symptoms represents sensitisation and not clinical disease and does not require treatment; therefore, ‘fishing expeditions’, in which multiple allergy tests are undertaken, are not recommended.

If IgE tests are unavailable, then a possible alternative is a trial of anti-allergy therapy using an intranasal corticosteroid or antihistamine or both.

Other tests measuring nasal patency and lung function may be helpful. The latter should always be done in persistent rhinitis or where there are any chest symptoms.

## Treatment

This aims at safe and effective symptom relief plus prevention of complications and disease progression. There is evidence that

outcomes are improved when evidence-based guidelines are applied.<sup>31</sup> The ARIA guidelines are currently the most widely used in Europe for AR treatment. A recent revision published in 2010 answers specific paediatric questions.<sup>32</sup> The EAACI guideline similarly suggests a stepwise therapeutic approach (figure 2).


Allergen avoidance, where possible, is recommended across the spectrum of disease severity. Those suffering from hay fever are asymptomatic outside the pollen season and symptoms in nose and eyes are reduced by nasal air filters,<sup>44</sup> and pet avoidance shows clear benefits; for house dust mite (HDM) reduction, individual measures gave equivocal results or lack of benefit.<sup>45</sup> FA rarely causes isolated rhinitis, but can be relevant where there are additional gastrointestinal problems, asthma and/or atopic eczema in toddlers.

Saline douching: This reduces symptoms of AR and improves the effect of intranasal steroids (INS)<sup>46</sup> and is effective in rhinosinusitis, where douching is guideline-recommended. Isotonic solutions are well tolerated, inexpensive, easy to use, with no evidence showing that regular, daily use adversely affects health. Other complementary therapies, such as homeopathy, acupuncture, butterbur and herbal medicines are not recommended.<sup>32</sup>

New-generation, non-sedating, antihistamines are recommended alone for mild to moderate AR and in combination with INS for moderate to severe disease.<sup>47</sup> Sedating antihistamines, all of which cause psychomotor retardation, should no longer be used.<sup>48</sup> Intranasal antihistamine alone is faster in onset (15 min, compared with 1 h) and is more effective on nasal symptoms than oral antihistamines<sup>49</sup> and is underused.

INS are superior, for all symptoms, (excluding eye symptoms for which they are equal) to antihistamines (oral or topical), to antileukotrienes and to the combination of antihistamine and antileukotriene in meta-analyses.<sup>50–52</sup> In particular, they are more effective in relieving nasal obstruction.<sup>50</sup> INS are recommended for moderate to severe disease; concerns relating to systemic effects are unfounded when the latest molecules are used, as their systemic bioavailability is extremely low (figure 3). Mometasone furoate and fluticasone propionate have been extensively studied in children, and no adverse effects on cortisol levels, the hypothalamic-pituitary-adrenal axis or growth

## Review



Nasonex Summary of Product Characteristics. 2011  
 Rhinocort: Summary of Product Characteristics. 2011  
 Beco-nase Summary of Product Characteristic 2011  
 Daley-Yates P et al Br J Clin Pharm Pharmacol  
 2001;Kariyawasam H and Scadding GK Journal of Asthma and Allergy 2011  
 Scadding GK Paediatric Drugs 2008

**Figure 3** Systemic bioavailability of intranasal steroids.


over 1 year have been detected,<sup>53 54</sup> unlike beclomethasone.<sup>55</sup> Fluticasone furoate, similarly, has low bioavailability, but the use of a double-dose 110 mcg per day has been associated with reduction in growth.<sup>56</sup> It is sensible to use the lowest dose of minimally bioavailable INS needed to maintain control and to administer this in the morning. 'Holidays' from treatment are ineffective in reduction of systemic effects and are liable to result in loss of control. Regular monitoring of growth is a sensitive marker for systemic effects and is sensible, particularly in children receiving corticosteroids to skin, lung and nose. The contribution to systemic effects is greater from skin than lung, with very little from the nose compared with skin or lung, so if growth is slowed, attempts to reduce those areas of treatment should be made first. Considering the adverse impact of uncontrolled rhinitis upon asthma, it may be possible to reduce or even stop inhaled corticosteroids once INS are started, since INS may reduce ocular symptoms and also bronchial hyper-reactivity, probably by reduction of inflammation-induced neural reflexes,<sup>57</sup> at which recently available molecules such as fluticasone furoate appear more consistently effective.<sup>58</sup> There is some evidence that INS can benefit asthma,<sup>17 59–61</sup> but randomised, controlled prospective trials are needed of the addition of long-term INS to regular inhaled corticosteroids, both for efficacy and safety. It is logical to treat AR and asthma as one with a corticosteroid inhaled via the nose. Attempts have been made,<sup>62 63</sup> but the practice has not yet been accepted.

It is vital to explain the safety and correct use of INS to children and their carers. The delay in onset of action should be explained, plus the need for regular treatment, how to administer the spray (figure 4) and linking use to a regular daily activity, such as teeth cleaning. Provision of contact details for advice should also aid concordance, since well-meaning uninformed relatives, friends and even primary care practitioners may state that temporary use is all that is allowed, or may switch the prescription to an older, cheaper, more bioavailable molecule.

Nasal biopsies do not show mucosal atrophy after 1 year of regular INS use in adults.<sup>64–66</sup> INS improve mucociliary clearance, and have shown benefits when used in acute rhinosinusitis, so continuation during viral colds is recommended.

A combined nasal spray containing fluticasone propionate and azelastine, an antihistamine, is under trial for use in children.

Systemic corticosteroid is available as a brief rescue therapy for very severe symptoms, the lowest dose for the shortest time should be accompanied by an INS. Depot corticosteroid



**Figure 4** (A) shows how to use a nasal spray so as to avoid the septum and spray as much of the lateral wall mucosa as possible, allowing subsequent mucociliary clearance to distribute the liquid all over the mucosa. (B) shows how nasal drops (superior for rhinosinusitis) should be used with the head completely upside down so that drops reach the ostiomeatal complex in the upper nose where sinuses drain and ventilate (adapted from Scadding *et al* [1] (with permission)).

injections are not recommended, as risk outweighs benefit.<sup>33 67</sup>

Leukotriene receptor antagonists have similar effects as oral antihistamines, with minimal additional benefit when the two are used in combination.<sup>33 68</sup> Use in children with seasonal AR, in preschool children with persistent AR and in children with AR and concomitant asthma, where concerns exist about the use of glucocorticoids, is suggested by ARIA 2010.

### ALLERGEN-SIT

At present, this is recommended for severe AR uncontrolled by pharmacotherapy.<sup>33</sup> There is good evidence in pollen-induced AR that SIT reduces symptoms in the nose and eyes.<sup>69–71</sup> In some trials, SIT improves allergic asthma.<sup>69–71</sup>

SIT can be subcutaneous or sublingual. Currently, only 1%–5% of European children suffering with AR are treated with SIT: with 75% receiving subcutaneous immunotherapy (SCIT) and 25% sublingual immunotherapy (SLIT).<sup>71</sup>

Benefits of therapy must always be weighed against risks. A recent review of UK SIT practice<sup>72</sup> confirmed a low incidence of severe adverse reactions and no fatalities, even though guidelines about excluding chronic asthma subjects were not observed, and patients with chronic asthma were treated with SIT. Adrenaline injections are still sometimes

needed for anaphylaxis in SCIT which, because it is not completely risk-free, must only be administered by those competent in its use.

SLIT in children shows efficacy in hay fever; HDM desensitisation gives variable results.<sup>73 74</sup> Local reactions are common with itching and swelling in mouth and throat, worst on first application and usually disappearing in 2 weeks, systemic reactions rarely occur and no fatalities have been reported. A meta-analysis of 49 SLIT studies, including 15 paediatric papers with over 1400 children, suggests that it is more effective in the paediatric group.<sup>75</sup>

SIT is the only treatment with a disease-modifying effect during the treatment period and in the years following. It appears to be capable of preventing the progression of rhinitis to asthma and new allergic sensitisations. The Prevention of Asthma Study demonstrated a reduction in cases of new asthma in children receiving 3 years of SCIT for hay fever at 3, 5 and 10 years, compared with those treated by pharmacotherapy alone,<sup>19</sup> but was not blinded. A double-blind, randomised trial is in progress to test the effectiveness of grass pollen SLIT in asthma prevention. Another is in progress comparing the efficacy of SCIT and SLIT. It seems likely that SIT administered correctly to appropriately selected patients is socioeconomically beneficial.<sup>75</sup>

Concordance with long-term therapy, like SIT, is challenging: good contact and communication are again necessary.

## CONCLUSION

AR is worth treating effectively, even when it is part of a myriad of allergic conditions; as the nose is the gateway to the respiratory tract, good rhinitis control can facilitate control of symptoms elsewhere. AR, if poorly controlled, results in troublesome symptoms and impacts on daily activities, quality of life and on other areas of the respiratory tract, such as ears, sinuses, throat and lungs. Possible causes for difficult-to-treat cases include doctor factors such as misdiagnosis and undertreatment or patient factors such as lack of concordance with therapy. This can be reduced by information leaflets on specific allergen avoidance, the correct technique in the application of nasal preparations (avoiding the septum and leaving in place rather than sniffing back) and explanation of potential side effects. Clearly written treatment plans are useful, particularly if several preparations are prescribed.

AR can sometimes copresent in association with other forms of rhinitis (or rhinosinusitis) leading to increased severity (mixed rhinitis). Associated comorbidities should be considered in the differential diagnosis in order to determine the most appropriate therapeutic approach. Allergen-SIT should be considered for patients with genuinely difficult-to-control allergic disease, despite guideline-directed pharmacotherapy properly taken. Grade A evidence via meta-analysis confirms its efficacy and it is unique in its ability to alter the long-term disease course.

Local primary care directives, such as use of the cheapest, significantly bioavailable INS and inhaled corticosteroids molecules, and minimal provision of SIT are misguided and liable to result in extra-disease manifestations, reduced quality of life, and will increase costs in the long term.<sup>76</sup>

**Competing interests** Research grants from GSK, ALK. Honoraria for articles, lectures/chairing and advisory boards: Astra Zeneca, Britannia Pharmaceuticals, Capnia, Church & Dwight, Circassia, GSK, Grupo Uriach, Meda, Merck, MSD, Ono Pharmaceuticals, Oxford Therapeutics, Sanofi-Aventis, UCB. Travel funding from Bayer, GSK.

**Provenance and peer review** Commissioned; externally peer reviewed.

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

## REFERENCES

- Scadding GK, Durham SR, Mirakian R, *et al.* BSACI guidelines for the management of allergic and non-allergic rhinitis. *Clin Exp Allergy* 2008;38:19–42.
- Bousquet J, Khaltaev N, Cruz AA, *et al.* Allergic Rhinitis and its Impact on Asthma (ARIA) 2008 update (in collaboration with the World Health Organization, GA(2) LEN and AllerGen). *Allergy* 2008;63(Suppl 86):8–160.
- Ait-Khaled N, Pearce N, Anderson HR, *et al.* Global map of the prevalence of symptoms of rhinoconjunctivitis in children: The International Study of Asthma and Allergies in Childhood (ISAAC) Phase Three. *Allergy* 2009;64:123–48.
- Bjorksten B, Clayton T, Ellwood P, *et al.* Worldwide time trends for symptoms of rhinitis and conjunctivitis: Phase III of the International Study of Asthma and Allergies in Childhood. *Pediatr Allergy Immunol* 2008;19:110–24.
- Burbach GJ, Heinzerling LM, Edenharter G, *et al.* GA(2)LEN skin test study II: clinical relevance of inhalant allergen sensitizations in Europe. *Allergy* 2009;64:1507–15.
- Bertelsen RJ, Carlsen KC, Carlsen KH. Rhinitis in children: co-morbidities and phenotypes. *Pediatr Allergy Immunol* 2010;21(4 Pt 1):612–12.
- Chawes BL, Bonnelykke K, Kreiner-Moller E, *et al.* Children with allergic and nonallergic rhinitis have a similar risk of asthma. *J Allergy Clin Immunol* 2010;126:567–7.
- Ponte EV, Franco R, Nascimento HF, *et al.* Lack of control of severe asthma is associated with co-existence of moderate-to-severe rhinitis. *Allergy* 2008;63:564–9.
- Masuda S, Fujisawa T, Katsumata H, *et al.* High prevalence and young onset of allergic rhinitis in children with bronchial asthma. *Pediatr Allergy Immunol* 2008;19:517–22.
- de Groot EP, Nijkamp A, Duiverman EJ, *et al.* Allergic rhinitis is associated with poor asthma control in children with asthma. *Thorax* 2012;67:582–7.
- Lasmar LM, Camargos PA, Ordonez A, *et al.* Prevalence of allergic rhinitis and its impact on the use of emergency care services in a group of children and adolescents with moderate to severe persistent asthma. *J Pediatr (Rio J)* 2007;83:555–61.
- Canonica GW, Compalati E. Minimal persistent inflammation in allergic rhinitis: implications for current treatment strategies. *Clin Exp Immunol* 2009;158:260–71.
- Cirillo I, Marseglia G, Klersy C, *et al.* Allergic patients have more numerous and prolonged respiratory infections than non-allergic subjects. *Allergy* 2007;62:1087–90.
- Murray CS, Poletti G, Kebabdzic T, *et al.* Study of modifiable risk factors for asthma exacerbations: virus infection and allergen exposure increase the risk of asthma hospital admissions in children. *Thorax* 2006;61:376–82.
- Burgess JA, Walters EH, Byrnes GB, *et al.* Childhood allergic rhinitis predicts asthma incidence and persistence to middle age: a longitudinal study. *J Allergy Clin Immunol* 2007;120:863–9.
- Kim SW, Han DH, Lee SJ, *et al.* Bronchial hyperresponsiveness in pediatric rhinitis patients: The difference between allergic and nonallergic rhinitis. *Am J Rhinol Allergy* 2013;27:63–8.
- Lohia S, Schlosser RJ, Soler ZM. Impact of intranasal corticosteroids on asthma outcomes in allergic rhinitis: a meta-analysis. *Allergy* 2013;68:569–79.
- Stelmach R, do Patrocinio TN, Ribeiro M, *et al.* Effect of treating allergic rhinitis with corticosteroids in patients with mild-to-moderate persistent asthma. *Chest* 2005;128:3140–7.
- Jacobsen L, Niggemann B, Dreborg S, *et al.* Specific immunotherapy has long-term preventive effect of seasonal and perennial asthma: 10-year follow-up on the PAT study. *Allergy* 2007;62:943–8.
- Silva CH, Silva TE, Morales NM, *et al.* Quality of life in children and adolescents with allergic rhinitis. *Braz J Otorhinolaryngol* 2009;75:642–9.
- Meltzer EO, Blaiss MS, Derebery MJ, *et al.* Burden of allergic rhinitis: results from the Pediatric Allergies in America survey. *J Allergy Clin Immunol* 2009;124(3 Suppl):S43–70.
- Roberts G, Mylonopoulou M, Hurley C, *et al.* Impairment in quality of life is directly related to the level of allergen exposure and allergic airway inflammation. *Clin Exp Allergy* 2005;35:1295–300.
- Emin O, Mustafa S, Nedim S. Psychological stress and family functioning in mothers of children with allergic rhinitis. *Int J Pediatr Otorhinolaryngol* 2009;73:1795–8.
- Santos CB, Pratt EL, Hanks C, *et al.* Allergic rhinitis and its effect on sleep, fatigue, and daytime somnolence. *Ann Allergy Asthma Immunol* 2006;97:579–86.
- Blaiss MS. Allergic rhinitis and impairment issues in schoolchildren: a consensus report. *Curr Med Res Opin* 2004;20:1937–52.
- Walker S, Khan-Wasti S, Fletcher M, *et al.* Seasonal allergic rhinitis is associated with a detrimental effect on examination performance in United Kingdom teenagers: case-control study. *J Allergy Clin Immunol* 2007;120:381–7.

## Review

- 27 Juniper EF, Guyatt GH, Dolovich J. Assessment of quality of life in adolescents with allergic rhinoconjunctivitis: development and testing of a questionnaire for clinical trials. *J Allergy Clin Immunol* 1994;93:413–23.
- 28 Vuurman EF, van Veggel LM, Uiterwijk MM, *et al.* Seasonal allergic rhinitis and antihistamine effects on children's learning. *Ann Allergy* 1993;71:121–6.
- 29 Keil T, Bockelbrink A, Reich A, *et al.* The natural history of allergic rhinitis in childhood. *Pediatr Allergy Immunol* 2010;21:962–9.
- 30 Rondon C, Canto G, Blanca M. Local allergic rhinitis: a new entity, characterization and further studies. *Curr Opin Allergy Clin Immunol* 2010;10:1–7.
- 31 Bousquet J, Lund VJ, Van Cauwenberge P, *et al.* Implementation of guidelines for seasonal allergic rhinitis: a randomized controlled trial. *Allergy* 2003;58:733–41.
- 32 Brozek JL, Bousquet J, Baena-Cagnani CE, *et al.* Allergic Rhinitis and its Impact on Asthma (ARIA) guidelines: 2010 revision. *J Allergy Clin Immunol* 2010;126:466–76.
- 33 Roberts G, Xatzipsalti M, Borrego LM, *et al.* Paediatric rhinitis: position paper of the European Academy of Allergy and Clinical Immunology. *Allergy* 2013;68:1102–16.
- 34 Vance G, Lloyd K, Scadding G, *et al.* The 'unified airway': the RCPC care pathway for children with asthma and/or rhinitis. *Arch Dis Child* 2011;96(Suppl 2):i10–14.
- 35 Greiner AN, Hellings PW, Rotiroi G, *et al.* Allergic Rhinitis. *Lancet* 2011;378:2112–22.
- 36 Scadding G. Non-surgical treatment of adenoidal hypertrophy: the role of treating IgE-mediated inflammation. *Pediatr Allergy Immunol* 2010;21:1095–106.
- 37 Fokkens WJ, Lund VJ, Mullol J, *et al.* EPOS 2012: European position paper on rhinosinusitis and nasal polyps 2012. A summary for otorhinolaryngologists. *Rhinology* 2012;50:1–12.
- 38 Alles R, Parikh A, Hawk L, *et al.* The prevalence of atopic disorders in children with chronic otitis media with effusion. *Pediatr Allergy Immunol* 2001;12:102–6.
- 39 Mari A, Ballmer-Weber BK, Vieths S. The oral allergy syndrome: improved diagnostic and treatment methods. *Curr Opin Allergy Clin Immunol* 2005;5:267–3.
- 40 Jauregui I, Davila I, Sastre J, *et al.* Validation of ARIA (Allergic Rhinitis and its Impact on Asthma) classification in a pediatric population: the PEDRIAL study. *Pediatr Allergy Immunol* 2011;22:388–92.
- 41 Chen CH, Lin YT, Wen CY, *et al.* Quantitative assessment of allergic shiners in children with allergic rhinitis. *J Allergy Clin Immunol* 2009;123:665–71, 671.
- 42 Smurthwaite L, Durham SR. Local IgE synthesis in allergic rhinitis and asthma. *Curr Allergy Asthma Rep* 2002;2:231–8.
- 43 Powe DG, Jagger C, Kleinjan A, *et al.* 'Entropy': localized mucosal allergic disease in the absence of systemic responses for atopy. *Clin Exp Allergy* 2003;33:1374–9.
- 44 O'Meara TJ, Sercombe JK, Morgan G, *et al.* The reduction of rhinitis symptoms by nasal filters during natural exposure to ragweed and grass pollen. *Allergy* 2005;60:529–32.
- 45 Eggleston PA. Methods and effectiveness of indoor environmental control. *Ann Allergy Asthma Immunol* 2001;87:44–7.
- 46 Li H, Sha Q, Zuo K, *et al.* Nasal saline irrigation facilitates control of allergic rhinitis by topical steroid in children. *ORL J Otorhinolaryngol Relat Spec* 2009;71:50–5.
- 47 Simons FE, Simons KJ. Histamine and H1-antihistamines: celebrating a century of progress. *J Allergy Clin Immunol* 2011;128:1139–50.
- 48 Church MK, Maurer M, Simons FER, *et al.* Risk of first-generation H1-antihistamines: a GA2LEN position paper. *Allergy* 2010;65:459–66.
- 49 Portnoy JM, Van Osdol T, Williams PB. Evidence-based strategies for treating allergic rhinitis. *Curr Allergy Asthma Rep* 2004;6:439–46.
- 50 Weiner JM, Abramson MJ, Puy RM. Intranasal corticosteroids versus oral H1 receptor antagonists in allergic rhinitis: systematic review of randomised controlled trials. *BMJ* 1998;317:1624–9.
- 51 Yáñez A, Rodrigo GJ. Intranasal corticosteroids versus topical H1 receptor antagonists for the treatment of allergic rhinitis: a systematic review with meta-analysis. *Ann Allergy Asthma Immunol* 2002;89:479–84.
- 52 Weinstein SF. Combination therapy in the treatment of allergic rhinitis. *Allergy Asthma Proc* 2002;23:1–3. Review.
- 53 Allen DB, Meltzer EO, Lemanske RF Jr, *et al.* No growth suppression in children treated with the maximum recommended dose of fluticasone propionate aqueous nasal spray for one year. *Allergy Asthma Proc* 2002;23:407–13.
- 54 Schenkel EJ, Skoner DP, Bronsky EA, *et al.* Absence of growth retardation in children with perennial allergic rhinitis after one year of treatment with mometasone furoate aqueous nasal spray. *Pediatrics* 2000;105:E22.
- 55 Skoner DP, Rachelefsky GS, Meltzer EO, *et al.* Detection of growth suppression in children during treatment with intranasal beclomethasone dipropionate. *Pediatrics* 2000;105:E23.
- 56 Lee L, Sterling R, Maspero J, *et al.* Growth velocity reduced with once-daily fluticasone furoate nasal spray in prepubescent children with perennial allergic rhinitis. *JACI Pract* 2014;2:421–7.
- 57 Baroody FM, Foster KA, Markaryan A, *et al.* Nasal ocular reflexes and eye symptoms in patients with allergic rhinitis. *Ann Allergy Asthma Immunol* 2008;100:194–9.
- 58 Keith P, Scadding G. Are intranasal corticosteroids all equally consistent in managing ocular symptoms of seasonal allergic rhinitis? *Curr Med Res Opin* 2009;25:2021–41.
- 59 Adams RJ, Fuhlbrigge AL, Finkelstein JA, *et al.* Intranasal steroids and the risk of emergency department visits for asthma. *J Allergy Clin Immunol* 2002;109:636–42.
- 60 Crystal-Peters J, Neslusan C, Crown WH, *et al.* Treating allergic rhinitis in patients with comorbid asthma: the risk of asthma-related hospitalizations and emergency department visits. *J Allergy Clin Immunol* 2002;109:57–62.
- 61 Corren J, Manning BE, Thompson SF, *et al.* Rhinitis therapy and the prevention of hospital care for asthma: a case-control study. *J Allergy Clin Immunol* 2004;113:415–19.
- 62 Pedersen W, Hjuler I, Bisgaard H, *et al.* Nasal inhalation of budesonide from a spacer in children with perennial rhinitis and asthma. *Allergy* 1998;53:383–7.
- 63 Camargos P, Ibiapina C, Lasmar L, *et al.* Obtaining concomitant control of allergic rhinitis and asthma with a nasally inhaled corticosteroid. *Allergy* 2007;62:310–16.
- 64 Holm AF, Fokkens WJ, Godthelp T, *et al.* A 1-year placebo-controlled study of intranasal fluticasone propionate aqueous nasal spray in patients with perennial allergic rhinitis: a safety and biopsy study. *Clin Otolaryngol Allied Sci* 1998;23:69–73.
- 65 Minshall E, Ghaffar O, Cameron L, *et al.* Assessment by nasal biopsy of long-term use of mometasone furoate aqueous nasal spray (Nasonex) in the treatment of perennial rhinitis. *Otolaryngol Head Neck Surg* 1998;118:648–54.
- 66 Pipkorn U, Pukander J, Suonpaa J, *et al.* Long-term safety of budesonide nasal aerosol: a 5.5-year follow-up study. *Clin Allergy* 1988;18:253–9.
- 67 Nasser SM, Ewan PW. Lesson of the week: Depot corticosteroid treatment for hay fever causing avascular necrosis of both hips. *BMJ* 2001;322:1589–91.
- 68 Wilson AM, O'Byrne PM, Parameswaran K. Leukotriene receptor antagonists for allergic rhinitis: a systematic review and meta-analysis. [Review] [44 refs]. *American Journal of Medicine* 2004;116:338–44.
- 69 Calderon MA, Gerth van Wijk R, *et al.* Perspectives on allergen-specific immunotherapy in childhood: An EAACI position statement. *Pediatr Allergy Immunol* 2012;23:300–6.
- 70 Walker SM, Durham SR, Till SJ, *et al.* Immunotherapy for allergic rhinitis. *Clin Exp Allergy* 2011;41:1177–200.
- 71 Bufe A, Roberts G. Specific immunotherapy in children. *Clin Exp Allergy* 2011;41:1256–62.
- 72 Vance GH, Goldring S, Warner JO, *et al.* A national audit of pollen immunotherapy for children in the United Kingdom: patient selection and programme safety. *Clin Exp Allergy* 2011;41:1313–23.
- 73 Hankin C, Cox L, Lang D, *et al.* Allergen immunotherapy and health care cost benefits for children with allergic rhinitis: a large-scale, retrospective, matched cohort study. *Ann Allergy Asthma Immunol* 2010;104:79–85.
- 74 Larenas-Linnemann D. Sublingual immunotherapy in children: complete and updated review supporting evidence of effect. *Curr Opin Allergy Clin Immunol* 2009;9:168–76.
- 75 Radulovic S, Calderon MA, Wilson D, *et al.* Sublingual immunotherapy for allergic rhinitis. *Cochrane Database Syst Rev* 2010;(12):CD002893.
- 76 <http://leicestershire.formulary.co.uk/chaptersSubDetails.asp?FormularySectionID=12&SubSectionRef=12.02.01&SubSectionID=A100>



## Optimal management of allergic rhinitis

Glenis K Scadding

*Arch Dis Child* 2015 100: 576-582 originally published online April 2, 2015  
doi: 10.1136/archdischild-2014-306300

---

Updated information and services can be found at:  
<http://adc.bmj.com/content/100/6/576>

---

*These include:*

### References

This article cites 74 articles, 6 of which you can access for free at:  
<http://adc.bmj.com/content/100/6/576#BIBL>

### Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

### Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

### Topic Collections

Articles on similar topics can be found in the following collections

- [Open access](#) (137)
- [Immunology \(including allergy\)](#) (1885)
- [Child health](#) (3664)
- [Asthma](#) (356)
- [Drugs: CNS \(not psychiatric\)](#) (462)
- [Ear, nose and throat/otolaryngology](#) (279)
- [Guidelines](#) (119)
- [Patients](#) (264)
- [TB and other respiratory infections](#) (613)

---

### Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>