

Features of the UK childcare environment and associations with preschooler's in-care physical activity

Authorship:

Kathryn R Hesketh¹ krh40@cam.ac.uk

Esther MF van Sluijs¹ ev234@medschl.cam.ac.uk

¹ MRC Epidemiology Unit & Centre for Diet and Activity Research (CEDAR)

University of Cambridge School of Clinical Medicine,

Box 285 Institute of Metabolic Science, Cambridge Biomedical Campus,

Cambridge CB2 0QQ

Corresponding author:

Dr Kathryn R. Hesketh

Email: kathryn.hesketh@ucl.ac.uk

Address: UCL Institute of Child Health

30 Guildford Street

London

WC1N1EH

Tel: +44 207 905 2320

Fax: +44 207 905 2793

Abstract: 198

Main article: 2027

Abbreviations

PA: Physical activity; **SED:** Sedentary; **LPA:** Light physical activity; **MVPA:** Moderate-to-vigorous physical activity; **ICC:** Intra-class correlation co-efficient; **SPACE:** Studying Physical Activity in preschool-aged Children and their Environment Study; **IMD:** Index of Multiple Deprivation; **EPAO:** Environment Policy Assessment and Observation; **Ofsted:** Office for Standards in Education

Highlights

- Little is known about what influences children's activity in UK childcare settings.
- In 30 settings, 201 3-4-year-olds provided valid in-care objective activity data.
- We assessed 23 potential care-provider, environmental and policy correlates.
- Few associations were observed with children's in-care sedentary/physical activity.
- UK preschool policies may be more conducive to individual activity preferences.

Abstract

Objective: Features of the childcare environment may influence children's in-care physical activity (PA). We assessed the association between UK preschool care-provider, environmental and policy factors and 3-4-year-olds' average daily in-care sedentary behaviour (SED) and PA.

Methods: In 2013, we used accelerometers to measure the in-care SED/ PA of 201 3-4-year-old children (51% female) in 30 preschools in Cambridgeshire, UK, (average wear time: (mean±SD) 4.2±1.3 week-days). We assessed the childcare environment using the Environment and Policy Assessment and Observation tool; demographic and carer information was taken from questionnaires. We used three-level mixed-effects regression analyses (adjusted for sex, in-care time and travel mode to care) to determine the association between childcare factors and children's in-care average daily minutes/hour spent SED, in light PA (LPA) and in moderate-to-vigorous PA (MVPA).

Results: Children spent 5.6±2.5 hours in care per day on average; clustering of PA within preschools was limited (ICCs: 0.003-0.05). Fully adjusted models showed that active opportunities were positively associated with children's in-care SED. No associations with in-care LPA and MVPA were observed.

Conclusion: Few care-provider, environmental and policy factors were associated with children's in-care activity. UK childcare policies advocating child-driven play, moving freely indoors and outdoors, may be more conducive to individual children's PA.

Keywords: Preschool-aged children, physical activity, childcare, sedentary, policy, environment

Background

As the time children spend in out-of-home care increases, the childcare environment is likely to exert a greater influence on young children's activity.¹ Guidelines for under-5s recommend 180 minutes of total activity daily,^{2,3} including light (LPA; e.g. crawling, walking) and moderate-to-vigorous physical activity (MVPA; e.g. running, jumping). Yet low levels of MVPA⁴ in combination with high levels of sedentary behaviour appear common during the childcare day.⁵

Much of the evidence regarding levels of preschool-aged children's activity in childcare comes from the USA and mainland Europe⁶ (where 'preschool' is defined as 2.5/3-5/6 years depending on country⁷). Positive associations with preschool-aged children's physical activity have been reported for fixed (e.g. climbing frames) and portable (e.g. wheeled) toys, the presence of natural elements (e.g. vegetation), and staff education, training and behaviour in the playground.⁶ In contrast, qualitative work suggests that factors including parental concerns about child safety and emphasis on educational outcomes⁸ may result in greater sedentary behaviour. The childcare day in the United States (US), and to a lesser extent in mainland Europe,^{9,10} tends to include structured periods of learning and recess. In the United Kingdom (UK), settings operate a free-flow policy where regardless of weather conditions children self-select activities, both inside and out, for the majority of the day. Understanding how these contextual differences and elements in the UK childcare environment influence preschoolers' physical activity may be beneficial to inform research and practitioners internationally.

This study therefore sought to determine whether elements in the interpersonal, environmental and policy domains are associated with UK 3-4-year-old children's sedentary behaviour and physical activity when in childcare.

Methods

Study Design and Recruitment

Data were from the “Studying Physical Activity in preschool-aged Children and their Environment (SPACE) Study”.¹¹ Both preschool (state-run education) and nursery (privately-run care) ‘settings’ were purposively recruited to enable comparison, as they are (usually) differentially funded, operate in different built environments and vary in the care provided (see Table 1). Recruitment and data collection took place in January-July 2013. Detailed information about setting and child recruitment has been published elsewhere.¹¹ Briefly, 88 settings in Cambridgeshire were approached to participate; 30 (34%) setting managers provided written consent. Within settings, preschool-aged children were eligible to participate (n=602) if they: were 3-4-years-old; would be present on the designated measurement day; were free from physical disability; and attended the setting for at least 9 hours per week. Parents/guardians provided written consent; children provided verbal assent prior to measurement. A minimum of 5 participating children per setting was required to ensure sufficient analytical power. The University of Cambridge Psychology Ethics Committee provided ethical approval for the study (Pre.2012.68).

Data collection

At settings, we fitted children with an Actiheart activity monitor (Cambridge Neurotechnology Ltd, UK), a combined lightweight heart-rate monitor and accelerometer, previously validated in preschool-aged children.¹² The unit was secured to the chest, and set

to record at 15-second epochs. Written instructions were sent home to the parents, together with a previously validated questionnaire¹³ designed to assess potential correlates of physical activity. We encouraged children to wear the monitor continuously for ≤ 7 days, including during water-based activity and sleep.

Outcome variables

Counts data from Actiheart monitors were downloaded and processed using STATA 13/SE. Childcare attendance during the measurement week was reported by parents using a specially designed open-ended question.¹¹ To reflect when children were most likely to be active and/or in care, we restricted data to between 7am and 6pm (maximum 660 minutes). Although children would plausibly be awake outside these hours, they were not, according to parental report, in care. We removed data periods of ≥ 100 minutes of zero-activity counts,¹⁴ and days with < 600 minutes of recording¹⁵ (average in-care days: (mean \pm SD) 4.2 \pm 1.3days). We applied a previously validated conversion factor,¹⁶ and used validated cut points¹⁷ to classify children's activity as sedentary (SED: < 38 Actigraph counts per 15s); LPA (≥ 38 -420); and MVPA (≥ 421).¹⁷ Each child's activity and location data were matched in 15-minute segments.¹¹ Only 'in care' segments were used in the present analyses; outcome measures were expressed as average daily minutes per hour spent SED, in LPA and MVPA.

Exposure measures

A trained researcher assessed the setting environment using the validated Environment Policy Assessment and Observation (EPAO) tool.¹⁸ Responses to questions across 8 physical activity sub-domains from the EPAO were scored from 0-2 and totalled within a given domain to a possible maximum of 20 points, yielding 8 physical activity subscale scores.¹⁹

An overall physical activity environment score (possible range 0-160, higher score indicates more supportive environment) was also calculated for each setting ('EPAO score').

Additional exposure variables were chosen based on prior evidence.⁶ The average time staff had spent at the setting and as a childcare provider was taken from the questionnaire completed by each carer and used to calculate averages for each setting. Setting managers reported daily minutes children spent in gross motor play (in categories: ≤ 60 minutes; 61-120 minutes; 121-180 minutes, > 180 minutes), and five rules relating to outside play: in light rain, heavy rain, snow, wet conditions and high UV/sun (allowed always; in special clothing; never). Each setting's Office for Standards in Education (Ofsted) rating (satisfactory, good/outstanding), given following independent external review by trained inspectors, was obtained from the Ofsted website (<https://www.gov.uk/government/organisations/ofsted>).

Statistical Analyses

All children with ≥ 2 valid week-days of accelerometry data were included in analyses (n=201), and a pre-defined significance level of $p < 0.05$ was used for all analyses. Descriptive statistics were calculated and compared by setting type using t-tests for normal, Mann-U Whitney for non-normal or χ^2 tests for categorical data.

Three-level hierarchical linear regression models were fitted, assessing the associations between childcare-related factors and children's daily average minutes per hour of in-care SED, LPA and MVPA (Level 1: in-care activity; Level 2: child; Level 3: setting). Univariable regression models were first conducted to assess the association between each exposure variable and children's activity. All variables significantly associated in univariable models were subsequently entered into a multivariable regression model. Variables were removed

from the adjusted model if they did not meet the pre-defined significance level. All analyses were adjusted for sex, daily hours spent in care and parent-reported travel mode to childcare.

Results

Thirty settings (15 preschools, 15 nurseries) provided valid observational and questionnaire data (Table 1). Area deprivation scores for participating settings did not differ from those who declined to participate (participating: median 8.3 (Range: 1-27); declined: 8.6 (2-35); Wilcoxon rank-sum test: $p=0.48$). Compared to care-providers, setting managers were on average older (44.8 (SD: 9.4) vs. 35.4 (12.0) years old) and had worked in childcare for longer (13.5 (8.7) vs. 8.1 (5.8) years).

The mean total EPAO physical activity environment score was 85.9 (SD: 11.6; Range: 58.9-110.2). Mean subscale scores ranged from 4.7 (3.9; 0-20) for physical activity training and education to 15.3 (3.8; 6.7-20) for Active Opportunities; the average subscale score was 10.7 (1.5; 7.4-13.8) across all 8 scales.

Associations between children's in-care activity and the preschool environment

In univariable analyses, four factors were associated with children's in-care SED; no factors were associated with children's in-care LPA and MVPA (Table 2). Only Active Opportunities remained significantly associated with SED in adjusted models. Children's in-care activity did not cluster within setting (intraclass correlation coefficients (ICCs): SED: 0.04; LPA: 0.003; MVPA: 0.05).

Table 1 Characteristics of participating settings by type

	All settings (n=30)	Nursery ^a (n=15)	Preschool ^b (n=15)
Interpersonal			
Children enrolled at setting ^c (mean(SD))	72 (52)	95 (58)	46 (28)
3-4 year-olds enrolled at setting (mean(SD))	44 (30)	49 (33)	38 (25)
Class composition (n (%))			
2-4 year olds	13 (43)	6 (40)	4 (27)
3-4 year olds	17 (57)	9 (60)	11 (73)
% Non-white children (mean (SD))	11.2 (13.6)	15.0 (17.7)	7.4 (6.6)
Government funded places (mean (SD))	33 (24)	27 (15)	37 (30)
Children per staff member ^d (mean (SD))	3.2 (7.1)	3.2 (9.0)	3.2 (5.6)
<i>Preschool Staff (all mean (SD))</i>			
Age in years	38.9 (8.5)	34.9 (7.9)	43.6 (6.7)
Years at setting	6.3 (3.4)	6.6 (3.7)	6.2 (3.3)
Years in childcare	9.7 (5.3)	8.9 (3.3)	10.8 (6.8)
Environmental			
Number of hours observed ^{**} (mean (SD))	7.1 (2.4)	9.1 (1.0)	5.1 (1.5)
Fixed equipment ^e (mean (SD))	4.8 (1.7)	5.0 (1.7)	4.6 (1.7)
Portable equipment ^e (mean (SD))	6.1 (1.5)	6.2 (1.6)	6.1 (1.5)
Reported time spent in GMP (n (%))			
0-60 minutes	4 (13)	2 (13)	2 (13)
61-120 minutes	8 (27)	2 (13)	6 (40)
121-180 minutes	7 (23)	3 (20)	4 (27)
>180 minutes	11 (37)	8 (53)	3 (20)

GMP: Gross Motor Play; a: Nursery: offers full day care (~7am-6pm) for children <1 year up to 4 years 11 months, usually privately run; b: offers sessional care (~9am-12noon and/or 12noon-3pm) for children between 2 years 9 months and 4 years 11 months old, usually state-run; c: Number of children enrolled at setting includes all children who attend on weekly basis, regardless of age and study eligibility; d: Calculated as a ratio: number of children in room /number of staff in room; e: refers to the average number of pieces of fixed/ portable play equipment visible at setting.

Significant difference by setting type: * $p < 0.05$; ** $p < 0.005$

Table 2 Associations between children's in-care activity and (elements in) the preschool environment

Exposure	Outcome [β (95% CI)]					
	SED		LPA		MVPA	
	Univariable	Multivariable	Univariable	Multivariable	Univariable	Multivariable
Interpersonal						
3-4 year-olds enrolled at setting (% of total)	-0.0 (-0.1, 0.0)	-	-0.0 (-0.0, 0.0)	-	0.0 (-0.1, 0.1)	-
Class composition (3-4yrs) (ref: 2-4yrs)	0.4 (-1.6, 2.4)	-	1.5 (-0.1, 3.2)	-	-1.8 (-4.6, 1.0)	-
Government funded places	-0.0 (-0.1, 0.0)	-	-0.0 (-0.0, 0.0)	-	-0.0 (-0.0, 0.0)	-
Children per staff member ^a	-4.3 (-14.7, 6.2)	-	3.7 (-4.4, 11.8)	-	-0.7 (-12.6, 11.3)	-
Staff mean age in years	-0.1 (-0.2, 0.1)	-	0.0 (-0.1, 0.1)	-	0.1 (-0.1, 0.2)	-
Staff mean years at setting	0.2 (-0.1, 0.6)	-	0.2 (-0.1, 0.5)	-	-0.4 (-0.9, 0.1)	-
Staff mean years in childcare	-0.1 (0.3, 0.1)	-	0.1 (-0.1, 0.2)	-	0.1 (-0.2, 0.3)	-
Staff Behaviour [§]						
Environmental						
Active opportunities [§]	1.9 (0.9, 2.9) ^{***}	1.9 (0.9, 2.9) ^{***}	-0.8 (-1.7, 0.1)	-	-1.1 (-2.6, 0.4)	-
Sedentary opportunities	0.1 (-1.4, 1.6)	-	-0.4 (-1.6, 0.9)	-	0.2 (-1.9, 2.2)	-
Fixed Equipment [§]	-0.1 (-0.5, 0.2)	-	-0.2 (-0.5, 0.1)	-	0.4 (-0.1, 0.9)	-
Portable Equipment [§]	-0.2 (-0.5, 0.1)	-	0.2 (-0.1, 0.4)	-	0.0 (-0.4, 0.5)	-
Sedentary environment [§]	0.2 (-0.1, 0.4)	-	-0.1 (-0.3, 0.2)	-	-0.1 (-0.5, 0.2)	-
Time allowed outside [¥] (%)	-0.0 (-0.1, 0.0)	-	-0.0 (-0.1, 0.0)	-	-0.0 (-0.1, 0.1)	-
Time children seated [¥] (%)	0.1 (-0.0, 0.2)	-	-0.0 (-0.1, 0.1)	-	-0.1 (-0.2, 0.1)	-
Reported time spent in GMP (ref: 0-60)						
61-120	1.2 (-2.0, 4.4)	-	-0.4 (-3.1, 2.3)	-	-0.7 (-5.3, 3.8)	-
121-180	3.1 (0.4, 7.0) [*]	ns	-1.4 (-4.2, 1.4)	-	-2.4 (-7.0, 2.2)	-
181+	3.4 (0.4, 6.4) [*]	ns	-1.1 (-3.6, 1.4)	-	-2.3 (-6.5, 1.9)	-

	Outcome β (95% CI)					
	SED		LPA		MVPA	
	Univariable	Multivariable	Univariable	Multivariable	Univariable	Multivariable
Policy						
Physical Activity Training and Education [§]	0.8 (-0.5, 2.1)	-	0.7 (-0.4, 1.8)	-	-1.3 (-3.1, 0.5)	-
Physical Activity Policies [§]	-1.3 (-2.8, 0.2)	-	1.1 (-0.2, 2.3)	-	0.3 (-1.8, 2.4)	-
Play outside in light rain: (ref: clothes)						
Always	-0.7 (-3.4, 1.9)	-	0.2 (-2.1, 2.4)	-	0.7 (-3.0, 4.4)	-
Play outside in heavy rain: (ref: with clothes)						
Always	-0.1 (-3.1, 2.9)	-	-0.1 (-2.5, 2.4)	-	0.1 (-4.1, 4.2)	-
Play outside in snow: (ref: with clothes)						
Always	2.1 (0.1, 4.0)*	ns	-0.9 (-2.6, 0.7)	-	-1.1 (-3.8, 1.6)	-
Play outside in sun: (ref: with clothes)						
Always	-1.2 (-5.3, 2.9)	-	2.9 (-3.9, 9.0)	-	-1.9 (-16.2, 12.3)	-
Ofsted score [¥] (ref: satisfactory)						
Good/Outstanding	1.4 (-1.3, 3.2)	-	-1.4 (-3.7, 0.9)	-	-0.1 (-4.0, 3.8)	-
Between child variance (Mean (Std Error))	4.50 (0.51)		3.82 (0.46)		7.59 (0.62)	
Within child variance (Mean (Std Error))	8.44 (0.26)		7.72 (0.23)		10.78 (0.33)	

β: Minutes of activity per hour in care; ^a Calculated as a ratio: number of children in room /number of staff in room; [§] Denotes EPAO subscale score used; [¥] from setting observation; all other variables taken from the setting questionnaire; ns: not significant in adjusted analyses; ‘-’: not entered into adjusted analyses; GMP: Gross Motor Play; All analyses adjusted for sex, daily hours in care and mode of travel to preschool: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Discussion

This is the first study to investigate associations between factors in the UK childcare environment and preschoolers' physical activity, showing that childcare variables explain little variation in children's activity. Although several interpersonal and environmental-level factors were associated with children's in-care sedentary behaviour in univariable analyses, only one remained in multivariable models. No factors were associated with in-care LPA and MVPA. This suggests the UK childcare environment may have a limited influence on children's activity, being conducive to children's individual activity preferences instead. How individual and unexplored social factors affect children's in-care activity now warrants further investigation, and may be useful when exploring ways to increase activity in lesser active children of this age.

Previous work conducted in the USA, using both direct observation and accelerometers to measure children's activity, showed that children's activity levels were primarily affected by the setting they attended.^{20,21} In contrast, children's in-care activity levels appeared not to cluster within settings here, with ICCs of 0.003-0.05 similar to those seen in a Danish study assessing preschoolers' objectively measured in-care activity.²² Variation in the childcare day may in part contribute to these differences. Structured periods of play, recess, and group teaching tends to occur in US and mainland European countries.⁹ For example, one study comparing differences between children's average activity in US and Swedish childcare centres found US children spent more time indoors, with greater MVPA observed when children were outdoors.⁹ In contrast, free-flow policies in the UK encourage children to select their own activities, both inside and out, for the majority of the day. A less structured childcare day may therefore result in the childcare environment exerting a smaller influence on UK children's activity. Given our findings, adoption of a less structured childcare day may therefore be one way for practitioners to positively influence young children's physical activity levels, and may be piloted relatively easily.

Additionally, our and the Danish study²² assessed children's individual-level daily in-care activity and used multi-level analyses to capture within-child fluctuations, which may better represent children's actual in-care activity levels. We identified larger fluctuations in within-child compared to between-child daily activity (when in care), which may reflect children's

self-selection of activities and UK childcare policies. As such, individual and social factors, may therefore be a stronger driver of children's in-care physical activity levels in the UK.^{19,23}

That few associations were found between childcare-related factors and children's in-care activity here may corroborate this. The EPAO has been used to assess childcare environments in our study, as well as in US¹⁹ and Dutch²³ studies, with similar average subscale scores seen (10.7 vs 10.2 in the US study¹⁹; not reported in Dutch study). Only the (unexpected) positive association between increased active opportunities and sedentary time remained significant in adjusted models. In contrast, in a Dutch cohort of 2-3 year olds, EPAO-assessed childcare active opportunities were positively associated with directly-observed higher intensity activity.²³ In the US, children in more supportive childcare environments were shown to have greater active and sedentary opportunities, spend more time in MVPA and less time sedentary.¹⁹ Notwithstanding the variation in outcome measures used, it is possible that differences in associations seen between these studies are indeed a result of cultural or operational differences in the childcare environment, which the EPAO was not designed to identify.

Strengths and Limitations

Previous studies assessing the influence of the childcare environment on children's activity have used direct observation or accelerometers to provide an aggregated (childcare-level) overview of children's physical activity levels.^{19,23} We used an objective measure to capture children's individual-level daily activity, which may reduce potential biases associated with direct observation. Staff were blinded to study aims to minimise bias and avoid behaviour change during the EPAO observation; no staff-related behaviours appeared to influence children's in-care activity here.

Children's actual in-care hours may have varied from those reported, resulting in misclassification of 'in-care' time; we adjusted for usual mode of transport to care to account for variation in actual / reported arrival time. Every effort was made to use accelerometry data reported to have occurred at the observed setting, but 7% of children attended two different settings during the measurement week and it was not possible to determine the participating setting. This may have attenuated the association between childcare factors and activity. However, post-hoc sensitivity analyses excluding these children did not alter the

overall conclusions. Finally, though heterogeneity in environment scores between settings is similar to those reported previously,¹⁹ insufficient variation in exposures may have contributed to the limited number of significant associations seen here.

Conclusion

This is the first work to assess the UK childcare physical activity environment and determine factors associated with children's in-care activity. Children's activity appeared not to cluster by setting, suggesting that the childcare environment may have a limited influence on children's in-care physical activity in the UK. This is supported by the finding that few investigated factors appear to be associated with children's in-care activity behaviour. Other locations or social groupings (e.g. parent-child groups) may prove more appropriate to facilitate and encourage activity amongst UK preschoolers.

Conflict of Interest Statement

The authors declare no conflict of interest.

Acknowledgements

We thank all children and their parents who participated in the SPACE study. In addition, we thank Kate Westgate and Stefanie Mayle from the physical activity technical team at the MRC Epidemiology Unit for their assistance in processing the accelerometer data, and members of the field team who conducted data collection. This work was conducted by the Medical Research Council [Unit Programme numbers MC_UU_12015/7; MC_UU_12015/4] and the Centre for Diet and Activity Research (CEDAR), a UKCRC Public Health Research Centre of Excellence. Funding from the British Heart Foundation, Economic and Social Research Council, Medical Research Council, National Institute for Health Research, and Wellcome Trust, under the auspices of the UK Clinical Research Collaboration, is gratefully acknowledged (CEDAR grant number: RES-590-28-0002). No financial disclosures were reported by the authors of this paper.

References

1. Ward DS, Vaughn A, McWilliams C, Hales D. Interventions for increasing physical activity at child care. *Med Sci Sport Exerc.* 2010;42(3):526–534.
2. Department of Health. *Start Active, Stay Active: A report on physical activity from the four home countries' Chief Medical Officers.* London, UK; 2011.
3. Tremblay MS, Leblanc AG, Carson V, et al. Canadian Physical Activity Guidelines for the Early Years (aged 0-4 years). *Appl Physiol Nutr Metab.* 2012;37(2):345–356.
4. Tucker P. The physical activity levels of preschool-aged children: A systematic review. *Early Child Res Q.* 2008;23(4):547–558.
5. Reilly JJ. Low levels of objectively measured physical activity in preschoolers in child care. *Med Sci Sport Exerc.* 2010;42(3):502–507.
6. Trost SG, Ward DS, Senso M. Effects of child care policy and environment on physical activity. *Med Sci Sport Exerc.* 2010;42(3):520–525.
7. The World Bank. Primary school starting age (years). 2013. Available at:<http://data.worldbank.org/indicator/SE.PRM.AGES>.
8. Copeland KA, Sherman SN, Kendeigh CA, Kalkwarf HJ, Saelens BE. Societal Values and Policies May Curtail Preschool Children's Physical Activity in Child Care Centers. *Pediatrics.* 2012;129(2):1–10.
9. Raustrop A, Pagels P, Boldemann C, Cosco N, Soderstrom M, Martensson F. Accelerometer Measured Level of Physical Activity Indoors and Outdoors During Preschool Time in Sweden and the United States. *J Phys Act Heal.* 2012;9(6):801–808.
10. Cardon GM, De Bourdeaudhuij IM. Are Preschool Children Active Enough?

- Objectively measured physical activity levels. *Res Q Exerc Sport*. 2008;79(3):326–332.
11. Hesketh KR, Griffin SJ, van Sluijs EMF. UK Preschool-aged children’s physical activity levels in childcare and at home: a cross-sectional exploration. *Int J Behav Nutr Phys Act*. 2015;12(1):123.
 12. Adolph AL, Puyau MR, Vohra FA, Nicklas TA, Zakeri IF, Butte NF. Validation of uniaxial and triaxial accelerometers for the assessment of physical activity in preschool children. *J Phys Act Heal*. 2012;9(7):944–953.
 13. McMinn AM, van Sluijs EM, Harvey NC, Cooper C, Inskip HM, Godfrey KM, Griffin SJ. Validation of a maternal questionnaire on correlates of physical activity in preschool children. *Int J Behav Nutr Phys Act*. 2009;6:81.
 14. Collings PJ, Brage S, Ridgway CL, Harvey NC, Godfrey KM, Inskip HM, Cooper C, Wareham NJ, Ekelund U. Physical activity intensity, sedentary time, and body composition in preschoolers. *Am J Clin Nutr*. 2013;97(5):1020–1028.
 15. Beets MW, Bornstein D, Dowda M, Pate RR. Compliance with national guidelines for physical activity in U.S. preschoolers: measurement and interpretation. *Pediatrics*. 2011;127(4):658–664.
 16. Ridgway CL, Brage S, Sharp SJ, Corder K, Westgate KL, van Sluijs EM, Goodyer IM, Hallal PC, Anderssen SA, Sardinha LB, Andersen LB, Ekelund U. Does birth weight influence physical activity in youth? A combined analysis of four studies using objectively measured physical activity. *PLoS One*. 2011;6(1):e16125.
 17. Pate RR, Almeida MJ, McIver KL, Pfeiffer KA, Dowda M. Validation and Calibration of an Accelerometer in Preschool Children. *Obesity*. 2006;14(11):2000–2006.
 18. Ward DS, Benjamin SE, Ammerman AS, Ball SC, Neelon BH, Bangdiwala SI.

Nutrition and physical activity in child care: results from an environmental intervention. *Am J Prev Med.* 2008;35(4):352–356.

19. Bower JK, Hales DP, Tate DF, Rubin DA, Benjamin SE, Ward DS. The Childcare Environment and Children's Physical Activity. *Am J Prev Med.* 2008;34(1):23–9.
20. Pate RR, Pfeiffer KA, Trost SG, Ziegler P, Dowda M. Physical activity among children attending preschools. *Pediatrics.* 2004;114(5):1258–63.
21. Pate R, McIver K, Dowda M, Brown W, Addy C. Directly Observed Activity Levels in Preschool Children. *J Sch Health.* 2008;78(8):438–444.
22. Groenholt Olesen L, Lund Kristensen P, Korsholm L, Froberg K. Physical Activity in Children Attending Preschools. *Pediatrics.* 2013;132:e1310–1318.
23. Gubbels JS, Kremers SP, van Kann DH, Stafleu A, Candel MJ, Dagnelie PC, Thijs C, de Vries NK. Interaction between physical environment, social environment, and child characteristics in determining physical activity at child care. *Heal Psychol.* 2011;30(1):84–90.