



## Original article

Scand J Work Environ Health 2017;43(1):5-14

doi:10.5271/sjweh.3587

### Job strain and informal caregiving as predictors of long-term sickness absence: A longitudinal multi-cohort study

by [Mortensen J](#), [Dich N](#), [Lange T](#), [Alexanderson K](#), [Goldberg M](#), [Head J](#), [Kivimäki M](#), [Madsen IEH](#), [Rugulies R](#), [Vahtera J](#), [Zins M](#), [Rod NH](#)

This study is the first to investigate the joint effects of job strain and informal caregiving on long-term sickness absence. The main finding was that informal caregiving responsibilities and/or high job strain predicted long-term sickness absence among women.

**Affiliation:** Department of Public Health, University of Copenhagen, Gothersgade 160, Copenhagen, Denmark. [jemo@sund.ku.dk](mailto:jemo@sund.ku.dk)

Refers to the following text of the Journal: [2015;41\(5\):421-507](#)

The following article refers to this text: [2017;43\(1\):1-96](#)

**Key terms:** [gender difference](#); [informal caregiving](#); [interaction](#); [job strain](#); [long-term sickness absence](#); [multi-cohort](#); [predictor](#); [sick leave](#); [sickness absence](#); [unpaid care](#); [women](#)

This article in PubMed: [www.ncbi.nlm.nih.gov/pubmed/27556905](http://www.ncbi.nlm.nih.gov/pubmed/27556905)

### Additional material

Please note that there is additional material available belonging to this article on the [Scandinavian Journal of Work, Environment & Health -website](#).



This work is licensed under a [Creative Commons Attribution 4.0 International License](#).

## Job strain and informal caregiving as predictors of long-term sickness absence: A longitudinal multi-cohort study

By Jesper Mortensen, MHS<sup>1</sup>, Nadya Dich, PhD,<sup>1</sup> Theis Lange, PhD,<sup>2</sup> Kristina Alexanderson, PhD,<sup>3</sup> Marcel Goldberg, PhD,<sup>4</sup> Jenny Head, MSc,<sup>5</sup> Mika Kivimäki, PhD,<sup>5</sup> Ida EH Madsen, PhD,<sup>6</sup> Reiner Rugulies, PhD,<sup>6</sup> Jussi Vahtera, PhD,<sup>7</sup> Marie Zins, MD,<sup>4</sup> Naja Hulvej Rod, DMSc<sup>1</sup>

Mortensen J, Dich N, Lange T, Alexanderson K, Goldberg M, Head J, Kivimäki M, Madsen IEH, Rugulies R, Vahtera J, Zins M, Rod NH. Job strain and informal caregiving as predictors of long-term sickness absence: A longitudinal multi-cohort study. *Scand J Work Environ Health*. 2017;43(1):5–14. doi:10.5271/sjweh.3587

**Objectives** The aim of this study was to investigate the individual, joint and interactive effects of job strain and informal caregiving on long-term sickness absence with special attention to gender differences.

**Methods** The study comprised a prospective cohort study of 6798 working adults from France, 14 727 from Finland, and 5275 from the UK. A total of 26 800 participants, age 52 (interquartile range 47–56) years participated in the study. Job strain was assessed using the demand–control model. Informal caregiving was defined as care for a sick, disabled, or elderly person. Long-term sickness absence spells defined as absence >14 consecutive days were registered during two years follow-up. We used recurrent-events Cox regression in random-effects meta-analyses.

**Results** A total of 12% men and 21% women had ≥1 long-term sickness absence spell. Among women, both high job strain [hazard ratio (HR) 1.08, 95% confidence interval (95% CI) 1.00–1.17] and informal caregiving (HR 1.13, 95% CI 1.04–1.23) were associated with a modestly higher risk of sickness absence. Women doubly exposed to high job strain and informal caregiving also showed a moderately higher risk of sickness absence (HR 1.20, 95% CI 1.03–1.41), but the excess risk was not more than expected from joint exposure to caregiving and job strain. Neither job strain nor informal caregiving predicted sickness absence for men.

**Conclusions** High job strain and informal caregiving predicted long-term sickness absence among women. However there was no noticeable interaction in the presence of both exposures.

**Key terms** gender difference; interaction; sick leave; unpaid care; women.

Informal caregiving, defined as unpaid assistance with activities of daily living for sick, disabled, or elderly relatives, contributes invaluablely to the well-being of populations in western countries and is becoming a key issue in many countries due to rapid population aging (1). Informal caregiving may entail benefits such as satisfaction and meaning for the caregiver (2). However, the combination of prolonged distress, physiological demands, financial strain and social isolation that sometimes accompanies informal caregiving may eventually compromise the caregivers' own health (3–6).

According to the role accumulation theory, having multiple roles from paid work and informal caregiving may carry positive effects for the caregiver (7) as one role may buffer stressors related to other roles (8). In contrast, the role strain theory hypothesizes that individuals who provide informal caregiving may be particularly vulnerable to its potential health consequences if they are simultaneously exposed to other stressors such as a high-strain job (9). This double exposure may lead to work–family conflicts introduced by the experience of role conflict (ie, being subject to conflicting

<sup>1</sup> Department of Public Health, University of Copenhagen, Copenhagen, Denmark.

<sup>2</sup> Department of Biostatistics, University of Copenhagen, Copenhagen, Denmark.

<sup>3</sup> Insurance Medicine, Karolinska Institutet, Stockholm, Sweden.

<sup>4</sup> INSERM, Population-based Epidemiological Cohorts Unit-UMS 11, and INSERM, UMR 1168, VIMA, Villejuif, France.

<sup>5</sup> Department of Epidemiology and Public Health, University College London, London, UK.

<sup>6</sup> National Research Centre for the Working Environment, Copenhagen, Denmark.

<sup>7</sup> Department of Public Health, University of Turku, Turku, Finland.

Corresponding author: Jesper Mortensen, Department of Public Health, University of Copenhagen, Gothersgade 160, Copenhagen, Denmark. [email: jemo@sund.ku.dk]

demands) or role overload (ie, having too much to do) (10) and eventually render the caregiver more vulnerable to negative health consequences.

Sickness absence has a large impact on employees, work sites, and society and provides a measure of work incapacity as a social consequence of morbidity (11). In two recent longitudinal studies from Australia and Belgium and a case–control study from Brazil, it has been shown that high job strain predicts long-term sickness absence (12–14). In contrast, one study only found an association for men in adjusted models (15), and another study found no association for either men or women in adjusted models (16). A few studies have also investigated the association between work–family conflicts and sickness absence (17–19). A large study including workers from 31 European countries found that work–life imbalance predict sickness absence (19), and a Finnish study showed that spillover from work into family life predicted a heightened rate of sickness absence spells (17). However, research on work–family conflicts has so far primarily focused on perceived imbalances between work and family life without distinguishing between different sources to this imbalance (20). Only few studies have investigated the combined effect of work and family factors on sickness absence (21–24). Two studies, based on the same occupational cohort, showed that the combination of high work and family demands predicted all-cause sickness absence and sickness absence due to mental health disorders (21, 22). In those studies, family demands were defined by the number of individuals the participant supported economically. Informal caregiving, however, involves both physically and mentally straining elements and is thus different from providing economic support. Looking specifically at informal caregiving, a study from Austria has shown that a time-based conflict between informal eldercare and paid work was related to the intended job change of female workers and that the intensity of caregiving provided was related to male workers' anticipated labor market exit (25). However, to our knowledge no studies have investigated the association between informal caregiving and sickness absence due to paid work.

Quantifying the separate and joint effects of job strain and informal caregiving on sickness absence will help in the identification of high-risk workers. Such information is important for policy-makers and employers to develop special need support and efficient preventive strategies (26).

Women are traditionally more engaged in informal caregiving than men, and it has been suggested that women may be especially vulnerable to the joint health effects of demanding work and life circumstances (27, 28). Furthermore, women in general still carry the largest household workload, and it has been shown that women have higher evening cortisol levels than men

(29), indicating that they may have difficulties relaxing and unwinding after work.

In this multi-cohort study, the objective was to investigate the joint effects of job strain and informal caregiving on long-term sickness absence with special attention to gender differences, using data from three large European occupational cohorts. Based on the role strain theory (9), we hypothesized that the joint effects of high job strain and informal caregiving on long-term sickness absence exceed the combination of their individual associations. We additionally hypothesized that women were at greater risk of sickness absence compared to men when exposed to both high job strain and informal caregiving.

## Methods

### Study participants

We used longitudinal data from the GAZEL cohort from France (30), the Finnish Public Sector Study (FPS) from Finland (31), and the Whitehall II study from the UK (32). These cohorts were chosen as they have detailed data on job strain, informal caregiving, and longitudinal register-based data on sickness absence.

The GAZEL cohort was established in 1989 and includes 20 625 employees of the French national gas and electricity company (response rate, approximately 75% annually).

FPS was established in 1997 and comprises all 151 901 public employees in ten towns and five hospital districts in Finland. A nested open cohort with 16 948 responders at baseline (baseline response rate, 70%), FPS includes data on psychosocial work and domestic factors, along with health outcomes.

Whitehall II was established in 1985 and includes 10 308 British civil servants from 20 London-based departments (baseline response rate, 73%).

Further information about each cohort can be found in the cohort profiles (30–32). We used data from year 2000 as baseline in GAZEL, year 2012 in FPS, and years 1991–1994 in Whitehall II, due to the availability of data on informal caregiving in the present years. We included individuals in paid work and excluded individuals on sick leave at the day of the baseline questionnaires. The present study comprised 6798 individuals from GAZEL, 14 727 individuals from FPS, and 5275 individuals from Whitehall II, a total of 26 800 individuals in paid work. Participants gave written consent to participate in the cohort studies, and all three cohorts have been approved by the respective human ethics committees. The data selection procedure is presented in a flow chart in the appendix ([www.sjweh.fi/index.php?page=data-repository](http://www.sjweh.fi/index.php?page=data-repository)).

## Measures

**Informal caregiving.** Informal caregiving was treated as a binary (yes/no) variable based on slightly different questions in each cohort. In GAZEL, individuals were asked if they provide regular care for an aged person (>65 years); in FPS individuals were asked if they provide care for a sick, disabled, or aged relative; and in Whitehall II individuals were asked if they provide care for an aged or disabled relative. To investigate the effect of weekly hours of informal caregiving, we applied a cut-off at >4 weekly hours, indicating caregiving of  $\geq 1$  hour (average) on each weekday. Information on weekly hours of caregiving was only available in FPS and Whitehall II.

**Job strain.** Job strain (33) was assessed using the Job Content Questionnaire in GAZEL and FPS and the Demand–Control Questionnaire in Whitehall II (34). High job strain was in accordance with the job strain model (33) defined as the combination of high job demands and low job control; all other combinations of job demands and job control was defined as no high strain (34). Job demands were assessed with five items in GAZEL and four items in Whitehall II and FPS, and encompass statements such as “My job requires working very fast”. High job demands were defined as a score above the median score of the specific study population. Job control (decision latitude) is comprised of two subscales: skill discretion and decision authority. Skill discretion was assessed with four items in all cohorts, and encompasses statements such as “My job requires a high level of skill”. Decision authority was assessed with two items in all cohorts, and encompasses statements such as “I have a lot of say about what happens on my job”. The two items in decision authority are weighted the same as the four items in skill discretion, in calculation of job control scores. Low job control was defined as a job control score below the median score of the specific study population (34). In GAZEL, job strain in 1999 was used as a proxy for information in 2000.

**Long-term sickness absence.** We used all-cause long-term sickness absence from paid work as the outcome, defined as spells of >14 consecutive days. Long-term sickness absence has shown to be a useful global measure of health differentials between employees (11, 35, 36), as opposed to shorter sickness absence spells, which may in some cases represent healthy coping behaviors to avoid serious morbidity (11, 35, 36). Furthermore, long-term spells were used as opposed to number of days absent, as spells  $\leq 14$  days are sometimes based on self-report in the three cohorts, and we wanted to prevent any reporting bias. Sickness absence information was from national registers in FPS and Whitehall II, and

company register in GAZEL (17, 22, 37). Recorded sickness absence was due to own morbidity and not absence due to, eg, maternity leave or care for a sick child. We used a follow-up period of two years in registration of long-term sickness absence. This period was chosen as longer follow-up might entail a weaker causal path between baseline exposures and later sickness absence.

## Confounders

We identified confounders for analyses using directed acyclic graphs (DAG) (38) (see appendix, [www.sjweh.fi/index.php?page=data-repository](http://www.sjweh.fi/index.php?page=data-repository)). Included variables were: age of children (0, 1–3, >3 years), occupational grade in three groups (low included low wage and manual laborers, intermediate included lower non-manual and midlevel managers, and high included upper non-manual workers and administrative staff), married/cohabiting, death of a close relative and/or divorce within the past 12 months,  $\geq 1$  spell of long-term sickness absence in the two years preceding baseline,  $\geq 1$  longstanding illness at baseline (diabetes, cardiovascular disease, cancer or respiratory disease).

## Statistical analysis

We used individual participant data in Cox proportional hazards random effects meta-analyses (39). This approach can be thought of as a type of multilevel model (with cohorts as cluster units) on the proportional hazard scale. Heterogeneity among cohort specific estimates were assessed with  $I^2$  statistics (40). A random effects model was chosen to allow for the true effect to vary between the included cohorts; eg, the effect size of informal caregiving on sickness absence could be greater when questions on caregiving encompass sick, disabled, or aged relatives in FPS and Whitehall II compared to only aged relatives in GAZEL.

Person-years at risk encompassed time until the occurrence of a long-term sickness absence spell or until censoring due to retirement, disability pension/incapacity benefit, death, or leaving the workplace. Sickness absence for spells  $\leq 14$  days were included in time-at-risk of long-term sickness absence based on the argument that one would still be at risk of long-term sickness absence during shorter spells of sickness absence.

To account for recurrent events of long-term sickness absence we used the Prentice, Williams and Peterson Total Time (PWP-TT) model (41), which is appropriate if the occurrence of the first event increases the likelihood of a recurrence. In PWP-TT, multiple events are ordered by stratification based on the prior number of events, such that all participants are at risk of an event in the first stratum, but only those with a prior event are at risk for a successive event (41). We truncated analy-

ses at two spells of long-term sickness absence during follow-up, due to small number of individuals with a greater number of spells.

We estimated hazard ratios (HR) and 95% confidence intervals (CI) with long-term sickness absence as the event of interest, for the individual associations with job strain and informal caregiving and the joint associations compared to those with no high strain and no informal caregiving as reference group (42). We then carried out similar analyses adjusted for the mentioned confounders. Confounders are presented in table 1 along with possible mediating lifestyle factors including smoking, alcohol, and overweight. We assessed additive interaction with the synergy index (SI) (43), which can be interpreted as the excess risk from double exposures when there is interaction relative to the risk from exposure without interaction, and further tested for multiplicative interaction. Analyses were conducted separately for men and women to address gender differences. It was not feasible to test formally for 3-way interaction with job strain, caregiving and gender due to lack of statistical power.

We carried out several sensitivity analyses to test the robustness of our results. In the first sensitivity analysis we used multiple imputation on the 14% of individuals with missing data, assuming that data were missing at random (44). We imputed missing data with a chained equation model and applied Rubin’s rules at the study level prior to meta-analysis (45). Results from the imputed analyses were compared to results from the complete case analyses to investigate selection bias introduced by excluding participants with missing data. Secondly, retirement may be a competing risk to long-term sickness absence and therefore we carried out a sensitivity analysis in which we excluded individuals >58 years at

baseline to account for retirement from the workforce due to the statutory pension age. Thirdly, by only looking at number of spells >14 days, individuals with very long spells may not have contributed with risk time following the first spell, as the first spell may have lasted the whole follow-up period. This may potentially underestimate the consequences of job strain and caregiving exposure on the risk of sickness absence. Therefore, we also made a sensitivity analysis where we only looked at time to the first event of long-term sickness absence. In a fourth analysis, we included weekly work hours as a confounding variable (<36 hours, 36-40 hours, >40 hours). This information was only available in Whitehall II. We also looked whether job strain and caregiving predicted the length of long-term sickness absence spells in the subpopulation of individuals with long-term spells. Lastly, we did a sensitivity analysis where we looked at job demands and job control separately to see if they were better predictors of sickness absence than the combined job strain measure. We used the ipdmetan statistical software package in STATA v13/IC for analyses. Ipdmetan performs two-stage individual participant-data meta-analysis using the inverse-variance method. The syntax used is “ipdmetan, study(studyID) : stcox...”, which fits the model command once within each level of study ID.

**Results**

In this population of 26 800 employees, there were 5946 spells of long-term sickness absence during 46 794 person-years. A total of 12% of male and 21% of female participants experienced ≥1 spell of long-term sickness absence within a 2-year period. When truncating at two

**Table 1.** Baseline characteristics according to cohort and gender. [FPS=Finnish Public Sector Study; M=median; IQR=interquartile range.]

	GAZEL: France (Baseline, year 2000)						FPS: Finland (Baseline, year 2012)						Whitehall II: UK (Baseline, year 1994)					
	Men (N=4546)			Women (N=2252)			Men (N=2999)			Women (N=11 728)			Men (N=3786)			Women (N=1489)		
	%	M	IQR	%	M	IQR	%	M	IQR	%	M	IQR	%	M	IQR	%	M	IQR
Age (years)		53	52–54		50	48–53		52	46–57		53	47–57		48	44–53		50	45–55
Married/cohabiting	92			71			82			74			82					63
Children <sup>a</sup>	95			87			84			86			58					30
Death of relative/divorce <sup>b</sup>	17			18			6			6			31					38
Low occupational grade	12			25			30			33			7					38
High alcohol intake	23			9			9			10			18					10
Smoking	16			16			17			12			13					17
Overweight	63			30			66			51			47					48
Previous long-term sickness absence	14			23			22			27			9					21
Longstanding illness (≥1)	11			9			25			26			11					12
High job strain	13			24			11			20			17					22
Informal caregiving	27			30			10			15			10					13

<sup>a</sup> Children in household in Whitehall II.

<sup>b</sup> In FPS, death of relative or divorce for the preceding part 2012 only (not one year previous to baseline).

years, the median length of the first event of long-term sickness absence spells was 29 days (15–436), 28 days (15–370), and 28 days (15–724) in Whitehall II, FPS and GAZEL, respectively. Clear gender differences were evident for marital status, with more men being married/cohabiting than women (table 1). Longstanding illness was most prevalent in FPS compared to GAZEL and Whitehall II. Both high job strain and informal caregiving were more prevalent for women (table 1), and informal caregiving was more prevalent in GAZEL with 27% of men and 30% of women providing care, compared to 10% and 15% in FPS and 10% and 13% in Whitehall II.

For men, neither informal caregiving nor high job strain were associated with a higher risk of long-term sickness absence in the unadjusted analyses and the multiple adjusted analyses, as seen in table 2. Among women, high job strain and informal caregiving were associated with long-term sickness absence in the unadjusted model, which was slightly attenuated in the multiple adjusted model, as seen in table 2.

Stratifying by hours of caregiving in a sub-analysis, caregiving >4 weekly hours was a risk factor for sickness absence among women in the unadjusted and multiple adjusted models (table 2). No associations were found between weekly hours of caregiving and long-term sickness absence for men in neither the unadjusted nor the multiple adjusted models.

Looking at the joint exposure, high job strain and no informal caregiving predicted long-term sickness absence for men in the unadjusted model with (HR 1.30, 95% CI 1.01–1.67). However, this was considerably attenuated in the multiple adjusted model with (HR 1.14, 95% CI 0.97–1.35), as seen in figure 1. For women, no high job strain and informal caregiving (HR 1.15, 95%

CI 1.01–1.29) was a predictor of long-term sickness absence in the multiple adjusted model, but the excess risk from exposure to both high job strain and informal caregiving was not more than expected (HR 1.20, 95% CI 1.03–1.41), ( $P=0.766$ ,  $SI=0.95$ ), as seen in figure 1. All estimates in the joint exposure analyses for both men and women were attenuated in the multiple adjusted models compared to the unadjusted models.


Sensitivity analyses using multiple imputations did not change results substantially (appendix), indicating that the partial missing data have not introduced any noteworthy selection bias. Furthermore, there was no noteworthy change in results from the sensitivity analysis using only first event of long-term sickness absence or the analysis with exclusion of individuals >58 years of age. Also, no noteworthy change in estimates was found in analyses on Whitehall II including weekly work hours. In the sensitivity analyses looking separately at job demands and job control, double exposure to both high demands or low control and informal caregiving were predictors of sickness absence, with fairly similar estimates as with the main analyses using the combined job strain model. Furthermore, for men, joint exposure from low control and informal caregiving was less than expected given the size of their individual effects ( $P=0.02$ ,  $SI=0.05$ ). Detailed results are not shown but can be obtained from the first author upon request.

## Discussion

We investigated the joint exposure of job strain and informal caregiving as predictors of long-term sickness

**Table 2.** Individuals associations between job strain, informal caregiving and long-term sickness absence (SA). [HR=hazard ratio;  $HR_{adj}$ =adjusted for age, married/cohabiting, children, occupational grade, death of relative and/or divorce, previous long-term sickness absence (SA), and longstanding illness; 95% CI=95% confidence interval.]

	Men							Women						
	Individuals with $\geq 1$ long-term SA spell	HR	95% CI	$I^2$ (%)	$HR_{adj}$	95% CI	$I^2$ (%)	Individuals with $\geq 1$ long-term SA spell	HR	95% CI	$I^2$ (%)	$HR_{adj}$	95% CI	$I^2$ (%)
High job strain														
No	1114	1.00			1.00			2350	1.00			1.00		
Yes	211	1.24	0.99–1.54	57	1.09	0.93–1.28	20	734	1.19	1.04–1.36	44	1.08	1.00–1.17	0
Informal caregiving														
No	1101	1.00	–		1.00			2572	1.00			1.00		
Yes	260	1.05	0.92–1.20	0	1.01	0.88–1.16	0	654	1.15	1.02–1.30	34	1.13	1.04–1.23	0
Weekly hours caregiving														
0	768	1.00			1.00			2283	1.00			1.00		
1–4	53	1.10	0.84–1.43	0	1.09	0.83–1.43	0	207	0.78	0.29–2.08	68	0.88	0.42–1.86	52
>4	25	0.89	0.55–1.45	36	0.80	0.42–1.53	64	205	1.25	1.09–1.44	0	1.19	1.04–1.36	0



**Figure 1.** Job strain and informal caregiving as predictors of recurrent events of long-term sickness absence during two years follow-up. Adjusted for age, married/cohabiting, children, occupational grade, death of relative and/or divorce, previous long-term sickness absence, and longstanding illness. Interaction between job strain and informal caregiving in meta-analysis: Multiplicative: Men (P=0.112), women (P=0.766). Additive: Men (synergy index=-0.43), women (synergy index=0.95).

absence. Informal caregiving responsibilities were common with prevalence's ranging between 10–30% across cohorts. With regards to sickness absence, 12% of men and 21% of women experienced  $\geq 1$  spell of long-term sickness absence during a 2-year period. We found that high job strain and informal caregiving were predictors of long-term sickness absence among women, and women jointly exposed to high job strain and informal caregiving had a 34% increased risk of sickness absence compared to women with no high strain and no informal caregiving. However, contrary to our hypothesis, being doubly exposed to informal caregiving and job strain was not associated with an excess risk of sickness absence compared to the expected risk from their individual effects. We found no associations between informal caregiving and job strain on sickness absence among men.

Informal caregiving as a predictor of long-term sickness absence among women may be a consequence of role overload ascribed to additional responsibilities in daily living (10). Women have been increasingly engaged in full-time employment, but still carry the largest household workload (23, 46), and might therefore be more likely than men to experience any health consequences due to the joint effects of paid work, household chores, and informal caregiving (47, 48). In line with this, previous studies have shown that women are more likely to take the role as primary caregiver (49–51), and more often assist with basic and instrumental activities of daily living

(49, 52). Thus, the additional role as informal caregiver may be more burdensome for women compared to men. A previous study using the GAZEL cohort (22), found no gender difference for individuals exposed to high work-family demands in risk of sickness absence. However, in that study family demands were defined as number of economically dependent family members, and informal caregiving may encompass a much greater emotional burden than having economically dependents. In line with our findings, a previous study from the GAZEL cohort also found tendencies for gender differences, but here, sickness absence was restricted to psychiatric disorders and family demands were also defined as number of economically dependents (21).

Job strain as a predictor of sickness absence in women is in line with findings from longitudinal studies in nurses (13, 53); and another longitudinal study covering a broader range of workers found that the strongest mediator in the association between gender and sickness absence was job strain (54). Also, a recent longitudinal population-based study on both men and women found support for the association between job strain and long-term sickness absence (12). In the latter study, long-term sickness absence was defined as absence >16 days, which is somewhat similar to our study. However, job strain was categorized in a variable with three groups: high, active or passive, and low strain. Thus, the reference category is different from our study. Other studies

have found high job demands (19) and low control (55) as predictors of sickness absence. One of the reasons why we chose to look at the combined job strain measure instead of demands and control separately was that it has been shown that job strain (even without positive interaction between job demands and control) predicts outcomes such as coronary heart disease better than low job control, when analyses are adjusted for socioeconomic status (56). Furthermore, sensitivity analyses on job demands and control did not change conclusions.

Even though we found a moderately higher risk of sickness absence among women who were exposed to both informal caregiving and job strain, this excess risk was not more than expected from their individual effects. These results are in line with recent findings showing no interactive effect of job strain and informal caregiving in relation to allostatic load (57), which is a biological measure of stress associated with poor health (58, 59). According to the role enhancement theory (7), a possible explanation for no interaction between high job strain and informal caregiving is that work may have acted as a buffer against stress from caregiving and vice versa (8). However, another possibility is a healthy caregiver effect, ie, those who undertake informal caregiving tasks may have more personal resources to begin with and are therefore also less likely to suffer any negative health consequences from informal caregiving. Based on this, we may have underestimated the effect of caregiving on sickness absence. It is also important to consider that individuals may have diverse experience with caregiving responsibilities. A Canadian study for example showed that >70% of caregivers were positive about the caregiving role (60); another study found that, despite difficulties, caregivers in general expressed great satisfaction with providing caregiving (61). However, other studies have shown that informal caregiving may compromise the caregivers' own health and interfere with working life (3–6). Unfortunately, we had no information on whether caregiving was perceived as burdensome in the present study. Hours of caregiving may be used as a proxy measure of workload, but our data on this variable was limited to Whitehall II, and we only found tendencies indicating that more hours of caregiving is associated with higher risk of sickness absence among women. Differentiating between those who experienced high and low caregiver burden in future studies may broaden our understanding of the relationship.

Informal caregiving was more prevalent in GAZEL compared to Whitehall II and FPS. This may seem counterintuitive as the question on informal caregiving in GAZEL only encompasses care for elderly relatives, whereas caregiving in the other cohorts also include care for sick and disabled relatives. However, a possible explanation for this discrepancy in prevalences is that France has a welfare system in which both family

and state have legal obligations regarding caregiving of disabled and elderly family members; whereas in Finland the state has clear obligations but families do not. In the UK, the rules of legal obligations for sick/disabled family members were somewhat unclear in the 1990s (ie, the baseline of this study) (62) compared to today. Still, the wording of the question in Whitehall and FPS might also have lead the participants to understand that only care of people with severe disabilities was included. Despite the differences in the prevalence of caregiving and the questions applied, we found low heterogeneity in the association between caregiving and sickness absence in the three cohorts. This association would most likely have been stronger if the question in the GAZEL study had also encompassed care for disabled/handicapped children and spouses. A limitation of this study is the different time periods used for baseline. Based on this, it cannot be ruled out that there have been changes/developments in work- and family-related policies or culture between the time periods that could have affected the outcome. We found some heterogeneity in the job strain and sickness absence association, showing a tendency for a time trend; with no association in Whitehall (1994–1996), a tendency for an increased risk in GAZEL (2000–2002), and a stronger tendency in FPS (2012–2014). This, it may be a time-trend where sickness absence has become easier and more socially acceptable. However, the discrepancy in cohort estimates may also be due to cultural differences.

Variables such as alcohol intake, smoking status, body mass index, and depression have been included as potential confounders in similar studies (21, 22, 63). We have chosen not to include these variables in our analysis as they may be mediators on the causal pathway between job strain and sickness absence rather than confounding factors. Thus, a meta-analysis has shown that work–family interference and family–work interference have been associated with depression and substance use/abuse as outcomes (64); and depression, which may be partly a consequence of exposure to work stressors, has been shown to be a highly contributory cause of sickness absence (65). A limitation in our analyses is that other potentially important covariates shown in the directed acyclic graph (DAG) ([www.sjweh.fi/index.php?page=data-repository](http://www.sjweh.fi/index.php?page=data-repository)) were not available, such as personality, spouse work status, household income and organizational changes at the workplace, which may be associated with informal caregiving, job strain and sickness absence (66, 67). However, household income is highly associated with being married and occupational grade (for which we adjusted), and the association between spouse work status and long-term sickness absence is likely to go through household income.

Sensitivity analyses on Whitehall II, including weekly work hours as a potential confounder, did not



change estimates. In line with this, there are some inconsistencies in findings from previous studies on the association between long work hours and sickness absence. A systematic review found that long work hours predict sickness absence and ill health (68), while another study found no association (69). A meta-analysis actually found that long working hours was a protective factor against sickness absence (19). The authors suggested the healthy worker effect as a plausible explanation and that some individuals with long working hours have highly demanding jobs and may feel pressure to be working even though they are sick (19).

To our knowledge, this is the first study to prospectively investigate the joint effects of job strain and informal caregiving on long-term sickness absence. Based on this large population of European workers, we have shown that informal caregiving is common among European workers and women with high job strain and/or informal caregiving responsibilities are at higher risk of long-term sickness absence compared to women with no high strain and no informal caregiving responsibilities. Presently, there may be unmeasured confounding from factors such as personality traits and organizational changes at workplaces.

### Acknowledgements

We gratefully acknowledge the work of the teams behind the GAZEL, FPS, and Whitehall II cohorts.

The Danish Work Environment Foundation financially supported the first author (grant no. 12-2013-03).

### References

- World Health Organization. Good health adds life to years Global brief for World Health Day 2012. Paris: WHO;2012. [Internet]. [cited 2015 Mar 8]. Available from: [http://apps.who.int/iris/bitstream/10665/70853/1/WHO\\_DCO\\_WHD\\_2012.2\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/70853/1/WHO_DCO_WHD_2012.2_eng.pdf)
- Pinquart M, Sörensen S. Differences between caregivers and noncaregivers in psychological health and physical health: A meta-analysis. *Psychol Aging*. 2003;18:250–67. <http://dx.doi.org/10.1037/0882-7974.18.2.250>
- Roepke SK, Mausbach BT, Patterson TL, Von Känel R, Ancoli-Israel S, Harmell AL, et al. Effects of Alzheimer caregiving on allostatic load. *J Health Psychol*. 2011 Jan;16:58–69. <http://dx.doi.org/10.1177/1359105310369188>
- von Känel R, Mausbach BT, Patterson TL, Dimsdale JE, Aschbacher K, Mills PJ, et al. Increased Framingham Coronary Heart Disease Risk Score in dementia caregivers relative to non-caregiving controls. *Gerontology*. 2008 Jan;54:131–7. <http://dx.doi.org/10.1159/000113649>
- Schulz R, Beach SR. Caregiving as a Risk Factor for Mortality. *JAMA*. 1999;282:2215–9. <http://dx.doi.org/10.1001/jama.282.23.2215>
- Christakis NA, Allison PD. Mortality after the Hospitalization of a Spouse. *N Eng J Med*. 2006;354:719–30. <http://dx.doi.org/10.1056/NEJMsa050196>
- Siebert SD. Toward a Theory of Role Accumulation. *Am Sociol Rev*. 1974;39:567–78. <http://dx.doi.org/10.2307/2094422>
- Chumbler NR, Pienta AM, Dwyer JW. The Depressive Symptomatology of Parent Care among the Near Elderly: The Influence of Multiple Role Commitments. *Res Aging*. 2004;26:330–51. <http://dx.doi.org/10.1177/0164027503262425>
- Goode WJ. A theory of role strain. *American sociological review*. 1960;25:483–496. <http://dx.doi.org/10.2307/2092933>
- Mastekaasa A. Parenthood, gender and sickness absence. *Soc Sci Med*. 2000;50:1827–42. [http://dx.doi.org/10.1016/S0277-9536\(99\)00420-7](http://dx.doi.org/10.1016/S0277-9536(99)00420-7)
- Kivimäki M, Head J, Ferrie JE, Shipley MJ, Vahtera J, Marmot MG. Sickness absence as a global measure of health: evidence from mortality in the Whitehall II prospective cohort study. *BMJ*. 2003;327:1–6. <http://dx.doi.org/10.1136/bmj.327.7411.364>
- Wang M-J, Mykletun A, Møyner EI, Øverland S, Henderson M, Stansfeld S, et al. Job strain, health and sickness absence: results from the Hordaland Health Study. *PLoS One*. 2014;9:e96025. <http://dx.doi.org/10.1371/journal.pone.0096025>
- Trybou J, Germonpre S, Janssens H, Casini A, Braeckman L, Bacquer D De, et al. Job-related stress and sickness absence among belgian nurses: A prospective study. *J Nurs Scholarsh*. 2014;46:292–301. <http://dx.doi.org/10.1111/jnu.12075>
- Silva-Junior JS da, Fischer FM. Long-Term Sickness Absence Due to Mental Disorders Is Associated with Individual Features and Psychosocial Work Conditions. *PLoS One*. 2014;9:e115885. <http://dx.doi.org/10.1371/journal.pone.0115885>
- Moreau M. Occupational stress and incidence of sick leave in the Belgian workforce: the Belstress study. *J Epidemiol Community Heal*. 2004;58:507–16. <http://dx.doi.org/10.1136/jech.2003.007518>
- Janssens H, Clays E, De Clercq B, Casini A, De Bacquer D, Kittel F, et al. The relation between psychosocial risk factors and cause-specific long-term sickness absence. *Eur J Public Health*. 2014;24:428–33. <http://dx.doi.org/10.1093/eurpub/cku009>
- Väänänen A, Kumpulainen R, Kevin MV, Ala-Mursula L, Kouvonen A, Kivimäki M, et al. Work-family characteristics as determinants of sickness absence: A large-scale cohort study of three occupational grades. *J Occup Health Psychol*. 2008;13:181–96. <http://dx.doi.org/10.1037/1076-8998.13.2.181>
- Jansen NWH, Kant IJ, van Amelsvoort LGPM, Kristensen TS, Swaen GMH, Nijhuis FJN. Work-family conflict as a risk factor

- for sickness absence. *Occup Environ Med.* 2006;63:488–94. <http://dx.doi.org/10.1136/oem.2005.024943>
19. Niedhammer I, Chastang JF, Sultan-Taieb H, Vermeulen G, Parent-Thirion A. Psychosocial work factors and sickness absence in 31 countries in Europe. *Eur J Public Health.* 2012;23:622–9. <http://dx.doi.org/10.1093/eurpub/cks124>
  20. Kossek EE, Baltes BB, Matthews R a. Innovative Ideas on How Work-Family Research Can Have More Impact. *Ind Organ Psychol.* 2011;4:426–32. <http://dx.doi.org/10.1111/j.1754-9434.2011.01367.x>
  21. Melchior M, Berkman LF, Niedhammer I, Zins M, Goldberg M. The mental health effects of multiple work and family demands. A prospective study of psychiatric sickness absence in the French GAZEL study. *Soc Psychiatry Psychiatr Epidemiol.* 2007;42:573–82. <http://dx.doi.org/10.1007/s00127-007-0203-2>
  22. Sabbath EL, Melchior M, Goldberg M, Zins M, Berkman LF. Work and family demands: predictors of all-cause sickness absence in the GAZEL cohort. *Eur J Public Health.* 2012;22:101–6. <http://dx.doi.org/10.1093/eurpub/ckr041>
  23. Väänänen A, Kevin MV, Ala-Mursula L, et al. The Double Burden of and Negative Spillover Between Paid and Domestic Work : Associations with Health Among Men and Women. *Women Heal.* 2008;40:1-18. [http://dx.doi.org/10.1300/J013v40n03\\_01](http://dx.doi.org/10.1300/J013v40n03_01)
  24. Ostlund G, Cedersund E, Hensing G, Alexanderson K. Domestic strain: a hindrance in rehabilitation? *Scand J Caring Sci.* 2004;18:49–56. <http://dx.doi.org/10.1111/j.1471-6712.2004.00254.x>
  25. Schneider U, Trukeschitz B, Mühlmann R, Ponocny I. "Do I stay or do I go?" – Job Change and Labor Market exit intentions of employees providing informal care to older adults. *Health Econ.* 2013;22:1230–49. <http://dx.doi.org/10.1002/hec.2880>
  26. National Institute for Occupational Safety and Health (NIOSH). Total Worker Health [Internet]. Atlanta: NIOSH; 2015. Available from: <http://www.cdc.gov/niosh/programs/totalworkerhealth/>
  27. Bonnaud S, Descatha A, Zins M, Buyck J-F, Ankri J. Is There Any Additional Psychological and/or Physical Job Constraint Associated With Informal Caregiving Status? Findings From the GAZEL Cohort Study. *J Occup Environ Med.* 2011;53:829–30. <http://dx.doi.org/10.1097/JOM.0b013e3182281178>
  28. Low CA, Thurston RC, Matthews KA. Psychosocial factors in the development of heart disease in women: current research and future directions. *Psychosom Med.* 2010;72:842–54. <http://dx.doi.org/10.1097/PSY.0b013e3181f6934f>
  29. Sjögren E, Leanderson P, Kristenson M. Diurnal saliva cortisol levels and relations to psychosocial factors in a population sample of middle-aged Swedish men and women. *Int J Behav Med.* 2006;13:193–200. [http://dx.doi.org/10.1207/s15327558ijbm1303\\_2](http://dx.doi.org/10.1207/s15327558ijbm1303_2)
  30. Goldberg M, Leclerc A, Bonenfant S, Chastang JF, Schmaus A, Kaniewski N, et al. Cohort profile: the GAZEL Cohort Study. *Int J Epidemiol.* 2007;36:32–9. <http://dx.doi.org/10.1093/ije/dyl247>
  31. Kouvonen A, Oksanen T, Vahtera J, Stafford M, Wilkinson R, Schneider J, et al. Low workplace social capital as a predictor of depression: the Finnish Public Sector Study. *Am J Epidemiol.* 2008;167:1143–51. <http://dx.doi.org/10.1093/aje/kwn067>
  32. Marmot M, Brunner E. Cohort Profile: the Whitehall II study. *Int J Epidemiol.* 2005;34:251–6. <http://dx.doi.org/10.1093/ije/dyh372>
  33. Karasek R, Theorell T. Healthy work. New York: Basic Books Inc; 1990.
  34. Fransson E, Nyberg S, Heikkilä K, Alfredsson L, Bacquer D, Batty G, et al. Comparison of alternative versions of the job demand-control scales in 17 European cohort studies: the IPD-Work consortium. *BMC Public Health.* 2012;12:62. <http://dx.doi.org/10.1186/1471-2458-12-62>
  35. Vahtera J, Pentti J, Kivimäki M. Sickness absence as a predictor of mortality among male and female employees. *J Epidemiol Community Health.* 2004;58:321–6. <http://dx.doi.org/10.1136/jech.2003.011817>
  36. Chevalier A, Luce D, Blanc C, Goldberg M. Sickness absence at the French National Electric and Gas Company. *Br J Ind Med.* 1987;44:101–10. <http://dx.doi.org/10.1136/oem.44.2.101>
  37. Jenny Head, Mika Kivimäki, Johannes Siegrist, Jane E. Ferrie, Jussi Vahtera, Martin J. Shipley, et al. Effort–reward imbalance and relational injustice at work predict sickness absence: The Whitehall II study. *J Psychosom Res.* 2007;63:433–40. <http://dx.doi.org/10.1016/j.jpsychores.2007.06.021>
  38. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology.* 1999; 10:37-48. <http://dx.doi.org/10.1097/00001648-199901000-00008>
  39. Dersimonian R, Laird N. Meta-Analysis in Clinical Trials. *Controlled Clinical Trials.* 1986;7:177–88.
  40. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ Br Med J.* 2003;327:557–60. <http://dx.doi.org/10.1136/bmj.327.7414.557>
  41. Amorim LDAF, Cai J. Modelling recurrent events: a tutorial for analysis in epidemiology. *Int J Epidemiol.* 2014;44:324–33. <http://dx.doi.org/10.1093/ije/dyu222>
  42. Vandembroucke JP, Von Elm E, Altman DG, Gøtzsche PC, Mulrow CD, Pocock SJ, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): Explanation and elaboration. *PLoS Med.* 2007;4 :1628–54. <http://dx.doi.org/10.1371/journal.pmed.0040297>
  43. de Mutsert R, Jager KJ, Zoccali C, Dekker FW. The effect of joint exposures: examining the presence of interaction. *Kidney Int.* 2009;75:677–81. <http://dx.doi.org/10.1038/ki.2008.645>
  44. Author. Multiple Imputation for Nonresponse in Surveys: Donald B. Rubin: Books [Internet]. [cited 2015 Mar 8]. Available from: <http://www.amazon.com/Multiple-Imputation-Nonresponse-Surveys-Donald/dp/0471655740>
  45. Burgess S, White IR, Resche-Rigon M, Wood AM. Combining multiple imputation and meta-analysis with individual participant data. *Stat Med.* 2013;32:4499–514. <http://dx.doi.org/10.1002/sim.5844>

46. Lundberg U, Frankenhaeuser M. Stress and workload of men and women in high-ranking positions. *J Occup Health Psychol.* 1999;4:142–51. <http://dx.doi.org/10.1037/1076-8998.4.2.142>
47. Eek F, Axmon A. Gender inequality at home is associated with poorer health for women. *Scand J Public Health.* 2014; 12:1–7.
48. Berntsson L, Lundberg U, Krantz G. Gender differences in work-home interplay and symptom perception among Swedish white-collar employees. *J Epidemiol Com Health.* 2006;60:1070–6. <http://dx.doi.org/10.1136/jech.2005.042192>
49. Allen SM. Gender differences in spousal caregiving and unmet need for care. *J Gerontol.* 1994;49:187–95. <http://dx.doi.org/10.1093/geronj/49.4.S187>
50. Ingersoll-Dayton B, Starrels ME, Dowler D. Caregiving for parents and parents-in-law: is gender important? *Gerontologist.* 1996;36:483–91. <http://dx.doi.org/10.1093/geront/36.4.483>
51. Neal MB, Ingersoll-Dayton B, Starrels ME. Gender and relationship differences in caregiving patterns and consequences among employed caregivers. *Gerontologist.* 1997;37:804–16. <http://dx.doi.org/10.1093/geront/37.6.804>
52. Kramer BJ, Kipnis S. Eldercare and work-role conflict: toward an understanding of gender differences in caregiver burden. *Gerontologist.* 1995;35:340–8. <http://dx.doi.org/10.1093/geront/35.3.340>
53. Bourbonnais R, Mondor M. Job strain and sickness absence among nurses in the province of Québec. *Am J Ind Med.* 2001;39:194–202. [http://dx.doi.org/10.1002/1097-0274\(200102\)39:2<194::AID-AJIM1006>3.0.CO;2-K](http://dx.doi.org/10.1002/1097-0274(200102)39:2<194::AID-AJIM1006>3.0.CO;2-K)
54. Casini A, Godin I, Clays E, Kittel F. Gender difference in sickness absence from work: A multiple mediation analysis of psychosocial factors. *Eur J Public Health.* 2013;23:635–42. <http://dx.doi.org/10.1093/eurpub/cks183>
55. North FM, Syme SL, Feeney a, Shipley M, Marmot M. Psychosocial work environment and sickness absence among British civil servants: the Whitehall II study. *Am J Public Health.* 1996;86:332–40. <http://dx.doi.org/10.2105/AJPH.86.3.332>
56. Kivimäki M, Nyberg ST, Kawachi I. Authors' reply: Calculation of population attributable risk should be based on robust estimates. *Scand J Work Environ Health.* 2015;41:506–7. <http://dx.doi.org/10.5271/sjweh.3518>
57. Dich N, Lange T, Head J, Rod NH. Work Stress, Caregiving, and Allostatic Load. *Psychosom Med.* 2015;77:539–47. <http://dx.doi.org/10.1097/PSY.000000000000191>
58. Seeman TE, McEwen BS, Rowe JW, Singer BH. Allostatic load as a marker of cumulative biological risk: MacArthur studies of successful aging. *PNAS.* 2001;98:4770–5. <http://dx.doi.org/10.1073/pnas.081072698>
59. Read S, Grundy E. Allostatic Load and Health in the Older Population of England. *Psychosom Med.* 2014;76:490–6. <http://dx.doi.org/10.1097/PSY.0000000000000083>
60. Cohen CA, Colantonio A, Vernich L. Positive aspects of caregiving: Rounding out the caregiver experience. *Int J Geriatr Psychiatry.* 2002;17:184–8. <http://dx.doi.org/10.1002/gps.561>
61. López J, López-Arrieta J, Crespo M. Factors associated with the positive impact of caring for elderly and dependent relatives. *Arch Gerontol Geriatr.* 2005;41:81–94. <http://dx.doi.org/10.1016/j.archger.2004.12.001>
62. Sundstrom G, Malmberg B, Castiello MS, del Barrio E, Castejon P, Tortosa M a, et al. Family Care for Elders in Europe: Policies and Practices. *Caregiving Context Cult Fam Soc Implic.* 2007;235:1–41.
63. Väänänen A, Kumpulainen R, Kevin M V, Ala-Mursula L, Kouvonen A, Kivimäki M, et al. Work-family characteristics as determinants of sickness absence: a large-scale cohort study of three occupational grades. *J Occup Health Psychol.* 2008;13:181–96. <http://dx.doi.org/10.1037/1076-8998.13.2.181>
64. Amstad FT, Meier LL, Fasel U, Elfering A, Semmer NK. A meta-analysis of work-family conflict and various outcomes with a special emphasis on cross-domain versus matching-domain relations. *J Occup Health Psychol.* 2011;16:151–69. <http://dx.doi.org/10.1037/a0022170>
65. Thomas CM, Morris S. Cost of depression among adults in England in 2000 service Cost of depression among adults in England in 2000. *Br J Psychiatry.* 2003;183:514–9. <http://dx.doi.org/10.1192/bjp.183.6.514>
66. Vahtera J, Kivimäki M, Pentti J, Linna A, Virtanen M, Virtanen P, et al. Organisational downsizing, sickness absence, and mortality: 10-town prospective cohort study. *BMJ.* 2004;328:1–5. <http://dx.doi.org/10.1136/bmj.37972.496262.0D>
67. Kivimäki M, Vahtera J, Koskenvuo M, Uutela A, Pentti J. Response of hostile individuals to stressful change in their working lives: test of a psychosocial vulnerability model. 1998;28:903–13.
68. Michie S, Williams S. Reducing work related psychological ill health and sickness absence: a systematic literature review. *Occup Environ Med.* 2003;60:3–9. <http://dx.doi.org/10.1136/oem.60.1.3>
69. Voss M, Floderus B, Diderichsen F. Physical, psychosocial, and organisational factors relative to sickness absence: a study based on Sweden Post. *Occup Environ Med.* 2001;58:178–84. <http://dx.doi.org/10.1136/oem.58.3.178>

Received for publication: 27 November 2015