

The heart of epilepsy: current views and future concepts

S. Shmuelly^{1,2}, M. van der Lende^{1,3}, R.J. Lamberts¹, J.W. Sander^{1,2}, R.D. Thijs^{1,2,3}

¹Stichting Epilepsie Instellingen Nederland - SEIN, Heemstede, the Netherlands

²NIHR University College London Hospitals Biomedical Research Centre, UCL Institute of Neurology, Queen Square, London WC1N 3BG and Epilepsy Society, Bucks SL9 0RJ, UK

³Department of Neurology, LUMC Leiden University Medical Centre, Leiden, the Netherlands

Corresponding author

Roland D. Thijs MD PhD

Stichting Epilepsie Instellingen Nederland - SEIN

PO Box 540

2130 AM Hoofddorp

The Netherlands

E-mail: rthijs@sein.nl

Telephone: +31(0)23 558 8948

Fax: +31(0)23 558 8159

OVERVIEW

ABSTRACT	2
INTRODUCTION	3
EPILEPSY AND CARDIAC ARRHYTHMIAS.....	4
EPILEPSY AND STRUCTURAL CARDIAC CONDITIONS	10
FUTURE CONCEPTS	12
ACKNOWLEDGEMENTS.....	14
DISCLOSURES.....	14
FUNDING	14
REFERENCES	16

ABSTRACT

Cardiovascular (CV) comorbidities are common in people with epilepsy. Several mechanisms explain why these conditions tend to co-exist including causal associations, shared risk factors and those resulting from epilepsy or its treatment

Various arrhythmias occurring during and after seizures have been described. Ictal asystole is the most common cause. The converse phenomenon, arrhythmias causing seizures, appears extremely rare and has only been reported in children following cardioinhibitory syncope. Arrhythmias in epilepsy may not only result from seizure activity but also from a shared genetic susceptibility. Various cardiac and epilepsy genes could be implicated but firm evidence is still lacking. Several antiepileptic drugs (AEDs) triggering conduction abnormalities can also explain the co-existence of arrhythmias in epilepsy.

Epidemiological studies have consistently shown that people with epilepsy have a higher prevalence of structural cardiac disease and a poorer CV risk profile than those without epilepsy. Shared CV risk factors, genetics and etiological factors can account for a significant part of the relationship between epilepsy and structural cardiac disease. Seizure activity may cause transient myocardial ischaemia and the Takotsubo syndrome. Additionally, certain AEDs may themselves negatively affect CV risk profile in epilepsy.

Here we discuss the fascinating borderland of epilepsy and cardiovascular conditions. The review focuses on epidemiology, clinical presentations and possible mechanisms for shared pathophysiology. It concludes with a discussion of future developments and a call for validated screening instruments and guidelines aiding the early identification and treatment of CV comorbidity in epilepsy.

Keywords: epilepsy; seizures; cardiac; cardiovascular; comorbidity; arrhythmia

INTRODUCTION

Well over 100 years ago, the occurrence of asystole during the course of an epileptic seizure was described: “He uttered a cry and was seen to be rubbing his hands together. His pulse was immediately examined for but was not palpable” [1]. Since then numerous associations between epilepsy and CV conditions have been identified, including this classical example of ictal asystole.

Co-existing conditions form an important part of the overall burden of epilepsy [2-5]. Several mechanisms of association between epilepsy and comorbid conditions have been described: associations can be explained by cause or effect, a shared risk factor may cause both conditions, or the mechanism of the association is unknown or spurious (i.e. coincidental) (Figure 1) [3, 5].

This review serves to discuss the fascinating borderland between epileptology and cardiology and focuses on the major developments over the last 25 years and on future developments. We use the comorbidity framework (Figure 1) [3] to review all cardiac conditions known, and alleged, to be linked to epilepsy. Associations with cardiac arrhythmias are discussed first, followed by an overview of all structural cardiac conditions related to epilepsy.




Figure 1. Mechanisms of association between epilepsy and comorbid conditions. Figure originates from Gaitatzis et al., 2012 [3].

EPILEPSY AND CARDIAC ARRHYTHMIAS

Various arrhythmias have been described, occurring during (ictal) or after (postictal) seizures. Sinus tachycardia is the most common ictal pattern, seen in up to 80% of all seizures [6] and in 82% of people with epilepsy [7], but usually without symptoms. The most frequent clinically relevant arrhythmia is ictal asystole, occurring in 0.318% (95% CI 0.316% to 0.320%) of people with refractory focal epilepsy admitted for video-EEG [8]. Ictal asystole, bradycardia and AV block predominantly occur in people with temporal lobe epilepsy (Table 1) [8]. Clinically, ictal asystole is characterised by sudden loss of tone during a dyscognitive seizure [9]. The circulatory pattern resembles vasovagal syncope with a transient, progressive and self-limiting slowing of the heart rate and decrease of blood pressure [9-11]. For many years, ictal asystole was thought to be a possible mechanism underlying sudden unexpected death in epilepsy (SUDEP). This appears to be unlikely: all but one reported case so far of ictal asystole were self-limiting [8]. In this one case successful resuscitation was started after 44 seconds of asystole and the event was classified as near-SUDEP [12]. The longest ictal asystole reported so far, however, lasted 96 seconds and appeared self-limiting [13]. Whether an event is classified as near-SUDEP or not will depend on interventions of medical personnel: prompt resuscitation in response to ictal asystole will likely lead to more classified as near-SUDEP cases. While there are no reports of fatal ictal asystole, it remains debatable whether ictal asystole can cause SUDEP.

The precise mechanism of ictal asystole is unknown. It may result from epileptic activity directly stimulating the central autonomic networks [6, 14]. For example, focal stimulation of parts of the limbic system (i.e. amygdala, cingulate gyrus) may provoke asystole [6, 15-17]. Alternatively, seizure-induced fear and catecholamine release [18] may evoke a vasovagal response causing cardioinhibition and vasodilation [19].

Ictal asystole is assumed to be self-limiting, but may cause falls and injuries due to seizure-induced syncope [20]. Proper trials are lacking but retrospective studies suggest that improving seizure control may prevent ictal asystole [21-23]. It also seems advisable to withdraw negative inotropic drugs and to consider the implantation of a loop recorder to

monitor possible future events in individuals in whom ictal asystole has been noted. If the asystolic episodes persist, cardiac pacemaker implantation should be considered to reduce the risk of trauma [20, 21, 23, 24].

In contrast to ictal asystole, postictal asystole is less common, associated with convulsive rather than focal (temporal lobe) seizures and has a higher fatality rate: 7 of 13 reported postictal asystole cases died from SUDEP [8]. All fatal cases had a convulsive seizure with immediate postictal generalised EEG suppression and a stuttering course of transient apnoea and asystole resulting in a terminal apnoea followed by a terminal asystole [25].

The mechanism underlying this sequence of postictal EEG suppression, apnoea, and terminal asystole has not yet been elucidated. Excessive inhibition causing brainstem depression might play a role [26]. Recent work in two animal models (mice carrying mutations in the *KCNA1* gene or the *SCN1A* gene) demonstrated that seizures initiated by direct cortical stimulation may evoke a spreading depression causing brain stem inhibition and cardiorespiratory collapse [27].

Another rare (post)ictal arrhythmia is ventricular tachycardia/ventricular fibrillation (VT/VF). So far three cases of postictal VT/VF leading to (near) SUDEP have been reported [8]. All VT/VF occurred directly following a convulsive seizure. No cardiac lesions were found in the case reports. There may be a publication bias, however, as cases with seizure-triggered VT/VF and cardiac lesions may not qualify as SUDEP and thus may be less likely to be reported. The mechanism of seizure-induced VT/VF is unclear. Convulsive seizures may exert proarrhythmogenic effects by triggering the sympathetic nervous system, as reflected by the peak in catecholamines and electrodermal activity [18, 28]. At the same time, convulsive seizures may increase cardiac oxygen deprivation by inducing sinus tachycardia [7] and respiratory impairment causing hypoxemia [29]. It has also been found that ECG-markers of sudden cardiac death such as QTc-lengthening and/or shortening [30, 31], and T-wave alternans are more prevalent [23] during and after convulsive seizures. The various factors might interact as seizure-related cardiac repolarization abnormalities appeared more frequent in seizures with ictal hypoxemia compared to those without [32].

Though seizure-induced VT/VF appears to be rare, a prospective community-based study of out-of-hospital cardiac arrests due to ECG-documented VT/VF showed that VT/VF risk in

those with epilepsy was three times as high as the general population [33]. A further analysis of those cases with epilepsy and VT/VF showed that most were not seizure-related, but rather occurred in the context of either pre-existing heart disease or as the immediate result of an acute myocardial infarction [34]. Pre-existing heart disease was a stronger predictor for VT/VF in people with epilepsy than markers of epilepsy severity. In a minority of cases, however, VT/VF was unexplained and a diagnosis of (near) SUDEP was established. It thus appears that sudden cardiac arrest and SUDEP are partially overlapping disease entities.

The increased risk of non-seizure related VF/VT episodes in people epilepsy may be explained by high cardiovascular comorbidity [3, 35]. People with epilepsy may have a propensity for sudden cardiac death as reduced heart rate variability, a measure of cardiac sympathovagal balance that is also a risk marker of sudden cardiac death, progressively worsens over time in people with refractory, but not in those with well-controlled, epilepsy [36]. In addition, other markers of sudden cardiac death such as early repolarization pattern and QTc-prolongation are more frequently found in the interictal ECGs of people with epilepsy than in those without epilepsy [37].

Seizure related arrhythmia	Reported in n cases	Associated seizure types	Reported in n cases	EEG seizure onset	Reported in n cases	SUDEP association
Ictal asystole	103	99% FDS 1% FAS	97	46% LT 31% RT 13% BT 10% Other	80	Unlikely
Postictal asystole	13	85% fbCS 15% FDS	13	20% LT 60% RT 20% Other	10	Likely, accompanied or preceded by PGES/apnea [Ryvlin et al., 2013]
Ictal bradycardia	25	100% FDS	8	52% LT 38% RT 10% Other	21	Unlikely
Ictal AV Block	11	90% FDS 10% FAS	10	73% LT 18% BT 10% Other	11	Unlikely
Postictal AV Block	2	100% fbCS	2	100% RT	1	Unlikely
Atrial Fibrillation	13	46% GTCS 46% fbCS 8% FDS	13	33% LT 33% Gen 33% Non loc	3	Unlikely
(Post)ictal ventricular fibrillation	3	100% GTCS	3	Insufficient data	0	Probable, but in a minority of cases

Table 1. Reported (post)ictal cardiac arrhythmias. FDS focal dyscognitive seizure; FAS focal autonomic seizure; fbCS focal seizure evolving to bilateral convulsive seizure; GTCS generalised tonic clonic seizure; LT left temporal; RT right temporal; BT bitemporal; Gen generalised; Non loc non-localising; PGES postictal generalized EEG suppression; *in people with refractory focal epilepsy admitted for a vEEG recording. For more details see van der Lende et al., 2016 [8].

Another mechanism explaining the association between arrhythmias and epilepsy is a shared genetic risk factor. A rapidly increasing number of genes potentially linking epilepsy to cardiac arrhythmias has been identified. Here we discuss some relevant examples; starting with the genes predominantly known for their cardiac functions and then the ‘epilepsy genes’.

Several genetic ion channel mutations are thought to be expressed in the brain as well as in the heart, and might thus cause seizures and cardiac arrhythmias. The first reported genetic link between epilepsy and cardiac arrhythmias was the discovery of cardiac sodium channel gene *SCN5A* in the brain [38]. Subsequently, more pathogenic variants in the long QT (LQT) gene family (i.e. *KCNQ1*, *KCNH2* and *SCN5A*) were associated with a “seizure phenotype” (e.g. self-reported diagnosis of epilepsy and AED use) [39-44]. Mice models indicated that other, non-LQT, cardiac channelopathy genes including *RYR2* (associated with catecholaminergic polymorphic ventricular tachycardia) [45], and *HCN1-4* [46, 47] potentially predispose to epilepsy.

Several postmortem studies suggest that the LQT and non-LQT cardiac gene mutations are more common in SUDEP victims [48-50]. As ictal recordings are lacking, it remains questionable whether the fatal events were caused by arrhythmias. The same applies to the identification of ‘epilepsy genes’ in the post-mortem cohorts [48, 49]. These mutations could be markers explaining epilepsy severity or a genetically mediated liability to fatal seizures. In certain epilepsy syndromes, SUDEP risk seems particularly high.

The most recognized example is the Dravet syndrome (DS), a severe epilepsy syndrome with high premature mortality, caused by *SCN1A* mutation [51]. In mutant *SCN1A* knock-out mice, postictal bradycardia and seizure-triggered ventricular fibrillation were recorded before a death resembling SUDEP [52, 53]. In DS subjects, markers associated with the risk of sudden cardiac death (decreased HRV and increased QT-dispersion) have been found [54, 55]. Ictal

proof is, however, lacking and is the subject of an ongoing study (ClinicalTrials.gov Identifier: NCT02415686).



Other less well studied examples of 'epilepsy genes' possibly mediating SUDEP risk include *KCNA1* and *SCN8A*. *KCNA1* is expressed in the vagal nerve as well as in the brain, and is associated with seizures, cardiac arrhythmias, vagal hyperexcitability and premature death in *KCNA1* null mice [56]. Mutations in this gene were found in a SUDEP case with epileptic encephalopathy and suspected cardiac arrhythmias [57].

A novel pathogenic *SCN8A* mutation was identified through whole-genome sequencing in a family affected by epileptic encephalopathy and SUDEP [58]. Before then, *SCN8A* mutations had only been linked to epilepsy in mice [59]. The *SCN8A* gene encodes a sodium channel that is expressed in heart and brain of mice and rats, and plays a role in excitation-contraction coupling, action potential propagation and pacemaking [60, 61].

We previously discussed how seizures may cause arrhythmias. Whether the converse phenomenon exists is a subject of controversy. The major complication is the fact that syncopal events are easily mistaken for epilepsy. Misdiagnosis of epilepsy is common, with reported rates of false positives of up to 71% , and syncope is the most misdiagnosed condition [Xu et al., 2016]. Rates of misdiagnosis of epilepsy vary from 23% in a community-based study of people with a primary diagnosis of epilepsy [62], to 42% in a group of people with seemingly refractory epilepsy [63]. Which is understandable as various symptoms and signs are seen in both conditions [11, 63-65]. Notably, jerking movements or signs indicative of cerebral standstill (complete flattening of the EEG) such as roving eye movements or stertorous breathing [11] are often interpreted as signs specific to epilepsy. Consequently, most seemingly overlapping presentations turn out **not** to be an isolated phenomenon of either syncope or epilepsy, if a proper investigation is performed (e.g. ictal recording of video, heart rate, blood pressure and EEG). Two large scale surveys of up to 2000 tilt-table tests failed to identify any adult case with syncopal-induced seizures [66, 67]. In children, however, a few cases have been reported with a cardioinhibitory reflex syncope followed by video-EEG documented clonic seizures [68-70]. The reason why this phenomenon only appears to affect children is unknown. It may be that the seizure threshold is lower in children (paralleling febrile seizures that also peak in childhood). Alternatively, the depth of

cerebral anoxia may be more profound in children as reflected by prolonged asystolic spells. For clinical management it is important to stress that syncope-induced seizures are extremely rare and probably only affect children. The diagnosis requires an ictal video-EEG recording.

Several AEDs, particularly those with sodium blocking properties are known to trigger conduction abnormalities or arrhythmias [71]. Atrioventricular (AV) conduction is the most frequent reported complication. ST changes, Brugada-like patterns, atrial fibrillation and QTc prolongation have also been reported but the association with AED treatment is less well established [72-86]. Most clinically relevant arrhythmias were related to AED overdose. Carbamazepine is, however, known to cause AV conduction blocks at low levels; this is almost exclusively reported in elderly women [77, 79, 87]. Rapid administration of phenytoin may also cause sinus arrest and hypotension; elderly people and those with pre-existing heart disease seem most vulnerable to these adverse effects. IV administration should, therefore, be undertaken slowly, with continuous cardiac monitoring [76, 83, 86, 88]. The above-mentioned AED effects do not seem to play a role in ictal arrhythmias. Nevertheless, it is important to take these effects into consideration in the selection of an AED and to monitor adverse effects closely especially in elderly people and those with cardiovascular comorbidities (Table 2).

Putative mechanisms of associations between epilepsy and cardiac arrhythmias		
Mechanisms of association	Conditions	References
<p>Causal</p>  <pre> graph LR CA[Condition A] --> E[Epilepsy] CB[Condition B] --> CA CA --> E </pre>	<p>Arrhythmias → Seizures</p>	<p>Battaglia et al., 1989; Horrocks et al., 2005; Stephenson et al., 2004</p>
<p>Shared risk factor</p>  <pre> graph LR Env[Environmental] --- J(()) Gen[Genetic] --- J Bio[Biological/Structural] --- J J --- Con[Condition] J --- E[Epilepsy] </pre>	<p>Genetics → Epilepsy and arrhythmias</p> <ul style="list-style-type: none"> - Important 'heart genes': <i>KCNQ1, KCNH2, SCN5A, RYR2</i> - Important 'epilepsy genes': <i>SCN1A, KCNA1, SCN8A</i> 	<p>Anderson et al., 2014; Auerbach et al., 2013; Auerbach et al., 2016; Aurlen et al., 2009; Heron et al., 2010; Bagnall et al., 2016; Benarroch, 2013; Delogu et al., 2011; Du et al., 2007; Glasscock et al., 2010; Goldman et al., 2016; Hartmann et al., 1999; Johnson et al., 2010; Kalume et al., 2013; Keller et al., 2009; Lehnart et al., 2008; Leu et al., 2015; Ludwig et al., 2003; Noujam et al., 2012; Papale et al., 2009; Partemi et al., 2013; Postma et al., 2005; Tu et al., 2011; Veeramah et al., 2012</p>


Resultant  <pre> graph LR Epilepsy --> AED Epilepsy --> Seizures AED --> Condition Seizures --> Condition </pre>	AED → Arrhythmias <ul style="list-style-type: none"> - Particularly carbamazepine, phenytoin and lacosamide 	Al Aloul et al., 2007; El-Menyar 2011; DeGiorgio et al., 2010; DeToledo et al., 2001; Feldman and Gidal, 2013; Guldiken et al., 2016; Ide and Kamijo, 2007; Kasarskis et al., 1992; Kaufman et al., 2012; Krause et al., 2011; Nizam et al., 2011; Randazzo et al., 1995; Takayanagi et al., 1998; Strimel et al., 2010; Swe et al., 2016; Zoneraich et al., 1976
	Seizures → Arrhythmias <ul style="list-style-type: none"> - Ictal: tachycardia, asystole, bradycardia and AV block. - Postictal: asystole, AV block, atrial flutter or fibrillation and ventricular fibrillation. 	Bardai et al., 2013; Chaila et al., 2010; Eggleston et al., 2014; Lanz et al., 2011; van der Lende et al., 2016; Sevcencu and Struijk, 2010

Table 2. Putative mechanisms of associations between epilepsy and cardiac arrhythmias. HRV heart rate variability; VT ventricular tachycardia; VF ventricular fibrillation; AED antiepileptic drugs.

EPILEPSY AND STRUCTURAL CARDIAC CONDITIONS

Epidemiological studies have consistently shown that people with epilepsy have a higher prevalence of structural cardiac disease than those without epilepsy [4, 5, 89-92].

Cardiovascular disease seems to be a significant contributor to the increased mortality in people with epilepsy, compared with the general population [93-95].

Shared cardiovascular risk factors can account for the relationship between epilepsy and heart disease, in addition to shared genetics and etiological factors. People with a history of epilepsy are more likely to be obese, physically inactive, and current smokers [90] and have a worse cardiovascular risk profile (i.e. hypertension, hypercholesterolemia, diabetes mellitus, stroke/TIA) than the general population [35, 90, 96, 97]. Unsurprisingly, people with epilepsy have higher rates of fatal and nonfatal cardio- and cerebrovascular disease than controls (mortality ratios up to 5.3 and morbidity ratio up to 7) [35, 98, 99]. The presence of cardiovascular disease (e.g. congestive heart failure and cardiac arrhythmias) was also associated with higher mortality risk in people with epilepsy [100].

Epilepsy treatment can also contribute to a poorer cardiovascular risk profile in epilepsy. Use of the enzyme-inducing AEDs phenytoin or carbamazepine may lead to elevated serological vascular risk markers (e.g. total cholesterol, LDL, homocysteine), and, thus, result in accelerated atherosclerosis [101-104]. Certain AEDs (e.g. valproic acid, carbamazepine) are

also known to cause weight gain and increase the risk of developing non-alcoholic fatty liver disease and metabolic syndrome, leading to further deterioration of the cardiovascular risk profile [102].

The co-occurrence of epilepsy and (congenital) heart disease, often accompanied by intellectual disability, may result from a multiple malformation syndrome: genetic defects may affect the development of both heart and brain, or abnormal cardiovascular function may lead to poor (intrauterine) brain growth [105].

CV disease can sometimes (indirectly) cause epilepsy through a predisposition to stroke [106, 107]. Stroke is a common risk factor for epilepsy and accounts for about a third of newly diagnosed seizures in people over the age of 60 years [107-110]. In particular, those with ischemic events with cortical involvement, cerebral hemorrhage (i.e. primary hemorrhage or hemorrhagic transformation of ischemic stroke) and early post-stroke seizures, have an increased risk of post-stroke epilepsy [107].

Seizure activity may not only induce arrhythmias but may also lead to structural cardiac changes [71, 111-113]. Epileptic seizures have been reported to provoke cardiac ischaemia via both acute and chronic effects on the heart (e.g. impaired heart rate variability, cardiac fibrosis, ST-segment depression and increased heart rate) [71, 114]. Transient myocardial ischaemia as indicated by ST-segment depression, was reported in a small-scale study in 40% of all 15 seizures [114]. Another study, however, failed to demonstrate troponin increases, suggesting that the reported ST changes do not usually cause myocardial damage [115].

Seizures are the second most frequent CNS condition known to induce the cardiomyopathy known as Takotsubo syndrome (TTS) [116]. TTS mimics myocardial infarction clinically, electrocardiographically and chemically [117]. It is characterized by acute onset of chest pain and dyspnoea, sometimes concomitant with palpitations, tiredness, oedema, fever, syncope, anxiety, nausea or vomiting [116]. The seizure type that most frequently causes TTS is the generalized tonic-clonic seizure [118, 119]. Seizures most likely trigger TTS by the stress-induced release of catecholamines [120]. This abundant catecholamine release may be a contributing factor in fatal status epilepticus [121]. A relationship between TTS and SUDEP, however, does not appear likely [116].




Putative mechanisms of associations between epilepsy and structural cardiac disease		
Mechanisms of association	Conditions	References
Causal 	Cardiac conditions, e.g. embolism and congenital cardiac abnormalities → Stroke	Attar et al., 2016; Ferlazzo et al., 2016; Gaitatzis et al. 2004
Shared risk factor 	Genetic → Malformation of cortical and cardiac development Shared cardiovascular risk factors → Stroke	Miller and Vogel, 1999 Centers for Disease Control and Prevention (CDC), 2013; Elliott et al., 2008; Gaitatzis et al., 2004; Kobau et al., 2008
Resultant 	AED → Arteriosclerosis AED → Weight gain, non-alcoholic fatty liver disease and metabolic syndrome Seizures → Transient myocardial ischaemia Seizures → Seizure-triggered Takotsubo syndrome (TTS)	Brodie et al., 2013; Katsiki et al., 2014; LoPinto and Mintzer, 2010; Mintzer et al., 2009 Katsiki et al., 2014 P-Codrea Tigaran et al., 2005; Schuele, 2009 Finsterer and Wahbi, 2014; Finsterer and Bersano, 2015; Lemke et al., 2008

Table 3. Putative mechanisms of associations between epilepsy and structural cardiac disease. AED antiepileptic drugs; TTS, Takotsubo syndrome.

FUTURE CONCEPTS

Significant progress has been made since the publication of Russel's case history: the complex interrelationship between epilepsy and cardiac conditions has been explored widely and this review aimed to capture all major discoveries made in this field (Table 2 and 3). Many discoveries of coexisting conditions were made by serendipity, and underlying mechanisms are yet to be uncovered. Treatment regimens are consequently often speculative and lack a personalized approach involving all comorbid conditions. As comorbidity gains recognition we now need to become better at noticing these symptom patterns. Today a substantial gap still remains between the specialties, but as we are now becoming aware of all overlapping syndromes epileptologists will increasingly need to improve their cardiac skills. Pattern recognition can be fostered by incorporating validated screening instruments and guidelines, aiding the early identification and treatment of cardiovascular comorbidity in epilepsy. Concomitantly, a fundamental change in the way clinicians think of epilepsy is crucial.

Epilepsy will soon be viewed as a collection of individual disorders that share a phenotype of an abnormal tendency for unprovoked epileptic seizures. The number of rare epilepsy syndromes with cardiac phenotypes will increase substantially. Epilepsy will be seen as a symptom-complex, and all comorbidities, even the most inconspicuous, should be considered as part of the stratification and phenotyping in people with epilepsy.

Cardiovascular comorbidities will provide insight into common mechanisms for epilepsy and give a window into common genetic predispositions. They may also provide important diagnostic clues. Channelopathies, for example, are increasingly identified in people with epilepsy. Genetic factors may explain both the epilepsy and the comorbid disorder(s), even in people with sporadic epilepsies [122]. Genome wide scanning will be widely available and drive the paradigm shift in epilepsy. Certain genes might be identified as contributing to SUDEP [48, 49], potentially allowing the development of individualised risk prevention strategies. Another major contributor to early identification of overlapping syndromes will be the development of new non-invasive tools to record heart function at home. The miniaturisation of sensors will favour long-term home-based recordings thus aiding the early identification of cardiac arrhythmias.

Advances in seizure detection will likely take off. ECG alone will help to detect a wide variety of seizures but lacks specificity. Combining ECG with other modalities including an accelerometry and electrodermal activity will likely improve accuracy and facilitate the widespread use of seizure detection devices in those with refractory epilepsy [123, 124].

Another unmet need relates to the treatment of epilepsy: many AEDs have proarrhythmogenic and arteriosclerogenic effects. Though non-pharmacological options exist, drug therapy is still the mainstay of epilepsy treatment and other options are usually only explored after AEDs have failed to successfully control seizures [125]. Many new AEDs have been launched in the last two decades, but have failed to improve the burden of side effects or substantially change prognosis for seizure control [126, 127]. With improved understanding of epileptogenesis, epigenetic determinants and pharmacogenomics comes the hope for better, disease-modifying or even curative pharmacological and non-pharmacological treatment strategies. Until then, comorbidity should be considered when prescribing AEDs.

The incorporation of neurocardiology into the paroxysmal spectrum will require a critical review of the epilepsy services. We need to validate new instruments to screen for cardiovascular conditions. Modern non-invasive long-term ECG devices may help screen for cardiac conditions and a cardiologist should review any relevant abnormalities. In cases where there is a relevant family history or abnormal ECG findings, a specialist cardiac assessment should be done. Identification and adequate treatment of cardiovascular disorders in epilepsy should therefore be an important part of epilepsy management.

Particular attention should be given to modifiable risk factors such as smoking, obesity, sedentary lifestyle, high cholesterol and hypertension. Physicians should screen for these risk factors in people with epilepsy, provide general health information and if necessary adjust AED treatment. Further studies are needed to improve risk profiling, thus allowing for screening in high risk individuals (with, for example, implantable loop recorders) and targeted interventions (e.g. defibrillators).

ACKNOWLEDGEMENTS

This work was partly done at the UCLH/UCL Comprehensive Bio-Medical Research Centre which received a proportion of funding from the Department of Health's NIHR Biomedical Research Centres funding scheme. We are grateful to Dr. GS Bell, for critically reviewing the manuscript.

DISCLOSURES

SS, MvdL and RJL report no disclosures. JWS reports personal fees from Lundbeck and Teva, grants and personal fees from UCB, Eisai, grants from GSK, WHO and Dutch National Epilepsy Fund, outside the submitted work; his current position is endowed by the Epilepsy Society, he is a member of the Editorial Board of the Lancet Neurology, and receives research support from the Marvin Weil Epilepsy Research Fund. RDT receives research support from the Dutch National Epilepsy Fund, NUTS Ohra Fund, Medtronic, and AC Thomson Foundation, and has received fees for lectures from Medtronic, UCB and GSK.

FUNDING

This work was supported by the Dutch National Epilepsy Fund [project number 15-10]; and Christelijke Vereniging voor de Verpleging van Lijders aan Epilepsie (The Netherlands).

REFERENCES

1. Russell AE. Cessation of the pulse during the onset of epileptic fits, with remarks on the mechanism of fits. *Lancet* 1906;168:152–4.
2. Forsgren L. Prevalence of epilepsy in adults in northern Sweden. *Epilepsia*. 1992;33(3):450-8. Epub 1992/05/01.
3. Gaitatzis A, Sisodiya SM, Sander JW. The somatic comorbidity of epilepsy: a weighty but often unrecognized burden. *Epilepsia*. 2012;53(8):1282-93. Epub 2012/06/14.
4. Kadima N KR, Zack M, Helmers S. Comorbidity in adults with epilepsy--United States, 2010. *MMWR Morb Mortal Wkly Rep*. 2013;62(43):849-53. Epub 2013/11/01.
5. Keezer MR, Sisodiya SM, Sander JW. Comorbidities of epilepsy: current concepts and future perspectives. *Lancet Neurol*. 2016;15(1):106-15. Epub 2015/11/10.
6. Sevcencu C, Struijk JJ. Autonomic alterations and cardiac changes in epilepsy. *Epilepsia*. 2010;51(5):725-37. Epub 2010/01/14.
7. Eggleston KS, Olin BD, Fisher RS. Ictal tachycardia: the head-heart connection. *Seizure*. 2014;23(7):496-505. Epub 2014/04/05.
8. van der Lende M, Surges R, Sander JW, Thijs RD. Cardiac arrhythmias during or after epileptic seizures. *J Neurol Neurosurg Psychiatry*. 2016;87(1):69-74. Epub 2015/06/04.
9. Schuele SU, Bermeo AC, Alexopoulos AV, Locatelli ER, Burgess RC, Dinner DS, et al. Video-electrographic and clinical features in patients with ictal asystole. *Neurology*. 2007;69(5):434-41. Epub 2007/08/01.
10. Tinuper P, Bisulli F, Cerullo A, Carcangiu R, Marini C, Pierangeli G, et al. Ictal bradycardia in partial epileptic seizures: Autonomic investigation in three cases and literature review. *Brain*. 2001;124(Pt 12):2361-71. Epub 2001/11/10.
11. van Dijk JG, Thijs RD, van Zwet E, Tannemaat MR, van Niekerk J, Benditt DG, et al. The semiology of tilt-induced reflex syncope in relation to electroencephalographic changes. *Brain*. 2014;137(Pt 2):576-85. Epub 2013/12/18.
12. Lanz M, Oehl B, Brandt A, Schulze-Bonhage A. Seizure induced cardiac asystole in epilepsy patients undergoing long term video-EEG monitoring. *Seizure*. 2011;20(2):167-72. Epub 2010/12/25.
13. Chaila E BJ, Tirupathi S, Delanty N. Ictal bradycardia and asystole associated with intractable epilepsy: a case series. *Br J Cardiol* 2010;17:245–8.
14. Leung H, Kwan P, Elger CE. Finding the missing link between ictal bradyarrhythmia, ictal asystole, and sudden unexpected death in epilepsy. *Epilepsy Behav*. 2006;9(1):19-30. Epub 2006/07/01.
15. Altenmuller DM, Zehender M, Schulze-Bonhage A. High-grade atrioventricular block triggered by spontaneous and stimulation-induced epileptic activity in the left temporal lobe. *Epilepsia*. 2004;45(12):1640-4. Epub 2004/12/02.
16. Oppenheimer SM, Gelb A, Girvin JP, Hachinski VC. Cardiovascular effects of human insular cortex stimulation. *Neurology*. 1992;42(9):1727-32. Epub 1992/09/01.
17. Pool JL, Ransohoff J. Autonomic effects on stimulating rostral portion of cingulate gyri in man. *J Neurophysiol*. 1949;12(6):385-92. Epub 1949/11/01.
18. Simon RP, Aminoff MJ, Benowitz NL. Changes in plasma catecholamines after tonic-clonic seizures. *Neurology*. 1984;34(2):255-7. Epub 1984/02/01.
19. Nilsson D, Sutton R, Melander O, Fedorowski A. Spontaneous vs nitroglycerin-induced vasovagal reflex on head-up tilt: Are there neuroendocrine differences? *Heart Rhythm*. 2016;13(8):1674-8. Epub 2016/07/28.
20. Moseley BD, Ghearing GR, Munger TM, Britton JW. The treatment of ictal asystole with cardiac pacing. *Epilepsia*. 2011;52(4):e16-e9.

21. Bestawros M, Darbar D, Arain A, Abou-Khalil B, Plummer D, Dupont WD, et al. Ictal asystole and ictal syncope: insights into clinical management. *Circ Arrhythm Electrophysiol.* 2015;8(1):159-64. Epub 2014/11/14.
22. Kohno R, Abe H, Akamatsu N, Benditt DG. Long-Term Follow-Up of Ictal Asystole in Temporal Lobe Epilepsy: Is Permanent Pacemaker Therapy Needed? *J Cardiovasc Electrophysiol.* 2016;27(8):930-6. Epub 2016/05/14.
23. Strzelczyk A, Cenusa M, Bauer S, Hamer HM, Mothersill IW, Grunwald T, et al. Management and long-term outcome in patients presenting with ictal asystole or bradycardia. *Epilepsia.* 2011;52(6):1160-7.
24. Duplyakov D, Golovina G, Lyukshina N, Surkova E, Elger CE, Surges R. Syncope, seizure-induced bradycardia and asystole: two cases and review of clinical and pathophysiological features. *Seizure.* 2014;23(7):506-11. Epub 2014/04/01.
25. Ryvlin P, Nashef L, Lhatoo SD, Bateman LM, Bird J, Bleasel A, et al. Incidence and mechanisms of cardiorespiratory arrests in epilepsy monitoring units (MORTEMUS): a retrospective study. *Lancet Neurol.* 2013;12(10):966-77. Epub 2013/09/10.
26. Massey CA, Sowers LP, Dlouhy BJ, Richerson GB. Mechanisms of sudden unexpected death in epilepsy: the pathway to prevention. *Nat Rev Neurol.* 2014;10(5):271-82. Epub 2014/04/23.
27. Aiba I, Noebels JL. Spreading depolarization in the brainstem mediates sudden cardiorespiratory arrest in mouse SUDEP models. *Sci Transl Med.* 2015;7(282):282ra46. Epub 2015/04/10.
28. Poh MZ, Loddenkemper T, Swenson NC, Goyal S, Madsen JR, Picard RW. Continuous monitoring of electrodermal activity during epileptic seizures using a wearable sensor. *Conf Proc IEEE Eng Med Biol Soc.* 2010;2010:4415-8. Epub 2010/11/26.
29. Bateman LM, Li CS, Seyal M. Ictal hypoxemia in localization-related epilepsy: analysis of incidence, severity and risk factors. *Brain.* 2008;131(Pt 12):3239-45. Epub 2008/10/28.
30. Surges R, Adjei P, Kallis C, Erhuero J, Scott CA, Bell GS, et al. Pathologic cardiac repolarization in pharmacoresistant epilepsy and its potential role in sudden unexpected death in epilepsy: a case-control study. *Epilepsia.* 2010;51(2):233-42.
31. Surges R, Scott CA, Walker MC. Enhanced QT shortening and persistent tachycardia after generalized seizures. *Neurology.* 2010;74(5):421-6.
32. Seyal M, Pascual F, Lee CY, Li CS, Bateman LM. Seizure-related cardiac repolarization abnormalities are associated with ictal hypoxemia. *Epilepsia.* 2011;52(11):2105-11. Epub 2011/09/13.
33. Bardai A, Lamberts RJ, Blom MT, Spanjaart AM, Berdowski J, van der Staal SR, et al. Epilepsy is a risk factor for sudden cardiac arrest in the general population. *PLoS One.* 2012;7(8):e42749.
34. Lamberts RJ, Blom MT, Wassenaar M, Bardai A, Leijten FS, de Haan GJ, et al. Sudden cardiac arrest in people with epilepsy in the community: Circumstances and risk factors. *Neurology.* 2015;85(3):212-8. Epub 2015/06/21.
35. Gaitatzis A, Carroll K, Majeed A, J WS. The epidemiology of the comorbidity of epilepsy in the general population. *Epilepsia.* 2004;45(12):1613-22. Epub 2004/12/02.
36. Suorsa E, Korpelainen JT, Ansakorpi H, Huikuri HV, Suorsa V, Myllyla VV, et al. Heart rate dynamics in temporal lobe epilepsy-A long-term follow-up study. *Epilepsy Res.* 2011;93(1):80-3. Epub 2010/11/26.
37. Lamberts RJ, Blom MT, Novy J, Belluzzo M, Seldenrijk A, Penninx BW, et al. Increased prevalence of ECG markers for sudden cardiac arrest in refractory epilepsy. *J Neurol Neurosurg Psychiatry.* 2015;86(3):309-13. Epub 2014/06/21.
38. Hartmann HA, Colom LV, Sutherland ML, Noebels JL. Selective localization of cardiac SCN5A sodium channels in limbic regions of rat brain. *Nat Neurosci.* 1999;2(7):593-5. Epub 1999/07/15.
39. Anderson JH, Bos JM, Cascino GD, Ackerman MJ. Prevalence and spectrum of electroencephalogram-identified epileptiform activity among patients with long QT syndrome. *Heart Rhythm.* 2014;11(1):53-7. Epub 2013/10/10.

40. Auerbach DS, McNitt S, Gross RA, Zareba W, Dirksen RT, Moss AJ. Genetic biomarkers for the risk of seizures in long QT syndrome. *Neurology*. 2016. Epub 2016/07/29.
41. Aurlien D, Leren TP, Tauboll E, Gjerstad L. New SCN5A mutation in a SUDEP victim with idiopathic epilepsy. *Seizure*. 2009;18(2):158-60. Epub 2008/08/30.
42. Heron SE, Hernandez M, Edwards C, Edkins E, Jansen FE, Scheffer IE, et al. Neonatal seizures and long QT syndrome: a cardiocerebral channelopathy? *Epilepsia*. 2010;51(2):293-6. Epub 2009/10/30.
43. Keller DI, Grenier J, Christe G, Dubouloz F, Osswald S, Brink M, et al. Characterization of novel KCNH2 mutations in type 2 long QT syndrome manifesting as seizures. *Can J Cardiol*. 2009;25(8):455-62. Epub 2009/08/12.
44. Partemi S, Cestele S, Pezzella M, Campuzano O, Paravidino R, Pascali VL, et al. Loss-of-function KCNH2 mutation in a family with long QT syndrome, epilepsy, and sudden death. *Epilepsia*. 2013;54(8):e112-6. Epub 2013/08/01.
45. Lehnart SE, Mongillo M, Bellinger A, Lindegger N, Chen BX, Hsueh W, et al. Leaky Ca²⁺ release channel/ryanodine receptor 2 causes seizures and sudden cardiac death in mice. *J Clin Invest*. 2008;118(6):2230-45. Epub 2008/05/17.
46. Benarroch EE. HCN channels: function and clinical implications. *Neurology*. 2013;80(3):304-10. Epub 2013/01/16.
47. Ludwig A, Budde T, Stieber J, Moosmang S, Wahl C, Holthoff K, et al. Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. *EMBO J*. 2003;22(2):216-24. Epub 2003/01/07.
48. Bagnall RD, Crompton DE, Petrovski S, Lam L, Cutmore C, Garry SI, et al. Exome-based analysis of cardiac arrhythmia, respiratory control, and epilepsy genes in sudden unexpected death in epilepsy. *Ann Neurol*. 2016;79(4):522-34. Epub 2015/12/26.
49. Leu C, Balestrini S, Maher B, Hernandez-Hernandez L, Gormley P, Hamalainen E, et al. Genome-wide Polygenic Burden of Rare Deleterious Variants in Sudden Unexpected Death in Epilepsy. *EBioMedicine*. 2015;2(9):1063-70. Epub 2015/10/27.
50. Tu E, Waterhouse L, Duflou J, Bagnall RD, Semsarian C. Genetic analysis of hyperpolarization-activated cyclic nucleotide-gated cation channels in sudden unexpected death in epilepsy cases. *Brain Pathol*. 2011;21(6):692-8. Epub 2011/05/28.
51. Shmueli S, Sisodiya SM, Gunning WB, Sander JW, Thijs RD. Mortality in Dravet syndrome: a review. *Epilepsy Behav*. In Press.
52. Auerbach DS, Jones J, Clawson BC, Offord J, Lenk GM, Ogiwara I, et al. Altered cardiac electrophysiology and SUDEP in a model of Dravet syndrome. *PLoS One*. 2013;8(10):e77843. Epub 2013/10/25.
53. Kalume F, Westenbroek RE, Cheah CS, Yu FH, Oakley JC, Scheuer T, et al. Sudden unexpected death in a mouse model of Dravet syndrome. *J Clin Invest*. 2013;123(4):1798-808. Epub 2013/03/26.
54. Delogu AB, Spinelli A, Battaglia D, Dravet C, De Nisco A, Saracino A, et al. Electrical and autonomic cardiac function in patients with Dravet syndrome. *Epilepsia*. 2011;52 Suppl 2:55-8. Epub 2011/04/08.
55. Ergul Y, Ekici B, Tatli B, Nisli K, Ozmen M. QT and P wave dispersion and heart rate variability in patients with Dravet syndrome. *Acta Neurol Belg*. 2013;113(2):161-6. Epub 2012/10/16.
56. Glasscock E, Yoo JW, Chen TT, Klassen TL, Noebels JL. Kv1.1 potassium channel deficiency reveals brain-driven cardiac dysfunction as a candidate mechanism for sudden unexplained death in epilepsy. *J Neurosci*. 2010;30(15):5167-75. Epub 2010/04/16.
57. Klassen TL, Bomben VC, Patel A, Drabek J, Chen TT, Gu W, et al. High-resolution molecular genomic autopsy reveals complex sudden unexpected death in epilepsy risk profile. *Epilepsia*. 2014;55(2):e6-12. Epub 2014/01/01.
58. Veeramah KR, O'Brien JE, Meisler MH, Cheng X, Dib-Hajj SD, Waxman SG, et al. De novo pathogenic SCN8A mutation identified by whole-genome sequencing of a family quartet affected by infantile epileptic encephalopathy and SUDEP. *Am J Hum Genet*. 2012;90(3):502-10. Epub 2012/03/01.

59. Papale LA, Beyer B, Jones JM, Sharkey LM, Tufik S, Epstein M, et al. Heterozygous mutations of the voltage-gated sodium channel SCN8A are associated with spike-wave discharges and absence epilepsy in mice. *Hum Mol Genet.* 2009;18(9):1633-41. Epub 2009/03/04.
60. Du Y, Huang X, Wang T, Han K, Zhang J, Xi Y, et al. Downregulation of neuronal sodium channel subunits Nav1.1 and Nav1.6 in the sinoatrial node from volume-overloaded heart failure rat. *Pflugers Arch.* 2007;454(3):451-9. Epub 2007/02/03.
61. Noujaim SF, Kaur K, Milstein M, Jones JM, Furspan P, Jiang D, et al. A null mutation of the neuronal sodium channel Nav1.6 disrupts action potential propagation and excitation-contraction coupling in the mouse heart. *FASEB J.* 2012;26(1):63-72. Epub 2011/09/29.
62. Scheepers B, Clough P, Pickles C. The misdiagnosis of epilepsy: findings of a population study. *Seizure.* 1998;7(5):403-6. Epub 1998/11/10.
63. Zaidi A, Clough P, Cooper P, Scheepers B, Fitzpatrick AP. Misdiagnosis of epilepsy: many seizure-like attacks have a cardiovascular cause. *J Am Coll Cardiol.* 2000;36(1):181-4. Epub 2000/07/18.
64. Grubb BP, Gerard G, Roush K, Temesy-Armos P, Elliott L, Hahn H, et al. Differentiation of convulsive syncope and epilepsy with head-up tilt testing. *Ann Intern Med.* 1991;115(11):871-6. Epub 1991/12/01.
65. Lempert T, Bauer M, Schmidt D. Syncope: a videometric analysis of 56 episodes of transient cerebral hypoxia. *Ann Neurol.* 1994;36(2):233-7. Epub 1994/08/01.
66. Blad H, Lamberts RJ, van Dijk GJ, Thijs RD. Tilt-induced vasovagal syncope and psychogenic pseudosyncope: Overlapping clinical entities. *Neurology.* 2015;85(23):2006-10. Epub 2015/11/13.
67. Mathias CJ, Deguchi K, Schatz I. Observations on recurrent syncope and presyncope in 641 patients. *Lancet.* 2001;357(9253):348-53. Epub 2001/02/24.
68. Battaglia A, Guerrini R, Gastaut H. Epileptic seizures induced by syncopal attacks. *Journal of Epilepsy* 1989;2:137-45.
69. Horrocks IA, Nechay A, Stephenson JB, Zuberi SM. Anoxic-epileptic seizures: observational study of epileptic seizures induced by syncopes. *Arch Dis Child.* 2005;90(12):1283-7. Epub 2005/09/15.
70. Stephenson J, Breningstall G, Steer C, Kirkpatrick M, Horrocks I, Nechay A, et al. Anoxic-epileptic seizures: home video recordings of epileptic seizures induced by syncopes. *Epileptic Disord.* 2004;6(1):15-9. Epub 2004/04/13.
71. Schuele SU. Effects of seizures on cardiac function. *Journal of Clinical Neurophysiology.* 2009;26(5):302-8.
72. Al Aloul B, Adabag AS, Houghland MA, Tholakanahalli V. Brugada pattern electrocardiogram associated with supratherapeutic phenytoin levels and the risk of sudden death. *Pacing Clin Electrophysiol.* 2007;30(5):713-5. Epub 2007/04/28.
73. DeGiorgio CM. Atrial flutter/atrial fibrillation associated with lacosamide for partial seizures. *Epilepsy and Behavior.* 2010;18(3):322-4.
74. El-Menyar A, Khan M, Al Suwaidi J, Eljerjawy E, Asaad N. Oxcarbazepine-induced resistant ventricular fibrillation in an apparently healthy young man. *Am J Emerg Med.* 2011;29(6):693 e1-3. Epub 2010/08/17.
75. Feldman AE, Gidal BE. QTc prolongation by antiepileptic drugs and the risk of torsade de pointes in patients with epilepsy. *Epilepsy Behav.* 2013;26(3):421-6. Epub 2012/12/12.
76. Guldiken B, Remi J, Noachtar S. Cardiovascular adverse effects of phenytoin. *J Neurol.* 2016;263(5):861-70. Epub 2015/12/10.
77. Ide A, Kamijo Y. Intermittent complete atrioventricular block after long term low-dose carbamazepine therapy with a serum concentration less than the therapeutic level. *InternMed.* 2007;46(9):627-9.
78. Ishizue N, Niwano S, Saito M, Fukaya H, Nakamura H, Igarashi T, et al. Polytherapy with sodium channel-blocking antiepileptic drugs is associated with arrhythmogenic ST-T abnormality in patients with epilepsy. *Seizure.* 2016;40:81-7. Epub 2016/07/03.

79. Kasarskis EJ, Kuo CS, Berger R, Nelson KR. Carbamazepine-induced cardiac dysfunction. Characterization of two distinct clinical syndromes. *Arch Intern Med*. 1992;152(1):186-91. Epub 1992/01/01.
80. Kaufman KR, Velez AE, Wong S, Mani R. Low-dose lacosamide-induced atrial fibrillation: Case analysis with literature review. *Epilepsy Behav Case Rep*. 2013;1:22-5. Epub 2013/01/01.
81. Krause LU, Brodowski KO, Kellinghaus C. Atrioventricular block following lacosamide intoxication. *Epilepsy Behav*. 2011;20(4):725-7. Epub 2011/03/18.
82. Nizam A, Mylavarapu K, Thomas D, Briskin K, Wu B, Saluja D, et al. Lacosamide-induced second-degree atrioventricular block in a patient with partial epilepsy. *Epilepsia*. 2011;52(10):e153-5. Epub 2011/08/02.
83. Randazzo DN, Ciccone A, Schweitzer P, Winters SL. Complete atrioventricular block with ventricular asystole following infusion of intravenous phenytoin. *J Electrocardiol*. 1995;28(2):157-9. Epub 1995/04/01.
84. Strimel WJ, Woodruff A, Cheung P, Kirmani BF, Stephen Huang SK. Brugada-like electrocardiographic pattern induced by lamotrigine toxicity. *Clin Neuropharmacol*. 2010;33(5):265-7. Epub 2010/09/25.
85. Swe T, Bhattarai B, Dufresne A. Type 1 Brugada pattern ECG due to supra-therapeutic phenytoin level. *BMJ Case Rep*. 2016;2016. Epub 2016/06/30.
86. Zoneraich S, Zoneraich O, Siegel J. Sudden death following intravenous sodium diphenylhydantoin. *Am Heart J*. 1976;91(3):375-7. Epub 1976/03/01.
87. Takayanagi K, Hisauchi I, Watanabe J, Maekawa Y, Fujito T, Sakai Y, et al. Carbamazepine-induced sinus node dysfunction and atrioventricular block in elderly women. *JpnHeart J*. 1998;39(4):469-79.
88. DeToledo JC, Lowe MR, Rabinstein A, Villaviza N. Cardiac arrest after fast intravenous infusion of phenytoin mistaken for fosphenytoin. *Epilepsia*. 2001;42(2):288. Epub 2001/03/10.
89. Elliott JO, Lu B, Shneker B, Charyton C, Layne Moore J. Comorbidity, health screening, and quality of life among persons with a history of epilepsy. *Epilepsy Behav*. 2009;14(1):125-9. Epub 2008/11/06.
90. Kobau R, Zahran H, Thurman DJ, Zack MM, Henry TR, Schachter SC, et al. Epilepsy surveillance among adults--19 States, Behavioral Risk Factor Surveillance System, 2005. *MMWR Surveill Summ*. 2008;57(6):1-20. Epub 2008/08/08.
91. Strine TW, Kobau R, Chapman DP, Thurman DJ, Price P, Balluz LS. Psychological distress, comorbidities, and health behaviors among U.S. adults with seizures: results from the 2002 National Health Interview Survey. *Epilepsia*. 2005;46(7):1133-9. Epub 2005/07/20.
92. Tellez-Zenteno JF, Matijevec S, Wiebe S. Somatic comorbidity of epilepsy in the general population in Canada. *Epilepsia*. 2005;46(12):1955-62. Epub 2006/01/06.
93. Ding D, Wang W, Wu J, Ma G, Dai X, Yang B, et al. Premature mortality in people with epilepsy in rural China: a prospective study. *Lancet Neurol*. 2006;5(10):823-7. Epub 2006/09/22.
94. Janszky I, Hallqvist J, Tomson T, Ahlbom A, Mukamal KJ, Ahnve S. Increased risk and worse prognosis of myocardial infarction in patients with prior hospitalization for epilepsy--the Stockholm Heart Epidemiology Program. *Brain*. 2009;132(Pt 10):2798-804. Epub 2009/09/01.
95. Neligan A, Bell GS, Johnson AL, Goodridge DM, Shorvon SD, Sander JW. The long-term risk of premature mortality in people with epilepsy. *Brain*. 2011;134(Pt 2):388-95. Epub 2011/02/01.
96. (CDC) CfDCaP. Comorbidity in adults with epilepsy--United States, 2010. *MMWR Morb Mortal Wkly Rep*. 2013;62(43):849-53. Epub 2013/11/01.
97. Elliott JO, Moore JL, Lu B. Health status and behavioral risk factors among persons with epilepsy in Ohio based on the 2006 Behavioral Risk Factor Surveillance System. *Epilepsy Behav*. 2008;12(3):434-44. Epub 2008/01/08.
98. Cockerell OC, Johnson AL, Sander JW, Hart YM, Goodridge DM, Shorvon SD. Mortality from epilepsy: results from a prospective population-based study. *Lancet*. 1994;344(8927):918-21. Epub 1994/10/01.

99. Nilsson L, Tomson T, Farahmand BY, Diwan V, Persson PG. Cause-specific mortality in epilepsy: a cohort study of more than 9,000 patients once hospitalized for epilepsy. *Epilepsia*. 1997;38(10):1062-8. Epub 1998/05/14.
100. St Germaine-Smith C, Liu M, Quan H, Wiebe S, Jette N. Development of an epilepsy-specific risk adjustment comorbidity index. *Epilepsia*. 2011;52(12):2161-7. Epub 2011/10/19.
101. Brodie MJ, Mintzer S, Pack AM, Gidal BE, Vecht CJ, Schmidt D. Enzyme induction with antiepileptic drugs: cause for concern? *Epilepsia*. 2013;54(1):11-27. Epub 2012/09/29.
102. Katsiki N, Mikhailidis DP, Nair DR. The effects of antiepileptic drugs on vascular risk factors: a narrative review. *Seizure*. 2014;23(9):677-84. Epub 2014/07/17.
103. Lopinto-Khoury C, Mintzer S. Antiepileptic drugs and markers of vascular risk. *Curr Treat Options Neurol*. 2010;12(4):300-8. Epub 2010/09/16.
104. Mintzer S, Skidmore CT, Abidin CJ, Morales MC, Chervoneva I, Capuzzi DM, et al. Effects of antiepileptic drugs on lipids, homocysteine, and C-reactive protein. *Ann Neurol*. 2009;65(4):448-56. Epub 2009/03/20.
105. Miller G, Vogel H. Structural evidence of injury or malformation in the brains of children with congenital heart disease. *Semin Pediatr Neurol*. 1999;6(1):20-6. Epub 1999/03/31.
106. Attar H, Sachdeva A, Sundararajan S. Cardioembolic Stroke in Adults With a History of Congenital Heart Disease. *Stroke*. 2016;47(5):e79-81. Epub 2016/04/14.
107. Ferlazzo E, Gasparini S, Beghi E, Sueri C, Russo E, Leo A, et al. Epilepsy in cerebrovascular diseases: Review of experimental and clinical data with meta-analysis of risk factors. *Epilepsia*. 2016;57(8):1205-14. Epub 2016/07/07.
108. Camilo O, Goldstein LB. Seizures and epilepsy after ischemic stroke. *Stroke*. 2004;35(7):1769-75. Epub 2004/05/29.
109. Forsgren L, Bucht G, Eriksson S, Bergmark L. Incidence and clinical characterization of unprovoked seizures in adults: a prospective population-based study. *Epilepsia*. 1996;37(3):224-9. Epub 1996/03/01.
110. Hauser WA, Annegers JF, Kurland LT. Incidence of epilepsy and unprovoked seizures in Rochester, Minnesota: 1935-1984. *Epilepsia*. 1993;34(3):453-68. Epub 1993/05/01.
111. Natelson BH, Suarez RV, Terrence CF, Turizo R. Patients with epilepsy who die suddenly have cardiac disease. *Arch Neurol*. 1998;55(6):857-60. Epub 1998/06/17.
112. Nei M, Sperling MR, Mintzer S, Ho RT. Long-term cardiac rhythm and repolarization abnormalities in refractory focal and generalized epilepsy. *Epilepsia*. 2012;53(8):e137-e40.
113. Tigaran S, Molgaard H, McClelland R, Dam M, Jaffe AS. Evidence of cardiac ischemia during seizures in drug refractory epilepsy patients. *Neurology*. 2003;60(3):492-5. Epub 2003/02/13.
114. S PCT, Dalager-Pedersen S, Baandrup U, Dam M, Vesterby-Charles A. Sudden unexpected death in epilepsy: is death by seizures a cardiac disease? *Am J Forensic Med Pathol*. 2005;26(2):99-105. Epub 2005/05/18.
115. Woodruff BK, Britton JW, Tigaran S, Cascino GD, Burritt MF, McConnell JP, et al. Cardiac troponin levels following monitored epileptic seizures. *Neurology*. 2003;60(10):1690-2. Epub 2003/05/29.
116. Finsterer J, Wahbi K. CNS disease triggering Takotsubo stress cardiomyopathy. *Int J Cardiol*. 2014;177(2):322-9. Epub 2014/09/13.
117. Finsterer J, Bersano A. Seizure-triggered Takotsubo syndrome rarely causes SUDEP. *Seizure*. 2015;31:84-7. Epub 2015/09/13.
118. Le Ven F, Pennec PY, Timsit S, Blanc JJ. Takotsubo syndrome associated with seizures: an underestimated cause of sudden death in epilepsy? *Int J Cardiol*. 2011;146(3):475-9. Epub 2011/01/05.
119. Lemke DM, Hussain SI, Wolfe TJ, Torbey MA, Lynch JR, Carlin A, et al. Takotsubo cardiomyopathy associated with seizures. *Neurocrit Care*. 2008;9(1):112-7. Epub 2008/03/19.
120. Szardien S, Mollmann H, Willmer M, Akashi YJ, Hamm CW, Nef HM. Mechanisms of stress (takotsubo) cardiomyopathy. *Heart Fail Clin*. 2013;9(2):197-205, ix. Epub 2013/04/09.

121. Manno EM, Pfeifer EA, Cascino GD, Noe KH, Wijdicks EF. Cardiac pathology in status epilepticus. *Ann Neurol*. 2005;58(6):954-7. Epub 2005/10/22.
122. Kasperaviciute D, Catarino CB, Chinthapalli K, Clayton LM, Thom M, Martinian L, et al. Uncovering genomic causes of co-morbidity in epilepsy: gene-driven phenotypic characterization of rare microdeletions. *PLoS One*. 2011;6(8):e23182. Epub 2011/08/23.
123. Ulate-Campos A, Coughlin F, Gainza-Lein M, Fernandez IS, Pearl PL, Loddenkemper T. Automated seizure detection systems and their effectiveness for each type of seizure. *Seizure*. 2016;40:88-101. Epub 2016/07/05.
124. van Andel J, Thijs RD, de Weerd A, Arends J, Leijten F. Non-EEG based ambulatory seizure detection designed for home use: What is available and how will it influence epilepsy care? *Epilepsy Behav*. 2016;57(Pt A):82-9. Epub 2016/03/02.
125. Duncan JS, Sander JW, Sisodiya SM, Walker MC. Adult epilepsy. *Lancet*. 2006;367(9516):1087-100. Epub 2006/04/04.
126. Loscher W, Schmidt D. Modern antiepileptic drug development has failed to deliver: ways out of the current dilemma. *Epilepsia*. 2011;52(4):657-78. Epub 2011/03/24.
127. Wassenaar M, van Heijl I, Leijten FS, van der Linden P, Uijl SG, Egberts AC, et al. Treatment of epilepsy in daily clinical practice: have outcomes improved over the past 10 years? *J Neurol*. 2013;260(11):2736-43. Epub 2013/07/31.