

Protocol

Schizosaccharomyces pombe Polysome Profile Analysis and RNA Purification

Dieter A. Wolf,^{1,4} Jürg Bähler,^{2,4} and Jo Ann Wise³

¹Degenerative Disease Program, Sanford Burnham Prebys Medical Discovery Institute, La Jolla, California 92037;

²Department of Genetics, Evolution & Environment, and UCL Cancer Institute, University College London, London WC1E 6BT, United Kingdom; ³Center for RNA Molecular Biology and Department of Biochemistry, Case Western Reserve University, Cleveland, Ohio 44106-4906

Polysome profile analysis is widely used by investigators studying the mechanism and regulation of translation. The method described here uses high-velocity centrifugation of whole cell extracts on linear sucrose gradients to separate 40S and 60S ribosomal subunits from 80S monosomes and polysomes. Cycloheximide is included in the lysis buffer to “freeze” polysomes by blocking translation. After centrifugation, the gradient is fractionated and RNA (and/or protein) is prepared from each fraction for subsequent analysis of individual species using northern or western blots. The entire RNA population in each fraction can be analyzed by hybridization to microarrays or by high-throughput RNA sequencing, and the proteins present can be identified by mass spectrometry analysis.

MATERIALS

It is essential that you consult the appropriate Material Safety Data Sheets and your institution’s Environmental Health and Safety Office for proper handling of equipment and hazardous material used in this protocol.

RECIPES: Please see the end of this protocol for recipes indicated by <R>. Additional recipes can be found online at <http://cshprotocols.cshlp.org/site/recipes>.

Reagents

Cycloheximide (100 mg/mL in H₂O) (optional; see Step 1)

Ethanol (85% [v/v] in water) (optional; see Step 18)

Ethanol (100%)

Ethylenediaminetetraacetic acid (EDTA) (500 mM) (optional; see Step 4.ii)

Guanidinium hydrochloride (8 M in double-distilled H₂O) (optional; see Step 16.iv)

Isopropanol

Polyribosomal lysis buffer <R> (for nonpuromycin-treated isolation procedures)

Polyribosomal lysis buffer with puromycin <R> (for puromycin-treated isolation procedures only)

Puromycin dihydrochloride (1 mM) (optional; see Step 4.i)

Puromycin is a tRNA mimic that causes polypeptide chain termination.

⁴Correspondence: dwolf@sbpdiscovery.org; j.bahler@ucl.ac.uk

From the Fission Yeast collection, edited by Iain M. Hagan, Antony M. Carr, Agnes Grallert, and Paul Nurse.

© 2017 Cold Spring Harbor Laboratory Press

Cite this protocol as *Cold Spring Harb Protoc*; doi:10.1101/pdb.prot091637

RNase I (Ambion) (optional; see Step 4.vi)
RNase A (optional; see Step 4.iv)
RNase T1 (Ambion) (optional; see Step 4.iv)
RNeasy kit (QIAGEN)
S. pombe cells ($0.5\text{--}1.0 \times 10^7$ cells/mL)
Sucrose gradient solutions (10%, 50%, and 55% [w/v], prepared in polyribosomal lysis buffer)
Water, RNase-free

Equipment

Beads, zirconia (0.5 mm diameter) (Biospec; Sigma-Aldrich)
Centrifuge, benchtop, equipped with swing-out rotor for 50-mL tubes
FastPrep device (MP Biomedicals)
Gradient fractionator (Brendel or ISCO)
Gradient maker (e.g., Biocomp or Sigma-Aldrich)
Microcentrifuge
Microcentrifuge tubes (flip-top, 1.5-mL)
Microcentrifuge tubes (screw-cap, 1.5-mL)
Microscope, phase contrast/bright field, equipped with 40 \times objective and 10 \times eyepieces
Needle (18-gauge)
Spectrophotometer (NanoDrop)
Tubes, glass (to fit the fraction collector)
Tubes (screw-cap, 50-mL) (Beckman)
Tubes, ultracentrifuge, polyallomer
Ultracentrifuge (e.g., Beckman Optima X), equipped with Sw40Ti or Sw41Ti rotors



METHOD

1. Harvest $0.3\text{--}0.7 \times 10^9$ mid-log phase cells by pouring the culture directly into 50-mL screw-cap centrifuge tubes on ice containing cycloheximide at a final concentration of 100 $\mu\text{g}/\text{mL}$. Centrifuge for 4–5 min at 670–1000g or 3 min at 2000g. Discard the culture fluid by decanting carefully.

Alternatively, add cycloheximide to the culture 5 min before harvesting to allow time for the inhibitor to work. In this case, pellet cells by centrifugation immediately (i.e., without additional added cycloheximide).

For studies in which polysome disruption involves treatment with puromycin (see Step 4.i), omit the cycloheximide treatment from Step 1, and use polyribosomal lysis buffer with puromycin throughout the protocol in place of the standard polyribosomal lysis buffer.

2. Wash cells with 0.9 mL of polyribosomal lysis buffer to remove residual media. Resuspend the pellet in 200 μL of ice-cold polyribosomal lysis buffer.
3. Transfer the cell suspension to prechilled 1.5-mL screw-cap microcentrifuge tube containing ~ 600 μL of zirconia beads. Lyse in a FastPrep device at level 6.0 for 18 sec (shorter times down to 13 sec can also be used).

Check an aliquot of the suspension using a microscope to ensure that at least 50%–70% of the cells have been lysed. If >30%–50% cells remain intact, repeat the FastPrep step.

4. (Optional) Disrupt polysomes as necessary using one of the following methods:

To induce polypeptide chain termination

- i. Treat lysate with 1 mM puromycin for 15 min.

Alternatively, add 1 mM puromycin to the cells for 1–10 min before disruption.

To dissociate ribosomal subunits

- ii. Add 25 μL of the 500 mM EDTA stock solution to 500 μL cell lysate, that is, to a final concentration of 25 mM.
- iii. Incubate for 15 min at room temperature.

To cleave RNA between individual ribosomes (Method 1)

- iv. Add RNase A and RNase T1 to final concentrations of 25 and 1000 U/mL, respectively, to the cell lysate.
- v. Incubate for 15 min at room temperature.

To cleave RNA between individual ribosomes (Method 2)

The efficacy of this procedure can vary with the batch of RNase and/or strain of yeast under investigation.

- vi. Add 750 U of RNase I to a cell lysate suspension with an OD_{260} of 40 units.
 - vii. Incubate for 30 min at room temperature.
5. After lysis (and any subsequent dissociative treatments), poke a hole in the bottom of the screw-cap microcentrifuge tube using an 18-gauge needle.
 6. Insert the perforated tube inside a 1.5-mL flip-top microcentrifuge tube. Separate the liquid from the beads in the upper tube by centrifugation at 5000g for 5 min at 4°C. Reserve the liquid collected in the flip-top tube.
 7. Add 250–350 μL of polyribosomal lysis buffer to the beads in the perforated tube. Repeat Step 6. Combine the flowthroughs from Steps 6 to 7.
 8. Clear the lysate by centrifugation at 20,000g for 15 min at 4°C.
 9. Add 2 μL of the lysate to 98 μL of dH_2O . Measure the OD_{260} of the diluted solution in a NanoDrop spectrophotometer.
 10. Prepare 11-mL linear (10%–50%) sucrose gradients.

Sucrose gradients can be stored in a cold room for up to 4 h at 4°C, with or without lysate layered on top. Do not store in a refrigerator; vibrations from the motor can disturb the gradient. See Pospíšek and Valásek (2013) for details on gradient separation.
 11. Dilute aliquots corresponding to 20–25 OD_{260} units to 350 μL with lysis buffer (with or without the protease inhibitor cocktail, heparin and RNasin). Layer onto the sucrose gradients.

When comparing polysome profiles from different cells, it is important to load the same number of OD units.
 12. Centrifuge the sucrose gradients at 4°C in a Beckman SW40Ti rotor at 39,000 rpm (~277,000g) for 150 min or a Beckman SW41Ti rotor at 35,000 rpm (~246,000g) for 160 min.

It is important to precool the centrifuge and all rotor parts.
 13. Remove gradients carefully to avoid disturbing the separation.
 14. Prewash the tubing through which the fractionated samples will pass with ethanol, then with water, and finally with sucrose solution.
 15. Fractionate each tube by upward displacement with 55% sucrose using a gradient fractionator:

The precise manipulations will vary depending on the exact combination of fractionator, UV monitor, and fraction collector. Some fractionators require a warming-up period.

Using a Brendel fractionator

- i. Set the flow rate at 800 $\mu\text{L}/\text{min}$.

Using an ISCO fractionator

- ii. Use a pump speed of 50%, 10 \times (~1.25 mL/min).

16. From each gradient, collect 12–13 fractions of 800–900 μ L each:

For protein isolation

- i. Collect fractions into empty glass tubes.

For RNA isolation (Method 1)

- ii. Collect fractions into tubes prefilled with 1.9–2.1 mL 100% isopropanol.

The final isopropanol concentration should be ~70%.

- iii. Mix each sample with the isopropanol. Precipitate overnight at -20°C .

For RNA isolation (Method 2)

- iv. Collect fractions into tubes prefilled with 2 mL of 8 M guanidinium hydrochloride.

The final concentration after collection should be 5.5 M.

- v. Add 1 volume of 100% ethanol. Mix by inversion. Precipitate overnight at -20°C .

17. After RNA precipitation (i.e., Step 16.iii or Step 16.v), centrifuge samples for 20 min at $>10,000g$ at 4°C . Discard the supernatant.

Alternatively, centrifuge at 3000g for 90 min.

18. (Optional) Wash pellets with 1 mL of 85% ethanol (see Protocol: **Preparation of Total RNA from Fission Yeast** [Bähler and Wise 2016] for tips to avoid losing the pellet).

19. Air-dry the RNA pellets for 20 min. Resuspend in 100 μ L of RNase-free water.

20. Purify RNA using a QIAGEN RNeasy kit according to the manufacturer's instructions.

RELATED INFORMATION

For additional information on the use of polysome profile analysis to study the mechanisms and regulation of translation, see Rallis et al. (2014). Procedures for RNA analysis by microarray hybridization are described in Lackner et al. (2012).

RECIPES

Polysomal Lysis Buffer

Tris-HCl (pH 7.5)	20 mM
KCl	50 mM
MgCl ₂	10 mM
Cycloheximide	100 μ g/mL
Dithiothreitol (DTT)	1 mM
cOmplete protease inhibitor cocktail tablets, EDTA-free (Roche)	2 tablets/100 mL culture
Heparin	0.2 mg/mL
RNasin (Promega)	1 μ L/mL
Phenylmethylsulfonyl fluoride (PMSF)	1 mM

Combine the first three ingredients. (If desired, a 10 \times stock solution containing 200 mM Tris-HCl, 500 mM KCl, and 100 mM MgCl₂ can be prepared and stored on a long-term basis.) Just before use, add the cycloheximide and DTT, as well as the protease inhibitor tablets, heparin, and RNasin (the latter three are optional but highly recommended). For processing meiotic cells, include PMSF.

Polysomal Lysis Buffer with Puromycin

Tris-HCl (pH 7.5)	20 mM
KCl	500 mM
MgCl ₂	2 mM
Puromycin	1 mM
Dithiothreitol (DTT)	1 mM
cOmplete protease inhibitor cocktail tablets, EDTA-free (Roche)	2 tablets/100 mL culture
Heparin	0.2 mg/mL
RNasin (Promega)	1 μL/mL
Phenylmethylsulfonyl fluoride (PMSF)	1 mM

Combine the first three ingredients. Just before use, add the puromycin and DTT, as well as the protease inhibitor tablets, heparin, and RNasin (the latter three are optional but highly recommended). For processing meiotic cells, include PMSF.

ACKNOWLEDGMENTS

D.A.W. was supported by the National Institutes of Health USA, J.B. was supported by the Wellcome Trust UK, and J.A.W. was supported by the National Science Foundation USA. This protocol is an amalgamation of procedures followed in the laboratories of Dieter Wolf and Jürg Bähler. The authors thank Charalampos Rallis (University College London, UK) for helpful comments.

REFERENCES

- Bähler J, Wise JA. 2016. Preparation of total RNA from fission yeast. *Cold Spring Harb Protoc* doi: 10.1101/pdb.prot091629.
- Lackner DH, Schmidt MW, Wu S, Wolf DA, Bähler J. 2012. Regulation of transcriptome, translation, and proteome in response to environmental stress in fission yeast. *Genome Biol* 13: R25.
- Pospíšek M, Valásek L. 2013. Polysome profile analysis—Yeast. *Methods Enzymol* 530: 173–181.
- Rallis C, López-Maury L, Georgescu T, Pancaldi V, Bähler J. 2014. Systematic screen for mutants resistant to TORC1 inhibition in fission yeast reveals genes involved in cellular ageing and growth. *Biol Open* 3: 161–171.





Cold Spring Harbor Protocols

***Schizosaccharomyces pombe* Polysome Profile Analysis and RNA Purification**

Dieter A. Wolf, Jürg Bähler and Jo Ann Wise

Cold Spring Harb Protoc; doi: 10.1101/pdb.prot091637

Email Alerting Service

Receive free email alerts when new articles cite this article - [click here](#).

Subject Categories

Browse articles on similar topics from *Cold Spring Harbor Protocols*.

[Analysis of Protein Expression in Cultured Cells](#) (44 articles)
[mRNA](#) (111 articles)
[Preparation of Cellular and Subcellular Extracts](#) (90 articles)
[RNA](#) (251 articles)
[RNA Purification](#) (56 articles)
[Subcellular Fractionation](#) (70 articles)
[Yeast](#) (219 articles)

To subscribe to *Cold Spring Harbor Protocols* go to:
<http://cshprotocols.cshlp.org/subscriptions>
