

Table 1 Demographic, diagnostic and risk profile features of the community-based (CBP), step-down (RT-CBP) and residential (RT) samples

	CBP (n=30)		RT-CBP (n=87)		RT (n=45)		Test of significance	OR (95% CI)		
	N	%	N	%	N	%		CBP v RT-CBP	CBP v RT	RT-CBP v RT
Female	20	66.7	69	79.3	36	80.0	$\chi^2_{(2)}=2.31, p=0.315$	0.52 (0.2, 1.32)	0.5 (0.17, 1.45)	0.95 (0.38, 2.36)
Single	18	60.0	61	71.1	31	68.9	$\chi^2_{(2)}=1.08, p=0.584$	0.63 (0.26, 1.52)	0.67 (0.25, 1.8)	1.05 (0.48, 2.32)
College Education	13	43.3	48	55.2	22	48.9	$\chi^2_{(2)}=1.27, p=0.531$	0.62 (0.26, 1.44)	0.73 (0.28, 1.88)	1.17 (0.56, 2.43)
Any Loss	16	53.3	38	43.7	23	51.1	$\chi^2_{(2)}=0.93, p=0.627$	1.47 (0.63, 3.41)	1.19 (0.46, 3.06)	0.81 (0.39, 1.67)
Any Sexual abuse	19	63.3	41	47.1	22	48.9	$\chi^2_{(2)}=2.42, p=0.298$	1.93 (0.81, 4.59)	1.8 (0.69, 4.71)	0.93 (0.45, 1.92)
Physical Abuse	17	56.7	43	49.4	24	53.3	$\chi^2_{(2)}=0.52, p=0.770$	1.33 (0.57, 3.11)	1.14 (0.44, 2.94)	0.85 (0.41, 1.77)
Avoidant PD	17	56.7	47	54.0	24	53.3	$\chi^2_{(2)}=0.09, p=0.957$	1.11 (0.47, 2.59)	1.14 (0.44, 2.94)	1.02 (0.49, 2.12)
Obsessive-compulsive PD	15	50.0	26	29.9	16	35.6	$\chi^2_{(2)}=3.96, p=0.138$	2.34 (0.99, 5.53)	1.81 (0.69, 4.71)	0.77 (0.35, 1.66)
Schizotypal PD	6	20.0	9	10.3	8	17.8	$\chi^2_{(2)}=2.36, p=0.307$	2.16 (0.69, 6.78)	1.15 (0.34, 3.82)	0.53 (0.18, 1.5)
Narcissistic PD	1	3.0	12	13.8	4	8.9	$\chi^2_{(2)}=4.84, p=0.089$	0.21 (0.02, 1.77)	0.35 (0.03, 3.45)	1.64 (0.49, 5.47)
Borderline PD	21	70.0	61	70.1	32	71.1	$\chi^2_{(2)}=0.02, p=0.992$	0.99 (0.39, 2.48)	0.94 (0.33, 2.65)	0.95 (0.42, 2.11)
Antisocial PD	3	10.0	9	10.3	4	8.9	$\chi^2_{(2)}=0.07, p=0.143$	0.96 (0.23, 3.87)	1.13 (0.23, 5.63)	1.18 (0.33, 4.12)
Clinical severity ^a (severe)	16	53.3	40	46.0	24	53.3	$\chi^2_{(2)}=0.87, p=0.647$	1.34 (0.57, 3.11)	1.00 (0.39, 2.56)	0.74 (0.35, 1.54)

	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>		<i>95% CI for the difference</i>		
Age	39.93	10.31	33.62	8.09	31.18	7.49	$F_{(2)}=10.14, p=0.000$	2.02, 10.61	3.97, 13.54	-1.28, 6.17
Average number of PD	3.87	1.91	3.60	1.97	3.60	1.76	$F_{(2)}=0.24, p=0.785$	-0.71, 1.24	-0.82, 1.35	-0.85, 0.84
GSI ^b	2.35	0.78	1.91	0.66	1.86	0.78	$F_{(2)}=5.02., p=0.008$	0.07, 0.80	0.08, 0.90	-0.26, 0.37

^a Clinical severity: calculated from the median number of Axis-II diagnoses each patient met criteria for, number of positive PD criteria met and intake general psychiatric distress scores

^b Brief Symptom Inventory General Severity Index