

Disclaimer: The manuscript and its contents are confidential,
intended for journal review purposes only, and not to be further
disclosed.

URL: <http://circimaging-submit.aha-journals.org>

Title: Mapping myocardial salvage index by extracellular volume
fraction: Are we there yet?

Manuscript number: CIRCCVIM/2017/006680

Author(s): Derek Hausenloy, University College London

**Mapping myocardial salvage index by extracellular volume fraction:
Are we there yet?**

Heerajnarain Bulluck PhD^{1,2,3}, Derek J Hausenloy MBChB PhD^{1,2,4,5,6,7}

¹The Hatter Cardiovascular Institute, Institute of Cardiovascular Science, University College London, UK

²The National Institute of Health Research University College London Hospitals Biomedical Research Centre, UK

³Papworth Hospital NHS Trust, Cambridge, UK

⁴Barts Heart Centre, St Bartholomew's Hospital, London, UK

⁵National Heart Research Institute Singapore, National Heart Centre Singapore

⁶Cardiovascular and Metabolic Disorders Program, Duke-National University of Singapore

⁷Yong Loo Lin School of Medicine, National University Singapore, Singapore

Corresponding author:

Professor Derek J Hausenloy
Cardiovascular & Metabolic Diseases Program
Duke-NUS Graduate Medical School Singapore
8 College Road,
Singapore 169857
Tel +65 65166719
Email derek.hausenloy@duke-nus.edu.sg

In ST-segment elevation myocardial infarction (STEMI) patients treated by primary percutaneous coronary intervention (PPCI), the myocardial salvage index (MSI) provides a more sensitive measure for assessing the efficacy of novel cardioprotective therapies, than an absolute reduction in myocardial infarct (MI) size. Knowledge of the MI size and the size of the area-at-risk (AAR) are pre-requisites for measuring the MSI, and both may be obtained by cardiovascular magnetic resonance (CMR) in reperfused STEMI patients, with the MSI shown to predict clinical outcomes following PPCI ¹. CMR is considered the gold standard imaging modality for quantifying MI size, and it can also delineate the edema-based AAR, with T2 and T1-mapping CMR emerging as the most robust techniques ², although no consensus has yet been reached.

In this issue of *Circulation: Cardiovascular Imaging*, Garg et al ³ report on a potentially novel approach for quantifying the AAR, chronic MI size, and MSI in a study of 50 STEMI patients reperfused by PPCI, based on extracellular volume fraction (ECV) maps from an acute CMR scan. The study derived specific ECV cut-off values on an acute CMR scan (performed at a median of 48 hours post-PPCI) in a subset of 10 patients, to delineate the AAR (when compared to T2-STIR imaging), and chronic MI size (when compared late gadolinium enhancement [LGE]) on a follow-up scan performed at 3 months post-PPCI). Using acute ECV cut-off values of >33% to delineate the AAR and >46% to delineate chronic MI size, they concluded that acute ECV maps could be used to reliably quantify AAR, chronic MI size, and MSI. Being able to accurately quantify MI size using a pixel-wise acute ECV map would be appealing, and could be potentially easily implemented in clinical practice with the wider availability of in-line, automated ECV map generation from the scanner, to improve workflow ⁴. However, there are several potential limitations to

consider concerning the use of acute ECV maps to detect the AAR, chronic MI size and MSI in reperfused STEMI patients.

Firstly, the study did not specifically compare the performance of ECV-derived MSI with conventional MSI (from LGE and T2-STIR), and whether acute ECV maps can be used to estimate the MSI remains to be demonstrated. The authors could have considered using the reference standard of manual delineation of AAR and MI size by experienced operators, rather than semi-automated thresholding methods, which are known to have their limitations^{5, 6}. The presence of microvascular obstruction (MVO) on the acute CMR scan is known to pseudo-normalize the ECV^{5, 7}, due to failure of the gadolinium chelate to penetrate areas of MVO, which could have affected MI size regression, thereby making it challenging to estimate chronic MI size from the acute ECV in patients with MVO. Finally, the study used $\leq 75\%$ transmural MI on the follow-up CMR scan to define viable myocardium, and an increase of $\geq 15\%$ in left ventricular (LV) end-systolic volume to define adverse LV remodeling, instead of the conventionally accepted definitions of $\leq 50\%$ for viability on CMR⁸, and $\geq 20\%$ increase in LV end-diastolic volume for adverse LV remodeling on echocardiography⁹. In this regard, we have recently proposed CMR-based definitions for assessing adverse LV remodeling following STEMI, which may be helpful for future CMR studies¹⁰. The study findings are also thought-provoking, and raise some interesting questions for the field.

Can acute ECV maps be used to accurately delineate the chronic MI size given the pathological differences that exist between acute and chronic MI?

It is well-recognized that acute MI size is dynamic, and acutely reduces in size within the first week^{11, 12}, and more chronically reduces over the first few months^{5, 11}. The regression in MI size represents the gradual resolution of myocardial edema,

intramyocardial hemorrhage (IMH) and MVO, and the gradual replacement of necrotic tissue with fibrosis in the chronic infarct ¹¹. Furthermore, LV remodeling leads to thinning of the infarcted territory, and compensatory LV hypertrophy of the adjacent and remote myocardium, resulting in the overall reduction in LGE mass observed in the chronic phase. The over-estimation of acute MI size by LGE on the acute CMR scan can be minimized by acquiring LGE images at least 15 minutes after contrast injection and undertaking the acute CMR scan at day 7 ¹³. In addition, acute MI size has recently been shown to be prognostic ¹⁴, and may help to reduce sample size in clinical cardioprotection studies in reperfused STEMI patients, thereby obviating the need to measure chronic MI size, which can be challenging to perform in clinical cardioprotection studies ⁶. Therefore, the differences between the pathologies underlying acute and chronic MI, may in part, have impacted on the study findings, and may have contributed to the substantial variability observed between acute MI size by ECV and chronic MI size by LGE, with the Bland-Altman analysis showing a bias of 1.9% and wide limits of agreement of $\pm 10.5\%$. This suggests that acute ECV maps may not yet be ready to measure chronic MI size following STEMI.

Can acute ECV maps be used to accurately delineate the AAR given that edema in salvaged myocardium can be both intracellular and interstitial following STEMI?

During acute myocardial ischemia, interruption of the blood supply to the myocardium disrupts cardiomyocyte Na^+/K^+ channel function, leading to increased trans-membrane Na^+ gradients and intracellular edema. Prolonged periods of ischemia can result in cardiomyocyte cell membrane rupture, thereby adding to the intracellular edema. Alterations in capillary permeability can promote interstitial edema, resulting in both intracellular and interstitial edema during acute myocardial ischemia.

Furthermore, reperfusion can exacerbate both interstitial and intracellular edema, and may also lead to extravasation of red blood cells¹⁵. Native T1-mapping, T2-mapping or T2-STIR CMR imaging detect both intracellular and interstitial edema, whereas acute ECV can only detect edema in the latter compartment¹⁶, and therefore areas of intracellular edema in the salvaged myocardium may be overlooked by acute ECV.

It may also be challenging for acute ECV maps to reliably differentiate salvaged myocardium within the AAR from remote myocardium outside the AAR, given that there can be overlap in acute ECV values between the salvaged and remote myocardium following STEMI. In this regard, Hammer-Hansen et al¹⁷ have recently reported an overlap in acute ECV values between salvaged versus remote myocardium (95% of the acute ECV values were 18 - 30% in the remote, and 28 - 51% in the salvaged myocardium) following STEMI. An overlap in acute ECV values between the salvaged and remote myocardium may also have affected the study by Garg et al³, with areas of acute ECV >33% observed in the remote myocardium (see figure 3, case 1). Furthermore, a previous study by the same research group had reported an acute ECV of 29±6% in the remote myocardium¹⁸, confirming the potential for overlap in acute ECV values between salvaged and remote myocardium. Again, this may explain, in part, the wide limits of agreement (±10.4%) observed in the current study, indicating a large variability in the observed values between acute ECV AAR and T2-STIR AAR³. In a previous study, we had observed limits of agreement of ±5.1% when comparing T1 and T2-mapping to detect the AAR in reperfused STEMI patients², and limits of agreement of ±10.4% observed with acute ECV maps for delineating T2-weighted AAR may be too wide for clinical application. This suggests that acute ECV maps may not yet be ready to measure AAR following STEMI.

In summary, performing a comprehensive CMR study in acutely reperfused STEMI patients can be very challenging, and Garg et al³ should be congratulated on their study investigating the potential role of acute ECV mapping for assessing AAR, chronic MI size and MSI. Although, multi-parametric CMR mapping has the real potential to provide valuable insights into the changes occurring in areas of MI, MVO, IMH, salvaged and remote myocardium in the setting of STEMI^{7, 19, 20}, more validation work is needed before acute ECV mapping, can be used to reliably assess AAR, MI size, and MSI in reperfused STEMI patients.

Acknowledgements

Derek Hausenloy is supported by the British Heart Foundation (FS/10/039/28270), Duke-National University Singapore Medical School, and the National Institute for Health Research University College London Hospitals Biomedical Research Centre. This research is supported by the Singapore Ministry of Health's National Medical Research Council under its Clinician Scientist-Senior Investigator scheme (NMRC/CSA-SI/0011/2017) and Collaborative Centre Grant scheme (NMRC/CGAug16C006). This research is supported by the Singapore Ministry of Education Academic Research Fund Tier 2 (MOE2016-T2-2-021).

Disclosures

The authors declare no disclosures

References

1. Eitel I, Desch S, Fuernau G, Hildebrand L, Gutberlet M, Schuler G and Thiele H. Prognostic significance and determinants of myocardial salvage assessed by cardiovascular magnetic resonance in acute reperfused myocardial infarction. *Journal of the American College of Cardiology*. 2010;55:2470-9.
2. Bulluck H, White SK, Rosmini S, Bhuvu A, Treibel TA, Fontana M, Abdel-Gadir A, Herrey A, Manisty C, Wan SM, Groves A, Menezes L, Moon JC and Hausenloy DJ. T1 mapping and T2 mapping at 3T for quantifying the area-at-risk in reperfused STEMI patients. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*. 2015;17:73.
3. Garg P, Broadbent D, Swoboda P, Foley JR, Fent GJ, Musa TA, Ripley DP, Erhayiem B, Dobson LE, McDiarmid A, Haaf P, Kidambi A, Geest RJVD, P. GJ and Plein S. Acute Infarct Extracellular Volume Mapping to Quantify Myocardial Area at Risk and Chronic Infarct Size on Cardiovascular Magnetic Resonance Imaging. *Circulation Cardiovascular imaging*. 2017.
4. Treibel TA, Fontana M, Maestrini V, Castelletti S, Rosmini S, Simpson J, Nasis A, Bhuvu AN, Bulluck H, Abdel-Gadir A, White SK, Manisty C, Spottiswoode BS, Wong TC, Piechnik SK, Kellman P, Robson MD, Schelbert EB and Moon JC. Automatic Measurement of the Myocardial Interstitium: Synthetic Extracellular Volume Quantification Without Hematocrit Sampling. *JACC Cardiovascular imaging*. 2016;9:54-63.
5. Bulluck H, Rosmini S, Abdel-Gadir A, Bhuvu AN, Treibel TA, Fontana M, Weinmann S, Sirker A, Herrey AS, Manisty C, Moon JC and Hausenloy DJ. Impact of microvascular obstruction on semiautomated techniques for quantifying acute and chronic myocardial infarction by cardiovascular magnetic resonance. *Open Heart*. 2016;3:e000535.
6. Bulluck H, Hammond-Haley M, Weinmann S, Martinez-Macias R and Hausenloy DJ. Myocardial Infarct Size by CMR in Clinical Cardioprotection Studies: Insights From Randomized Controlled Trials. *JACC Cardiovascular imaging*. 2017;10:230-240.
7. Bulluck H, Rosmini S, Abdel-Gadir A, White SK, Bhuvu AN, Treibel TA, Fontana M, Gonzalez-Lopez E, Reant P, Ramlall M, Hamarneh A, Sirker A, Herrey AS, Manisty C, Yellon DM, Kellman P, Moon JC and Hausenloy DJ. Automated Extracellular Volume Fraction Mapping Provides Insights Into the Pathophysiology of Left Ventricular Remodeling Post-Reperfused ST-Elevation Myocardial Infarction. *Journal of the American Heart Association*. 2016;5.
8. Kim RJ, Wu E, Rafael A, Chen EL, Parker MA, Simonetti O, Klocke FJ, Bonow RO and Judd RM. The use of contrast-enhanced magnetic resonance imaging to identify reversible myocardial dysfunction. *The New England journal of medicine*. 2000;343:1445-53.
9. Bolognese L, Cerisano G, Buonamici P, Santini A, Santoro GM, Antoniucci D and Fazzini PF. Influence of infarct-zone viability on left ventricular remodeling after acute myocardial infarction. *Circulation*. 1997;96:3353-9.

10. Bulluck H, Go YY, Crimi G, Ludman AJ, Rosmini S, Abdel-Gadir A, Bhuva AN, Treibel TA, Fontana M, Pica S, Raineri C, Sirker A, Herrey AS, Manisty C, Groves A, Moon JC and Hausenloy DJ. Defining left ventricular remodeling following acute ST-segment elevation myocardial infarction using cardiovascular magnetic resonance. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*. 2017;19:26.
11. Engblom H, Hedstrom E, Heiberg E, Wagner GS, Pahlm O and Arheden H. Rapid initial reduction of hyperenhanced myocardium after reperfused first myocardial infarction suggests recovery of the peri-infarction zone: one-year follow-up by MRI. *Circulation Cardiovascular imaging*. 2009;2:47-55.
12. Carrick D, Haig C, Ahmed N, Rauhalammi S, Clerfond G, Carberry J, Mordi I, McEntegart M, Petrie MC, Eteiba H, Hood S, Watkins S, Lindsay MM, Mahrous A, Welsh P, Sattar N, Ford I, Oldroyd KG, Radjenovic A and Berry C. Temporal Evolution of Myocardial Hemorrhage and Edema in Patients After Acute ST-Segment Elevation Myocardial Infarction: Pathophysiological Insights and Clinical Implications. *Journal of the American Heart Association*. 2016;5:Advance online publication.
13. Jablonowski R, Engblom H, Kanski M, Nordlund D, Koul S, van der Pals J, Englund E, Heiberg E, Erlinge D, Carlsson M and Arheden H. Contrast-Enhanced CMR Overestimates Early Myocardial Infarct Size: Mechanistic Insights Using ECV Measurements on Day 1 and Day 7. *JACC Cardiovascular imaging*. 2015;8:1379-89.
14. Stone GW, Selker HP, Thiele H, Patel MR, Udelson JE, Ohman EM, Maehara A, Eitel I, Granger CB, Jenkins PL, Nichols M and Ben-Yehuda O. Relationship Between Infarct Size and Outcomes Following Primary PCI: Patient-Level Analysis From 10 Randomized Trials. 2016.
15. Garcia-Dorado D and Oliveras J. Myocardial oedema: a preventable cause of reperfusion injury? *Cardiovascular research*. 1993;27:1555-63.
16. Bulluck H, Maestrini V, Rosmini S, Abdel-Gadir A, Treibel TA, Castelletti S, Bucciarelli-Ducci C, Manisty C and Moon JC. Myocardial T1 mapping. *Circulation journal : official journal of the Japanese Circulation Society*. 2015;79:487-94.
17. Hammer-Hansen S, Bandettini WP, Hsu LY, Leung SW, Shanbhag S, Mancini C, Greve AM, Kober L, Thune JJ, Kellman P and Arai AE. Mechanisms for overestimating acute myocardial infarct size with gadolinium-enhanced cardiovascular magnetic resonance imaging in humans: a quantitative and kinetic study. *European heart journal cardiovascular Imaging*. 2016;17:76-84.
18. Kidambi A, Motwani M, Uddin A, Ripley DP, McDiarmid AK, Swoboda PP, Broadbent DA, Musa TA, Erhayiem B, Leader J, Croisille P, Clarysse P, Greenwood JP and Plein S. Myocardial Extracellular Volume Estimation by CMR Predicts Functional Recovery Following Acute MI. *JACC Cardiovascular imaging*. 2016.
19. Bulluck H, Rosmini S, Abdel-Gadir A, Bhuva AN, Treibel TA, Fontana M, Gonzalez-Lopez E, Ramlall M, Hamarneh A, Sirker A, Herrey AS, Manisty C, Yellon DM, Moon JC and Hausenloy DJ. Diagnostic Performance of T1 and T2 Mapping to Detect Intramyocardial Hemorrhage in Reperfused ST-Segment Elevation Myocardial Infarction (STEMI) Patients. *Journal of magnetic resonance imaging : JMRI*. 2017.
20. Bulluck H, Rosmini S, Abdel-Gadir A, White SK, Bhuva AN, Treibel TA, Fontana M, Ramlall M, Hamarneh A, Sirker A, Herrey AS, Manisty C, Yellon DM, Kellman P, Moon JC and Hausenloy DJ. Residual Myocardial Iron Following

Intramyocardial Hemorrhage During the Convalescent Phase of Reperfused ST-Segment-Elevation Myocardial Infarction and Adverse Left Ventricular Remodeling. *Circulation Cardiovascular imaging*. 2016;9.

Disclaimer: The manuscript and its contents are confidential, intended for journal review purposes only, and not to be further disclosed.