

**Intracerebral hemorrhage as a manifestation of cerebral hyperperfusion syndrome after carotid revascularization: Systematic review and meta-analysis**

Authors:

Pedro Abreu<sup>1</sup>

Jerina Nogueira<sup>1</sup>

Filipe Brogueira Rodrigues<sup>2,3,4</sup>

Ana Nascimento<sup>5</sup>

Mariana Carvalho<sup>5</sup>

Ana Marreiros<sup>1</sup>

Hipólito Nzwalo<sup>1</sup>

<sup>1</sup>Department of Biomedical Sciences and Medicine, University of Algarve, Portugal

<sup>2</sup>Clinical Pharmacology Unit, Instituto de Medicina Molecular, Lisboa, Portugal

<sup>3</sup>Laboratory of Clinical Pharmacology and Therapeutics, Faculty of Medicine, University of Lisbon, Lisbon, Portugal

<sup>4</sup>Huntington's Disease Centre, Institute of Neurology, University College London, London, United Kingdom

<sup>5</sup>Department of Neurology, Centro Hospitalar do Algarve, Portugal

**Corresponding Author :**

Hipólito Nzwalo, MD, MSc

Departement of Biomedical Sciences and Medicine,

University of Algarve, Campus da Penha, 8005-139 Faro, Portugal.

E-mail address: nzwalo@gmail.com

## **Abstract**

**Background:** Intracerebral hemorrhage (ICH) in the context of cerebral hyperperfusion syndrome (CHS) is an uncommon but potentially lethal complication after carotid revascularization for carotid occlusive disease. Information about its incidence, risk factors and fatality is scarce. Therefore, we aimed to perform a systematic review and meta-analysis focusing on the incidence, risk factors and outcomes of ICH in the context of CHS after carotid revascularization.

**Methods:** We searched on PubMed and EBSCO host for all studies published in English about CHS in the context of carotid revascularization. Two reviewers independently assessed each study for eligibility based on predefined criteria. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed and PROSPERO register was made (register number: CRD42016033190), including the pre-specified protocol.

**Results:** Forty-one studies involving 28956 participants were deemed eligible and included in our analysis. The overall quality of the included studies was fair. The pooled frequency of ICH in the context of CHS was 38% (95% CI: 26% to 51%, I<sup>2</sup>=84%, 24 studies) and the pooled case-fatality of ICH after CHS was 51% (95% CI: 32% to 71%, I<sup>2</sup>=77%, 17 studies). When comparing carotid angioplasty with stenting (CAS) with carotid endarterectomy (CEA), post-procedural ICH in the context of CHS was less frequent in CEA. ICH following CHS occurred less often in large series and was rare in asymptomatic patients. The most common risk factors were periprocedural hypertension and ipsilateral severe stenosis.

**Conclusions:** ICH as a manifestation of CHS is rare, more frequent after CAS, and associated with poor prognosis. Periprocedural control of hypertension can reduce its occurrence.

## **Keywords**

“cerebral hyperperfusion syndrome”, “intracerebral hemorrhage”, “carotid endarterectomy”, “carotid angioplasty”, “carotid revascularization”

## **1-Introduction**

Ischemic strokes is a major cause of morbidity and mortality worldwide. (12, 33, 53) Carotid artery disease contributes to 18-29% of all cases.(59, 66) Carotid invasive interventions such as carotid endarterectomy (CEA) and carotid angioplasty with stenting (CAS) are widely used to treat carotid artery disease.(6, 26, 62) One potentially severe complication is intracerebral hemorrhage (ICH) in the context of cerebral hyperperfusion syndrome (CHS).(37) Clinical manifestations of CHS are diverse and include symptoms such as throbbing headaches, confusion, focal neurological deficits, partial or generalized seizures, among others.(25, 27, 60) Diagnostic exams such computerized tomography scan (CT), magnetic resonance imaging (MRI), transcranial Doppler (TCD), single-photon emission computerized tomography (SPECT) and positron emission tomography (PET) can be used to confirm or exclude this diagnosis.(3, 11, 24, 60) The pathophysiology behind CHS is still poorly understood.(41) Three mechanisms have been associated with the syndrome: failure of brain vessels' auto-regulatory mechanisms to adapt to the sudden and deregulated increase in cerebral blood flow after carotid revascularization in long standing hypoperfused brains due to severe stenosis/obstruction(2, 36, 48, 54, 60); baroreflex disturbances secondary to carotid revascularization (58); disturbances in the trigeminovascular reflex.(35, 60) Despite its severity, the knowledge about the frequency, risk factors and prognosis of ICH in the context of CHS is scarce. Previous reviews did not specifically addressed the occurrence of ICH in the context of CHS.(32, 41) Therefore, we set up to perform a systematic review of the existent studies and meta-analytically estimate the frequency of ICH after CHS and its case-fatality.

## **2-Material and Methods**

### **2.1-Protocol and registration**

This systematic review was registered at PROSPERO database (CRD42016033190) and written in accordance with PRISMA guidelines.(31)

### **2.2-Eligibility criteria**

Primary studies involving patients submitted to carotid revascularization (CAS or CEA) due to carotid occlusive disease were included. Studies with CEA/CAS preformed for

other specific conditions, case reports and animal studies; and studies in which the definition and frequency of ICH related CHS was not described were excluded.

### **2.3-Information sources**

The search process was performed using the search engines PubMed and EBSCOHost (1986 to January 2016). Databases accessed via EBSCOhost include MEDLINE, sciencedirect, academic one file, J-stage, general one file, OAlster, expanded academic ASAP, China/Asia on demand, SciELO, Scitech conect, MedicLatina and Korean studies information study system. Only full-text English-written publications were included.

### **2.4-Search strategy and study selection**

The Mesh terms “cerebral hyperperfusion syndrome”, “complications”, “carotid revascularization”, “endarterectomy” and “carotid angioplasty” were used to retrieve relevant literature. Studies were selected by two independent investigators. A consensus between the authors was used to resolve any disagreements about the inclusion of specific studies.

### **2.6-Data collection process**

Studies were analysed by two independent investigators. A consensus between the authors was used to resolve any disagreements about the inclusion of specific studies.

### **2.7-Data items**

The following items were extracted: type surgical procedure; frequency of CHS with ICH; associated risk factors, outcome (case-fatality, morbidity rates).

### **2.8-Studies Risk of bias**

The National institutes of Health (NIH) tools were used (15) for quality assessment (supplement). Two reviewers performed the assessment independently. Discrepancies in the classifications were discussed and agreement achieved.

### **2.11-Planned methods of analysis**

Qualitative analysis with quantitative description including all selected studies was performed whenever applicable. To address the risk factors; and for the meta-analysis of frequency and case-fatality of ICH after CHS, only large studies ( $\geq 100$  patients) were

considered. This arbitrary threshold was selected to minimize the effects of substantial variability in the diagnostic criteria, time of evaluation and populations included in our study. We used Stata/SE 14.0 software for conducting the analysis and to derive forest plots. Random-effects meta-analysis weighted by the inverse-variance method was performed to estimate pooled frequency and 95% confidence intervals (CI). Heterogeneity was assessed with the I<sup>2</sup> test. We used a random-effects model as substantial heterogeneity between studies results was expected. The limit for statistical significance was established at 0.05.

### **3-Results**

#### **3.1-Study selection**

Initial search yielded a total of 545 manuscripts (423 publications at EBSCOhost and 122 publications at PubMed) (Figure 1). After extraction removal of duplicates and studies not fulfilling our eligibility criteria, 41 studies were included in the final analysis (Figure 1). Reasons for study exclusions were documented at supplement - 1.

#### **3.2-Study characteristics**

A total of 28,956 participants were included, with studies sample sizes ranging from 26 to 4,494 participants (Table 1). Eighteen studies (44%) defined both CHS and ICH at methods section. (1, 2, 7-9, 16, 19, 20, 22, 24, 29, 34, 42, 44, 46, 47, 61, 65) The frequency of ICH after carotid intervention in studies with less than 100 participants ranged from 0% to 4.44%. (4, 8, 14, 17, 18, 21, 22, 24, 40, 46, 48, 64) In studies with 101 to 1000 participants the range was from 0% to 2.21%. (1, 2, 5, 7, 9, 10, 13, 19, 20, 23, 29, 30, 34, 39, 42, 43, 51, 55, 57, 65) In studies that included 1001 or more participants a range from 0.09% to 0.6% was found. (16, 44, 45, 47, 49, 50, 52, 61, 63) With regard to the quality evaluation, the majority were rated as "good" (1, 2, 5, 7, 8, 18, 22, 29, 30, 34, 50, 65) or fair (4, 9, 10, 13, 16, 17, 19-21, 23, 24, 39, 40, 42-49, 51, 55-57, 63, 64), and three were "poor". (14, 52, 61) The risk of bias was considerable in most studies (Supplement 2). (4, 9, 10, 13, 16, 17, 19-21, 23, 24, 39, 42-45, 47, 49, 55, 64) The pooled frequency of ICH in the context of CHS was 38% (95% CI: 26% to 51%, I<sup>2</sup>=84%) (Figure 2)

#### **3.3-Risk factors**

Table 2 resumes the risk factors found in larger studies ( $\geq 100$  patients). Table 1 shows that when comparing CEA and CAS, the frequency of ICH in the context of CHS in large studies was higher after CAS, ranging from 0.28% to 4.05%. (1, 7, 9, 19, 30, 39, 42, 47,

49, 55, 57, 65) In CEA varied from 0% to 2.15%.(2, 5, 9, 10, 13, 16, 18, 20, 23, 29, 34, 43-45, 47, 61) Also, 73% of the studies involving CAS had a frequency of ICH in the context of CHS above 0.5%.(1, 7, 9, 19, 30, 39, 42, 47, 49, 55, 57, 65) In CEA, these frequencies occurred in only 26% of the cases.(2, 5, 9, 10, 13, 16, 18, 20, 23, 29, 34, 43-45, 47, 61) The “ICH to CHS proportion” has higher after CAS in comparison to CES: 7 out of 11 CAS studies (63.6%) had 50% or more hemorrhagic CHS (range 0-100%).(2, 5, 9, 10, 13, 16, 18, 20, 23, 29, 34, 43-45, 47, 61) In CEA, 5 out of 12 studies (41.6%) had 50% or more cases of hemorrhagic CHS (range 0-80%).(2, 5, 9, 10, 13, 16, 18, 20, 23, 29, 34, 43-45, 47, 61) Post procedure ICH in asymptomatic patients was addressed in 9 large studies (1, 7, 10, 13, 47, 49, 55, 57, 61) and occurred in three.(47, 57, 61) Overall, periprocedural hypertension is the most frequent risk factor, being documented in 4 studies.(16, 20, 47, 50) Three studies mention ipsilateral severe stenosis as risk factor.(50, 51, 63) Younger age was considered a risk factor in 2 studies but denied as a risk factor in other 2 studies.(16, 47, 51, 63)

### **3.4-Outcomes**

The mortality of ICH related to CHS ranged from 0% to 100%.(1, 8, 9, 16, 20, 21, 23, 30, 39, 40, 42, 45, 47, 49, 51, 55, 57, 61, 63-65). In large studies the mortality was  $\geq 50\%$  in more than half of the studies (range 0 to 100%).(1, 10, 17, 21, 24, 31, 39, 42, 45, 47, 49, 51, 55, 57, 61, 63, 65) The pooled case-fatality of ICH after CHS was 51% (95% CI: 32% to 71%,  $I^2=77\%$ ) (Figure 3).

### **4-Discussion**

This is the first systematic review and meta-analysis addressing the frequency, risk factors and outcome of ICH in the context of CHS. Despite being discussed since 1981(54), no consensual definition exists for CHS and its pathophysiology is still to be elucidated.(28, 60) We found a high variation in the frequency of ICH in the context of CHS, with a range of 0% to 4.44%.(1, 2, 4, 5, 7-10, 13, 14, 16-24, 29, 30, 34, 39, 40, 42-52, 55, 57, 61, 63-65) The overall case-fatality associated to ICH in the context of CHS was high.(1, 8, 10, 17, 21, 22, 24, 31, 39, 40, 42, 45, 47, 49, 51, 55, 57, 61, 63-65) Of notice, the two larger studies reported associated mortality varying from 25.93% to 57.14%(16, 47). Important variation exists regarding mortality rates when considering all studies information (0 to 100%). However, this may be explained by the inclusion of

different studies designs, samples sizes and the classification criteria used for ICH in the context of CHS . In large studies, the frequency of ICH was higher after CAS in comparison to CEA and was rare after asymptomatic carotid disease. The higher “ICH to CHS proportion” post CAS CHS further supports the notion that patients with CHS after CAS are at increased risk of ICH.(38) The mandatory use of double antiplatelet therapy in CAS could contribute to this finding. Indeed, the use of antithrombotics was associated with the occurrence of post CAS ICH in two small studies and in one large study based on administrative data.(38) Periprocedural hypertension and ipsilateral severe stenosis were the commonest risk factors described.(16, 20, 47, 50, 51, 63) This data is relevant and stresses the importance of pre and post carotid revascularization blood pressure control, particularly in patients with severe stenosis. One frequent bias found in the included studies was the lack of definition for CHS with ICH. However, the requirement of brain imaging for ICH diagnosis may minimize the impact of this bias for the overall comparison. The use of different methodologies and size discrepancies between the studies can also explain the variation in the frequency of ICH in the context of CHS.(1, 2, 4, 5, 7-10, 13, 14, 16-24, 29, 30, 34, 39, 40, 42-52, 55, 57, 61, 63-65) The lack of information regarding associated factors such as use of antithrombotics, time interval from ischemic event to revascularization procedure, and presence of chronic white matter alterations represent a limitation when evaluating the occurrence of ICH in the context of CHS.

## **5-Conclusion**

This systematic review and meta-analysis showed that ICH in the context of CHS is rare in large series, occurs more frequently after CHS secondary to CAS and than post CAS, and is generally associated with high case-fatality rate. The main risk factors are periprocedural hypertension and ipsilateral severe stenosis. Further studies to better describe the contribution of other risk factors are needed.

**Funding** - No funding was received for this research

**Conflict of interest:** FBR is supported by CHDI Foundation and by the EHDN. All authors certify that they are no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers'

bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

This article does not contain any studies with human participants performed by any of the authors.

## **8-References**

1. Abou-Chebl A, MD; Yadav, J.S., M.D; Reginelli, J.P., M.D; Bajzer, C., M.D.; Bhatt, D, M.D.; Krieger, D.W., MD (2004) Intracranial Hemorrhage and Hyperperfusion Syndrome Following Carotid Artery Stenting. *Journal of the American College of Cardiology* 43:6
2. Ascher EM, N.; Schutzer, R.W.; Kallakuri, S.; Jacob, T.; Hingorani, A.P.; (2003) Cerebral hyperperfusion syndrome after carotid endarterectomy: Predictive factors and hemodynamic changes. *Journal of Vascular Surgery* 37:9
3. Benzel ECH, K.D. (1991) Factors associated with postoperative hypertension complicating carotid endarterectomy. *Acta Neurochirurgica* 112:5
4. Bonaldi GA, L.; Baruzzi, F; Biroli, F.; Facchinettu, A.; Fachinetti, P.; Lunghi, A; Terraneo, F. (2005) ANGIOPLASTY AND STENTING OF THE CERVICAL CAROTID BIFURCATION UNDER FILTER PROTECTION: A PROSPECTIVE STUDY IN A SERIES OF 53 PATIENTS. *Journal of neuroradiology*:9
5. Borst GJM, F.L.; Van de Pavoordt, H.D.W.M.; Mauser, H.W.; Kelder, J.C.; Ackerstaff, R.G.A. (2001) Stroke From Carotid Endarterectomy: When and How to Reduce Perioperative Stroke Rate? *European journal of Vascular and Endovascular Surgery*:6
6. Bradac OM, M.; Kramar, F.; Netuka, D.; Ostry, S.; Charvat, F.; Lacman, J.; Benes, V. (2014) Carotid endarterectomy and carotid artery stenting: changing paradigm during 10 years in a high-volume centre. *Acta Neurochirurgica* 156:8



7. Brantley HPK, J.L.; Hugh, M.B.; Mendelsohn, F.O. (2009) Hyperperfusion syndrome following carotid artery stenting: the largest single-operator series to date. *Journal of Invasive Cardiology*:4
8. Buczek JK, M.; Kobayashi, A.; Bialek, P.; Czlonkowska, A. (2013) Hyperperfusion syndrome after carotid endarterectomy and carotid stenting. *Cerebrovascular Diseases*:7
9. Coutts SBH, M.D.; Hu, W.; Sutherland, G.R.; (2003) Hyperperfusion syndrome: toward a stricter definition. *Neurosurgery*:8
10. Dalman JEB, I.C.M.; Moll, F.L.; Leusink, J.A.; Ackerstaff (1999) Transcranial doppler monitoring during carotid endarterectomy helps to identify patients at risk of postoperative hyperperfusion. *European Journal of Vascular and Endovascular Surgery*:8
11. Deruty RM, C.; Pelissou-Guyotat, I.; Lapras, C. (1991) The Carotid Endarterectomy: Experience with 260 cases and discussion of the indications. *Acta Neurochirurgica* 112:7
12. Go ASM, D.; Roger, V.L.; Benjamin, E.J.; Berry, J.D.; Blaha, M.J. (2014) Heart disease and stroke statistics e 2014 update: a report from the American Heart Association. *Circulation* 129:28
13. Gosseti BM, O.; Guericchio, R.; Irace, L.; Benedetti-Valentini, F. (1997) Transcranial doppler in 178 patients before, during and after carotid endarterectomy. *Journal of Neuroimaging*:4
14. Gupta AKP, S.; Unnikrishnan, M.; Vattoth, S.; Krishnamoorthy, T.; Kesavadas, C. (2005) HYPERPERFUSION SYNDROME AFTER SUPRAAORTIC VESSEL INTERVENTIONS AND BYPASS SURGERY. *Journal of Neuroradiology* 32:7
15. 15. National Institute of Health: Study quality assessment Tools [<https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-reduction/tools>] Last Updated April 2014.
16. Henderson RDP, T.G.; Piepgras, D.G.; Wijdicks, E.F.M. (2001) Mechanisms of intracerebral hemorrhage after carotid endarterectomy. *Journal of Neurosurgery* 95:6
17. Henry MG, L.; Rjagopal, S.; Rath, P.C.; Henry, I.; Hugel, M (2005) Bilateral Carotid Angioplasty and Stenting. *Catheterization and Cardiovascular Interventions*, Wiley-Liss 64:8

18. Hosoda KK, T.; Shibata, Y.; Kamei, M.; Kidoguchi, K.; Koyama, J.; Fujita, S.; Tamaki, N. (2001) Cerebral vasoreactivity and internal carotid artery flow help to identify patients at risk for hyperperfusion after carotid endarterectomy. *Stroke*:7
19. Iko MA, H.; Go, Y.; Nakai, K.; Tsutsumi, M.; Yu, I.; Mizokami, T.; Sakamoto, K.; Inoue, R.; Mitsutake, T.; Eto, A.; Hanada, H.; Kazekawa, K. (2014) Treatment outcomes of carotid artery stenting with two types of distal protection filter device. *SpringerPlus* 3:7
20. Jansen CS, A.M.; Moli, F.L.; Vermeulen, F.E.E.; Hamerlijnck, R.P.H.M.; Gijin, J.V.; Ackerstaff, R.G.A. (1994) Prediction of intracerebral hemorrhage after carotid endarterectomy by clinical criteria and intraoperative transcranial doppler monitoring: results of 233 operations. *European journal of Vascular and Endovascular Surgery*:6
21. Kablak-Ziembicka AP, T.; Pieniazek, P.; Musialek, P.; Motyl, R.; Moczulski, Z.; Tracz, W (2006) Assessment of Flow Changes in the Circle of Willis After Stenting for Severe Internal Carotid Artery Stenosis. *Journal of Endovascular Therapy* 13:10
22. Kaku YY, S.; Kokuzawa, J. (2004) Factors predictive of cerebral hyperperfusion after carotid angioplasty and stent placement. *American journal of Neuroradiology*:6
23. Karapanayiotides TM, R.; Devuyt, G.; Piechowski-Jozwiak, B.; Dewarrat, A.; Ruchat, P.; Segesser, L.V.; Bogousslavsky (2004) Postcarotid Endarterectomy Hyperperfusion or Reperfusion Syndrome. *Stroke* 36:7
24. Katano HM, M.; Sakurai, K.; Miyachi, S.; Yamada, K (2012) Reevaluation of collateral pathways as escape routes from hyperemia/hyperperfusion following surgical treatment for carotid stenosis. *Acta Neurochirurgica* 154:12
25. Kayahara TT, R.; Kikkawa, Y.; Take, Y.; Kurita, H. (2015) Hyperperfusion syndrome after aneurysm surgery: a case report. *Acta Neurochirurgica* 157:3
26. Khattar NKF, R.M.; Chaer, R.A.; Avgerinos, E.D.; Kretz, E.S.; Balzer, J.R.; Crammond, D.J.; Habeych, M.H.; Thirumala, P.D. (2016) Perioperative stroke after carotid endarterectomy: etiology and implications. *Acta Neurochirurgica* 158:7
27. Kim DYK, B.M.; Park, H.; Chung, J. (2012) Retinal hemorrhage as an initial sign of cerebral hyperperfusion syndrome after carotid stenting. *Acta Neurochirurgica* 154:3
28. Kozar SJ, M. (2014) Hyperperfusion and intracranial haemorrhage after carotid angioplasty with stenting – latest review. *Signa vitae* 9:6
29. Lai ZL, B.; Chen, Y.; Ni, L.; Liu, C. (2015) Prediction of cerebral hyperperfusion syndrome with velocity blood pressure index. *Chinese Medical Journal* 128:7

30. Li SML, D.; Ling, F.; Miao, Z.R.; Wang, M.L. (2005) Carotid Artery Stenting: Experience of a Single Institute in China. *Interventional Neuroradiology* 11:8
31. Liberati AA, D.G.; Tetzlaff, J.; Mulrow, C.; Gotzsche, P.C.; Ionannidis, J.P.A.; Clarke, M.; Devereaux, P.J.; Kleijnen, J.; Moher, D. (2009) The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *Annals of Internal Medicine* 151:30
32. Lieb MS, U.; Hines, G.L. (2012) Cerebral hyperperfusion syndrome after carotid intervention: a review *Cardiology in review*:6
33. Luengo-Fernandez RG, A.M.; Bull, L.; Cuthbertson, F.; Rothwell, P.M. (2013) Oxford Vascular Study. Quality of life after TIA and stroke: ten-year results of the oxford vascular study. *Neurology* 81:95
34. Maas MBK, C.J.; Hirsch, J.A.; Jaff, M.R.; Rordorf, G.A. (2013) Clinical risk predictors for cerebral hyperperfusion syndrome after carotid endarterectomy. *Journal of Neurology, Neurosurgery and Psychiatry* 84:5
35. Macfarlane RM, M.A.; Sakas, D.E.; Tasdemiroglu, E.; Wei, E.P.; Kontos, H.A. (1991) The role of neuroeffector mechanisms in cerebral hyperperfusion syndromes. *Journal of Neurosurgery* 1991:11
36. Magee TRD, A.H.; Baird, R.N.; Horrocks, M. (1992) Transcranial Doppler measurement before and after carotid endarterectomy. *Journal of the Royal College of Surgeons of Edinburgh* 37:2
37. McCabe DB, M.; Clifton, A. (1999) Fatal cerebral reperfusion hemorrhage after carotid stenting. *Stroke* 30:4
38. McDonald RJH, J.C.; Kallmes, D.F. (2011) Intracranial hemorrhage is much more common after carotid stenting than after endarterectomy. Evidence from the national inpatient sample. *Stroke*:6
39. Mohammadian RS, B.; Mansourizadeh, R.; Mohammadian, F.; Nasiri, B.; Haririan, S. (2012) Unprotected carotid artery stenting: complications in 6 months follow-up. *Journal of neuroradiology* 54:7
40. Morrish WG, S.; Douen, A.; Gordon, C.; Hu, W.; Farb, R.; Kalapos, P.; Wee, R.; Hudon, M.; Agbi, C.; Richard, M. (2000) Intracranial hemorrhage after stenting and angioplasty of extracranial carotid stenosis. *American journal of Neuroradiology*:6

41. Moulakakis KGM, S.N.; Sfyroeras, G.S.; Andrikopoulos V. (2009) Hyperperfusion syndrome after carotid revascularization. *Journal of Vascular Surgery* 49:9
42. Narita SA, H.; Nagata, S; Tstutsumi, M.; Kouhei, N.; Yoshida, H.; Matsumoto, Y.; Hamaguchi, S.; Etoh, H.; Sakamoto, K.; Inoue, R.; Kazekawa, K. (2013) Intraoperative Prediction of Hemorrhagic Cerebral Hyperperfusion Syndrome After Carotid Artery Stenting. *Journal of Stroke and Cerebrovascular Diseases* 22:5
43. Naylor ARE, J.; Thompson, M.M.; London, N.J.M.; Abbott, R.J.; Cherryman, G.; Bell, P.R.F. (2003) Seizures After Carotid Endarterectomy: Hyperperfusion, Dysautoregulation or Hypertensive Encephalopathy? *European journal of Vascular and Endovascular Surgery* 26:6
44. Newman JEA, M.; Sharpe, R.; Bown, M.J.; Sayers, R.D.; Naylor, A.R. (2013) Changes in Middle Cerebral Artery Velocity after Carotid Endarterectomy do not Identify Patients at High-risk of Suffering Intracranial Haemorrhage or Stroke due to Hyperperfusion Syndrome. *European Society for Vascular Surgery*:10
45. Nicholas GGH, H.; Gee, W.; Reed, J.F. (1993) The cerebral hyperperfusion syndrome: Diagnostic value of ocular pneumoplethysmography. *Journal of Vascular Surgery* 17:6
46. Ogasawara KI, T.; Kobayashi, M.; Endo, H.; Yoshida, K.; Fukuda, T.; Terasaki, K.; Ogawa, A. (2005) Cerebral hyperperfusion following carotid endarterectomy: diagnostic utility of intraoperative transcranial doppler ultrasonography compared with single-photon emission computed tomography study. *American journal of Neuroradiology*:6
47. Ogasawara KMD, Ph.D.; Sakai, N. M.D., Ph.D.; Kuroiwa, T, M.D., Ph.D.; Hosoda, K, M.D., Ph.D.; Iihara MD, Ph.D.; Toyoda, K, M.D., Ph.D.; Sakai, Chiaki, M.D., Ph.D.; Nagata, I., M.D., Ph.D.; Ogawa, A., M.D., Ph.D. (2007) Intracranial hemorrhage associated with cerebral hyperperfusion syndrome following carotid endarterectomy and carotid artery stenting: retrospective review of 4494 patients. *Journal of Neurosurgery*:7
48. Ogasawara. K.; Yukawa HK, M., et al. (2003) Prediction and monitoring of cerebral hyperperfusion after carotid endarterectomy by using single-photon emission computerized tomography scanning. *Journal of Neurosurgery* 99:7
49. Pieniazek PT, L.; Musialek, P.; Kablak-Ziembicka, A.; Przewlocki, T.; Motyl, R.; Dzierwa, K.; Paluszek, P.; Hlawaty, M.; Zmudka, K.; Podolec, P. (2012) Carotid artery

stenting according to the 'tailored-CAS' algorithm is associated with a low complication rate at 30 days: data from the TARGET-CAS study. *Kardiologia Polska* 70:9

50. Piepgras DM, M.K.; Sundt, T.; Yanagihara, T.; Mussman, L.M. (1988) Intracerebral hemorrhage after carotid endarterectomy. *Journal of Neurosurgery* 68:5

51. Schroeder TS, H.; Boesen, J.; Laursen, H.; Sorensen, P.S. (1987) Intracerebral hemorrhage after carotid endarterectomy. *European journal of Vascular and Endovascular Surgery*:10

52. Solomon RAL, M.; Quest, D.Q.; Correll, J. (1986) Incidence and etiology of intracerebral hemorrhage following carotid endarterectomy. *Journal of Neurosurgery*:6

53. Sprigg NS, J.; Fox, L.; Berge, E.; Whynes, D.; Bath, P.M. (2013) Efficacy of Nitric Oxide in Stroke Investigators. Very low quality of life after acute stroke: data from the efficacy of nitric oxide in stroke trial. *Stroke* 44:62

54. Sundt TMJS, F.W.; Piepgras, D.G.; Kearns, T.P.; Messick, J.M. Jr; O'Fallon, W.M. (1981) Correlation of cerebral blood flow and electroencephalographic changes during carotid endarterectomy: with results of surgery and hemodynamics of cerebral ischemia. *Mayo Clinic Proceedings* 56:11

55. Tan GS-SP, C.C. (2009) Cerebral hyperperfusion syndrome post-carotid artery stenting. *Journal of Medical Imaging and Radiation Oncology* 53:7

56. Terada TT, M.; Matsumoto, H.; Masuo, O.; Tsumoto, T.; Yamaga, H.; Ohura, Y.; Itakura, T.; (2006) HEMORRHAGIC COMPLICATIONS AFTER ENDOVASCULAR THERAPY FOR ATHEROSCLEROTIC INTRACRANIAL ARTERIAL STENOSES. *Neurosurgery* 59:9

57. Tietke MWKK, T.; Alfke, K.; Riedel, C.; Rohr, A.; Jensen, U.; Zimmermann, P.; Stingele, R.; Jansen, O. (2009) Complication rate in unprotected carotid artery stenting with closed-cell stents. *Neuroradiology*:8

58. Timmers HJLMW, W.; Karemaker, J.M.; Lenders, J.W.M. (2004) Baroreflex failure: a neglected type of secondary hypertension. *Netherlands journal of Medicine* 62:5

59. Urbach DR (2005) Measuring quality of life after surgery. *Surgical Innovation* 12:5

60. van Mook WNKAR, R.J.M.W.; Schurink, G.W.; van Oostenbrugge, R.J.; Mess, W.H.; Hofman, P.A.M.; de Leeuw, P.W. (2005) Cerebral hyperperfusion syndrome. *Lancet Neurology* 4:12

61. Wagner WHC, D.V.; Farber, A.; Levin, P.M.; Cohen, J.L. (2005) Hyperperfusion syndrome after carotid endarterectomy. *Annals of vascular surgery*:8

62. Wallaert JBN, B.W.; Stone, D.H.; Powell, R.J.; Brown, J.R.; Cronenwett, J.L.; Googney, P.P.; (2016) Physician specialty and variation in carotid revascularization technique selected for Medicare patients. *Journal of Vascular Surgery* 63:9
63. Wilson PVA, A.D. (2005) The incidence of Ischaemic stroke versus intracerebral hemorrhage after carotid endarterectomy: a review of 2452 cases. *Annals of vascular surgery*:4
64. Wu Y-MW, H-F.; Chen, Y-L.; Wong, M-C.; Toh, C-H. (2011) Carotid stenting of asymptomatic and symptomatic carotid artery stenoses with and without the use of a distal embolic protection device. *Acta cardiologica* 66:6
65. yoshie TU, T.; Takada, T.; Nogoshi, S.; Fukano, T.; Hasegawa, Y. (2016) Prediction of cerebral hyperperfusion syndrome after carotid artery stenting by CT perfusion imaging with acetazolamide challenge. *Neuroradiology* 58:7
66. Zhu CZN, J.W. (1990) Role of carotid stenosis in ischemic stroke. *Stroke* 21:4