

Chromatic Roots are Dense in the Whole Complex Plane

ALAN D. SOKAL[†]

Department of Physics, New York University, 4 Washington Place, New York, NY 10003, USA
(e-mail: sokal@nyu.edu)

Received 29 December 2000; revised 12 August 2003

I show that the zeros of the chromatic polynomials $P_G(q)$ for the generalized theta graphs $\Theta^{(s,p)}$ are, taken together, dense in the whole complex plane with the possible exception of the disc $|q - 1| < 1$. The same holds for their dichromatic polynomials (alias Tutte polynomials, alias Potts-model partition functions) $Z_G(q, v)$ outside the disc $|q + v| < |v|$. An immediate corollary is that the chromatic roots of not-necessarily-planar graphs are dense in the whole complex plane. The main technical tool in the proof of these results is the Beraha–Kahane–Weiss theorem on the limit sets of zeros for certain sequences of analytic functions, for which I give a new and simpler proof.

1. Introduction

1.1. (Di)chromatic polynomials and the Potts model

The polynomials studied in this paper arise independently in graph theory and in statistical mechanics. It is appropriate, therefore, to begin by explaining each of these contexts. Specialists in these fields are warned that they will find at least one (and perhaps both) of these summaries excruciatingly boring; they can skip them.

Let $G = (V, E)$ be a finite undirected graph¹ with vertex set V and edge set E . For each positive integer q , let $P_G(q)$ be the number of ways that the vertices of G can be assigned ‘colours’ from the set $\{1, 2, \dots, q\}$ in such a way that adjacent vertices always receive different colours. It is not hard to show (see below) that $P_G(q)$ is the restriction to \mathbb{Z}_+ of a polynomial in q . This (obviously unique) polynomial is called the *chromatic*

[†] This research was supported in part by US National Science Foundation grants PHY-9520978, PHY-9900769 and PHY-0099393. Some of the work took place during a Visiting Fellowship at All Souls College, Oxford, where it was supported in part by Engineering and Physical Sciences Research Council grant GR/M 71626 and aided by the warm hospitality of John Cardy and the Department of Theoretical Physics.

¹ In this paper a ‘graph’ is allowed to have loops and/or multiple edges unless explicitly stated otherwise.

polynomial of G , and can be taken as the *definition* of $P_G(q)$ for arbitrary real or complex values of q .²

The chromatic polynomial was introduced in 1912 by Birkhoff [16]. The original hope was that study of the real or complex zeros of $P_G(q)$ might lead to an analytic proof of the Four-Colour Conjecture [74, 91], which states that $P_G(4) > 0$ for all loopless planar graphs G . To date this hope has not been realized, although combinatoric proofs of the Four-Colour Theorem have been found [1, 2, 3, 88, 114]. Even so, the zeros of $P_G(q)$ are interesting in their own right and have been extensively studied. Most of the available theorems concern real zeros [17, 120, 133, 134, 135, 57, 136, 115, 42, 116], but there have been some theorems on complex zeros [12, 8, 9, 128, 21, 22, 23, 24, 25, 111, 112, 26] as well as numerical studies and computations for special families [54, 10, 12, 45, 4, 5, 83, 100, 101, 102, 103, 89, 104, 90, 117, 105, 106, 107, 108, 109, 97, 99, 15, 30, 31, 32, 33, 34, 35, 36, 37, 38, 13, 14, 110, 98, 92, 59, 60].

A more general polynomial can be obtained as follows. Assign to each edge $e \in E$ a real or complex weight v_e . Then define

$$Z_G(q, \{v_e\}) = \sum_{\sigma} \prod_{e \in E} [1 + v_e \delta(\sigma(x_1(e)), \sigma(x_2(e)))], \tag{1.1}$$

where the sum runs over all maps $\sigma: V \rightarrow \{1, 2, \dots, q\}$, the δ is the Kronecker delta, and $x_1(e), x_2(e) \in V$ are the two endpoints of the edge e (in arbitrary order). It is not hard to show (see below) that $Z_G(q, \{v_e\})$ is the restriction to $q \in \mathbb{Z}_+$ of a polynomial in q and $\{v_e\}$. If we take $v_e = -1$ for all e , this reduces to the chromatic polynomial. If we take $v_e = v$ for all e , this defines a two-variable polynomial $Z_G(q, v)$ that was introduced implicitly by Whitney [130, 131, 132] and explicitly by Tutte [118, 119]; it is known variously (modulo trivial changes of variable) as the *dichromatic polynomial*, the *dichromate*, the *Whitney rank function* or the *Tutte polynomial* [129, 11].³

In statistical mechanics, (1.1) is known as the partition function of the q -state Potts model. In the Potts model [80, 138, 139], an ‘atom’ (or ‘spin’) at site $x \in V$ can exist in any one of q different states (where q is an integer ≥ 1). The *energy* of a configuration is the sum, over all edges $e \in E$, of 0 if the spins at the two endpoints of that edge are unequal and $-J_e$ if they are equal. The *Boltzmann weight* of a configuration is then $e^{-\beta H}$, where H is the energy of the configuration and $\beta \geq 0$ is the inverse temperature. The *partition function* is the sum, over all configurations, of their Boltzmann weights. Clearly this is just a rephrasing of (1.1), with $v_e = e^{\beta J_e} - 1$. A coupling J_e (or v_e) is called *ferromagnetic* if $J_e \geq 0$ ($v_e \geq 0$) and *antiferromagnetic* if $-\infty \leq J_e \leq 0$ ($-1 \leq v_e \leq 0$).

To see that $Z_G(q, \{v_e\})$ is indeed a polynomial in its arguments (with coefficients that are in fact 0 or 1), we proceed as follows. In (1.1), expand out the product over $e \in E$,

² See [81, 84] for excellent reviews on chromatic polynomials, and [39] for an extensive bibliography.

³ The Tutte polynomial $T_G(x, y)$ is conventionally defined as [129, p. 45] [11, pp. 73, 101]

$$T_G(x, y) = \sum_{E' \subseteq E} (x - 1)^{k(E') - k(E)} (y - 1)^{|E'| + k(E') - |V|},$$

where $k(E')$ is the number of connected components in the subgraph (V, E') . Comparison with (1.2) below yields

$$T_G(x, y) = (x - 1)^{-k(E)} (y - 1)^{-|V|} Z_G((x - 1)(y - 1), y - 1).$$

and let $E' \subseteq E$ be the set of edges for which the term $v_e \delta(\sigma(x_1(e)), \sigma(x_2(e)))$ is taken. Now perform the sum over configurations σ : in each connected component of the subgraph (V, E') the spin value $\sigma(x)$ must be constant, and there are no other constraints. Therefore,

$$Z_G(q, \{v_e\}) = \sum_{E' \subseteq E} q^{k(E')} \prod_{e \in E'} v_e, \tag{1.2}$$

where $k(E')$ is the number of connected components (including isolated vertices) in the subgraph (V, E') . The expansion (1.2) was discovered by Birkhoff [16] and Whitney [130] for the special case $v_e = -1$ (see also Tutte [118, 119]); in its general form it is due to Fortuin and Kasteleyn [62, 46] (see also [43]). We take (1.2) as the *definition* of $Z_G(q, \{v_e\})$ for arbitrary complex q and $\{v_e\}$.

In statistical mechanics, a very important role is played by the complex zeros of the partition function. This arises as follows [141]. Statistical physicists are interested in *phase transitions*, namely in points where one or more physical quantities (e.g., the energy or the magnetization) depend nonanalytically (in many cases even discontinuously) on one or more control parameters (e.g., the temperature or the magnetic field). Now, such nonanalyticity is manifestly impossible in (1.1)/(1.2) for any finite graph G . Rather, phase transitions arise only in the *infinite-volume limit*. That is, we consider some countably infinite graph $G_\infty = (V_\infty, E_\infty)$ – usually a regular lattice, such as \mathbb{Z}^d with nearest-neighbour edges – and an increasing sequence of finite subgraphs $G_n = (V_n, E_n)$. It can then be shown (under modest hypotheses on the G_n) that the (*limiting*) *free energy per unit volume*

$$f_{G_\infty}(q, v) = \lim_{n \rightarrow \infty} |V_n|^{-1} \log Z_{G_n}(q, v) \tag{1.3}$$

exists for all *nondegenerate physical* values of the parameters,⁴ namely either

- (a) q integer ≥ 1 and $-1 < v < \infty$ (using (1.1): see, e.g., [55, Section I.2]), or
- (b) q real > 0 and $0 \leq v < \infty$ (using (1.2): see [53, Theorem 4.1] and [52, 96]).

This limit $f_{G_\infty}(q, v)$ is in general a continuous function of v ; but it can fail to be a real-analytic function of v , because complex singularities of $\log Z_{G_n}(q, v)$ – namely, complex zeros of $Z_{G_n}(q, v)$ – can approach the real axis in the limit $n \rightarrow \infty$. Therefore, the possible points of physical phase transitions are precisely the real limit points of such complex zeros (see Theorem 3.1). As a result, theorems that constrain the possible location of complex zeros of the partition function are of great interest. In particular, theorems guaranteeing that a certain complex domain is free of zeros are often known as *Lee–Yang theorems*.⁵

In summary, both graph theorists and statistical mechanics are interested in the zeros of $Z_G(q, v)$. Graph theorists most often fix $v = -1$ and look for zeros in the complex q -plane, while statistical mechanics most often fix q real > 0 (usually but not always

⁴ Here ‘physical’ means that the weights are nonnegative, so that the model has a probabilistic interpretation; and ‘nondegenerate’ means that we exclude the limiting cases $v = -1$ in (a) and $q = 0$ in (b), which cause difficulties due to the existence of configurations having zero weight.

⁵ The first such theorem, concerning the behaviour of the ferromagnetic Ising model at complex magnetic field, was proved by Lee and Yang [64] in 1952. A partial bibliography (up to 1980) of generalizations of this result can be found in [65].

an integer) and look for zeros in the complex v -plane. But these inclinations are not hard-and-fast; both groups have seen the value of investigating the more general two-variable problem. The analysis in this paper will, in fact, be an illustration of the value of doing so, even if one is ultimately interested in a specific one-variable specialization of $Z_G(q, v)$.

1.2. A Lee–Yang theorem for chromatic polynomials?

Let me now review some known facts about the *real* zeros of the chromatic polynomial $P_G(q)$, in order to motivate some conjectures concerning the *complex* zeros.

(1) It is not hard to show that for any loopless graph G with n vertices, $(-1)^n P_G(q) > 0$ for real $q < 0$ [84]. It is then natural to ask whether the absence of negative real zeros might be the tip of the iceberg of a Lee–Yang theorem: that is, might there exist a *complex* domain D containing $(-\infty, 0)$ that is zero-free for all P_G ? One’s first guess is that the half-plane $\operatorname{Re} q < 0$ might be zero-free [45]. This turns out to be false: for about a decade, examples have been known of loopless graphs G that have chromatic roots with slightly negative real part [5, 83, 104, 107, 110, 24, 25, 92]; and examples have very recently been constructed with arbitrarily negative real part [105, equation (3.12)]. Nevertheless, it is not ruled out that some smaller domain $D \supset (-\infty, 0)$ might be zero-free.

(2) For any loopless *planar* graph G , Birkhoff and Lewis [17] proved in 1946 that $P_G(q) > 0$ for real $q \geq 5$;⁶ we now know that $P_G(4) > 0$ [3, 88]; and it is very likely true (though not yet proved as far as I know) that $P_G(q) > 0$ also for $4 < q < 5$. Thus, it is natural to conjecture that there might exist a *complex* domain D containing $(4, \infty)$ (or $(5, \infty)$) that is zero-free for all planar P_G . One’s first guess might be that $\operatorname{Re} q > 4$ works. This again turns out to be false: examples are known of loopless planar graphs G that have chromatic roots with real part as large as ≈ 4.2 [5, 60]. Nevertheless, it is not ruled out that some smaller domain $D \supset (4, \infty)$ might be zero-free.

As with most of my conjectures, these two are false; but what is interesting is that they are *utterly, spectacularly false*, for I can prove:

Theorem 1.1. *There is a countably infinite family of planar (in fact, series-parallel) graphs whose chromatic roots are, taken together, dense in the entire complex q -plane with the possible exception of the disc $|q - 1| < 1$.*

As far as I know, it was until now an open question whether the closure of the set of all chromatic roots (of all graphs taken together) even has nonzero two-dimensional Lebesgue measure. Theorem 1.1 answers this question in a most spectacular way.

The graphs arising in Theorem 1.1 are ‘generalized theta graphs’ $\Theta^{(s,p)}$ obtained by parallel-connecting p chains each of which has s edges in series.⁷ Theorem 1.1 is in fact a corollary of the following more general result for the two-variable polynomials $Z_G(q, v)$.

⁶ See also [136, Theorem 1] and [115, Theorem 3.1 ff.] for alternative proofs of a more general result.

⁷ More generally, the generalized theta graph Θ_{s_1, \dots, s_p} consists of end-vertices x, y connected by p internally disjoint paths of lengths $s_1, \dots, s_p \geq 1$ [26]. The graphs arising in Theorem 1.1 thus correspond to the special case $s_1 = \dots = s_p = s$. See Section 2.3 below for a computation of the Potts-model partition function for an arbitrary Θ_{s_1, \dots, s_p} .

Theorem 1.2. Fix complex numbers q_0, v_0 satisfying $|v_0| \leq |q_0 + v_0|$. Then, for each $\epsilon > 0$, there exist $s_0 \in \mathbb{N}$ and a map $p_0 : \mathbb{N} \cap [s_0, \infty) \rightarrow \mathbb{N}$ such that, for all $s \geq s_0$ and $p \geq p_0(s)$,

- (a) if $v_0 \neq 0$, then $Z_{\Theta^{(s,p)}}(\cdot, v_0)$ has a zero in the disc $|q - q_0| < \epsilon$,
- (b) $Z_{\Theta^{(s,p)}}(q_0, \cdot)$ has a zero in the disc $|v - v_0| < \epsilon$.

(Setting $v_0 = -1$, Theorem 1.2(a) implies Theorem 1.1.)

As Jason Brown pointed out to me (why didn't I notice it myself?), an immediate corollary of Theorem 1.1 is as follows.

Corollary 1.3. There is a countably infinite family of (not-necessarily-planar) graphs whose chromatic roots are, taken together, dense in the entire complex q -plane.

Indeed, it suffices to consider the union of the two families $\Theta^{(s,p)}$ and $\Theta^{(s,p)} + K_2$ (the latter is the join of $\Theta^{(s,p)}$ with the complete graph on two vertices) and to recall that $P_{G+K_n}(q) = q(q-1) \cdots (q-n+1)P_G(q-n)$.

It is an open question whether the chromatic roots of planar (or perhaps even series-parallel) graphs are dense in the disc $|q - 1| < 1$. I have some partial results on this question, but since they are not yet definitive, I shall report them elsewhere.

The methods of this paper actually prove a result stronger than Theorem 1.2, namely:

Theorem 1.4.

- (a) Fix a complex number $v_0 \neq 0$. Then, for each $\epsilon > 0$ and $R < \infty$, there exist $s_0 \in \mathbb{N}$ and a map $p_0 : \mathbb{N} \cap [s_0, \infty) \rightarrow \mathbb{N}$ such that for all $s \geq s_0$ and $p \geq p_0(s)$, the zeros of $Z_{\Theta^{(s,p)}}(\cdot, v_0)$ come within ϵ of every point in the region $\{q \in \mathbb{C} : |q + v_0| \geq |v_0| \text{ and } |q| \leq R\}$.
- (b) Fix a complex number q_0 . Then, for each $\epsilon > 0$ and $R < \infty$, there exist $s_0 \in \mathbb{N}$ and a map $p_0 : \mathbb{N} \cap [s_0, \infty) \rightarrow \mathbb{N}$ such that, for all $s \geq s_0$ and $p \geq p_0(s)$, the zeros of $Z_{\Theta^{(s,p)}}(q_0, \cdot)$ come within ϵ of every point in the region $\{v \in \mathbb{C} : |v| \leq |q_0 + v| \text{ and } |v| \leq R\}$.

I thank Roberto Fernández for posing the question of whether something like Theorem 1.4 might be true.

1.3. Sketch of the proof

The intuition behind Theorem 1.2 is based on recalling the rules for parallel and series combination of Potts edges (see Section 2 for details):

$$\begin{aligned} \text{Parallel: } v_{\text{eff}} &= v_1 + v_2 + v_1 v_2 && \text{(mnemonic: } 1 + v \text{ multiplies),} \\ \text{Series: } v_{\text{eff}} &= v_1 v_2 / (q + v_1 + v_2) && \text{(mnemonic: } v / (q + v) \text{ multiplies).} \end{aligned}$$

In particular, if $0 < |v / (q + v)| < 1$, then putting a large number s of edges in series drives the effective coupling v_{eff} to a small (but nonzero) number; moreover, by small perturbations of v and/or q we can give v_{eff} any phase we please. But then, by putting a large number p of such chains in parallel, we can make the resulting v_{eff} lie anywhere in the complex plane we please. In particular, we can make v_{eff} equal to $-q$, which causes the partition function $Z_{\Theta^{(s,p)}}$ to be zero.

To convert this intuition into a proof, I employ a complex-variables result due to Beraha, Kahane and Weiss [6, 7, 8, 9], which I slightly generalize as follows. Let D be a domain (connected open set) in \mathbb{C} , and let $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_m$ ($m \geq 2$) be analytic functions on D , none of which is identically zero. For each integer $n \geq 0$, define

$$f_n(z) = \sum_{k=1}^m \alpha_k(z) \beta_k(z)^n. \tag{1.4}$$

We are interested in the zero sets

$$\mathcal{Z}(f_n) = \{z \in D : f_n(z) = 0\} \tag{1.5}$$

and in particular in their limit sets as $n \rightarrow \infty$:

$$\liminf \mathcal{Z}(f_n) = \{z \in D : \text{every neighbourhood } U \ni z \text{ has a nonempty intersection with all but finitely many of the sets } \mathcal{Z}(f_n)\}, \tag{1.6}$$

$$\limsup \mathcal{Z}(f_n) = \{z \in D : \text{every neighbourhood } U \ni z \text{ has a nonempty intersection with infinitely many of the sets } \mathcal{Z}(f_n)\}. \tag{1.7}$$

Let us call an index k *dominant at z* if $|\beta_k(z)| \geq |\beta_l(z)|$ for all l ($1 \leq l \leq m$); and let us write

$$D_k = \{z \in D : k \text{ is dominant at } z\}. \tag{1.8}$$

Then the limiting zero sets can be completely characterized as follows.

Theorem 1.5. *Let D be a domain in \mathbb{C} , and let $\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_m$ ($m \geq 2$) be analytic functions on D , none of which is identically zero. Let us further assume a ‘no-degenerate-dominance’ condition: there do not exist indices $k \neq k'$ such that $\beta_k \equiv \omega \beta_{k'}$ for some constant ω with $|\omega| = 1$ and such that $D_k (= D_{k'})$ has nonempty interior. For each integer $n \geq 0$, define f_n by*

$$f_n(z) = \sum_{k=1}^m \alpha_k(z) \beta_k(z)^n.$$

Then $\liminf \mathcal{Z}(f_n) = \limsup \mathcal{Z}(f_n)$, and a point z lies in this set if and only if either

- (a) *there is a unique dominant index k at z , and $\alpha_k(z) = 0$, or*
- (b) *there are two or more dominant indices at z .*

Note that case (a) consists of isolated points in D , while case (b) consists of curves (plus possibly isolated points where all the β_k vanish simultaneously). Beraha–Kahane–Weiss considered the special case of Theorem 1.5 in which the f_n are polynomials satisfying a linear finite-order recurrence relation, and they assumed a slightly stronger nondegeneracy condition. Their theorem is all we really need to prove Theorems 1.2 and 1.4, but the general result is more natural and its proof is no more difficult. Indeed, my proof (see Section 3) is quite a bit simpler than the original proof of Beraha–Kahane–Weiss [7] (though also less powerful in that it gives no information on the *rate* of convergence of the zeros of f_n to their limiting set).

The next step is to notice that the dichromatic polynomial for the generalized theta graph $\Theta^{(s,p)}$ has precisely the form (1.4) with $m = 2$:

$$Z_{\Theta^{(s,p)}}(q, v) = \frac{[(q + v)^s + (q - 1)v^s]^p + (q - 1)[(q + v)^s - v^s]^p}{q^{p-1}} \tag{1.9}$$

(see Section 2 for the easy calculation). It follows from Theorem 1.5 that when $p \rightarrow \infty$ at fixed s , the zeros of $Z_{\Theta^{(s,p)}}$ accumulate where⁸

$$\left| 1 + (q - 1) \left(\frac{v}{q + v} \right)^s \right| = \left| 1 - \left(\frac{v}{q + v} \right)^s \right|. \tag{1.10}$$

I then use the following lemma to handle the limit $s \rightarrow \infty$.

Lemma 1.6. *Let F_1, F_2, G be analytic functions on a disc $|z| < R$ satisfying $|G(0)| \leq 1$ and $G \not\equiv \text{constant}$. Then, for each $\epsilon > 0$, there exists $s_0 < \infty$ such that for all integers $s \geq s_0$ the equation*

$$|1 + F_1(z)G(z)^s| = |1 + F_2(z)G(z)^s| \tag{1.11}$$

has a solution in the disc $|z| < \epsilon$.

Theorems 1.2 and 1.4 are an almost immediate consequence (see Section 5 for details).

1.4. Plan of this paper

The plan of this paper is as follows. In Section 2 I discuss some identities satisfied by the Potts-model partition function $Z_G(q, \{v_e\})$; in particular, I show what happens when a 2-rooted graph (G, x, y) is inserted in place of an edge e_* in some other graph H . As a special case, I obtain the well-known rules for series and parallel combination of Potts edges, which allow one to compute in a straightforward way the Potts-model partition function $Z_G(q, \{v_e\})$ for any series-parallel graph G . Specializing further, I derive the formula (1.9) for the dichromatic polynomial $Z_G(q, v)$ of the generalized theta graphs $\Theta^{(s,p)}$. In Section 3 I prove some theorems – which seem to me of considerable interest in their own right – concerning the limit sets of zeros for certain sequences of analytic functions. As a corollary, I obtain a simple proof of Theorem 1.5. In Section 4 I prove a strengthened version of Lemma 1.6. In Section 5 I complete the proof of Theorems 1.2 and 1.4. In Section 6 I digress to study the *real* chromatic roots of the graphs $\Theta^{(s,p)}$. In Section 7 I discuss some variants of the construction employed in this paper. In Section 8 I discuss some open questions.

The two appendices provide further examples of the power of the identities derived in Section 2. In Appendix A I give a simple proof of the Brown–Hickman [25] theorem on chromatic roots of large subdivisions. In Appendix B I extend Thomassen’s [115] construction concerning the chromatic roots of 2-degenerate graphs.

⁸ For Theorems 1.2(a) and 1.4(a), the no-degenerate-dominance condition of Theorem 1.5 is satisfied whenever $v_0 \neq 0$. For Theorems 1.2(b) and 1.4(b), the no-degenerate-dominance condition is satisfied whenever $q_0 \neq 0$; but if $q_0 = 0$, then $Z_{\Theta^{(s,p)}}(q_0, \cdot)$ is identically zero and the assertion is trivially true.

2. Some identities for Potts models

In this section I discuss some identities for Potts-model partition functions and use them to calculate $Z_G(q, \{v_e\})$ for generalized theta graphs. There are two alternative approaches to proving such identities: one is to prove the identity directly for complex q , using the Fortuin–Kasteleyn representation (1.2); the other is to prove the identity first for *positive integer* q , using the spin representation (1.1), and then to extend it to complex q by arguing that two polynomials (or rational functions) that coincide at infinitely many points must be equal. The latter approach is perhaps less elegant, but it is often simpler or more intuitive.

2.1. Restricted Potts-model partition functions for 2-rooted graphs

Let $G = (V, E)$ be a finite graph, and let x, y be distinct vertices of G . We define $G \bullet xy$ to be the graph in which x and y are contracted to a single vertex. (NB: If G contains one or more edges xy , then these edges are *not* deleted, but become loops in $G \bullet xy$.) There is a canonical one-to-one correspondence between the edges of G and the edges of $G \bullet xy$; for simplicity (though by slight abuse of notation) we denote an edge of G and the corresponding edge of $G \bullet xy$ by the same letter. In particular, we can apply a given set of edge weights $\{v_e\}_{e \in E}$ to both G and $G \bullet xy$.

Let us now define

$$Z_G^{(x \leftrightarrow y)}(q, \{v_e\}) = \sum_{\substack{E' \subseteq E \\ E' \text{ connects } x \text{ to } y}} q^{k(E')} \prod_{e \in E'} v_e, \tag{2.1}$$

$$Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}) = \sum_{\substack{E' \subseteq E \\ E' \text{ does not connect } x \text{ to } y}} q^{k(E')} \prod_{e \in E'} v_e. \tag{2.2}$$

Note that $Z_G^{(x \leftrightarrow y)}(q, \{v_e\})$ and $Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\})$ are polynomials in q and $\{v_e\}$: the lowest-order contribution to $Z_G^{(x \leftrightarrow y)}$ (resp. $Z_G^{(x \not\leftrightarrow y)}$) is of order at least q (resp. at least q^2); the highest-order contribution to $Z_G^{(x \leftrightarrow y)}$ is of order at most $q^{|V| - \text{dist}(x,y)}$, where $\text{dist}(x, y)$ is the length of the shortest path from x to y using edges having $v_e \neq 0$ (if no such path exists, then $Z_G^{(x \leftrightarrow y)}$ is identically zero); and the highest-order contribution to $Z_G^{(x \not\leftrightarrow y)}$ is of order exactly $q^{|V|}$ (with coefficient 1, coming from the term $E' = \emptyset$). From (1.2) we have trivially

$$Z_G(q, \{v_e\}) = Z_G^{(x \leftrightarrow y)}(q, \{v_e\}) + Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}) \tag{2.3}$$

and almost as trivially

$$Z_{G \bullet xy}(q, \{v_e\}) = Z_G^{(x \leftrightarrow y)}(q, \{v_e\}) + q^{-1} Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}). \tag{2.4}$$

Remark. Let $G + xy$ denote the graph G with an extra edge xy added. Then it is not hard to see that (1.2) also implies

$$Z_{G+xy}(q, \{v_e\}, v_{xy}) = (1 + v_{xy}) Z_G^{(x \leftrightarrow y)}(q, \{v_e\}) + \left(1 + \frac{v_{xy}}{q}\right) Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}), \tag{2.5}$$

which equals $Z_G + v_{xy}Z_{G \bullet xy}$ in agreement with the deletion–contraction formula. In particular, when $v_{xy} = -1$ we have

$$Z_{G+xy}(q, \{v_e\}, v_{xy} = -1) = \frac{q-1}{q} Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}), \tag{2.6}$$

which equals $Z_G - Z_{G \bullet xy}$.

Now let q be an integer ≥ 1 , and define the restricted partition function

$$Z_{G,x,y}(q, \{v_e\}; \sigma_x, \sigma_y) = \sum_{\substack{\sigma: V \rightarrow \{1, \dots, q\} \\ \sigma(x) = \sigma_x \\ \sigma(y) = \sigma_y}} \prod_{e \in E} [1 + v_e \delta(\sigma(x_1(e)), \sigma(x_2(e)))], \tag{2.7}$$

where $\sigma_x, \sigma_y \in \{1, \dots, q\}$ and the sum runs over all maps $\sigma: V \rightarrow \{1, \dots, q\}$ satisfying $\sigma(x) = \sigma_x$ and $\sigma(y) = \sigma_y$. We then have the following refinement of the Fortuin–Kasteleyn identity (1.2).

Proposition 2.1.

$$Z_{G,x,y}(q, \{v_e\}; \sigma_x, \sigma_y) = A_{G,x,y}(q, \{v_e\}) + B_{G,x,y}(q, \{v_e\}) \delta(\sigma_x, \sigma_y), \tag{2.8}$$

where

$$A_{G,x,y}(q, \{v_e\}) = q^{-2} Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}), \tag{2.9a}$$

$$B_{G,x,y}(q, \{v_e\}) = q^{-1} Z_G^{(x \leftrightarrow y)}(q, \{v_e\}) \tag{2.9b}$$

are polynomials in q and $\{v_e\}$, whose degrees in q are

$$\deg A = |V| - 2, \tag{2.10a}$$

$$\deg B \leq |V| - 1 - \text{dist}(x, y), \tag{2.10b}$$

where $\text{dist}(x, y)$ is the length of the shortest path from x to y using edges having $v_e \neq 0$ (if no such path exists, then $B = 0$). Moreover, (2.9a/b) define the unique functions $A_{G,x,y}$ and $B_{G,x,y}$ that are polynomials in q and satisfy (2.8).

First proof. In (2.7), expand out the product over $e \in E$, and let $E' \subseteq E$ be the set of edges for which the term $v_e \delta(\sigma(x_1(e)), \sigma(x_2(e)))$ is taken. Now perform the sum over configurations $\{\sigma(z)\}_{z \in V \setminus \{x,y\}}$: in each connected component of the subgraph (V, E') the spin value $\sigma(z)$ must be constant. In particular, in each component containing x and/or y , the spins must all equal the specified value σ_x and/or σ_y ; in all other components, the spin value is free. Therefore,

$$Z_{G,x,y}(q, \{v_e\}; \sigma_x, \sigma_y) = \begin{cases} q^{-2} Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}) & \text{if } \sigma_x \neq \sigma_y, \\ q^{-2} Z_G^{(x \not\leftrightarrow y)}(q, \{v_e\}) + q^{-1} Z_G^{(x \leftrightarrow y)}(q, \{v_e\}) & \text{if } \sigma_x = \sigma_y. \end{cases} \tag{2.11}$$

Finally, the numbers $A_{G,x,y}$ and $B_{G,x,y}$ in (2.8) are uniquely defined for each integer $q \geq 2$; and any polynomials that coincide with (2.9a/b) at all integers $q \geq 2$ must coincide everywhere. □

Second proof. For each integer q , the S_q permutation symmetry of the Potts model implies that

$$Z_{G,x,y}(q, \{v_e\}; \sigma_x, \sigma_y) = A + B\delta(\sigma_x, \sigma_y) \tag{2.12}$$

for some numbers A and B depending on G, x, y, q and $\{v_e\}$; moreover, the numbers A and B are obviously unique when $q \geq 2$. But then, summing (2.12) over σ_x, σ_y without and with the constraint $\sigma_x = \sigma_y$, we get

$$Z_G = q^2A + qB, \tag{2.13a}$$

$$Z_{G \bullet xy} = qA + qB. \tag{2.13b}$$

Hence

$$A = \frac{Z_G - Z_{G \bullet xy}}{q(q-1)} = q^{-2}Z_G^{(x \leftrightarrow y)}, \tag{2.14a}$$

$$B = \frac{qZ_{G \bullet xy} - Z_G}{q(q-1)} = q^{-1}Z_G^{(x \leftrightarrow y)} \tag{2.14b}$$

by virtue of (2.3) and (2.4). □

Remark. Extensions of Proposition 2.1 to k -rooted graphs (for any $k \geq 2$) can also be derived [140, 79]. I thank Fred Wu for bringing reference [140] to my attention.

Let us now consider inserting the 2-rooted graph (G, x, y) in place of an edge e_* in some other graph H , and let us call the resulting graph H^G . We can trivially rewrite (2.8)/(2.9) as

$$Z_{G,x,y}(q, \{v_e\}; \sigma_x, \sigma_y) = A_{G,x,y}(q, \{v_e\}) \left[1 + \frac{B_{G,x,y}(q, \{v_e\})}{A_{G,x,y}(q, \{v_e\})} \delta(\sigma_x, \sigma_y) \right], \tag{2.15a}$$

$$= A_{G,x,y}(q, \{v_e\}) [1 + v_{\text{eff},G,x,y}(q, \{v_e\}) \delta(\sigma_x, \sigma_y)], \tag{2.15b}$$

where

$$v_{\text{eff}} \equiv v_{\text{eff},G,x,y}(q, \{v_e\}) = \frac{B_{G,x,y}(q, \{v_e\})}{A_{G,z,y}(q, \{v_e\})} = \frac{qZ_G^{(x \leftrightarrow y)}(q, \{v_e\})}{Z_G^{(x \leftrightarrow y)}(q, \{v_e\})} \tag{2.16}$$

is a rational function of q and $\{v_e\}$. We then see that (G, x, y) acts within the graph H as a single edge with effective weight v_{eff} , provided that the partition function Z_H is multiplied by an overall prefactor $A_{G,z,y}(q, \{v_e\})$. More precisely,

$$Z_{H^G}(q, \{v_e\}_{e \in (E(H) \setminus e_*) \cup E(G)}) = A_{G,x,y}(q, \{v_e\}_{e \in E(G)}) Z_H(q, \{v_e\}_{e \in E(H) \setminus e_*}, v_{\text{eff}}), \tag{2.17}$$

where $v_{\text{eff}} \equiv v_{\text{eff},G,x,y}(q, \{v_e\}_{e \in E(G)})$ replaces v_{e_*} as an argument of Z_H . This follows from Proposition 2.1 whenever q is an integer ≥ 1 ; and the corresponding identity then holds for all q , because both sides are rational functions of q that agree at infinitely many points. It is also worth noting that the ‘transmissivity’ $t_{\text{eff}} \equiv v_{\text{eff}}/(q + v_{\text{eff}})$ is given by the simple formula

$$t_{\text{eff}} = \frac{Z_G^{(x \leftrightarrow y)}(q, \{v_e\})}{Z_G}. \tag{2.18}$$

The most general version of this construction appears to be the following [137]. Let $H = (V, E)$ be a finite undirected graph, and let \tilde{H} be a directed graph obtained by assigning an orientation to each edge of H . For each edge $e \in E$, let $G_e = (V_e, E_e, x_e, y_e)$ be a 2-rooted finite undirected graph (so that $x_e, y_e \in V_e$ with $x_e \neq y_e$) equipped with edge weights $\{v_{\tilde{e}}\}_{\tilde{e} \in E_e}$. We denote by \mathbf{G} the family $\{G_e\}_{e \in E}$, and we denote by $\tilde{H}^{\mathbf{G}}$ the undirected graph obtained from H by replacing each edge $e \in E$ with a copy of the corresponding graph G_e , attaching x_e to the tail of e and y_e to the head. Its edge set is thus $\mathbf{E} = \bigcup_{e \in E} E_e$ (disjoint union). We then have the following result.

Proposition 2.2. *Let $H = (V, E)$ and $\{G_e\}_{e \in E}$ be as above. Suppose that*

$$\begin{aligned} Z_{G_e, x_e, y_e}(q, \{v_{\tilde{e}}\}_{\tilde{e} \in E_e}; \sigma_{x_e}, \sigma_{y_e}) \\ = A_{G_e, x_e, y_e}(q, \{v_{\tilde{e}}\}_{\tilde{e} \in E_e}) + B_{G_e, x_e, y_e}(q, \{v_{\tilde{e}}\}_{\tilde{e} \in E_e}) \delta(\sigma_{x_e}, \sigma_{y_e}), \end{aligned} \tag{2.19}$$

and define

$$v_{\text{eff}, e} \equiv \frac{B_{G_e, x_e, y_e}(q, \{v_{\tilde{e}}\}_{\tilde{e} \in E_e})}{A_{G_e, x_e, y_e}(q, \{v_{\tilde{e}}\}_{\tilde{e} \in E_e})}. \tag{2.20}$$

Then

$$Z_{\tilde{H}^{\mathbf{G}}}(q, \{v_{\tilde{e}}\}_{\tilde{e} \in \mathbf{E}}) = \left(\prod_{e \in E} A_{G_e, x_e, y_e}(q, \{v_{\tilde{e}}\}_{\tilde{e} \in E_e}) \right) \times Z_H(q, \{v_{\text{eff}, e}\}_{e \in E}). \tag{2.21}$$

In particular, $Z_{\tilde{H}^{\mathbf{G}}}$ does not depend on the orientations of the edges of H .

A special case of this construction is the subdivision of edges [85, 86, 82, 25], discussed further in Appendix A.

2.2. Parallel and series connection

Important special cases of Proposition 2.1 concern parallel and series connection.

The case of parallel connection is almost trivial: Let G consist of edges e_1, \dots, e_n (with corresponding weights v_1, \dots, v_n) in parallel between the same pair of vertices x, y . Then

$$\prod_{i=1}^n [1 + v_i \delta(\sigma_x, \sigma_y)] = 1 + \left[\prod_{i=1}^n (1 + v_i) - 1 \right] \delta(\sigma_x, \sigma_y), \tag{2.22}$$

so that the parallel edges are equivalent to a single edge with effective weight

$$v_{\text{eff}} = \prod_{i=1}^n (1 + v_i) - 1 \tag{2.23}$$

or equivalently

$$1 + v_{\text{eff}} = \prod_{i=1}^n (1 + v_i). \tag{2.24}$$

That is, for parallel connection, ‘ $1 + v$ multiplies’.

The case of series connection is a bit less trivial. Let G be a path P_n consisting of edges e_1, \dots, e_n (with weights v_1, \dots, v_n) between end-vertices 0 and n . Then

$$\sum_{\sigma_1, \dots, \sigma_{n-1}=1}^q \prod_{i=1}^n [1 + v_i \delta(\sigma_{i-1}, \sigma_i)] = A + B \delta(\sigma_0, \sigma_n) \tag{2.25}$$

with

$$A = q^{-1} \left[\prod_{i=1}^n (q + v_i) - \prod_{i=1}^n v_i \right], \tag{2.26a}$$

$$B = \prod_{i=1}^n v_i. \tag{2.26b}$$

This formula can easily be proved by induction on n ; or it can alternatively be derived from (2.14) by remembering that

$$Z_G = Z_{P_n} = q \prod_{i=1}^n (q + v_i), \tag{2.27}$$

$$Z_{G \bullet 0n} = Z_{C_n} = \prod_{i=1}^n (q + v_i) + (q - 1) \prod_{i=1}^n v_i \tag{2.28}$$

(formulae that can themselves be proved by induction). In particular, the series edges are equivalent to a single edge with effective weight

$$v_{\text{eff}} \equiv \frac{B}{A} = \frac{q \prod_{i=1}^n v_i}{\prod_{i=1}^n (q + v_i) - \prod_{i=1}^n v_i} \tag{2.29}$$

or equivalently

$$\frac{v_{\text{eff}}}{q + v_{\text{eff}}} = \prod_{i=1}^n \frac{v_i}{q + v_i}. \tag{2.30}$$

That is, for series connection, ‘ $v/(q + v)$ multiplies’. But one should not forget the overall prefactor A given by (2.26).

Remarks. (1) Note that applying series reduction (2.29)/(2.30) to a chromatic polynomial (all $v_i = -1$) leads to edge weights v_{eff} that are in general unequal and $\neq -1$. This is further evidence of the value of studying the general Potts-model partition function $Z_G(q, \{v_e\})$ with not-necessarily-equal edge weights $\{v_e\}$. In particular, the series-reduction formulae allow an easy proof of the Brown–Hickman theorem on chromatic roots of large subdivisions [25]: see Appendix A.

(2) The foregoing formulae form the basis for a linear-time (*i.e.*, $O(|V| + |E|)$) algorithm for computing the Potts-model partition function $Z_G(q, \{v_e\})$ for a series-parallel graph G at any fixed set of numbers q and $\{v_e\}$, and hence a quadratic-time ($O(|V|^2 + |V||E|)$) algorithm for computing $Z_G(q, \{v_e\})$ as a polynomial in q for fixed $\{v_e\}$.⁹ (One needs, as a

⁹ This is in a computational model where the elementary arithmetic operations (+, −, ×, ÷) are assumed to take a time of order 1 irrespective of the size of their arguments.

subroutine, a linear-time algorithm for recognizing whether a graph is series-parallel and, if it is, computing a decomposition tree of series-parallel reductions: see [127, Sections 2.3 and 3.3].) This is essentially the approach used, for the special case of the chromatic polynomial, in [29]. More generally, Noble [73] has shown how to compute, in linear time, the dichromatic polynomial $Z_G(q, v)$ at any point (q, v) for graphs of bounded tree-width (the case of series-parallel graphs corresponds to tree-width 2); this algorithm presumably extends to handling the general Potts-model partition function at any point $(q, \{v_e\})$. For general graphs, by contrast – or even for general *planar* graphs of maximum degree 4 – even the problem of determining whether G is 3-colourable (*i.e.*, whether $P_G(3) \neq 0$) is NP-complete [47, p. 191].

2.3. Potts-model partition function of generalized theta graphs

Consider a generalized theta graph Θ_{s_1, \dots, s_p} consisting of end-vertices x, y connected by p internally disjoint paths of lengths $s_1, \dots, s_p \geq 1$. Label the edges e_{ij} ($1 \leq i \leq p, 1 \leq j \leq s_i$) and let v_{ij} be the corresponding weights. Applying the series-connection identity (2.25)/(2.26) on each of the p paths, and then applying the parallel-connection identity (2.22), we obtain the restricted partition function

$$Z_{\Theta_{s_1, \dots, s_p}, x, y}(q, \{v_{ij}\}; \sigma_x, \sigma_y) = A + B\delta(\sigma_x, \sigma_y) \tag{2.31}$$

with

$$A = q^{-p} \prod_{i=1}^p \left[\prod_{j=1}^{s_i} (q + v_{ij}) - \prod_{j=1}^{s_i} v_{ij} \right], \tag{2.32a}$$

$$B = q^{-p} \left\{ \prod_{i=1}^p \left[\prod_{j=1}^{s_i} (q + v_{ij}) + (q - 1) \prod_{j=1}^{s_i} v_{ij} \right] - \prod_{i=1}^p \left[\prod_{j=1}^{s_i} (q + v_{ij}) - \prod_{j=1}^{s_i} v_{ij} \right] \right\}. \tag{2.32b}$$

In particular, summing (2.31) over σ_x and σ_y without constraint, we obtain $Z_{\Theta_{s_1, \dots, s_p}} = q^2 A + qB$ and hence the following result.

Proposition 2.3. *The Potts-model partition function for the generalized theta graph Θ_{s_1, \dots, s_p} with edge weights $\{v_{ij}\}_{1 \leq i \leq p, 1 \leq j \leq s_i}$ is*

$$\begin{aligned} & Z_{\Theta_{s_1, \dots, s_p}}(q, \{v_{ij}\}) \\ &= q^{-(p-1)} \left\{ \prod_{i=1}^p \left[\prod_{j=1}^{s_i} (q + v_{ij}) + (q - 1) \prod_{j=1}^{s_i} v_{ij} \right] + (q - 1) \prod_{i=1}^p \left[\prod_{j=1}^{s_i} (q + v_{ij}) - \prod_{j=1}^{s_i} v_{ij} \right] \right\}. \end{aligned} \tag{2.33}$$

In particular, when $v_{ij} = v$ for all i, j , we have the dichromatic polynomial

$$Z_{\Theta_{s_1, \dots, s_p}}(q, v) = q^{-(p-1)} \left\{ \prod_{i=1}^p [(q + v)^{s_i} + (q - 1)v^{s_i}] + (q - 1) \prod_{i=1}^p [(q + v)^{s_i} - v^{s_i}] \right\}. \tag{2.34}$$

And when also $s_1 = \dots = s_p = s$, we have

$$Z_{\Theta_{(s,p)}}(q, v) = q^{-(p-1)} \{ [(q + v)^s + (q - 1)v^s]^p + (q - 1) [(q + v)^s - v^s]^p \}. \tag{2.35}$$

Remarks. (1) Here is an alternate derivation of (2.31)/(2.32) and (2.33): If in Θ_{s_1, \dots, s_p} we contract x to y , we obtain p cycles of lengths s_i joined at a single vertex, so that

$$\begin{aligned} Z_{\Theta_{s_1, \dots, s_p} \bullet xy}(q, \{v_{ij}\}_{1 \leq i \leq p, 1 \leq j \leq s_i}) &= q^{-(p-1)} \prod_{i=1}^p Z_{C_{s_i}}(q, \{v_{ij}\}_{1 \leq j \leq s_i}) \\ &= q^{-(p-1)} \prod_{i=1}^p \left[\prod_{j=1}^{s_i} (q + v_{ij}) + (q - 1) \prod_{j=1}^{s_i} v_{ij} \right]. \end{aligned} \tag{2.36}$$

If, on the other hand, we add to Θ_{s_1, \dots, s_p} an extra edge xy with $v_{xy} = -1$, we obtain p cycles of lengths $s_i + 1$ joined along a single $v = -1$ edge, so that

$$\begin{aligned} Z_{\Theta_{s_1, \dots, s_p} + xy}(q, \{v_{ij}\}_{1 \leq i \leq p, 1 \leq j \leq s_i}, v_{xy} = -1) &= [q(q - 1)]^{-(p-1)} \prod_{i=1}^p Z_{C_{s_i+1}}(q, \{v_{ij}\}_{1 \leq j \leq s_i}, v_{xy} = -1) \\ &= [q(q - 1)]^{-(p-1)} \prod_{i=1}^p \left[(q - 1) \prod_{j=1}^{s_i} (q + v_{ij}) - (q - 1) \prod_{j=1}^{s_i} v_{ij} \right] \\ &= q^{-(p-1)} (q - 1) \prod_{i=1}^p \left[\prod_{j=1}^{s_i} (q + v_{ij}) - \prod_{j=1}^{s_i} v_{ij} \right]. \end{aligned} \tag{2.37}$$

The addition–contraction formula $Z_G = Z_{G \bullet xy} + Z_{G + xy}(v_{xy} = -1)$ then gives (2.33); and using (2.14) we obtain (2.32).

(2) In Appendix B I use formula (2.32) to give an extension (and simple proof) of Thomassen’s [115, Theorem 3.9] construction concerning the chromatic roots of 2-degenerate graphs.

3. Limit sets of zeros for certain sequences of analytic functions

Let (f_n) be a sequence of analytic functions on a domain $D \subset \mathbb{C}$, and let (a_n) be a sequence of positive real numbers such that $(|f_n|^{a_n})$ are uniformly bounded on compact subsets of D . We are interested in the zero sets $\mathcal{Z}(f_n)$ and in particular in their limit sets as $n \rightarrow \infty$. We shall relate these sets to the existence and behaviour of the limit

$$u(z) = \lim_{n \rightarrow \infty} a_n \log |f_n(z)|. \tag{3.1}$$

We begin with a warm-up result.

Theorem 3.1. *Let D be a domain in \mathbb{C} , and let $x_0 \in D \cap \mathbb{R}$. Let (f_n) be analytic functions on D , and let (a_n) be positive real constants such that $(|f_n|^{a_n})$ are uniformly bounded on compact subsets of D . Assume further that:*

- (a) for each $x \in D \cap \mathbb{R}$, $f_n(x)$ is real and > 0 ,
- (b) for each $x \in D \cap \mathbb{R}$, $\lim_{n \rightarrow \infty} a_n \log f_n(x) \equiv u(x)$ exists and is finite,
- (c) u is not real-analytic at x_0 .

Then $\liminf \mathcal{L}(f_n) \ni x_0$: that is, for all n sufficiently large, there exist zeros z_n^* of f_n such that $\lim_{n \rightarrow \infty} z_n^* = x_0$.

Theorem 3.1 is well known to workers in mathematical statistical mechanics: it is the contrapositive of an observation going back to Yang and Lee [141] concerning the genesis of phase transitions (see, e.g., [50, Theorem 4.1, p. 51]). We give its proof later in this section.

The main result of this section, Theorem 3.2, is similar in nature to Theorem 3.1: the difference is that, by taking the logarithm only of the absolute value of $f_n(z)$, we need not worry *a priori* whether $f_n(z) \neq 0$ (since $\log 0 = -\infty$ is a legitimate value), nor need we worry about potential ambiguities in the branch of the logarithm.

Theorem 3.2. *Let D be a domain in \mathbb{C} , and let $z_0 \in D$. Let (f_n) be analytic functions on D , and let (a_n) be positive real constants such that $(|f_n|^{a_n})$ are uniformly bounded on compact subsets of D . Suppose that there does not exist a neighbourhood $U \ni z_0$ and a function v on U that is either harmonic or else identically $-\infty$ such that $\liminf_{n \rightarrow \infty} a_n \log |f_n(z)| \leq v(z) \leq \limsup_{n \rightarrow \infty} a_n \log |f_n(z)|$ for all $z \in U$. Then $z_0 \in \liminf \mathcal{L}(f_n)$: that is, for all n sufficiently large, there exist zeros z_n^* of f_n such that $\lim_{n \rightarrow \infty} z_n^* = z_0$.*

The proofs of Theorems 3.1 and 3.2 depend crucially on the concept of a normal family of analytic functions [72, 95], which we now define.

Definition 1. A family \mathcal{F} of analytic functions on a domain $D \subset \mathbb{C}$ is said to be normal (in D) if, for every sequence $(f_n) \subset \mathcal{F}$, there exists a subsequence (f_{n_k}) that converges, uniformly on compact subsets of D , either to a (finite-valued) analytic function or else to the constant function ∞ .

(Some authors omit the possibility of convergence to ∞ from the definition of normality. But the foregoing definition is more useful for our purposes.) An easy covering and diagonalization argument [95, Theorem 2.1.2] shows that normality is a local property: \mathcal{F} is normal in D if and only if, for each $z \in D$, there exists an open disc $U \ni z$ in which \mathcal{F} is normal.

The first key result concerning normal families is Montel’s (1907) theorem [95, pp. 35–36]: if the family \mathcal{F} is uniformly bounded on compact subsets of D , then it is normal. (In this case, of course, convergence to ∞ is impossible.) We will need a slight extension of this result [95, Example 2.3.9]. For each set $A \subset \mathbb{C}$, let $\mathcal{F}_{\notin A}$ be the family of analytic functions whose values avoid A :

$$\mathcal{F}_{\notin A} = \{f \text{ analytic on } D : f[D] \cap A = \emptyset\}. \tag{3.2}$$

We then have the following.

Proposition 3.3. *If $A \subset \mathbb{C}$ has nonempty interior, then $\mathcal{F}_{\notin A}$ is normal.*

Remarks. (1) Since normality is a local property, this result can be strengthened as follows: for a family \mathcal{F} to be normal, it suffices that for each $z \in D$ there exist an open disc $U \ni z$ and a set $A_U \subset \mathbb{C}$ with nonempty interior, such that $f[U] \cap A_U = \emptyset$ for all $f \in \mathcal{F}$.

(2) A result vastly stronger than Proposition 3.3 is true: Montel's *Critère fondamental* (1912) states that $\mathcal{F}_{\neq A}$ is normal as soon as A contains *at least two points*. Detailed accounts of this deep result can be found in [72, 95]. But we will not need it.

Proof of Proposition 3.3. Suppose that A contains the disc $\{w : |w - a| < \epsilon\}$. For each $f \in \mathcal{F}_{\neq A}$, define $\tilde{f}(z) = 1/[f(z) - a]$. Then the family $\tilde{\mathcal{F}} \equiv \{\tilde{f} : f \in \mathcal{F}_{\neq A}\}$ is uniformly bounded by $1/\epsilon$, hence normal by Montel's theorem. So let $(f_n) \subset \mathcal{F}_{\neq A}$, and consider the corresponding sequence $(\tilde{f}_n) \subset \tilde{\mathcal{F}}$. By normality of $\tilde{\mathcal{F}}$, there exists a subsequence (\tilde{f}_{n_k}) that converges, uniformly on compact subsets of D , to an analytic function g . But since the \tilde{f}_{n_k} are nonvanishing on D , Hurwitz's theorem [87, p. 262] tells us that g is either identically zero or else nonvanishing. It is straightforward to show that, in the former case, the corresponding subsequence (f_{n_k}) converges, uniformly on compact subsets of D , to the constant function ∞ ; and that in the latter case, (f_{n_k}) converges, uniformly on compact subsets of D , to the analytic function $f = a + 1/g$. (For each compact $K \subset D$, we have $\inf_{z \in K} |g(z)| \equiv \delta > 0$, so for all sufficiently large k we have $|\tilde{f}_{n_k}| \geq \delta/2$ everywhere on K . Now apply the uniform continuity of the function $w \mapsto 1/w$ on the set where $|w| \geq \delta/2$.) \square

We also need the following well-known lemma, which expresses the main idea underlying the Vitali–Porter theorem.

Lemma 3.4. *Let \mathcal{F} be a normal family of analytic functions on a domain $D \subset \mathbb{C}$, and suppose that the sequence $(f_n) \subset \mathcal{F}$ converges (pointwise either to a complex number or to ∞) on a set $S \subset D$ having at least one accumulation point in D . Then (f_n) converges, uniformly on compact subsets of D , either to a (finite-valued) analytic function or else to the constant function ∞ .*

Proof. Define $f(z) = \lim_{n \rightarrow \infty} f_n(z)$ for $z \in S$. Now, for any subsequence $\mathcal{S} = (f_{n_k})$, there exists (by normality) a subsubsequence $\mathcal{S}' = (f_{n_{k_j}})$ that converges, uniformly on compact subsets of D , either to a (finite-valued) analytic function $g_{\mathcal{S}'}$ or else to the constant function $g_{\mathcal{S}'} \equiv \infty$. Moreover, we must have $g_{\mathcal{S}'} \upharpoonright S = f$ for all \mathcal{S}' . Therefore, we either have $f \equiv \infty$, in which case $g_{\mathcal{S}'} \equiv \infty$ for all \mathcal{S}' ; or else f is everywhere finite-valued, in which case the $g_{\mathcal{S}'}$ are all equal to the *same* analytic function g (since S is a determining set for analytic functions on D). It follows that (f_n) converges, uniformly on compact subsets of D , either to the constant function ∞ (in the first case) or to the analytic function g (in the second). \square

We are now ready to prove Theorem 3.1.

Proof of Theorem 3.1. Suppose the contrary, *i.e.*, suppose that there exists an $\epsilon > 0$ and an infinite sequence $n_1 < n_2 < \dots$ such that none of the functions f_{n_k} has a zero in the set $D_\epsilon = \{z \in \mathbb{C} : |z - x_0| < \epsilon\}$; let us also take ϵ small enough so that $\overline{D_\epsilon} \subset D$. Then, since D_ϵ is simply connected, $\log f_{n_k}$ is analytic in D_ϵ (we take the branch that is real on $D_\epsilon \cap \mathbb{R}$), and $u_k \equiv a_{n_k} \log f_{n_k}$ satisfies $\lim_{k \rightarrow \infty} u_k(x) = u(x)$ for $x \in D_\epsilon \cap \mathbb{R}$. Moreover, by hypothesis the functions $\operatorname{Re} u_k$ are uniformly bounded above on D_ϵ , so it follows from Proposition 3.3 that the (u_k) are a normal family on D_ϵ . Lemma 3.4 then implies that the sequence (u_k) converges on D_ϵ to an analytic function \tilde{u} that extends u . But this contradicts the hypothesis that u is not real-analytic at x_0 . \square

Let us next recall that a real-valued function u on a domain $D \subset \mathbb{C} \simeq \mathbb{R}^2$ is called *harmonic* if it is twice continuously differentiable and satisfies Laplace’s equation

$$\Delta u \equiv \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0, \tag{3.3}$$

where we write $z = x + iy$. The key facts (see, *e.g.*, [28, pp. 122–124]) are: the real part of every analytic function is harmonic; and conversely, if the domain D is simply connected, then every harmonic function on D is the real part of some analytic function.

One can define normality for families of harmonic functions [95, Section 5.4], as follows.

Definition 2. A family \mathcal{H} of harmonic functions on a domain $D \subset \mathbb{C}$ is said to be *normal* (in D) if, for every sequence $(u_n) \subset \mathcal{H}$, there exists a subsequence (u_{n_k}) that converges, uniformly on compact subsets of D , either to a (finite-valued) harmonic function or else to the constant function $+\infty$ or $-\infty$.

A covering and diagonalization argument analogous to [95, Theorem 2.1.2] shows that normality is a local property: \mathcal{H} is normal in D if and only if, for each $z \in D$, there exists an open disc $U \ni z$ in which \mathcal{H} is normal.

There is a sufficient condition for normality analogous to Proposition 3.3, as follows.

Proposition 3.5. *Let \mathcal{H} be a family of harmonic functions on a domain D , which are uniformly bounded above on compact subsets of D . Then \mathcal{H} is normal: that is, for every sequence $(u_n) \subset \mathcal{H}$, there exists a subsequence (u_{n_k}) that converges, uniformly on compact subsets of D , either to a (finite-valued) harmonic function or else to the constant function $-\infty$.*

Proof. Since normality is a local property, it suffices to prove the proposition when D is an open disc U . (The only property of U we will really use is its simple connectedness.) So let (u_n) be a sequence of harmonic functions in U that is uniformly bounded above on compact subsets of U . Choose analytic functions f_n on U such that $\operatorname{Re} f_n = u_n$, and let $F_n = \exp(f_n)$. The functions F_n are nonvanishing, and they are uniformly bounded on compact subsets of U . By Montel’s (1907) theorem, there exists a subsequence (F_{n_k}) that converges, uniformly on compact subsets of U , to an analytic function F . By Hurwitz’s theorem, F is either identically zero or else nonvanishing. In the former case,

$u_{n_k} \equiv \operatorname{Re} \log F_{n_k} = \log |F_{n_k}|$ tends to $-\infty$ uniformly on compact subsets of U . In the latter case, u_{n_k} tends to the harmonic function $u_\infty \equiv \operatorname{Re} \log F = \log |F|$ uniformly on compact subsets of U . (For each compact $K \subset U$, we have $\inf_{z \in K} |F(z)| \equiv \delta > 0$ and $\sup_{z \in K} |F(z)| \equiv M < \infty$, so for all sufficiently large k we have $\delta/2 \leq |F_{n_k}| \leq 2M$ everywhere on K . Now apply the uniform continuity of the log function on $[\delta/2, 2M]$.) \square

Remarks. (1) For a slightly different proof, see [95, Theorems 5.4.2 and 5.4.3].

(2) Montel’s proof [72, Section 23] is *not* valid: if a subsequence of $(f_n \equiv u_n + iv_n)$ tends to ∞ , it does not follow that the corresponding subsequence of $(|u_n|)$ also tends to ∞ ; it could be that $(|v_n|)$ does so instead (see, e.g., [95, p. 184]).

Proof of Theorem 3.2. Suppose the contrary, *i.e.*, suppose that there exists an $\epsilon > 0$ and an infinite sequence $n_1 < n_2 < \dots$ such that none of the functions f_{n_k} has a zero in the set $D_\epsilon = \{z \in \mathbb{C} : |z - z_0| < \epsilon\} \subset D$. Then each function $u_k \equiv a_{n_k} \log |f_{n_k}|$ is harmonic on D_ϵ ; by hypothesis, the functions u_k are uniformly bounded above on compact subsets of D_ϵ ; so it follows from Proposition 3.5 that the (u_k) are a normal family on D_ϵ . Therefore, there exists a subsequence (u_{k_l}) that converges, uniformly on compact subsets of D_ϵ , to a function v that is either harmonic on D_ϵ or else identically $-\infty$. But this contradicts the hypothesis of the theorem. \square

Remark. An argument closely related to that of Theorems 3.1 and 3.2 was used previously, in a special context, by Borwein, Chen and Dilcher [20, pp. 79 and 82]. I thank Karl Dilcher for informing me of this article.

We now proceed to the proof of Theorem 1.5. We have already defined

$$D_k = \{z : k \text{ is dominant at } z\}. \tag{3.4}$$

Let us further define the ‘good sets’

$$V_k = \{z : k \text{ is the unique dominant index at } z, \text{ and } \alpha_k(z) \neq 0\} \tag{3.5}$$

and the ‘bad sets’

$$W_k = \{z : k \text{ is the unique dominant index at } z, \text{ and } \alpha_k(z) = 0\}, \tag{3.6}$$

$$X = \{z : \beta_1(z) = \beta_2(z) = \dots = \beta_m(z) = 0\}, \tag{3.7}$$

$$Y_{kl} = \{z : k \text{ and } l \ (k \neq l) \text{ are dominant indices at } z \text{ (there may be others)} \\ \text{and } |\beta_k(z)| > 0\}. \tag{3.8}$$

Clearly all the sets (3.5)–(3.8) are disjoint, except that some of the Y_{kl} may overlap at points where there are three or more dominant indices. We shall write $V = \bigcup_{k=1}^m V_k$, $W = \bigcup_{k=1}^m W_k$ and $Y = \bigcup_{1 \leq k < l \leq m} Y_{kl}$; we obviously have the disjoint decomposition $D = V \cup W \cup X \cup Y$. Theorem 1.5 amounts to the statement that

$$\liminf \mathcal{L}(f_n) = \limsup \mathcal{L}(f_n) = W \cup X \cup Y. \tag{3.9}$$

We begin by collecting some simple facts about these sets.

Lemma 3.6. *Under the hypotheses of Theorem 1.5:*

- (a) *Each D_k is closed in D . In particular, each ∂D_k has empty interior.*
- (b) *$D_k = V_k \cup W_k \cup X \cup (\bigcup_{l \neq k} Y_{kl})$, and the latter four sets are disjoint.*
- (c) *For $k \neq l$, $D_k \cap D_l = X \cup Y_{kl}$. In particular, each set $X \cup Y_{kl}$ is closed.*
- (d) *For $k \neq l$, $Y_{kl} \subset \partial D_k \cap \partial D_l$. In particular, each Y_{kl} has empty interior.*
- (e) *Each V_k is open.*
- (f) *Each $V_k \cup W_k$ is open. Moreover, $V_k \cup W_k \subset D_k^\circ \subset V_k \cup W_k \cup X$, where $^\circ$ denotes interior.*
- (g) *W has no limit points in D . Moreover, for each $z \in W_k$ there exists $\epsilon > 0$ such that the punctured neighbourhood $\{z' : 0 < |z' - z| < \epsilon\}$ is contained in V_k .*
- (h) *X has no limit points in D .*
- (i) *$W \cup X \cup Y$ is a closed set with empty interior, so that V is a dense open subset of D .*
- (j) *$W \cup X \cup Y = \bigcup_{k=1}^m \partial V_k$.*
- (k) *$Y = \bigcup_{1 \leq k < l \leq m} (Y_{kl} \cap \partial V_k \cap \partial V_l)$.*
- (l) *Y is dense-in-itself.*

Proof. (a)–(c) are trivial.

(d) Let $z_0 \in Y_{kl} \subset D_k \cap D_l$, so that $|\beta_k(z_0)| = |\beta_l(z_0)| > 0$. Then β_k/β_l is analytic in a neighbourhood $U \ni z_0$. If β_k/β_l equals a constant ω , then the no-degenerate-dominance condition implies that $D_k (= D_l)$ must have empty interior, so trivially $z_0 \in \partial D_k = \partial D_l$. If, on the other hand, β_k/β_l is nonconstant, then the open mapping theorem implies that β_k/β_l must take values both inside and outside the unit circle in every neighbourhood of z_0 . The former points belong to D_k^c , and the latter to D_l^c ; so $z_0 \in \partial D_k \cap \partial D_l$.

(e) is trivial.

(f) $V_k \cup W_k$ is clearly open, hence contained in D_k° . On the other hand, by (d), $D_k^\circ \cap Y_{kl} = \emptyset$ for each $l \neq k$, so $D_k^\circ \subset V_k \cup W_k \cup X$.

(g) and (h) are trivial consequences of the assumption that none of the α_k or β_k are identically zero.

(i) We use the fact (valid in arbitrary topological spaces [44, Exercise 1.3.D]) that if A is closed, then for every set B we have $(A \cup B)^\circ = (A \cup B^\circ)^\circ$. In particular, if A and B have empty interior and at least one of them is closed, then $A \cup B$ has empty interior. Applying this to $A = X$ and $B = Y_{kl}$, we conclude from (d) and (h) that $X \cup Y_{kl}$ has empty interior. Moreover, by (c), $X \cup Y_{kl}$ is closed. Further repeated applications, using (g), then lead to the conclusion that $W \cup X \cup Y$ is a closed set with empty interior.

(j) By (i), V is open and dense, so $W \cup X \cup Y = V^c = \partial V$. On the other hand, $\partial(\bigcup_{k=1}^m V_k) = \bigcup_{k=1}^m \partial V_k$ for any finite collection of disjoint open sets in any topological space [44, Theorem 1.3.2(iii) and Exercise 1.3.A].

(k) Let $z_0 \in Y$, and let S be the set of indices that are dominant at z_0 . Let U be an open neighbourhood of z_0 , chosen small enough so that no index in S^c is dominant at any point of U . By (i), $U \cap V$ is dense in U . I claim that there must exist at least two distinct indices $k \in S$ for which $U \cap V_k$ is nonempty. For suppose that there is only one such

index, so that $U \cap V_k$ is dense in U . Then $U \subset D_k$ and $z_0 \in Y_{kl}$ for some $l \neq k$ ($l \in S$). Now, shrinking U if necessary so that β_k is nonvanishing in U , we have $|\beta_l/\beta_k| \leq 1$ in U , with $|\beta_l(z_0)/\beta_k(z_0)| = 1$; so by the maximum modulus theorem we must have $\beta_l \equiv \omega\beta_k$ for some constant ω with $|\omega| = 1$, contrary to the no-degenerate-dominance condition. Hence there are indices $k \neq l$ ($k, l \in S$) such that $U \cap V_k$ and $U \cap V_l$ are both nonempty. Since this holds for arbitrarily small $U \ni z_0$, and since there are finitely many pairs k, l , we must have $z_0 \in Y_{kl} \cap \partial V_k \cap \partial V_l$ for some pair $k \neq l$.

(I) Let $z_0 \in Y$, and let U be any connected open neighbourhood of z_0 , chosen small enough so that $U \cap (W \cup X) = \emptyset$; and let $U' = U \setminus \{z_0\}$. If it were true that $U' \cap Y = \emptyset$, then we would have $U' \subset \bigcup_{k=1}^m V_k$. But since the V_k are open and disjoint, and U' is connected, U' would have to be contained in one set V_k ; but this contradicts the fact (k) that every point in Y has nearby points in at least two sets V_k and V_l . □

Remarks. (1) It is not necessarily true that $Y_{kl} \subset \partial V_k \cap \partial V_l$. For example, let $D = \mathbb{C} \setminus \{0\}$ and $m = 3$, and let $\beta_1(z) = z, \beta_2(z) = 1/z, \beta_3(z) = 1, \alpha_1(z) = \alpha_2(z) = \alpha_3(z) = 1$. Then $Y_{12} = Y_{13} = Y_{23} = \partial V_1 = \partial V_2 = \text{unit circle}$, but $V_3 = \emptyset$.

(2) Y cannot have isolated points, but the individual sets Y_{kl} can: though in a neighbourhood of $z_0 \in Y_{kl}$ there certainly exist nearby points where $|\beta_k| = |\beta_l|$, such points may fail to belong to Y_{kl} because the indices k, l fail to remain dominant. *Example:* $\beta_1(z) = 1 + z, \beta_2(z) = 1 - z, \beta_3(z) = 1$. Then $Y_{12} = \text{imaginary axis}$, but $Y_{13} = Y_{23} = \{0\}$.

(3) In this paper we will not use parts (j) and (l) of Lemma 3.6, but they are useful facts to know in applications.

Let us now define

$$u_n(z) = \frac{1}{n} \log |f_n(z)| = \frac{1}{n} \operatorname{Re} \log f_n(z), \tag{3.10}$$

which is well-defined everywhere on D provided that we give it the value $-\infty$ at the zeros of f_n . Clearly, u_n is a continuous map from D into $\mathbb{R} \cup \{-\infty\}$, and is a harmonic function on $D \setminus \mathcal{Z}(f_n)$.

We can compute $\lim_{n \rightarrow \infty} u_n(z)$ at nearly every point $z \in D$. Let us define

$$S(z) = \{k : \alpha_k(z) \neq 0\}, \tag{3.11}$$

$$\tilde{\beta}(z) = \begin{cases} \max\{|\beta_k(z)| : k \in S(z)\} & \text{if } S(z) \neq \emptyset, \\ 0 & \text{if } S(z) = \emptyset, \end{cases} \tag{3.12}$$

$$T(z) = \{k \in S(z) : |\beta_k(z)| = \tilde{\beta}(z)\}. \tag{3.13}$$

The next lemma then follows easily from the definition of f_n .

Lemma 3.7. *Under the hypotheses of Theorem 1.5:*

(a) *For $z \in V_k, \lim_{n \rightarrow \infty} u_n(z) = \log |\beta_k(z)| > -\infty$, and the convergence is uniform on compact subsets of V_k .*

- (b) More generally, if $T(z) = \{k\}$, then $\lim_{n \rightarrow \infty} u_n(z) = \log |\beta_k(z)|$, though this may equal $-\infty$.
- (c) For $z \in X$, $u_n(z) = -\infty$ for all n , so that $\lim_{n \rightarrow \infty} u_n(z) = -\infty$. More generally, this occurs whenever $\tilde{\beta}(z) = 0$ (i.e., when $\alpha_k(z)\beta_k(z) = 0$ for all k).
- (d) For all $z \in D$, $\limsup_{n \rightarrow \infty} u_n(z) \leq \log \tilde{\beta}(z)$. In particular $\limsup_{n \rightarrow \infty} u_n(z) < \log |\beta_k(z)|$, for $z \in W_k$.

Remark. The behaviour of the sequence $\{u_n(z)\}$ is subtle at points z where two or more dominant indices compete, i.e., where $|T(z)| \geq 2$ and $\tilde{\beta}(z) > 0$. But we will not need this information.

Proof of Theorem 1.5. If $z_0 \in V_k$, then choose $\epsilon > 0$ so that $\overline{D}_\epsilon \equiv \{z : |z - z_0| \leq \epsilon\} \subset V_k$; it follows that there exist constants $\delta > 0$ and $M < \infty$ such that

$$|\alpha_k(z)| \geq \delta, \tag{3.14a}$$

$$\left| \frac{\alpha_l(z)}{\alpha_k(z)} \right| \leq M \quad \text{for all } l \neq k, \tag{3.14b}$$

$$|\beta_k(z)| \geq \delta, \tag{3.14c}$$

$$\left| \frac{\beta_l(z)}{\beta_k(z)} \right| \leq 1 - \delta \quad \text{for all } l \neq k, \tag{3.14d}$$

for all $z \in \overline{D}_\epsilon$. Then

$$|f_n(z)| \geq |\alpha_k(z)| |\beta_k(z)|^n - \sum_{l: l \neq k} |\alpha_l(z)| |\beta_l(z)|^n \tag{3.15a}$$

$$\geq \delta^{n+1} [1 - (m - 1)M(1 - \delta)^n] \tag{3.15b}$$

for all $z \in \overline{D}_\epsilon$. Therefore, f_n is nonvanishing on \overline{D}_ϵ for all sufficiently large n , so that $z_0 \notin \limsup \mathcal{Z}(f_n)$.

If $z_0 \in W_k$, we have $\limsup_{n \rightarrow \infty} u_n(z_0) \leq \log \tilde{\beta}(z_0) < \log |\beta_k(z_0)|$ by Lemma 3.7(d). On the other hand, by Lemmas 3.6(g) and 3.7(a) there exists a punctured neighbourhood of z_0 on which $\lim_{n \rightarrow \infty} u_n(z) = \log |\beta_k(z)|$; and this quantity tends to $\log |\beta_k(z_0)|$ as $z \rightarrow z_0$. Therefore, there does not exist a neighbourhood $U \ni z_0$ and a continuous function $v : U \rightarrow \mathbb{R} \cup \{-\infty\}$ satisfying $\liminf_{n \rightarrow \infty} u_n(z) \leq v(z) \leq \limsup_{n \rightarrow \infty} u_n(z)$ for all $z \in U$. In particular, we can apply Theorem 3.2 and conclude that $z_0 \in \liminf \mathcal{Z}(f_n)$.

If $z_0 \in X$, we have $f_n(z_0) = 0$ for all n , so trivially $z_0 \in \liminf \mathcal{Z}(f_n)$.

Finally, if $z_0 \in Y$, Lemma 3.6(k) implies that there exist indices $k \neq l$ such that $z_0 \in Y_{kl} \cap \partial V_k \cap \partial V_l$. So let U be any open neighbourhood of z_0 , chosen small enough so that β_k and β_l are nonvanishing on U ; then the sets $U_k \equiv U \cap V_k$ and $U_l \equiv U \cap V_l$ are nonempty. Lemma 3.7(a) implies that

$$\lim_{n \rightarrow \infty} u_n(z) = \begin{cases} \log |\beta_k(z)| & \text{for } z \in U_k, \\ \log |\beta_l(z)| & \text{for } z \in U_l. \end{cases} \tag{3.16}$$

Now there cannot exist a harmonic function v on U that coincides with $\lim_{n \rightarrow \infty} u_n(z)$ on $U_k \cup U_l$: for if there did, then by the uniqueness of harmonic continuation we would

have $|\beta_k(z)| = |\beta_l(z)| = e^{v(z)}$ on U , hence $\beta_k \equiv \omega\beta_l$ for some constant ω with $|\omega| = 1$, contrary to the no-degenerate-dominance condition. Therefore, by Theorem 3.2, we have $z_0 \in \liminf \mathcal{L}(f_n)$. \square

4. Proof of a strong form of Lemma 1.6

I will prove the following strengthened version of Lemma 1.6.

Theorem 4.1. *Let F_1, F_2, G be analytic functions on a domain $D \subset \mathbb{C}$, with $G \neq \text{constant}$. For each integer $s \geq 0$, define*

$$\mathcal{U}_s = \{z \in D : |1 + F_1(z)G(z)^s| = |1 + F_2(z)G(z)^s|\}. \tag{4.1}$$

Then $\liminf \mathcal{U}_s = \limsup \mathcal{U}_s = \{z \in D : |G(z)| \leq 1\} \cup \{z \in D : |F_1(z)| = |F_2(z)|\}$.

In the proof of Theorem 4.1, I will need the following elementary consequence of Taylor’s theorem with remainder.

Lemma 4.2. *Let w be a complex number. Then $|\log(1 + w) - w| \leq |w|^2$ provided that $|w| \leq 1/4$, where \log denotes the principal branch.*

Proof of Theorem 4.1. It is easy to see that $\limsup \mathcal{U}_s \subset \{z \in D : |G(z)| \leq 1\} \cup \{z \in D : |F_1(z)| = |F_2(z)|\}$: for if $z_0 \in D$ with $|G(z_0)| > 1$ and $|F_1(z_0)| > |F_2(z_0)|$ (say), then we can find a neighbourhood $U \ni z_0$ on which $|G(z)| \geq 1 + \delta$ and $|F_1(z)| - |F_2(z)| \geq \delta$, for some $\delta > 0$; but then we have $|1 + F_1(z)G(z)^s| > |1 + F_2(z)G(z)^s|$ for all s large enough so that $\delta(1 + \delta)^s > 2$.

So it suffices to prove that

- (a) $\liminf \mathcal{U}_s \supset \{z \in D : |G(z)| \leq 1\}$ and
- (b) $\liminf \mathcal{U}_s \supset \{z \in D : |G(z)| > 1 \text{ and } |F_1(z)| = |F_2(z)|\}$.

(a) Assume that $z_0 \in D$ with $|G(z_0)| \leq 1$. Fix any $\epsilon > 0$ small enough so that $D_\epsilon \equiv \{z : |z - z_0| < \epsilon\} \subset D$. Since G is nonconstant, by the open mapping theorem we can find $z_1 \in D_\epsilon$ such that $|G(z_1)| < 1$ and $G'(z_1) \neq 0$; and we can then find an open set $U \ni z_1$ with $U \subset \overline{U} \subset D_\epsilon$, and a number $\rho < 1$, such that G is a homeomorphism of U onto $G[U] \subset \{w : |w| < \rho\}$. If $F_1 \equiv F_2$, the claim we want to prove is trivial; so assume $F_1 \neq F_2$ and choose an open set $V \subset U$ such that

$$(F_1 - F_2)[V] \subset \left\{ w : |w| \geq \delta \text{ and } \varphi_1 - \frac{\pi}{4} \leq \arg w \leq \varphi_1 + \frac{\pi}{4} \right\} \tag{4.2}$$

for some $\delta > 0$ and some φ_1 . Since $\overline{V} \subset D_\epsilon$, we have

$$M \equiv \sup_{z \in V} [|F_1(z)| + |F_2(z)|] < \infty. \tag{4.3}$$

Since $G[V]$ is an open subset of $\{w : |w| < \rho\}$, it contains an arc $A_{r,\theta_1,\theta_2} \equiv \{w = re^{i\theta} : \theta_1 \leq \theta \leq \theta_2\}$ for some $0 < r < \rho$ and $\theta_1 < \theta_2$. Let $C = U \cap G^{-1}[A_{r,\theta_1,\theta_2}]$; since $G \upharpoonright U$

is a homeomorphism, we know that C is a topological arc (and in particular a connected set) contained in V , satisfying $G[C] = A_{r,\theta_1,\theta_2}$. Now take $s_1 = 2\pi/(\theta_2 - \theta_1)$. It follows that for each integer $s \geq s_1$, the set $G^s[C] \equiv \{G(z)^s : z \in C\}$ is exactly the circle $\{w : |w| = r^s\}$.

Now choose s_2 so that $M\rho^{s_2} \leq 1/4$. Then, for any integer $s \geq s_2$, the function

$$L_s(z) = \log[1 + F_1(z)G(z)^s] - \log[1 + F_2(z)G(z)^s] \tag{4.4}$$

(where ‘log’ denotes the principal branch) is analytic on V . Furthermore, by Lemma 4.2 we have

$$|L_s(z) - [F_1(z) - F_2(z)]G(z)^s| \leq 2(Mr^s)^2 \tag{4.5}$$

for $z \in C$.

For each integer $s \geq s_1$, there exist points $z_s^+, z_s^- \in C$ such that $G(z_s^\pm)^s = \pm r^s e^{-i\varphi_1}$, and hence

$$\pm \operatorname{Re}\{[F_1(z_s^\pm) - F_2(z_s^\pm)] G(z_s^\pm)^s\} \geq \frac{\delta r^s}{\sqrt{2}}. \tag{4.6}$$

For $s \geq \max(s_1, s_2)$ we then have

$$\pm \operatorname{Re} L_s(z_s^\pm) \geq \frac{\delta r^s}{\sqrt{2}} - 2M^2 r^{2s}. \tag{4.7}$$

Now choose $s_0 \geq \max(s_1, s_2)$ large enough so that $2M^2 r^{s_0} < \delta/\sqrt{2}$. It follows that for $s \geq s_0$ we have $\pm \operatorname{Re} L_s(z_s^\pm) > 0$. Since L_s is continuous and C is connected, it follows that there exists $z_s^{(0)} \in C$ such that $\operatorname{Re} L_s(z_s^{(0)}) = 0$. This completes the proof of assertion (a).

(b) Assume that $z_0 \in D$ with $|G(z_0)| > 1$ and $|F_1(z_0)| = |F_2(z_0)|$. If $F_1(z_0) = F_2(z_0) = 0$, the claim is trivial (we have $z_0 \in \mathcal{U}_s$ for all s); so assume that $|F_1(z_0)| = |F_2(z_0)| > 0$. Fix any $\epsilon > 0$ small enough so that $D_\epsilon \equiv \{z : |z - z_0| < \epsilon\} \subset D$ and

$$|G(z)| \geq 1 + \delta, \tag{4.8a}$$

$$|F_1(z)| \geq \delta, \tag{4.8b}$$

$$|F_2(z)| \geq \delta, \tag{4.8c}$$

for all $z \in D_\epsilon$, for some $\delta > 0$. We then consider two cases.

Case 1: $F_1 \equiv \omega F_2$ for some $\omega \in \mathbb{C}$ (obviously $|\omega| = 1$).

We need to show that for all s sufficiently large, there exists $z \in D_\epsilon$ such that

$$|1 + F_1(z)^{-1}G(z)^{-s}| = |1 + F_2(z)^{-1}G(z)^{-s}|. \tag{4.9}$$

But this follows from part (a) applied to the functions $1/F_1$, $1/F_2$ and $1/G$.

Case 2: There does not exist $\omega \in \mathbb{C}$ such that $F_1 \equiv \omega F_2$.

In this case the function

$$f(z) = F_1(z_0)F_2(z) - F_2(z_0)F_1(z) \tag{4.10}$$

has a zero at $z = z_0$ but is not identically vanishing. So it follows from Rouché’s theorem that for all sufficiently large s , the equation

$$F_1(z_0)F_2(z) - F_2(z_0)F_1(z) + [F_1(z_0) - F_2(z_0)]G(z)^{-s} = 0 \tag{4.11}$$

has a solution $z \in D_\epsilon$. But this means that

$$\frac{F_1(z) + G(z)^{-s}}{F_1(z_0)} = \frac{F_2(z) + G(z)^{-s}}{F_2(z_0)}, \tag{4.12}$$

hence that $z \in \mathcal{U}_s$. □

5. Completion of the proof of Theorem 1.4

We can now put together all the ingredients to complete the proof of Theorem 1.4 (and hence also of the weaker Theorem 1.2) along the lines sketched in Section 1.3. We shall give the details for part (a); the proof of (b) is almost identical, *mutatis mutandis*.

Fix a complex number $v_0 \neq 0$, and fix $\epsilon > 0$ and $R < \infty$. Since the open region $D \equiv \{q \in \mathbb{C} : |q + v_0| > |v_0| \text{ and } |q| < R\}$ has compact closure $\bar{D} \equiv \{q \in \mathbb{C} : |q + v_0| \geq |v_0| \text{ and } |q| \leq R\}$, we can choose finitely many points $q_1, \dots, q_n \in D$ such that each point $q \in \bar{D}$ lies within a distance $\epsilon/3$ of at least one q_i .

Now, for each i ($1 \leq i \leq n$), apply Lemma 1.6 (or the stronger Theorem 4.1) with $z = q - q_i$, $F_1(z) = q - 1$, $F_2(z) = -1$, $G(z) = v_0/(q + v_0)$. (The hypothesis $v_0 \neq 0$ guarantees that $G \neq \text{constant}$, and the hypothesis $q_i \in D$ guarantees that $|G(0)| < 1$.) We conclude that, for each i , there exists an integer $s_{0,i} < \infty$ such that for all integers $s \geq s_{0,i}$ the equation

$$\left| 1 + (q - 1) \left(\frac{v_0}{q + v_0} \right)^s \right| = \left| 1 - \left(\frac{v_0}{q + v_0} \right)^s \right| \tag{5.1}$$

(cf. (1.10)) has a solution in the disc $|q - q_i| < \epsilon/3$. Let $s_0 = \max_{1 \leq i \leq n} s_{0,i}$.

Now fix any $s \geq s_0$, and let q'_1, \dots, q'_n be solutions of (5.1) satisfying $|q'_i - q_i| < \epsilon/3$ and $q'_i \neq 0$ for all i . Apply Theorem 1.5 (or the original Beraha–Kahane–Weiss theorem) with $m = 2$, $n = p$, $z = q$, $\alpha_1(z) = 1$, $\alpha_2(z) = q - 1$, $\beta_1(z) = (q + v_0)^s + (q - 1)v_0^s$, $\beta_2(z) = (q + v_0)^s - v_0^s$. (The nondegeneracy condition holds because $v_0 \neq 0$.) We conclude that the zeros of

$$q^{p-1} Z_{\Theta^{(s,p)}}(q, v) = [(q + v_0)^s + (q - 1)v_0^s]^p + (q - 1)[(q + v_0)^s - v_0^s]^p \tag{5.2}$$

(cf. (1.9)) accumulate (in \liminf as well as \limsup sense) at each point of the solution set of (5.1). In particular, for each i , there exists a zero of (5.2) in the disc $|q - q'_i| < \min(\epsilon/3, |q'_i|)$. (Since this disc does not contain 0, the given q must be a zero of $Z_{\Theta^{(s,p)}}(q, v_0)$.) Set $p_0 = \max_{1 \leq i \leq n} p_{0,i}$.

We have shown that for each $p \geq p_0$, there exist zeros q''_1, \dots, q''_n of $Z_{\Theta^{(s,p)}}(\cdot, v_0)$ such that each point $q \in \bar{D}$ lies within distance ϵ of at least one q''_i . This completes the proof.

6. Real chromatic roots of the graphs $\Theta^{(s,p)}$

Though this paper is primarily concerned with the *complex* chromatic roots of the graphs $\Theta^{(s,p)}$ (and more generally, with the complex roots of their dichromatic polynomials $Z_{\Theta^{(s,p)}}(q, v)$), let me digress to discuss their *real* chromatic roots. The chromatic polynomial

$P_{\Theta^{(s,p)}}(q)$ is obtained by specializing (1.9) to $v = -1$:

$$P_{\Theta^{(s,p)}}(q) = \frac{[(q-1)^s + (q-1)(-1)^s]^p + (q-1)[(q-1)^s - (-1)^s]^p}{q^{p-1}}. \tag{6.1}$$

It is useful to write $y = 1 - q$, so that

$$F(y) \equiv (-1)^{sp+1} q^{p-1} P_{\Theta^{(s,p)}}(q) = y(y^s - 1)^p - (y^s - y)^p. \tag{6.2}$$

Recall that $\Theta^{(s,p)}$ has $n \equiv (s-1)p + 2$ vertices. I write $P_G^{(j)}(q)$ to denote the j th derivative of $P_G(q)$. The real chromatic roots of the graphs $\Theta^{(s,p)}$ are then fully characterized by following result.

Proposition 6.1. *Fix integers $s, p \geq 2$ and set $n = (s-1)p + 2$. Then:*

- (a) $(-1)^{n+j} P_{\Theta^{(s,p)}}^{(j)}(q) > 0$ for $q < 0$ and $0 \leq j \leq n$.
- (b) $P_{\Theta^{(s,p)}}(0) = 0$ and $(-1)^n P'_{\Theta^{(s,p)}}(0) = (s-1)^p - s^p < 0$.
- (c) $(-1)^n P_{\Theta^{(s,p)}}(q) < 0$ for $0 < q < 1$.
- (d) $P_{\Theta^{(s,p)}}(1) = 0$ and $(-1)^n P'_{\Theta^{(s,p)}}(1) = 1 > 0$.
- (e) If n is even (i.e., s is odd or p is even or both), then $P_{\Theta^{(s,p)}}(q) > 0$ for $q > 1$.
- (e') If n is odd (i.e., s is even and p is odd), then there exists $q_*^{(s,p)} \in [1.4301597, \dots, 2)$ such that:

- (e'_1) $P_{\Theta^{(s,p)}}(q) < 0$ for $1 < q < q_*^{(s,p)}$;
- (e'_2) $P_{\Theta^{(s,p)}}(q_*^{(s,p)}) = 0$ and $P'_{\Theta^{(s,p)}}(q_*^{(s,p)}) > 0$; and
- (e'_3) $P_{\Theta^{(s,p)}}(q) > 0$ and $P'_{\Theta^{(s,p)}}(q) > 0$ for $q > q_*^{(s,p)}$.

Here $1.4301597, \dots (= q_*^{(2,3)})$ is shorthand for the unique real root of $q^3 - 5q^2 + 10q - 7 = 0$. Moreover,

- (e'_a) $q_*^{(s,p)}$ is a strictly increasing function of s and p ;
- (e'_b) $\lim_{s \rightarrow \infty} q_*^{(s,p)} = 2$ for each p ; and
- (e'_c) $\lim_{p \rightarrow \infty} q_*^{(s,p)} = q_*^{(s,\infty)}$, where $q_*^{(s,\infty)} \in [\frac{3}{2}, 2)$ is the unique nonzero real root of $2(q-1)^s + q - 2 = 0$.

The signs of $P_G(q)$ and its derivatives in (a)–(d) are of course general facts about the chromatic polynomial of any 2-connected graph [133, Theorem 2], but it is amusing to verify them explicitly for $\Theta^{(s,p)}$.

Proof. (a) We have

$$F(y) = (y-1)(y^s - 1)^p + [(y^s - 1)^p - (y^s - y)^p] \tag{6.3a}$$

$$= (y-1) \left[(y^s - 1)^p + \sum_{k=0}^{p-1} (y^s - 1)^k (y^s - y)^{p-1-k} \right], \tag{6.3b}$$

so clearly F and all its derivatives are nonnegative for $y \geq 1$. If we now write $P_{\Theta^{(s,p)}}(q) = \sum_{k=1}^n a_k^{(s,p)} q^k$, we see that $(-1)^{n-k} a_k^{(s,p)} = F^{(k+p-1)}(1)/(k+p-1)! \geq 0$; and of course $a_n^{(s,p)} = 1 > 0$. The claims follow.

(b) An easy calculation.

(c) For $0 < y < 1$, we have $(1 - y^s)/(y - y^s) > 1/y > 1$ and hence $[(1 - y^s)/(y - y^s)]^p > 1/y$, so that $(-1)^p F(y) > 0$.

(d) An easy calculation.

(e) For $y < 0$ and p even, we manifestly have $F(y) < 0$. For $y = -z < 0$ and s odd, we have $(-1)^{p+1} F(y) = z(z^s + 1)^p + (z^s - z)^p$. For $z \geq 1$ this is manifestly > 0 ; for $0 < z < 1$ it is $> z - z^p > 0$ as well.

(e') For $z = q - 1 > 0$ with s even and p odd, we can write $P_{\Theta(s,p)}(q) = (1 + z)G(z)^p [1 - H(z)^p]$ where

$$G(z) = \frac{z + z^s}{1 + z}, \tag{6.4a}$$

$$H(z) = \frac{1 - z^s}{z^{(p-1)/p}(1 + z^{s-1})}. \tag{6.4b}$$

Now $H(z)$ is a strictly decreasing function of z on $(0, \infty)$, which runs from $+\infty$ at $z = 0$ to 0 at $z = 1$ to $-\infty$ at $z = +\infty$; therefore, the equation $H(z) = 1$ has a unique solution $z_*^{(s,p)} \in (0, \infty)$, which lies in fact in $(0, 1)$. Moreover, $H(z)$ is a strictly increasing function of s and p when $z \in (0, 1)$, so $z_*^{(s,p)}$ is a strictly increasing function of s and p ; and it is easy to see that it has properties (e'_b) and (e'_c). Finally, $G(z)$ is a strictly increasing function of z on $(0, \infty)$, so $P'_{\Theta(s,p)}(q) > 0$ for $q \geq q_*^{(s,p)}$. \square

Remarks. (1) For any n -vertex connected graph G and any vertex v of G , the quantity $T_G(1, 0) = (-1)^{n+1} P'_G(0)$ counts the acyclic orientations of G in which v is the unique source ([49, Theorem 7.3], [27, Proposition 6.3.18], [48]). Taking v to be one of the end-vertices of Θ_{s_1, \dots, s_p} , it is easy to show that this quantity equals $\prod_{i=1}^p s_i - \prod_{i=1}^p (s_i - 1)$, exactly as computed from the chromatic polynomial.

(2) For any n -vertex connected graph $G \neq K_2$ and any edge $e = \langle uv \rangle$ of G , the quantity $[\partial T_G(x, y) / \partial x](0, 0) = (-1)^n P'_G(1)$ counts the acyclic orientations of G in which u is the unique source and v is the unique sink ([49, Theorem 7.2], [27, Exercise 6.35], [48]). It also equals half the number of totally cyclic orientations of G (i.e., orientations in which every edge of G belongs to some directed cycle) in which every directed cycle uses e ([49, Theorem 8.2], [27, Proposition 6.2.12 and Example 6.3.29]). Taking u to be one of the end-vertices of Θ_{s_1, \dots, s_p} and v one of its neighbours, it is easy to see that there is exactly one orientation of the first kind (and it is acyclic), and two of the second.

(3) In part (e) we *cannot* assert that $P_{\Theta(s,p)}(q)$ is increasing for $q > 1$. Indeed, for many pairs (s, p) – including $(2, 4)$, $(2, 6)$, $(4, 6)$, $(5, 6)$, $(5, 7)$, ... – there is an interval $1 < q_1 < q < q_2 < 2$ where $P'_{\Theta(s,p)}(q) < 0$.

(4) In part (e'), $P_{\Theta(s,p)}(q)$ is not necessarily convex on $(1, 2)$: for example, for $(s, p) = (2, 7)$ there are inflection points at $q \approx 1.282916, 1.405642$. However, numerical calculations for small s, p suggest that $P_{\Theta(s,p)}(q)$ is convex on the interval $[q_*^{(s,p)}, \infty)$. Indeed, it appears that *all* the derivatives of $P_{\Theta(s,p)}(q)$ are strictly positive at $q = q_*^{(s,p)}$, except that for $s = 2$ and $p \geq 7$ the $(n - 1)$ st derivative is < 0 .

Though I am unable to say much about the monotonicity or convexity of $P_{\Theta(s,p)}(q)$ for $1 < q < 2$ (cf. Remarks (3) and (4) above), the situation simplifies dramatically for $q \geq 2$ for all series-parallel graphs. We begin with the following definition.

Definition 3. Let P be a degree- n polynomial with real coefficients, and let $a \in \mathbb{R}$. We say that P is *strongly nonnegative* (resp. *strongly positive*) at a in case $P^{(j)}(a) \geq 0$ (resp. $P^{(j)}(a) > 0$) for all $0 \leq j \leq n$.

Note that if P is strongly nonnegative at a and is not identically zero, then P is strongly positive on (a, ∞) ; and conversely, if P is strongly nonnegative on (a, ∞) , then it is also strongly nonnegative at a . Note also that if P and Q are strongly nonnegative (resp. strongly positive) at a , then so are $P + Q$ and PQ . I suspect that strong positivity is the appropriate concept for many (though perhaps not all) theorems asserting upper zero-free intervals for chromatic polynomials [17, 136, 115]. Here, at any rate, is one example.

Proposition 6.2. Let $G = (V, E)$ be a finite graph equipped with edge weights $\{v_e\}_{e \in E}$; let x, y be distinct vertices of G ; and let

$$Z_{G,x,y}(q, \{v_e\}; \sigma_x, \sigma_y) = A_{G,x,y}(q, \{v_e\}) + B_{G,x,y}(q, \{v_e\}) \delta(\sigma_x, \sigma_y) \tag{6.5}$$

be its restricted partition function. Suppose that the two-terminal graph (G, x, y) is series-parallel,¹⁰ and that $v_e \geq -1$ for all edges $e \in E$. Then A and $A + B$ are strongly nonnegative at $q = 2$.

Corollary 6.3. Let G be a series-parallel graph equipped with edge weights $\{v_e\}_{e \in E}$ satisfying $v_e \geq -1$ for all edges e . Then $Z_G(q, \{v_e\})/q$ (and hence also $Z_G(q, \{v_e\})$) is strongly nonnegative at $q = 2$. In particular, $P_G(q) > 0$ for $q > 2$.

The proof of Proposition 6.2 is an easy application of the formulae (2.23) and (2.26) for parallel and series connection, which can be rewritten as

$$((A_1, B_1), (A_2, B_2)) \xrightarrow{\text{parallel}} (A^{\text{par}}, B^{\text{par}}) \equiv (A_1A_2, A_1B_2 + A_2B_1 + B_1B_2), \tag{6.6}$$

$$((A_1, B_1), (A_2, B_2)) \xrightarrow{\text{series}} (A^{\text{ser}}, B^{\text{ser}}) \equiv (A_1B_2 + A_2B_1 + qA_1A_2, B_1B_2). \tag{6.7}$$

The first step is the following.

Lemma 6.4. If $A_1, A_1 + B_1, A_2$ and $A_2 + B_2$ are all strongly nonnegative at $q = 2$, then so are $A^{\text{par}}, A^{\text{par}} + B^{\text{par}}, A^{\text{ser}}$ and $A^{\text{ser}} + B^{\text{ser}}$.

Proof. A trivial computation:

$$A^{\text{par}} = A_1A_2, \tag{6.8a}$$

$$A^{\text{par}} + B^{\text{par}} = (A_1 + B_1)(A_2 + B_2), \tag{6.8b}$$

$$A^{\text{ser}} = A_1(A_2 + B_2) + A_2(A_1 + B_1) + (q - 2)A_1A_2, \tag{6.8c}$$

$$A^{\text{ser}} + B^{\text{ser}} = (A_1 + B_1)(A_2 + B_2) + (q - 1)A_1A_2. \tag{6.8d}$$

□

¹⁰ Note that this is *stronger* than requiring simply that G be series-parallel; we require that G with the two fixed terminals x and y be transformable to a single edge by a sequence of series and parallel reductions.

Proposition 6.2 then follows by induction, starting from the fact that it holds when G is a single edge $\langle xy \rangle$ (for which $A = 1$ and $A + B = 1 + v_e \geq 0$). Corollary 6.3 for 2-connected series-parallel graphs G then follows from Proposition 6.2 by choosing some pair of vertices between which the graph is series-parallel, and using $Z_G/q = qA + B = (A + B) + (q - 1)A$; it then holds for general series-parallel graphs by factorization into 2-connected components.

Remark. Thomassen [115, Theorem 3.4] shows that for any integer $k \geq 2$, the set M_k of real chromatic roots of graphs of tree-width $\leq k$ consists of 0, 1 and a dense subset of the interval $(32/27, k]$. Indeed, Thomassen’s argument (which is based on [115, Theorem 3.3]) actually proves the stronger result that $P_G(q)/q$ (and hence also $P_G(q)$) is strongly nonnegative at $q = k$. The case $k = 2$ corresponds to series-parallel graphs.

7. Variants of the construction

7.1. Adding an edge to $\Theta^{(s,p)}$

I recently proved that if G is a graph of maximum degree Δ , then all the chromatic roots of G lie in the disc $|q| < 7.963907\Delta$ [111, Corollary 5.3 and Proposition 5.4].¹¹ Moreover, if *all but one* of the vertices of G have degree $\leq \Delta$, then the chromatic roots of G are still bounded, namely they lie in the disc $|q| < 7.963907\Delta + 1$ [111, Corollary 6.4]. The graphs $\Theta^{(s,p)}$ show that no analogous theorem can hold if ‘all but one’ is replaced by ‘all but two’ (not even when $\Delta = 2$ and G is series-parallel), for in that case the chromatic roots can spread over the whole complex plane (except perhaps the disc $|q - 1| < 1$).

There is at least one instance in which the presence of several vertices of large degree cannot cause large chromatic roots, namely when those vertices form an N -clique within which every pair of vertices is connected by a $v = -1$ edge. In that case, if the remaining vertices of G have degree $\leq \Delta$, then all the chromatic roots of G lie in the disc $|q| < 7.963907\Delta + N$ [111, Theorem 6.3(b)].

The condition here that the clique edges have $v = -1$ is crucial. Indeed, consider the graph $\Theta^{(s,p)} + xy$ (where x and y are the end-vertices of $\Theta^{(s,p)}$) in which the edge xy is assigned a weight v_{xy} and all the other edges have weight v . The corresponding Potts-model partition function is easily calculated from (2.5), (2.8) and (2.31)/(2.32):

$$Z_{\Theta^{(s,p)}+xy}(q, v, v_{xy}) = \frac{(1 + v_{xy})[(q + v)^s + (q - 1)v^s]^p + (q - 1)[(q + v)^s - v^s]^p}{q^{p-1}}, \tag{7.1}$$

which reduces to (2.35) when $v_{xy} = 0$ and to (2.37) when $v_{xy} = -1$. Provided that $v_{xy} \neq -1$, Theorem 1.5 still applies and the zeros accumulate as $s, p \rightarrow \infty$ in the entire region where $|v/(q + v)| \leq 1$.

¹¹ This result is, in fact, the specialization to chromatic polynomials of a more general bound on the zeros of the Potts-model partition function $Z_G(q, \{v_e\})$ throughout the ‘complex antiferromagnetic regime’, *i.e.*, when $|1 + v_e| \leq 1$ for all edges e . The other two results of [111] cited later in this subsection are likewise specializations of bounds valid throughout the complex antiferromagnetic regime. See [111] for details.

7.2. The dual construction

Let $C_n^{(r)}$ be the cycle C_n with each edge replaced by r edges in parallel. Note that the dual graph of $\Theta^{(s,p)}$ is $C_p^{(s)}$, and *vice versa*. The dichromatic polynomial of $C_n^{(r)}$ is easily computed from the dichromatic polynomial (2.28) of C_n together with the parallel reduction formula (2.22):

$$Z_{C_n^{(r)}}(q, v) = [q + (1 + v)^r - 1]^n + (q - 1)[(1 + v)^r - 1]^n. \tag{7.2}$$

It follows from Theorem 1.5 that when $n \rightarrow \infty$ at fixed r , the zeros of $Z_{C_n^{(r)}}$ accumulate where

$$|1 + (q - 1)(1 + v)^{-r}| = |1 - (1 + v)^{-r}|. \tag{7.3}$$

Here the $G \not\equiv$ constant condition of Lemma 1.6 means that we cannot fix v and use q as the variable. However, we can fix $q = q_0 \neq 0$ and use v as the variable;¹² we then have the following theorem.

Theorem 7.1. *Fix complex numbers q_0, v_0 satisfying $q_0 \neq 0$ and $|1 + v_0| \geq 1$. Then, for each $\epsilon > 0$, there exist numbers $r_0 < \infty$ and $n_0(r) < \infty$ such that, for all $r \geq r_0$ and $n \geq n_0(r)$, the dichromatic polynomial $Z_{C_n^{(r)}}(q_0, \cdot)$ has a zero in the disc $|v - v_0| < \epsilon$.*

We can also slice the (q, v) -space in other ways; for example, we can fix $u_0 \neq 0$ and take $v = q/u_0$.

Theorem 7.2. *Fix nonzero complex numbers q_0, u_0 satisfying $|u_0| \leq |q_0 + u_0|$. Then, for each $\epsilon > 0$, there exist numbers $r_0 < \infty$ and $n_0(r) < \infty$ such that for all $r \geq r_0$ and $n \geq n_0(r)$, the polynomial $Z_{C_n^{(r)}}(q, q/u_0)$ has a zero in the disc $|q - q_0| < \epsilon$.*

In particular, if $|q_0 - 1| \geq 1$, then the flow polynomial $F_{C_n^{(r)}}(q) = (-1)^{nr} q^{-n} Z_{C_n^{(r)}}(q, -q)$ has a zero in the disc $|q - q_0| < \epsilon$.

Since the $G \not\equiv$ constant condition forbids us to use q here as the variable with v fixed, these results are somewhat less interesting (at least for direct application to chromatic polynomials) than Theorem 1.2. However, they become useful when we insert nontrivial 2-rooted graphs in place of the edges of $C_n^{(r)}$. But we defer this construction to a separate paper.

8. Discussion

It now appears that there are three *quite distinct* theories of chromatic roots:

- (a) the theory of *integer* chromatic roots (*i.e.*, the combinatorial theory of the chromatic number $\chi(G)$);

¹² Note that the no-degenerate-dominance condition of Theorem 1.5 is satisfied provided that the fixed value of q is nonzero.

- (b) the theory of *real* chromatic roots (i.e., the theory of zero-free and zero-dense intervals for various classes of graphs [17, 133, 134, 135, 57, 136, 115, 42, 116]); and
 (c) the theory of *complex* chromatic roots.¹³

Woodall [133, Theorem 8] has shown that the complete bipartite graphs K_{n_1, n_2} , in the limit $n_2 \rightarrow \infty$ with n_1 fixed, have *real* chromatic roots arbitrarily close to all the integers from 2 through $\lfloor n_1/2 \rfloor$, even though their only *integer* chromatic roots are of course 0 and 1. More generally, the joins $G = K_{n_1, n_2} + K_n$, which have chromatic number $\chi(G) = n + 2$, have real chromatic roots arbitrarily close to all the integers from $n + 2$ through $n + \lfloor n_1/2 \rfloor$, even though their only integer chromatic roots are 0, 1, \dots , $n + 1$. This suggests that the real roots of the chromatic polynomial have nothing much to do with the integer roots. In particular, the real roots of $P_G(q)$ cannot be bounded in terms of the chromatic number $\chi(G)$ alone.

Likewise, the results of the present paper show that the generalized theta graphs $\Theta^{(s,p)}$ have *complex* chromatic roots arbitrarily close to any point in $\mathbb{C} \setminus \{|q - 1| < 1\}$, even though their only *real* chromatic roots are 0, 1 and $q_*^{(s,p)} \in [1.4301597, \dots, 2)$ and their only integer roots are 0 and 1. This suggests that the complex roots of the chromatic polynomial have nothing much to do with the real or integer roots.

In particular, planar graphs (such as $\Theta^{(s,p)}$) can have complex chromatic roots densely surrounding the point 4.¹⁴ This suggests that it is unlikely that there will ever be a ‘complex-analysis’ proof of the Four-Colour Theorem: quite simply, $q = 4$ behaves very differently from complex values of q arbitrarily close to 4. On the other hand, it is conceivable that there may some day be a ‘real-analysis’ proof of the Four-Colour Theorem in the same sense that Birkhoff and Lewis [17, pp. 413–415] (see also [136, 115]) provided a ‘real-analysis’ proof of the Five-Colour Theorem: namely, it may be possible to prove that $P_G(q) > 0$ for real $q \geq 4$, for all loopless planar graphs G . Indeed, Birkhoff and Lewis [17, p. 413] have conjectured that $P_G(q)/[q(q-1)(q-2)] - (q-3)^{n-3}$ is strongly nonnegative (see Definition 3) at $q = 4$ for all loopless planar graphs G with n vertices.

We conclude with a few disconnected remarks.

(1) Let $\mathcal{G}_k^<$ denote the class of graphs G such that every subcontraction (= minor) of G has a vertex of degree $\leq k$. Woodall [136, Question 1] asked whether all the (complex) chromatic roots of $G \in \mathcal{G}_k^<$ are bounded in modulus by k . For $k \geq 2$, the answer is no,

¹³ There is also a (small) fourth theory, namely the ‘number-theoretic’ (or ‘algebraic’) theory of chromatic roots, which exploits the fact that the chromatic polynomial is a monic polynomial with *integer* coefficients. I include here the proof that the generalized Beraha numbers $B_n^{(k)} = 4 \cos^2(k\pi/n)$ for $n = 5, 7, 8, 9$ and $n \geq 11$, with k coprime to n , are never chromatic roots [92] (see [123] for earlier related results); Tutte’s golden-ratio theorems for plane triangulations [120, 121, 123]; and speculations concerning the accumulation of chromatic roots at Beraha numbers $B_n = 4 \cos^2(\pi/n)$ for certain sequences of planar graphs [10, 122, 123, 124, 125, 5, 69, 70, 71, 93, 94, 61, 63, 113, 126, 58, 68, 67, 66, 92, 59, 60]. This theory is, however, as yet rather undeveloped.

¹⁴ The fact that planar graphs can have chromatic roots arbitrarily close to 4 was established already two decades ago in the seminal paper of Beraha and Kahane [8]. They considered $4 \times n$ strips of the triangular lattice with periodic boundary conditions in the ‘short’ direction (and extra sites at both ends of the ‘long’ direction, but this is inessential) and found that the chromatic roots accumulate as $n \rightarrow \infty$ on a curve passing through $q = 4$.

since

$$\text{generalized theta graphs} \subset \text{series-parallel graphs} = \mathcal{G}_2^<, \tag{8.1}$$

and we have shown that the chromatic roots of generalized theta graphs $\Theta^{(s,p)}$ are unbounded. (In fact, the family $\Theta^{(2,p)} \simeq K_{2,p}$ already has unbounded chromatic roots: see [26].)

(2) Read and Royle [83, pp. 1027–1028] have observed empirically that, for cubic graphs with a fixed number of vertices, the graphs with high girth tend to contribute the chromatic roots with smallest real part. It would be valuable to obtain a better understanding of this phenomenon. A first conjecture might be that there is a universal lower bound on the real part of a chromatic root in terms of the girth, but this is false: the graphs $\Theta^{(3,p)}$ all have girth = circumference = 6, but their chromatic roots accumulate as $p \rightarrow \infty$ on the hyperbola

$$(\text{Im } q)^2 - (\text{Re } q - \frac{3}{2})^2 = \frac{1}{4} \tag{8.2}$$

(set $s = 3$ and $v = -1$ in (1.10)), which contains points with arbitrarily negative real part.

(3) The construction employed in this paper is based on concatenating 2-rooted subgraphs (here single edges, but that can be generalized) in order to create a larger 2-rooted subgraph with some desired value of the effective coupling v_{eff} . The situation may thus be very different for 3-connected graphs, in which 2-rooted subgraphs of more than a single edge cannot occur. It is thus reasonable to ask: What is the closure of the set of chromatic roots of 3-connected graphs? The answer, as Bill Jackson has pointed out to me, is ‘still the whole complex plane’. To see this, consider the graphs $\Theta^{(s,p)} + K_n$ for fixed n and varying s, p : they are $(n + 2)$ -connected, and their chromatic roots (taken together) are dense in the region $|q - (n + 1)| \geq 1$. In particular, considering both n and $n + 2$, the chromatic roots are dense in the whole complex plane. So even arbitrarily high connectedness does not, by itself, stop the chromatic roots from being dense in the whole complex plane.

On the other hand, the graphs $\Theta^{(s,p)} + K_n$ are non-planar whenever $n \geq 1$ and $p \geq 3$. This suggests posing a more restricted question.

Question 8.1. *What is the closure of the set of chromatic roots for 3-connected planar graphs?*

Here the answer may well be much smaller than \mathbb{C} or $\mathbb{C} \setminus \{|q - 1| < 1\}$. But it will certainly not be a bounded set, for as Bill Jackson has pointed out to me, the bipyramids $C_n + \bar{K}_2$ are 4-connected plane triangulations, but their chromatic roots are unbounded [84, 100, 102].

(4) The chromatic roots of generalized theta graphs have been further studied in [105, 24, 26]. In particular, Shrock and Tsai [105, Section 3] plot the limiting set (1.10) of chromatic roots as $p \rightarrow \infty$ for fixed $s = 3, 4, 5$. Brown, Hickman, Sokal and Wagner [26] prove that the chromatic roots of Θ_{s_1, \dots, s_p} are bounded in magnitude by $[1 + o(1)] p / \log p$,

uniformly in the path lengths s_1, \dots, s_p ; moreover; they prove that for $\Theta^{(2,p)} \simeq K_{2,p}$ this bound is sharp.

Acknowledgements

I wish to thank Jason Brown, Roberto Fernández, Walter Hayman, Bill Jackson, Henry McKean, Criel Merino, Robert Shrock, Dave Wagner, Norman Weiss, Dominic Welsh and Douglas Woodall for valuable conversations and/or correspondence. I especially wish to thank Norman Weiss for helpful comments on a previous draft of this manuscript. Finally, I wish to thank Herbert Spohn for suggesting, circa 1993, that I study Potts models on hierarchical versions of the graphs $\Theta^{(s,p)}$ [51, 40, 41, 56, 75, 76, 77, 78, 142, 18]; though that work is still in progress, it directly inspired the present paper.

I also wish to thank an anonymous referee for comments on the first version of this paper that led to improvements (I hope) in the exposition.

Appendix A: Simple proof of the Brown–Hickman theorem on chromatic roots of large subdivisions

In this appendix I use the formula (2.25)/(2.26) for series reduction of Potts edges to give a simple proof of the Brown–Hickman [25] theorem on chromatic roots of large subdivisions. This proof gives a further illustration of the utility of considering the general Potts-model partition function $Z_G(q, \{v_e\})$ with not-necessarily-equal edge weights v_e , even if one is ultimately interested in the dichromatic polynomial $Z_G(q, v)$ or the chromatic polynomial $P_G(q)$.

Lemma A1. *Let G be a graph of n vertices and m edges, and let all the edge weights satisfy $|v_e| \leq \delta$. Then all the roots of $Z_G(q, \{v_e\})$ lie in the disc $|q| \leq \max(C, C^{1/(n-1)})$ where $C = (1 + \delta)^m - 1$.*

Proof. From (1.2) we have $Z_G(q, \{v_e\}) = \sum_{k=1}^n a_k q^k$ with $a_n = 1$ and $\sum_{k=1}^{n-1} |a_k| \leq (1 + \delta)^m - 1 = C$. Then

$$|Z_G(q, \{v_e\})| \geq |q|^n - \sum_{k=1}^{n-1} |a_k| |q|^k \geq |q|^n - \begin{cases} C|q|^{n-1} & \text{if } |q| \geq 1 \\ C|q| & \text{if } |q| \leq 1 \end{cases}. \tag{A.1}$$

So $Z_G \neq 0$ if $|q| > \max(C, C^{1/(n-1)})$. □

Remark. This bound is far from being quantitatively optimal; much better bounds, which depend on the maximum degree Δ but not on n or m , can be obtained using the methods of [111]. But Lemma A1 is good enough for our present purposes, as we do not seek any uniformity in G .

Given a finite graph $G = (V, E)$ and a family of integers $\mathbf{k} = \{k_e\}_{e \in E} \geq 1$, we define $G^{\mathbf{k}}$ to be the graph in which the edge e of G is subdivided into k_e edges in series.

Theorem A2. *Let G be a finite graph, let v be a complex number, and let $\epsilon > 0$. Then there exists $K < \infty$ such that $k_e \geq K$ for all edges e implies that all the roots of $Z_{G^k}(q, v)$ lie in the disc $|q + v| \leq (1 + \epsilon)|v|$.*

Proof. Let G have n vertices and m edges. The claim is trivial if $v = 0$, so assume that $v \neq 0$. By the series reduction formula (2.25)/(2.26), we have

$$Z_{G^k}(q, v) = \left(\prod_{e \in E} \frac{(q + v)^{k_e} - v^{k_e}}{q} \right) \times Z_G(q, \{v_{e,\text{eff}}\}), \tag{A.2}$$

where

$$v_{e,\text{eff}} = \frac{q v^{k_e}}{(q + v)^{k_e} - v^{k_e}}. \tag{A.3}$$

The zeros of the prefactor in (A.2) all lie on the circle $|q + v| = |v|$. So let us assume that $|(q + v)/v| = R > 1 + \epsilon$; we shall show that $Z_G(q, \{v_{e,\text{eff}}\}) \neq 0$ provided that K is chosen sufficiently large (depending only on ϵ, n, m and $|v|$). We have $\epsilon < R - 1 \leq |q/v| \leq R + 1$ and hence

$$|v_{e,\text{eff}}| \leq \frac{R + 1}{R^K - 1} |v| < \frac{2 + \epsilon}{(1 + \epsilon)^K - 1} |v| \equiv \delta \tag{A.4}$$

for all edges e (since $(R + 1)/(R^K - 1)$ is a decreasing function of R for $R > 1$). It follows from Lemma A1 that $Z_G(q, \{v_{e,\text{eff}}\}) \neq 0$ provided that we choose K large enough so that $\delta \leq (1 + \alpha)^{1/m} - 1$ where $\alpha = \min[\epsilon|v|, (\epsilon|v|)^{1/(n-1)}]$. □

The Brown–Hickman theorem is the special case $v = -1$.

Appendix B: Chromatic roots of 2-degenerate graphs

Let us recall that the *degeneracy number* of a graph G is defined by

$$D(G) = \max_{H \subseteq G} \delta(H), \tag{B.1}$$

where the max runs over all subgraphs H of G , and $\delta(H)$ is the minimum degree of H . A graph G is said to be *k-degenerate* if $D(G) \leq k$. Equivalently, G is *k-degenerate* if there exists an ordering x_1, x_2, \dots, x_n of the vertex set of G such that

$$\deg_{G \upharpoonright \{x_1, x_2, \dots, x_j\}}(x_j) \leq k \tag{B.2}$$

for $1 \leq j \leq n$, where $G \upharpoonright \{x_1, x_2, \dots, x_j\}$ denotes the induced subgraph.

It is an elementary fact [19, Theorem V.1] that all of the *integer* chromatic roots of G lie in the interval $[0, D(G)]$, i.e., the chromatic number $\chi(G)$ is at most $D(G) + 1$. On the other hand, Thomassen [115, Theorem 3.9] has shown that the *real* chromatic roots cannot be bounded in terms of the degeneracy number: indeed, there are 2-degenerate graphs with arbitrarily large real chromatic roots. Here we shall use Proposition 2.2 and equation (2.32) to give a simple proof of the formula underlying (a slight generalization of) Thomassen’s construction, and to exhibit its exact range of applicability.

One sees immediately from (2.16)/(2.32) that the 2-rooted generalized theta graph $\Theta^{(s,p)}$ (rooted at the end-vertices) has an effective coupling

$$v_{\text{eff}}(q) = \left(\frac{C_s(q)}{A_s(q)} \right)^p - 1 \tag{B.3}$$

where

$$A_s(q) = \frac{(q-1)^s - (-1)^s}{q}, \tag{B.4}$$

$$C_s(q) = \frac{(q-1)^s + (q-1)(-1)^s}{q}. \tag{B.5}$$

So let $Q_s \subset \mathbb{C}$ be the set on which $|1 + v_{\text{eff}}(q)| > 1$, i.e.,

$$Q_s = \{q \in \mathbb{C} : |(q-1)^s + (q-1)(-1)^s| > |(q-1)^s - (-1)^s|\}, \tag{B.6}$$

so that in particular

$$Q_2 = \{q : \text{Re } q > \frac{3}{2}\}, \tag{B.7a}$$

$$Q_3 = \{q : (\text{Im } q)^2 - (\text{Re } q - \frac{3}{2})^2 > \frac{1}{4}\}. \tag{B.7b}$$

We then have the following easy extension of Thomassen’s result.

Theorem B1. *Let G be a loopless graph, and let s be an integer ≥ 2 . Then there exist 2-degenerate (hence 3-colourable) graphs $\{G'(p)\}_{p=1}^\infty$ and an integer $k \geq 0$ such that:*

- (a) $\lim_{p \rightarrow \infty} C_s(q)^{-kp} P_{G'(p)}(q) = P_G(q)$ uniformly on compact subsets of Q_s .
- (b) If G is connected, then so are the $G'(p)$.
- (c) If s is even and G is bipartite, then so are the $G'(p)$.
- (d) If G is embeddable on a surface S , then so are the $G'(p)$.

In particular:

- (i) If $q_0 \in Q_s$ is a zero of P_G , then for all $\epsilon > 0$ and all sufficiently large p (depending on ϵ) $P_{G'(p)}$ has a zero in the disc $|q - q_0| < \epsilon$.
- (ii) If $q_0 \in Q_s \cap \mathbb{R}$ is a zero of P_G of odd multiplicity, then for all $\epsilon > 0$ and all sufficiently large p (depending on ϵ) $P_{G'(p)}$ has a zero in the interval $(q_0 - \epsilon, q_0 + \epsilon)$.

Proof. We can assume without loss of generality that $G = (V, E)$ has no multiple edges (this is just to simplify the notation). For each $x \in V$, define

$$V'_x = \begin{cases} \{x\} & \text{if } x \text{ is an isolated vertex of } G, \\ \{(x, y) : xy \in E\} & \text{otherwise,} \end{cases} \tag{B.8}$$

and let $V' = \bigcup_{x \in V} V'_x$. Let $E' = \{(x, y)(y, x) : xy \in E\}$, and let E'' be a set of edges (with vertex set V') chosen so that the vertex sets of the connected components of the graph (V', E'') are precisely the sets V'_x . (For example, we could connect the vertices of each V'_x by a path or a cycle or a complete graph.) Now let $G' = (V', E' \cup E'')$, and let $Z(q, v)$ be the Potts-model partition function $Z_{G'}(q, \{v_e\})$ with weights $v_e = -1$ for $e \in E'$ and $v_e = v$

for $e \in E''$. It then follows immediately from (1.2) that

$$Z(q, v) = \sum_{k=0}^{|E''|} v^k p_k(q) \tag{B.9}$$

with highest-order coefficient

$$p_{|E''|}(q) = P_G(q). \tag{B.10}$$

Now let $G'(p)$ be obtained from G' by replacing each edge in E'' by the 2-rooted graph $\Theta^{(s,p)}$ (rooted at the end-vertices). By construction, $G'(p)$ is 2-degenerate, hence 3-colourable. Moreover, $G'(p)$ is connected if G is, and $G'(p)$ is bipartite if G is and s is even. Finally, if G is embedded on a surface S , then $G'(p)$ can also be taken to be embedded on S , by drawing the vertex $(x, y) \in V'_x$ on the edge xy very near to x , and by choosing E'' to be a path (or cycle) connecting the vertices of each V'_x in cyclic order as one circles x .

It follows immediately from Proposition 2.2 and equation (2.32) that

$$P_{G'(p)}(q) = A_s(q)^{|E''|} Z(q, v_{\text{eff}}(q)) \tag{B.11}$$

where $v_{\text{eff}}(q)$ is given by (B.3). In particular, for $q \in Q_s$ we have $v_{\text{eff}}(q) \rightarrow \infty$ as $p \rightarrow \infty$, so that

$$\lim_{p \rightarrow \infty} C_s(q)^{-|E''|} P_{G'(p)}(q) = P_G(q). \tag{B.12}$$

Moreover, the convergence is clearly uniform on compact subsets of Q_s .

Claim (i) then follows immediately from (a) using Hurwitz’s theorem [87, p. 262] and the fact that P_G is not identically zero. To prove claim (ii), choose $\epsilon > 0$ small enough so that $P_G(q_0 - \epsilon)$ and $P_G(q_0 + \epsilon)$ have opposite signs, and choose p_0 large enough so that $C_s(q_0 \pm \epsilon)^{-|E''|} P_{G'(p)}(q_0 \pm \epsilon)$ has the same sign as $P_G(q_0 \pm \epsilon)$ for all $p \geq p_0$; then use the intermediate value theorem. □

Remarks. (1) Thomassen [115, Theorem 3.9] considers the special case $s = 2$ and restricts attention to real $q > 2$. He establishes (B.12) for integer $q > 2$ by counting colourings, and then asserts that ‘this also holds’ for real $q > 2$. The assertion is correct but requires a bit of argument; Proposition 2.2 provides a convenient way of filling in the missing details, and determining the exact set Q_s on which (B.12) holds.

(2) One consequence of this result is that the upper zero-free interval for 3-colourable (or 2-degenerate) planar graphs is the same as that for all planar graphs (unless the largest chromatic roots have even multiplicity, which seems unlikely). In commenting on this corollary, Thomassen [115, p. 506] asserted that there exist planar graphs with real chromatic roots arbitrarily close to 4; but this assertion apparently arises from a misunderstanding of the Beraha–Kahane [8] theorem that $4_{\text{periodic}} \times n_{\text{free}}$ triangular lattices have complex chromatic roots arbitrarily close to 4. In fact I do not know of any planar graphs with real chromatic roots arbitrarily close to 4. A study of triangular-lattice strips [60] has thus far found chromatic roots up to ≈ 3.64 ; and Baxter’s work [5] suggests that sufficiently wide and long pieces of the triangular lattice will have real chromatic roots up to at least the Beraha number $B_{14} = 4 \cos^2(\pi/14) \approx 3.801938$ (see also [60] for further discussion). Finally, Douglas Woodall (private communication) has found a 23-vertex

plane triangulation with a chromatic root at ≈ 3.811475 . But I do not know of any planar graphs with chromatic roots larger than this. It is therefore conceivable that the upper zero-free interval for planar graphs extends *below* $q = 4$. I thank Carsten Thomassen and Douglas Woodall for correspondence concerning these questions.

References

- [1] Appel, K. and Haken, W. (1977) Every planar map is four colorable: Part 1, Discharging. *Illinois J. Math.* **21** 429–490.
- [2] Appel, K., Haken, W. and Koch, J. (1977) Every planar map is four colorable: Part 2, Reducibility. *Illinois J. Math.* **21** 491–567.
- [3] Appel, K. and Haken, W. (1989) Every planar map is four colorable. *Contemp. Math.* **98**. American Mathematical Society, Providence RI.
- [4] Baxter, R. J. (1986) q colourings of the triangular lattice. *J. Phys. A* **19** 2821–2839.
- [5] Baxter, R. J. (1987) Chromatic polynomials of large triangular lattices. *J. Phys. A* **20** 5241–5261.
- [6] Beraha, S., Kahane, J. and Weiss, N. J. (1975) Limits of zeroes of recursively defined polynomials. *Proc. Nat. Acad. Sci. USA* **72** 4209.
- [7] Beraha, S., Kahane, J. and Weiss, N. J. (1978) Limits of zeroes of recursively defined families of polynomials. In *Studies in Foundations and Combinatorics*, Vol. 1 of *Advances in Mathematics Supplementary Studies* (G.-C. Rota, ed.), Academic Press, New York, pp. 213–232.
- [8] Beraha, S. and Kahane, J. (1979) Is the four-color conjecture almost false? *J. Combin. Theory Ser. B* **27** 1–12.
- [9] Beraha, S., Kahane, J. and Weiss, N. J. (1980) Limits of chromatic zeros of some families of maps. *J. Combin. Theory Ser. B* **28** 52–65.
- [10] Berman, G. and Tutte, W. T. (1969) The golden root of a chromatic polynomial. *J. Combin. Theory* **6** 301–302.
- [11] Biggs, N. (1993) *Algebraic Graph Theory*, 2nd edn, Cambridge University Press, Cambridge/New York.
- [12] Biggs, N. L., Damerell, R. M. and Sands, D. A. (1972) Recursive families of graphs. *J. Combin. Theory Ser. B* **12** 123–131.
- [13] Biggs, N. (1999) The chromatic polynomial of the $3 \times n$ toroidal square lattice. London School of Economics CDAM Research Report LSE-CDAM-99-05.
- [14] Biggs, N. (2001) A matrix method for chromatic polynomials. *J. Combin. Theory Ser. B* **82** 19–29.
- [15] Biggs, N. and Shrock, R. (1999) $T = 0$ partition functions for Potts antiferromagnets on square lattice strips with (twisted) periodic boundary conditions. *J. Phys. A* **32** L489–L493, cond-mat/0001407 at xxx.lanl.gov.
- [16] Birkhoff, G. D. (1912) A determinantal formula for the number of ways of coloring a map. *Ann. Math.* **14** 42–46.
- [17] Birkhoff, G. D. and Lewis, D. C. (1946) Chromatic polynomials. *Trans. Amer. Math. Soc.* **60** 355–451.
- [18] Bleher, P. M. and Lyubich, M. Yu. (1991) Julia sets and complex singularities in hierarchical Ising models. *Commun. Math. Phys.* **141** 453–474.
- [19] Bollobás, B. (1998) *Modern Graph Theory*, Springer, New York.
- [20] Borwein, P. B., Chen, W. and Dilcher, K. (1995) Zeros of iterated integrals of polynomials. *Canad. J. Math.* **47** 65–87.
- [21] Brenti, F., Royle, G. F. and Wagner, D. G. (1994) Location of zeros of chromatic and related polynomials of graphs. *Canad. J. Math.* **46** 55–80.
- [22] Brown, J. I. (1998) On the roots of chromatic polynomials. *J. Combin. Theory Ser. B* **72** 251–256.

- [23] Brown, J. I. (1998) Chromatic polynomials and order ideals of monomials. *Discrete Math.* **189** 43–68.
- [24] Brown, J. I. and Hickman, C. A. (2002) On chromatic roots with negative real part. *Ars Combinatorica* **63** 211–221.
- [25] Brown, J. I. and Hickman, C. A. (2002) On chromatic roots of large subdivisions of graphs. *Discrete Math.* **242** 17–30.
- [26] Brown, J. I., Hickman, C. A., Sokal, A. D. and Wagner, D. G. (2001) On the chromatic roots of generalized theta graphs. *J. Combin. Theory Ser. B* **83** 272–297, math.CO/0012033 at xxx.lanl.gov.
- [27] Brylawski, T. and Oxley, J. (1992) The Tutte polynomial and its applications. In *Matroid Applications* (N. White, ed.), Vol. 40 of *Encyclopedia of Mathematics and its Applications*, Cambridge University Press, Cambridge, pp. 123–225.
- [28] Cartan, H. (1963) *Elementary Theory of Analytic Functions of One or Several Complex Variables*, Addison-Wesley, Reading, MA. Reprinted by Dover, New York (1995).
- [29] Chandrasekharan, N. and Hannehalli, S. (1994) Efficient algorithms for computing the matching and chromatic polynomials for series-parallel graphs. *J. Combin. Math. Combin. Comput.* **15** 19–32.
- [30] Chang, S.-C. and Shrock, R. (2000) Exact Potts model partition function on strips of the triangular lattice. *Physica A* **286** 189–238, cond-mat/0004181 at xxx.lanl.gov.
- [31] Chang, S.-C. and Shrock, R. (2000) Ground state entropy of the Potts antiferromagnet with next-nearest-neighbor spin-spin couplings on strips of the square lattice. *Phys. Rev. E* **62** 4650–4664, cond-mat/0005236 at xxx.lanl.gov.
- [32] Chang, S.-C. and Shrock, R. (2001) Ground state entropy of the Potts antiferromagnet on triangular lattice strips. *Ann. Phys.* **290** 124–155, cond-mat/0004129 at xxx.lanl.gov.
- [33] Chang, S.-C. and Shrock, R. (2001) Ground state entropy of the Potts antiferromagnet on strips of the square lattice. *Physica A* **290** 402–430, cond-mat/0004161 at xxx.lanl.gov.
- [34] Chang, S.-C. and Shrock, R. (2001) $T = 0$ partition functions for Potts antiferromagnets on lattice strips with fully periodic boundary conditions. *Physica A* **292** 307–345, cond-mat/0007491 at xxx.lanl.gov.
- [35] Chang, S.-C. and Shrock, R. (2001) Structural properties of Potts model partition functions and chromatic polynomials for lattice strips. *Physica A* **296** 131–182, cond-mat/0005232 at xxx.lanl.gov.
- [36] Chang, S.-C. and Shrock, R. (2001) Exact Potts model partition functions on strips of the honeycomb lattice. *Physica A* **296** 183–233, cond-mat/0008477 at xxx.lanl.gov.
- [37] Chang, S.-C. and Shrock, R. (2001) Exact Potts model partition functions on wider arbitrary-length strips of the square lattice. *Physica A* **296** 234–288, cond-mat/0011503 at xxx.lanl.gov.
- [38] Chang, S.-C. and Shrock, R. (2001) Exact partition function for the Potts model with next-nearest neighbor couplings on strips of the square lattice. *Int. J. Mod. Phys. B* **15** 443–478, cond-mat/0007505 at xxx.lanl.gov.
- [39] Chia, G. L. (1997) A bibliography on chromatic polynomials. *Discrete Math.* **172** 175–191.
- [40] Derrida, B., De Seze, L. and Itzykson, C. (1983) Fractal structure of zeros in hierarchical models. *J. Statist. Phys.* **33** 559–569.
- [41] Derrida, B., Itzykson, C. and Luck, J. M. (1984) Oscillatory critical amplitudes in hierarchical models. *Commun. Math. Phys.* **94** 115–132.
- [42] Edwards, H., Hierons, R. and Jackson, B. (1998) The zero-free intervals for characteristic polynomials of matroids. *Combin. Probab. Comput.* **7** 153–165.
- [43] Edwards, R. G. and Sokal, A. D. (1988) Generalization of the Fortuin–Kasteleyn–Swendsen–Wang representation and Monte Carlo algorithm. *Phys. Rev. D* **38** 2009–2012.
- [44] Engelking, R. (1977) *General Topology*, PWN (Polish Scientific Publishers), Warszawa.
- [45] Farrell, E. J. (1980) Chromatic roots: Some observations and conjectures. *Discrete Math.* **29** 161–167.

- [46] Fortuin, C. M. and Kasteleyn, P. W. (1972) On the random-cluster model, I: Introduction and relation to other models. *Physica* **57** 536–564.
- [47] Garey, M. R. and Johnson, D. S. (1979) *Computers and Intractability: A Guide to the Theory of NP-Completeness*, Freeman, San Francisco.
- [48] Gebhard, D. D. and Sagan, B. E. (2000) Sinks in acyclic orientations of graphs. *J. Combin. Theory Ser. B* **80** 130–146.
- [49] Greene, C. and Zaslavsky, T. (1983) On the interpretation of Whitney numbers through arrangements of hyperplanes, zonotopes, non-Radon partitions, and orientations of graphs. *Trans. Amer. Math. Soc.* **280** 97–126.
- [50] Griffiths, R. B. (1972) Rigorous results and theorems. In *Phase Transitions and Critical Phenomena*, Vol. 1 (C. Domb and M. S. Green, eds), Academic Press, London/New York, pp. 7–109.
- [51] Griffiths, R. B. and Kaufman, M. (1982) Spin systems of hierarchical lattices. Introduction and thermodynamic limit. *Phys. Rev. B* **26** 5022–5032.
- [52] Grimmett, G. (1978) The rank polynomials of large random lattices. *J. London Math. Soc.* **18** 567–575.
- [53] Grimmett, G. (1995) The stochastic random-cluster process and the uniqueness of random-cluster measures. *Ann. Probab.* **23** 1461–1510.
- [54] Hall, D. W., Siry, J. W. and Vanderslice, B. R. (1965) The chromatic polynomial of the truncated icosahedron. *Proc. Amer. Math. Soc.* **16** 620–628.
- [55] Israel, R. B. (1979) *Convexity in the Theory of Lattice Gases*, Princeton University Press, Princeton, NJ.
- [56] Itzykson, C. and Luck, J. M. (1985) Zeroes of the partition function for statistical models on regular and hierarchical lattices. In *Critical Phenomena: 1983 Brasov School Conference* (V. Ceaşescu, G. Costache and V. Georgescu, eds), Birkhäuser, Boston/Basel/Stuttgart, pp. 45–82.
- [57] Jackson, B. (1993) A zero-free interval for chromatic polynomials of graphs. *Combin. Probab. Comput.* **2** 325–336.
- [58] Jackson, D. M. (1994) The lattice of noncrossing partitions and the Birkhoff–Lewis equations. *European J. Combin.* **15** 245–250.
- [59] Jacobsen, J. L. and Salas, J. (2001) Transfer matrices and partition-function zeros for antiferromagnetic Potts models, II: Extended results for square-lattice chromatic polynomial. *J. Statist. Phys.* **104** 701–723, cond-mat/0011456 at xxx.lanl.gov.
- [60] Jacobsen, J. L., Salas, J. and Sokal, A. D. (2003) Transfer matrices and partition-function zeros for antiferromagnetic Potts models, III: Triangular-lattice chromatic polynomial. *J. Statist. Phys.* **112** 921–1017, cond-mat/0204587 at xxx.lanl.gov.
- [61] Jaeger, F. (1991) Even subgraph expansions for the flow polynomial of cubic plane maps. *J. Combin. Theory Ser. B* **52** 259–273.
- [62] Kasteleyn, P. W. and Fortuin, C. M. (1969) Phase transitions in lattice systems with random local properties. *J. Phys. Soc. Japan* **26** (Suppl.) 11–14.
- [63] Kauffmann, L. and Saleur, H. (1993) An algebraic approach to the planar coloring problem. *Commun. Math. Phys.* **152** 565–590.
- [64] Lee, T. D. and Yang, C. N. (1952) Statistical theory of equations of state and phase transitions, II: Lattice gas and Ising model. *Phys. Rev.* **87** 410–419.
- [65] Lieb, E. H. and Sokal, A. D. (1981) A general Lee–Yang theorem for one-component and multicomponent ferromagnets. *Commun. Math. Phys.* **80** 153–179.
- [66] Maillard, J.-M. (1997) Hyperbolic Coxeter groups, symmetry group invariants for lattice models in statistical mechanics, and the Tutte–Beraha numbers. *Math. Comput. Modelling* **26** 169–225.
- [67] Maillard, J.-M. and Rollet, G. (1994) Hyperbolic Coxeter groups for triangular Potts models. *J. Phys. A* **27** 6963–6986.

- [68] Maillard, J.-M., Rollet, G. and Wu, F. Y. (1994) Inversion relations and symmetry groups for Potts models on the triangular lattice. *J. Phys. A* **27** 3373–3379.
- [69] Martin, P. P. (1987) The Potts model and the Beraha numbers. *J. Phys. A* **20** L399–L403.
- [70] Martin, P. P. (1989) Block spin transformations in the operator formulation of two-dimensional Potts models. *J. Phys. A* **22** 3991–4005.
- [71] Martin, P. P. (1991) *Potts Models and Related Problems in Statistical Mechanics*, World Scientific, Singapore.
- [72] Montel, P. (1927) *Leçons sur les Familles Normales de Fonctions Analytiques et leurs Applications*, Gauthier-Villars, Paris. Reprinted by Chelsea Publishing Co., New York (1974).
- [73] Noble, S. D. (1998) Evaluating the Tutte polynomial for graphs of bounded tree-width. *Combin. Probab. Comput.* **7** 307–321.
- [74] Ore, O. (1967) *The Four Color Problem*, Academic Press, New York.
- [75] Peitgen, H.-O., Prüfer, M. and Richter, P. H. (1985) Phase transitions and Julia sets. In *Lotka–Volterra-Approach to Cooperation and Competition in Dynamic Systems* (W. Ebeling and M. Peschel, eds), Akademie-Verlag, Berlin, pp. 81–102.
- [76] Peitgen, H.-O. and Richter, P. H. (1985) The Mandelbrot set in a model for phase transitions. In *Arbeitstagung Bonn 1984* (F. Hirzebruch, J. Schwermer and S. Suter, eds), Vol. 1111 of *Lecture Notes in Mathematics*, Springer, Berlin/Heidelberg/New York/Tokyo, pp. 111–134.
- [77] Peitgen, H.-O. and Richter, P. H. (1986) *The Beauty of Fractals: Images of Complex Dynamical Systems*, Springer, Berlin/Heidelberg/New York/Tokyo, pp. 45–49, 129–146, 194–195.
- [78] Peitgen, H.-O. and Richter, P. H. (1986) Fraktale und die Theorie der Phasenübergänge. *Physikalische Blätter* **42**(1) 9–22.
- [79] Polin, M. (2002) Private communication.
- [80] Potts, R. B. (1952) Some generalized order-disorder transformations. *Proc. Cambridge Philos. Soc.* **48** 106–109.
- [81] Read, R. C. (1968) An introduction to chromatic polynomials. *J. Combin. Theory* **4** 52–71.
- [82] Read, R. C. (2003) Chain polynomials of graphs. *Discrete Math.* **265** 213–235.
- [83] Read, R. C. and Royle, G. F. (1991) Chromatic roots of families of graphs. In *Graph Theory, Combinatorics, and Applications: Proceedings of the Sixth Quadrennial International Conference on the Theory and Applications of Graphs, Western Michigan University, 1988* Vol. 2 (Y. Alavi et al., eds), Wiley, New York, pp. 1009–1029.
- [84] Read, R. C. and Tutte, W. T. (1988) Chromatic polynomials. In *Selected Topics in Graph Theory 3* (L. W. Beineke and R. J. Wilson, eds), Academic Press, London, pp. 15–42.
- [85] Read, R. C. and Whitehead, E. G. Jr. (1999) Chromatic polynomials of homeomorphism classes of graphs. *Discrete Math.* **204** 337–356.
- [86] Read, R. C. and Whitehead, E. G. Jr. (2002) The Tutte polynomial for homeomorphism classes of graphs. *Discrete Math.* **243** 267–272.
- [87] Remmert, R. (1991) *Theory of Complex Functions*, Springer, New York.
- [88] Robertson, N., Sanders, D. P., Seymour, P. D. and Thomas, R. (1997) The four-colour theorem. *J. Combin. Theory Ser. B* **70** 2–44.
- [89] Roček, M., Shrock, R. and Tsai, S.-H. (1998) Chromatic polynomials for families of strip graphs and their asymptotic limits. *Physica A* **252** 505–546, cond-mat/9712148 at xxx.lanl.gov.
- [90] Roček, M., Shrock, R. and Tsai, S.-H. (1998) Chromatic polynomials for $J(\prod H)$ strip graphs and their asymptotic limits. *Physica A* **259** 367–387, cond-mat/9807106 at xxx.lanl.gov.
- [91] Saaty, T. L. and Kainen, P. C. (1977) *The Four-Color Problem: Assaults and Conquest*, McGraw-Hill, New York.
- [92] Salas, J. and Sokal, A. D. (2001) Transfer matrices and partition-function zeros for antiferromagnetic Potts models, I: General theory and square-lattice chromatic polynomial. *J. Statist. Phys.* **104** 609–699, cond-mat/0004330 at xxx.lanl.gov.

- [93] Saleur, H. (1990) Zeroes of chromatic polynomials: A new approach to Beraha conjecture using quantum groups. *Commun. Math. Phys.* **132** 657–679.
- [94] Saleur, H. (1991) The antiferromagnetic Potts model in two dimensions: Berker–Kadanoff phase, antiferromagnetic transition, and the role of Beraha numbers. *Nucl. Phys. B* **360** 219–263.
- [95] Schiff, J. L. (1993) *Normal Families*, Springer, New York.
- [96] Seppäläinen, T. (1998) Entropy for translation-invariant random-cluster measures. *Ann. Probab.* **26** 1139–1178.
- [97] Shrock, R. (1999) $T = 0$ partition functions for Potts antiferromagnets on Möbius strips and effects of graph topology. *Phys. Lett. A* **261** 57–62, cond-mat/9908323 at xxx.lanl.gov.
- [98] Shrock, R. (2000) Exact Potts model partition functions on ladder graphs. *Physica A* **283** 388–446, cond-mat/0001389 at xxx.lanl.gov.
- [99] Shrock, R. (2001) Chromatic polynomials and their zeros and asymptotic limits for families of graphs. *17th British Combinatorial Conference (Canterbury, 1999)*, *Discrete Math.* **231** 421–446, cond-mat/9908387 at xxx.lanl.gov.
- [100] Shrock, R. and Tsai, S.-H. (1997) Asymptotic limits and zeros of chromatic polynomials and ground state entropy of Potts antiferromagnets. *Phys. Rev. E* **55** 5165–5178, cond-mat/9612249 at xxx.lanl.gov.
- [101] Shrock, R. and Tsai, S.-H. (1997) Families of graphs with chromatic zeros lying on circles. *Phys. Rev. E* **56** 1342–1345, cond-mat/9703249 at xxx.lanl.gov.
- [102] Shrock, R. and Tsai, S.-H. (1997) Families of graphs with $W_r(\{G\}, q)$ functions that are nonanalytic at $1/q = 0$. *Phys. Rev. E* **56** 3935–3943, cond-mat/9707096 at xxx.lanl.gov.
- [103] Shrock, R. and Tsai, S.-H. (1997) Lower bounds and series for the ground state entropy of the Potts antiferromagnet on Archimedean lattices and their duals. *Phys. Rev. E* **56** 4111–4124, cond-mat/9707306 at xxx.lanl.gov.
- [104] Shrock, R. and Tsai, S.-H. (1998) Ground state entropy of Potts antiferromagnets on homeomorphic families of strip graphs. *Physica A* **259** 315–348, cond-mat/9807105 at xxx.lanl.gov.
- [105] Shrock, R. and Tsai, S.-H. (1998) Ground state degeneracy of Potts antiferromagnets: Cases with noncompact W boundaries having multiple points at $1/q = 0$. *J. Phys. A* **31** 9641–9665, cond-mat/9810057 at xxx.lanl.gov.
- [106] Shrock, R. and Tsai, S.-H. (1999) Ground state degeneracy of Potts antiferromagnets: Homeomorphic classes with noncompact W boundaries. *Physica A* **265** 186–223, cond-mat/9811410 at xxx.lanl.gov.
- [107] Shrock, R. and Tsai, S.-H. (1999) Ground state entropy of the Potts antiferromagnet on cyclic strip graphs. *J. Phys. A* **32** L195–L200, cond-mat/9903233 at xxx.lanl.gov.
- [108] Shrock, R. and Tsai, S.-H. (1999) Ground-state degeneracy of Potts antiferromagnets on two-dimensional lattices: Approach using infinite cyclic strip graphs. *Phys. Rev. E* **60** 3512–3515, cond-mat/9910377 at xxx.lanl.gov.
- [109] Shrock, R. and Tsai, S.-H. (1999) Ground state entropy of Potts antiferromagnets on cyclic polygon chain graphs. *J. Phys. A* **32** 5053–5070, cond-mat/9905431 at xxx.lanl.gov.
- [110] Shrock, R. and Tsai, S.-H. (2000) Exact partition functions for Potts antiferromagnets on cyclic lattice strips. *Physica A* **275** 429–449, cond-mat/9907403 at xxx.lanl.gov.
- [111] Sokal, A. D. (2001) Bounds on the complex zeros of (di)chromatic polynomials and Potts-model partition functions. *Combin. Probab. Comput.* **10** 41–77, cond-mat/9904146 at xxx.lanl.gov.
- [112] Sokal, A. D. (2002) A remark on the roots of chromatic polynomials. In preparation.
- [113] Temperley, H. N. V. (1993) New representations of the Temperley–Lieb algebra with applications. In *Low-Dimensional Topology and Quantum Field Theory: Cambridge, 1992* (H. Osborn, ed.), Vol. 315 of *NATO Advanced Science Institutes Series B: Physics*, Plenum, New York, pp. 203–212.

- [114] Thomas, R. (1998) An update on the four-color theorem. *Notices Amer. Math. Soc.* **45** 848–859.
- [115] Thomassen, C. (1997) The zero-free intervals for chromatic polynomials of graphs. *Combin. Probab. Comput.* **6** 497–506.
- [116] Thomassen, C. (2000) Chromatic roots and hamiltonian paths. *J. Combin. Theory Ser. B* **80** 218–224.
- [117] Tsai, S.-H. (1998) End graph effects on chromatic polynomials for strip graphs of lattices and their asymptotic limits. *Physica A* **259** 349–366, cond-mat/9807107 at xxx.lanl.gov.
- [118] Tutte, W. T. (1947) A ring in graph theory. *Proc. Cambridge Philos. Soc.* **43** 26–40.
- [119] Tutte, W. T. (1954) A contribution to the theory of chromatic polynomials. *Canad. J. Math.* **6** 80–91.
- [120] Tutte, W. T. (1970) On chromatic polynomials and the golden ratio. *J. Combin. Theory* **9** 289–296.
- [121] Tutte, W. T. (1970) The golden ratio in the theory of chromatic polynomials. *Ann. New York Acad. Sci.* **175** 391–402.
- [122] Tutte, W. T. (1974) Chromatic sums for rooted planar triangulations, V: Special equations. *Canad. J. Math.* **26** 893–907.
- [123] Tutte, W. T. (1975) Chromials. In *Studies in Graph Theory* (D. R. Fulkerson, ed.), Part II: *Studies in Mathematics*, #12, Mathematical Association of America, Washington, pp. 361–377.
- [124] Tutte, W. T. (1982) Chromatic solutions. *Canad. J. Math.* **34** 741–758.
- [125] Tutte, W. T. (1982) Chromatic solutions, II. *Canad. J. Math.* **34** 952–960.
- [126] Tutte, W. T. (1993) The matrix of chromatic joins. *J. Combin. Theory Ser. B* **57** 269–288.
- [127] Valdes, J., Tarjan, R. E. and Lawler, E. L. (1982) The recognition of series parallel digraphs. *SIAM J. Comput.* **11** 298–313.
- [128] Wakelin, C. D. and Woodall, D. R. (1992) Chromatic polynomials, polygon trees, and outerplanar graphs. *J. Graph Theory* **16** 459–466.
- [129] Welsh, D. J. A. (1993) *Complexity: Knots, Colourings, and Counting*, Vol. 186 of *London Mathematical Society Lecture Note Series*, Cambridge University Press, Cambridge/New York.
- [130] Whitney, H. (1932) A logical expansion in mathematics. *Bull. Amer. Math. Soc.* **38** 572–579.
- [131] Whitney, H. (1932) On the coloring of graphs. *Ann. Math.* **33** 688–718.
- [132] Whitney, H. (1933) A set of topological invariants for graphs. *Amer. J. Math.* **55** 231–235.
- [133] Woodall, D. R. (1977) Zeros of chromatic polynomials. In *Combinatorial Surveys: Proceedings of the Sixth British Combinatorial Conference* (P. J. Cameron, ed.), Academic Press, London, pp. 199–223.
- [134] Woodall, D. R. (1992) An inequality for chromatic polynomials. *Discrete Math.* **101** 327–331.
- [135] Woodall, D. R. (1992) A zero-free interval for chromatic polynomials. *Discrete Math.* **101** 333–341.
- [136] Woodall, D. R. (1997) The largest real zero of the chromatic polynomial. *Discrete Math.* **172** 141–153.
- [137] Woodall, D. R. (2002) Tutte polynomial expansions for 2-separable graphs. *Discrete Math.* **247** 201–213.
- [138] Wu, F. Y. (1982) The Potts model. *Rev. Mod. Phys.* **54** 235–268. Erratum **55** (1983) 315.
- [139] Wu, F. Y. (1984) Potts model of magnetism (invited). *J. Appl. Phys.* **55** 2421–2425.
- [140] Wu, F. Y., King, C. and Lu, W. T. (1999) On the rooted Tutte polynomial. *Ann. Inst. Fourier (Grenoble)* **49** 1103–1114.
- [141] Yang, C. N. and Lee, T. D. (1952) Statistical theory of equations of state and phase transitions, I: Theory of condensation. *Phys. Rev.* **87** 404–409.
- [142] Yung Qin and Yang, Z. R. (1991) Diamond-type hierarchical lattices for the Potts antiferromagnet. *Phys. Rev. B* **43** 8576–8582.