

Supplementary table 1. Drop-out analysis comparing those not available to those who completed the hypomania symptoms questionnaire			
Predictors	Completed questionnaire N (%)	Did not complete questionnaire N (%)	Non-response versus response OR (95% CI)
Gender			
Female	2182 (30.2%)	5037 (69.8%)	[reference]
Male	1188 (15.6%)	6447 (84.4%)	0.43 (0.39, 0.46)
Birthweight			
>2499g	3020 (23.1%)	1063 (76.9%)	[reference]
<2500g	153 (18.8%)	659 (81.2%)	0.77 (0.65, 0.93)
Family adversity (Mean score)	3.42 (3.62)	4.24 (4.22)	0.95 (0.94, 0.96)
Childhood abuse			
No	2928 (26.9%)	7976 (73.1%)	[reference]
Yes	363 (26.7%)	996 (73.3%)	0.99 (0.87, 1.13)
Maternal hitting			
No	1615 (27.8%)	4184 (72.2%)	[reference]
Yes	1486 (27.5%)	3908 (72.5%)	0.99 (0.91, 1.07)
Maternal shouting			
No	1732 (28.6%)	4317 (71.4%)	[reference]
Yes	1369 (26.5%)	3789 (73.5%)	0.90 (0.83, 0.98)

OR=Odds Ratio; CI=Confidence Intervals; Significant results presented in bold type face. Family adversity assessed with the family adversity index which comprises 18 items (see Winsper et al. 2012 for more details).