

1. Ali, S., & Ripley, S. D. (1971). Handbook of the birds of India and Pakistan: together with those of Nepal, Sikkim, Bhutan and Ceylon. Vol. 6, Cuckoo-shrikes to babaxes: synopsis nos. 1064-1271, colour plates 65-72. Oxford University Press.
2. Ansari, M. A. R. (1957) Studies on ischnoceran Mallophaga infesting birds in the Panjab. In Proceedings of Seventh Pakistan Science Conference, pp. 42-62
3. Atkinson, C. T., Utzurum, R. C., Seamon, J. O., Savage, A. F., & Lapointe, D. A. (2006). Hematozoa of forest birds in American Samoa-evidence for a diverse, indigenous parasite fauna from the South Pacific. *Pacific Conservation Biology*, 12(3), 229-237.
4. Azin, F., Nosrati, SM., & Amini, H. (2008). Occurrence of Red-vented Bulbul *Pycnonotus cafer* on Kish Island, Northeastern Persian Gulf Iran. *Podoces* 3: 105-107.
5. Bates, J. H., Spotswood, E. N., & Russel, J. C. (2014). Foraging behavior and habitat partitioning in sympatric invasive birds in French Polynesia. *Notornis*, 61(1), 35-42.
6. Beckon, W. N. (1993). The effect of insularity on the diversity of land birds in the Fiji islands: implications for refuge design. *Oecologia*, 94, 318-329.
7. Berger, A. J. (1972). Hawaiian birdlife. University Press of Hawai'i. 1st edition. 270 pp.
8. Berger, A. J. (1975). Red-whiskered and Red-vented bulbuls on Oahu. *Elepaio*, 36(2), 16-19.
9. Berger, A. J. (1981). Hawaiian birdlife. University Press of Hawai'i. 2nd edition. 260 pp.
10. Bergman, D. L., Chandler, M. D., & Locklear, A. (2000). The economic impact of invasive species to wildlife services' cooperators. In L. Clark, J. Hone, J. Shrivik, K. VerCauteren, R. Watkins & J. Yoder (Eds.), *Human conflicts with wildlife: economic considerations* (pp. 169-178). Fort Collins, Colorado: National Wildlife Research.
11. Bhatt, D., & Kumar, A. (2001). Foraging ecology of red-vented bulbul *Pycnonotus cafer* in Haridwar, India. *Forktail*, 109-109.
12. Blanvillain, C., Salducci, J. M., Tutururai, G., & Maeura, M. (2003). Impact of introduced birds on the recovery of the Tahiti Flycatcher (*Pomarea nigra*), a critically endangered forest bird of Tahiti. *Biological Conservation*, 109(2), 197-205.
13. Blanvillain et all, 2013. Mission d'enquête sur la contamination de l'avifaune par salmonella enteritidis et d'autre pathogènes des volailles. C. Blanvillain, S. Saavedra and T. Withers. Rapport SOP/DIREN 23pp
14. Boughton, D. C., Boughton, R. B., & Volk, J. (1938). Avian hosts of the genus *Isospora* (Coccidiida). *Ohio Journal of Science*. 38(3):149-163
15. Brochier, B., Vangeluwe, D., & van den Berg, T. (2010). Alien invasive birds. *Rev. sci. tech. Off. int. Epiz.*, 29(2), 217-226.

16. Brooks, D. M. (2013). Ecology, behavior, and reproduction of an introduced population of Red-vented Bulbuls (*Pycnonotus cafer*) in Houston, Texas. *The Wilson Journal of Ornithology*, 125(4), 800-808.
17. Brown, J. H. (1989). Patterns, Modes and Extents of Invasions by Vertebrates. In J. A. Drake, H. A. Mooney, F. Di Castri, R. H. Groves, F. J. Kruger, M. Rejmanek & M. Williamson (Eds.), *Biological Invasions: a Global Perspective* (pp. 85-109). Chichester: John Wiley & Sons Ltd.
18. Bruner, A. (1979). Red-vented Bulbul now in Tahiti. *Elepaio*, 40(6), 92.
19. Caplong, P. & Barjon, F. (2010). Le Bulbul à ventre rouge, une star méconnue en Nouvelle Calédonie. *La Calédonie Agricole*. 123 :22-25
20. Carlson, E. A. (1974). The avifauna of Tonga. *Peace Corps Tonga*.
21. Clapp, R. B., & Sibley, F. C. (1966). Notes on the birds of Tutuila, American Samoa. *Notornis*, 13(3), 157-164.
22. Clunie, F. (1976). Red-vented Bulbul uses Vanikoro Broadbill nest. *Notornis*, 23(3), 263-263
23. Cummings, J. L., Mason, J. R., Otis, D. L., Davis, J. E., & Ohashi, T. J. (1994). Evaluation of methiocarb, ziram, and methyl anthranilate as bird repellents applied to dendrobium orchids. *Wildlife Society Bulletin*, 633-638.
24. Daigneault, A., Brown, P., & Australia, S. N. (2013, February). Invasive species management in the Pacific using survey data and benefit-cost analysis. In *57th Australian Agricultural and Resource Economic Conference*. Sydney, Australia.
25. Delacour, J., 1943a. A revision of the genera and species of the family Pyconotidae (Bulbuls).— *Zoologica* 28(1): 17-28
26. Dhondt. 1976a. Bird Notes From The Kingdom Of Tonga. *Notomis* 23:4-7
27. Dhondt, A. (1976). Bird Observations in Western Samoa. *Notornis* 23: 29-43.
28. Dhondt, A. A. (1977). Breeding and postnuptial molt of the red-vented bulbul in Western Samoa. *The Condor*, 79(2), 257-260.
29. Dickinson & Dekker (2002). A preliminary review of the Pyconotidae. *Zool. Verh. Leiden* 340
30. Dickinson et al. (2002) Types of the Pyconotidae. *Zool. Verh. Leiden* 340
31. Duncan, R. P., Bomford, M., Forsyth, D. M., & Conibear, L. (2001). High predictability in introduction outcomes and the geographical range size of introduced Australian birds: a role for climate. *Journal of Animal Ecology*, 70(4), 621-632.
32. John E. duPont. (1972). Notes from Western Samoa, including the description of a new parrot-finch (*Erythrura*). *The Wilson Bulletin*, 375-376.
33. Fox, L. (2011) Red vented bulbul can alter ecosystems. *The Maui News*, 01/09/2011, C6
34. Freifeld, H. B. (1999). Habitat relationships of forest birds on Tutuila Island, American Samoa. *Journal of Biogeography*, 26(6), 1191-1213.

35. Fuong, H., Keeley, K. N., Bulut, Y., & Blumstein, D. T. (2014). Heterospecific alarm call eavesdropping in nonvocal, white-bellied copper-striped skinks, *Emoia cyanura*. *Animal Behaviour*, 95, 129-135.
36. Gill B.J., Hunt G.R. & Sirgouant S. (1995). – Red-Vented Bulbul (*Pycnonotus Cafer*) In New Caledonia. *Notornis*, 42, 214-215
37. Gill, B.J. 1999. A myna increase - notes on introduced mynas (*Acridotheres*) and bulbuls (*Pycnonotus*) in Western Samoa. *Notornis* 46: 268-269.
38. Gorman, M. L. (1975). HABITATS OF THE LAND-BIRDS OF VITI LEVU, FIJI ISLANDS. *Ibis*, 117(2), 152-161.
39. Gregory, G. (2005). The birds of the State of Kuwait. George Gregory, 219 pp
40. Hannecart, F., & Letocart, Y. (1980). Oiseaux de Nouvelle Calédonie et des Loyautés. Editions Cardinalis.
41. Hirshfield, E., & King, R. (1992). The status of some escaped species of birds in Bahrain. *The Phoenix*, 9, 11-13.
42. Holyoak, D.T.; Thibault, J.C. 1984. Contribution à l'étude des oiseaux de Polynésie Orientale. Mémoires du Muséum National d'Histoire Naturelle Série A, Zoologie: 62-183.
43. Islam, K., & Williams, R. N. (2000). Red-vented Bulbul(*Pycnonotus cafer*) and Red-whiskered Bulbul(*Pycnonotus jocosus*). *The Birds of North America*, (520), 20.
44. Jarvi, S. I., Farias, M. E., Baker, H., Freifeld, H. B., Baker, P. E., Van Gelder, E., ... & Atkinson, C. T. (2003). Detection of avian malaria (*Plasmodium* spp.) in native land birds of American Samoa. *Conservation Genetics*, 4(5), 629-637.
45. Jennings, M. C. (2004). Breeding birds in Central Arabia 1978-2003. *Sandgrouse*, 26(1), 35-47.
46. Jennings, M. C. (2010). Atlas of the breeding birds of Arabia. *Fauna of Arabia*, 25, 1-751.
47. Khamis, A. (2010). Alien species in Bahrain: the overseen threats. *BNHS Newsletter*, 1, 20-26
48. Khan, M. A. R. (1993). The introduced but naturalized avifauna of the United Arab Emirates. *Journal of the Bombay Natural History Society*, 90(3), 437-445.
49. KUMSCHICK, S., BLACKBURN, T. M., & RICHARDSON, D. M. (2015). Managing alien bird species: Time to move beyond “100 of the worst” lists?. *Bird Conservation International*, 1-10.
50. Kumar, A. (2004). Acoustic communication in the Red-vented Bulbul *Pycnonotus cafer*. *Anais da Academia Brasileira de Ciências*, 76(2), 350-358.
51. Kumar, A. (2010). Communication value of displays and postures in Red-vented Bulbul *Pycnonotus cafer* (Aves: Pycnonotidae). *Journal of Threatened Taxa*, 2(6), 919-929
52. Lendon, A. (1952). Bulbuls in Melbourne. *Emu*, 52(1), 67-68.

53. Lockwood, J. L., Moulton, M. P., & Anderson, S. K. (1993). Morphological Assortment and the Assembly of Communities of Introduced Passeriforms on Oceanic Islands: Tahiti Versus Oahu. *The American Naturalist*, 141(3), 398-408.
54. Long, J. L. (1981). *Introduced birds of the world: The worldwide history, distribution and influence of birds introduced to New environments*. New York: Universe Books
55. Long, J. L., & Mawson, P. R. (1991). Species of introduced birds in mediterranean Australia. In R. Groves & F. di Castri (Eds.), *Biogeography of Mediterranean Invasions*: Cambridge Univeristy Press.
56. Loope, L. L., Howarth, F. G., Kraus, F., & Pratt, T. K. (2001). Newly emergent and future threats of alien species to pacific birds and ecosystems. *Studies in Avian Biology*, 22, 291-304.
57. Lowe, S., Browne, M., Boudjelas, S., De Poorter, M. (2000) 100 of the World's Worst Invasive Alien Species A selection from the Global Invasive Species Database. Published by The Invasive Species Specialist Group (ISSG) a specialist group of the Species Survival Commission (SSC) of the World Conservation Union (IUCN), 12pp. First published as special lift-out in Aliens 12, December 2000. Updated and reprinted version: November 2004.
58. Marathe, S. (1989). Fly-catching bulbuls. *Newsletter for Birdwatchers* 29 (9&10): 10–11.
59. Martin-Albarracin, V. L., Amico, G. C., Simberloff, D., & Nuñez, M. A. (2015). Impact of Non-Native Birds on Native Ecosystems: A Global Analysis. *PloS one*, 10(11).
60. HOBCROFT, I. A. W. & Hobcroft, D. (2005). The further spread of introduced birds in Samoa. *Notornis*, 52, 16-20.
61. McLain, D. K., Moulton, M. P., & Redfearn, T. P. (1995). Sexual selection and the risk of extinction of introduced birds on oceanic islands. *Oikos*, 74(1), 27-34.
62. Merlin, M. D., & Juvik, J. O. (1985). Bird protection in Western Samoa. *Oryx*, 19(2), 97-103.
63. Metzdorf N., & Brescia F. (2008) Impact De L'avifaune Et Des Roussettes Sur Les Productions Fruitières En Nouvelle-Calédonie. Etude Préliminaire. Etat Des Lieux et Recommandations. Institut Agronomique Néo-Caledonien. Rapport D'étude N°3/2008. 34 Pp. Unpublished Report.
64. MEYER, J.-Y. 1993. Dispersion de *Miconia calvescens* par les oiseaux dans les îles de la Société. Pp. 27-42 in Séminaire Manu, Connaissance et Protection des Oiseaux. Société d'Ornithologie de Polynésie, 9-10-12 novembre 1993, Punaauia, Tahiti.
65. Meyer, J.-Y. 1996. Status Of *Miconia Calvescens* (Melastomataceae), A Dominant Invasive Tree In The Society Islands (French Polynesia). *Pacific Science* 50(1):66-7
66. Monnet, C., Thibault, J. C., & Varney, A. (1993). Stability and changes during the twentieth century in the breeding landbirds of Tahiti (Polynesia). *Bird Conservation International*, 3(04), 261-280.
67. Moulton, M. P., & Pimm, S. L. (1983). The introduced Hawaiian avifauna: biogeographic evidence for competition. *The American Naturalist*, 121(5).

68. Moulton, M. P., & Pimm, S. L. (1987). Morphological assortment in introduced Hawaiian passerines. *Evolutionary Ecology*, 1(2), 113-124.
69. Moulton, M. P. (1985). Morphological similarity and coexistence of congeners: an experimental test with introduced Hawaiian birds. *Oikos*, 44(2), 301-305.
70. Moulton, M. P. (1993). The All-or-None Pattern in Introduced Hawaiian Passeriforms: The Role of Competition Sustained. *The American Naturalist*, 141(1), 105-119.
71. Moulton, M. P., & Scioli, M. E. T. (1986). Range Sizes and Abundances of Passerines Introduced to Oahu, Hawaii. *Journal of Biogeography*, 13(4), 339-344.
72. Moulton, M. P., & Sanderson, J. G. (1997). Predicting the fates of Passeriform introductions on oceanic islands. *Conservation Biology*, 11(2), 552-558.
73. Muensch, A. J., Leininger, P. D., Werth, D. E., Fawks, A. M., & Thomas, S. M. (2006). The Anoles of Coconut Island, Kane'ohe Bay, O'ahu, Hawai'i. *Iguana*, 13(3), 198-206.
74. Nation, B., Nation, H., & Hooper, A. (1997). Birds new to Qatar. *Sandgrouse*, 19, 56-62.
75. Parham, B. E. V. (1955). Birds as pests in Fiji. *Journal of Fijian Agriculture* 25 :9-14
76. Pedersen, T. & Aspinall, S. J. (compilers) 2015. EBRC Annotated Checklist of the birds of the United Arab Emirates.
77. Pernetta JC, Watling D. 1978. The introduced and native terrestrial vertebrates of Fiji. *Pac Sci* 32(3): 223-244.
78. Prajapati, S. H., Patel, C. D., Parmar, R. V. & Patel, M. I. 2011. Breeding performance of Red-vented Bulbul (*Pycnonotus cafer*). *Life Sciences Leaflets*. 11:298–304
79. Pratt, H. D., Bruner, P. L., & Berrett, D. G. (1987). A field guide to the birds of Hawaii and the tropical Pacific (p. 409). Princeton, NJ: Princeton University Press.
80. Price, R. D. (1975). The Menacanthus eurysternus complex (Mallophaga: Menoponidae) of the Passeriformes and Piciformes (Aves). *Annals of the Entomological Society of America*, 68(4), 617-622.
81. Pyle, R. L. (2002). Checklist of the birds of Hawaii. *Elepaio*, 62(6), 135-148.
82. Rana, R. S., Chand, J., & Patyal, S. K. (2014). Evaluation of Birds as Predators of Insect Pests Infesting Cauliflower. *Journal of Community Mobilization and Sustainable Development*, 9(1), 18-22.
83. Richardson, C. (1992). Escapes and introductions in the United Arab Emirates. *The Phoenix*, 9, 13-15.
84. Rinke, D. (1987). The avifauna of 'Eua and its off-shore Islet Kalau, Kingdom of Tonga. *Emu*, 87, 26-34.
85. Roy, R. N., & Guha, B. C. (1958). Production of experimental scurvy in a bird species. *Nature*, 182, 319

86. Saavedra, S. 2012. First control campaign for Common myna and Red vented bulbul on Tahiti. Internal Report, SOP Manu, 25 pp.
87. Saavedra, S. 2013. Second control campaign for Common myna and Red vented bulbul on Tahiti. Internal Report, SOP Manu, 15 pp.
88. Sherley, G. (2000). Invasive species in the Pacific: a technical review and draft regional strategy. Samoa: South Pacific Regional Environment Programme.
89. Sherman, J. A., & Fall, P. L. (2010). Observations on feeding frequencies among native and exotic birds and fruit bats at *Erythrina variegata* and *Dysoxylum* trees on American Samoa. Altered ecologies: fire, climate and human influence on terrestrial landscapes. Australian National University Press, Canberra, 101-116.
90. Simberloff, D., & Boecklen, W. (1991). Patterns of extinction in the introduced Hawaiian avifauna: a reexamination of the role of competition. *The American Naturalist*, 138(2), 300-327.
91. Slater, P. 1974. A Field Guide To Australian Birds: Passerines. Rigby Ltd, Adelaide, Australia.
92. Spotswood, E. N., Meyer, J. Y., & Bartolome, J. W. (2012). An invasive tree alters the structure of seed dispersal networks between birds and plants in French Polynesia. *Journal of Biogeography*, 39(11), 2007-2020.
93. Stimson, J., & Berman, M. (1990). Predator induced colour polymorphism in *Danaus plexippus* L.(Lepidoptera: Nymphalidae) in Hawaii. *Heredity*, 65(3), 401-406.
94. Stimson, J., & Kasuya, M. (2000). Decline in the frequency of the white morph of the monarch butterfly (*Danaus plexippus plexippus* L., Nymphalidae) on Oahu, Hawaii. *JOURNAL-LEPIDOPTERISTS SOCIETY*, 54(1), 29-32.
95. Stuart, C., & Stuart, T. (1999). Birds of Africa: from seabirds to seed-eaters. MIT Press.
96. Thouzeau-Fonseca C. (2013). Contribution à la gestion d'un oiseau introduit envahissant en Nouvelle-Calédonie, le bulbul à ventre rouge (*Pycnonotus cafer* L.) : étude des modalités de piégeage et du comportement alimentaire en milieu naturel. Nouvelle-Calédonie : IAC Production : Angers : Agrocampus Ouest, 2013 . - 67p.+annexes
97. Thibault, J. C., Martin, J. L., Penloup, A., & Meyer, J. Y. (2002). Understanding the decline and extinction of monarchs (Aves) in Polynesian Islands. *Biological Conservation*, 108(2), 161-174.
98. Trail, P. W. (1994). The phenology of rainforest plants on Tutuila Island, American Samoa. American Samoa Government Department of Marine and Wildlife Resources Biological Report Series, (58).
99. Trail, P. W. (1994). Distribution and status of mynas in American Samoa. *Elepaio*, 54(4).
100. Turbott, E. G. (1956). Bulbuls in Auckland. *Notornis*, 6(7), 185-192
101. Vander Velde, N. (2002). The Red-vented Bulbul has come to Micronesia. *Aliens*, 16, 13-14.

102. Vijayan, V. S. (1980). Breeding biology of bulbuls, *Pycnonotus cafer* and *Pycnonotus luteolus* (Class: Aves, Family: Pycnonotidae) with special reference to their ecological isolation. Journal of the Bombay Natural History Society, 75, 1090-1117.
103. Walker, R., (2008). The Red-vented Bulbul - « Superbird »? Elepaio,, 68 (9), pp. 73-74.
104. Watling, D. (1977). The ecology of the Red-vented Bulbul in Fiji. Thesis. University of Cambridge, Cambridge, United Kingdom. (en cours d'obtention)
105. Watling, D. (1978). Observations on the naturalised distribution of the red-vented bulbul in the pacific, with special reference to the fiji islands. Notornis 25 : 109-117.
106. Watling, D. (1979). Bulbul gets a clean bill. New Scientist, 81(1147), 963-965.
107. Watling, D. (1983). The breeding biology of the Red-vented Bulbul *Pycnonotus cafer* in Fiji. Emu, 83(3), 173-180.
108. Williams, R N. 1983. Bulbul introductions on Oahu. The Elepaio, 43, 89–90.
109. Williams, G. (2011). 100 Alien Invaders. Bradt Travel Guides, 2011, 160 pp
110. Williams, R. N., & Giddings, L. V. (1984). Differential range expansion and population growth of bulbuls in Hawaii. The Wilson Bulletin, 647-655.
111. Wood, C. A., & Wetmore, A. (1926). A collection of birds from the Fiji Islands. Ibis, 68(1), 91-136.
112. Zia, U., Ansari, M. S., Akhter, S., & Rakha, B. A. (2014). breeding biology of red vented bulbul (*pycnonotus cafer*) in the area of rawalpindi/islamabad. Journal of Animal and Plant Sciences, 24(2), 656-659.