



Deriving a clinical prediction rule to target sexual healthcare to women attending British General Practices



N.L. Edelman^{a,b,*}, J.A. Cassell^a, C.H. Mercer^c, S.A. Bremner^a, C.I. Jones^a, A. Gersten^d, R.O. deVisser^e

^a Brighton & Sussex Medical School, United Kingdom

^b University of Brighton, United Kingdom

^c University College London, United Kingdom

^d Pavilion General Practice, United Kingdom

^e University of Sussex, United Kingdom

ARTICLE INFO

Keywords:

Primary care
Sexually transmitted infections
Women
Contraception
Sexual behaviour
Primary prevention
Sexual healthcare
Reproductive healthcare

ABSTRACT

Some women attending General Practices (GPs) are at higher risk of unintended pregnancy (RUIP) and sexually transmitted infections (STI) than others. A clinical prediction rule (CPR) may help target resources using psychosocial questions as an acceptable, effective means of assessment. The aim was to derive a CPR that discriminates women who would benefit from sexual health discussion and intervention.

Participants were recruited to a cross-sectional survey from six GPs in a city in South-East England in 2016. On arrival, female patients aged 16–44 years were invited to complete a questionnaire that addressed psychosocial factors, and the following self-reported outcomes: 2+ sexual partners in the last year (2PP) and RUIP. For each sexual risk, psychosocial questions were retained from logistic regression modelling which best discriminated women at risk using the C-statistic. Sensitivity and specificity were established in consultation with GP staff.

The final sample comprised $N = 1238$ women. 2PP was predicted by 11 questions including age, binge-drinking weekly, ever having a partner who insulted you often, current smoking, and not cohabiting (C-statistic = 0.83, sensitivity = 73% and specificity = 77%). RUIP was predicted by 5 questions including sexual debut < 16 years, and emergency contraception use in the last 6 months (C-statistic = 0.70, sensitivity = 69% and specificity = 57%).

2PP was better discriminated than RUIP but neither to a clinically-useful degree. The finding that different psychosocial factors predicted each outcome has implications for prevention strategies. Further research should investigate causal links between psychosocial factors and sexual risk.

1. Introduction

In Britain, General Practices (GP) act as the gatekeeper to specialist secondary services, and are staffed by Practice Nurses and General Practitioners (akin to Family Physicians in the United States). In England 58,969,634 people (approximately 90% of the resident population) are currently registered with a GP (NHS Digital, 2018) from whom they can also obtain a range of interventions directly. Britain's sexual health guidance and policy (DH, 2001, 2013) recommends GPs as sites for provision of sexual health interventions. This widens the availability of testing for sexually transmitted infections (STIs) and contraception; to a broad population who are likely to vary in need for those interventions to a greater degree than those attending specialist

contraception and sexual health (CASH) services (Cassell et al., 2006). Without an evidence-based means of targeting these interventions to women in GPs, resources may be wasted offering interventions unnecessarily (Fairley, 2016). Opportunities may also be missed to offer STI testing and contraception to high-risk individuals presenting for unrelated problems.

Within CASH clinics, a sexual history is the standard approach to determining appropriate intervention. This is resource-intensive during GP appointments for unrelated concerns and may feel unacceptable to some GP attenders (Edelman et al., 2013; Define, 2008), leading to possible under-report (DiClemente, 2016). Guidance recommends STI screening in GP settings only for target populations such as men who have sex with men, and for symptomatic individuals. The absence of

* Corresponding author at: University of Brighton, United Kingdom.
E-mail address: N.Edelman@brighton.ac.uk (N.L. Edelman).

Table 1
Candidate predictors (prospective CPR items).

Item wording	Response options	Source
How old are you?	<ul style="list-style-type: none"> • 15 years or less • Between 16 and 24 years • Between 25 and 34 years • Between 35 and 44 years • 45 years or older 	Natsal-3 wording, categories developed by authors
Thinking about where you are living now, which statement best describes your circumstances?	<ul style="list-style-type: none"> • I am renting or living rent-free (including living with parents or staying with friends) • I own my own home (including mortgage, shared ownership or bought outright) 	Natsal-3 response options with different piloted question wording
Did you live more or less continuously with <u>both of your natural (birth parents)</u> at home until you were 14?	<ul style="list-style-type: none"> • Yes • No • Prefer not to answer 	Natsal-3 with underline instead of lengthy explanatory text
How often is each of the following kinds of support available to you if you need it: Someone to help if you're confined to bed Someone to take you to the doctor if you need it Someone to prepare your meals if you're unable to do it yourself Someone to help with daily chores if you're sick	<ul style="list-style-type: none"> • None of the time • A little of the time • Some of the time • Most of the time • All of the time 	Canadian Community Health Survey (systematic review)
To what extent is the statement 'I have high self-esteem' true for you	<ul style="list-style-type: none"> • Not very true of me • Somewhat untrue of me • Neither untrue nor true of me • Somewhat true of me • Very true of me 	Robins' Single Item Self-Esteem Scale (SISE)
In the last 12 months have you received treatment from a health professional for depression?	<ul style="list-style-type: none"> • Yes • No • Prefer not to answer 	Natsal-3 with slightly adapted wording (as this was presented in a card)
How strongly do you agree with the statement 'Having a partner at all times is important to me'?	<ul style="list-style-type: none"> • Strongly agree • Agree • Disagree • Strongly disagree 	Atlanta Centre for Disease Control (CDC) study
How often do you have 6 or more units of alcohol on one occasion?	<ul style="list-style-type: none"> • Daily or almost daily • Weekly or almost weekly • Monthly • Less than monthly • Never • Prefer not to answer 	Natsal-3 (also an item in the validated Fast Alcohol Screening Test)
Do you smoke cigarettes at all nowadays?	<ul style="list-style-type: none"> • Yes I smoke cigarettes or roll-ups • Yes I smoke e-cigarettes • No • Prefer not to answer 	Natsal-3 with additional e-cigarette category developed following PPI
Have you <u>ever</u> taken any non-prescribed, illicit or illegal drugs, including legal highs?	<ul style="list-style-type: none"> • Yes • No • Don't know • Prefer not to answer 	Natsal-3 with wording adapted to specify that legal highs included following PPI
At present are you....	<ul style="list-style-type: none"> • Living as a couple with a partner or spouse • In a steady relationship but not living together • In a casual relationship • Single • Prefer not to answer 	Natsal-3
Please rate how emotionally satisfying your current relationship is, or how emotionally satisfying your most recent relationship was if you are currently single	<ul style="list-style-type: none"> • Extremely satisfying • Very satisfying • Moderately satisfying • Slightly satisfying • Not at all satisfying • Prefer not to answer 	Testa et al., 2005 (not systematic review)
Please indicate how strongly you agree with the following statement: My partner tells me who I can spend time with My partner does what he wants even if I don't want him to	<ul style="list-style-type: none"> • Strongly agree • Agree • Disagree • Strongly disagree • Prefer not to answer 	Atlanta CDC study
<u>During</u> your current or most recent relationship did your partner ever have sexual intercourse with anyone besides you	<ul style="list-style-type: none"> • No definitely not • I don't think so • It's quite likely • Yes, definitely • Prefer not to answer 	Testa et al., 2005 (not systematic review)
Have you ever been in a relationship with a partner who... Insulted or talked down to you often? Shouted or swore at you often? Threatened you with harm sometimes? Physically hurt you sometimes?	<ul style="list-style-type: none"> • Yes • No • Prefer not to answer 	HITS domestic violence tool (GP in place of Australian Women's Health Survey tool in systematic review)

(continued on next page)

Table 1 (continued)

Item wording	Response options	Source
How old were you when you <i>first</i> had sexual intercourse with someone of the opposite sex (including experiences you may not have wanted or that happened at an early age)?	<ul style="list-style-type: none"> • Under 16 years old • 16 years or older • I've never had sexual intercourse with someone of the opposite sex • Prefer not to answer 	Natsal-3
The man I most recently had sex with is 5 or more years older than me	<ul style="list-style-type: none"> • True • Probably true • I have no idea • Probably not true • Not true 	Natsal-3
In the last six months have you used emergency contraception at all?	<ul style="list-style-type: none"> • Yes • No • Prefer not to answer 	Adapted from the Contessa study
In the last six months have you taken a pregnancy test because you thought you might be pregnant?	<ul style="list-style-type: none"> • Yes • No • Prefer not to answer 	Adapted from the Contessa study

incentives for contraception and chlamydia screening in Britain and elsewhere may further demotivate opportunistic questioning by GP staff, who may struggle to initiate sexual health discussions (Wakley, 2005; Cassell et al., 2013). Interventions to address this have had limited impact (Macdowall et al., 2010; Town et al., 2015). Together this evidence suggests that a brief, acceptable, bespoke tool to target sexual healthcare could benefit GP staff and patients.

Clinical prediction rules (CPRs) identify risk of current or future adverse outcomes in individuals (Armstrong and Eborall, 2012) using several patient characteristics (Falasinnu et al., 2014a), to inform decisions about whether to offer interventions (Adams and Leveson, 2012). Many existing sexual health risk assessment tools and risk scores are CPRs (Falasinnu et al., 2014b). Most comprise sexual behavioural and socio-demographic 'known factors' (Haukoos et al., 2012; Gaydos et al., 2015) and focus on STI risk (Falasinnu et al., 2014b). However, no CPRs have been developed to identify women at risk of STIs and/or risk of unintended pregnancy (RUIP) in primary care, and few have used psychosocial questions (Falasinnu et al., 2014b).

Psychosocial factors are increasingly being explored alongside socio-demographic and sexual-behavioural predictors of sexual health, reflecting growing recognition of the need to address the social determinants of sexual health (DH, 2013). A systematic review and survey analysis identified several psychosocial factors associated with sexual risk in general population surveys of women, which may be usefully deployed as questions in a CPR for women attending GPs (Edelman et al., 2015; Edelman et al., 2017). That work suggests a psychosocial CPR should focus on identifying women experiencing recent potential risk of:

- STI acquisition through multiple partnerships in the last year (the primary outcome in this study)
- RUIP (desire to avoid pregnancy and not consistently using contraception in last six months)
- STI acquisition (through a potentially infected partner).

The latter outcome was included in the hope of identifying a novel population at risk through their partner's behaviour (Mittal et al., 2012). Re-infection from the same partner is believed to be a key factor in the re-infection rates for Chlamydia trachomatis, estimated at 20% (cumulative risk) (Walker et al., 2012). Together these outcomes facilitate CPR use for primary prevention (contraception and promotion of condom use) and for secondary prevention (pregnancy and STI testing and treatment).

The aim of this study was to identify the best combination of psychosocial questions to form an acceptable CPR to target sexual health intervention in General Practices to women of reproductive age. To do this we addressed the following research question: What combination of

acceptable psychosocial and socio-demographic questions discriminates best those women experiencing multiple partnerships, potential risk of STIs through partner and risk of unintended pregnancy?

2. Methods

We undertook a cross-sectional quantitative survey across GPs in a city in South-East England between April and September 2016. All female GP attenders aged 16–44 years were eligible to participate: we included women who did not report any male sexual partners in the last year so that we could assess the discriminatory power of the CPR.

2.1. Exposures

We identified psychosocial questions for inclusion in the survey questionnaire from a variety of sources based on the findings of preliminary studies (Edelman et al., 2015; Edelman et al., 2017; Robins et al., 2001; Gao and Chen, 2011; Raiford et al., 2009; Testa et al., 2005; Wellings et al., 2015; Sherin et al., 1998). These are listed in full in Table 1, and included questions on age group and housing tenure (renting or living rent-free versus home ownership) which remained predictive of multiple partnerships after adjustment for psychosocial questions in a preliminary study (Edelman et al., 2017). We chose psychosocial items that were brief and had few response options so that they would be easy to self-score and therefore suitable for self-completion in the CPR. We privileged items that were more common to ensure adequate prediction (i.e. applying the rationale that rarer exposures would lead to the identification of fewer women). We added a 'prefer not to answer' option to exposures that might be deemed unacceptable in order to measure this.

2.2. Outcomes

We developed a model for each of the following outcomes, which were designed to represent recent histories that would warrant sexual health discussion and possible intervention.

1. Report of 2+ male sexual partners in the last year (2PP) - indicating possible need for sexual health advice and STI testing. This outcome was measured using a single item 'In the last year, how many men have you had sexual intercourse with? *By sexual intercourse we mean a man's penis in a woman's vagina, mouth or anus*'.
2. Report of 2PP and/or risk of STI through a male partner (i.e. that the participant perceived that their most recent male sexual partner had had other sexual partners in the last year and had not always used condoms with those other partners).
3. Report of risk of unintended pregnancy in the last six months

(RUIP), indicating a possible need for contraception. At the time of development there was no suitable existing measure. Therefore we constructed a composite measure from an item measuring contraception use (Wellings et al., 2013a) and an item measuring desire to avoid pregnancy (Miller et al., 2013), adapting each to report retrospectively on the last six months.

These outcomes were limited to heterosexual experiences on the basis that women who only have sex with other women are at considerably lower risk for STI acquisition than other women (Everett, 2013). Nonetheless, the exclusion criteria did not incorporate women who self-identified as lesbian, as their exclusion from sexual health research is a growing concern as sexual health moves away from a disease-focused biomedical model (Wellings and Johnson, 2013). In addition, evidence suggests that some women who identify as lesbian may also report recent sexual activity with men (Everett, 2013) and may therefore experience the outcomes of interest.

2.3. Data collection

Data collection was designed to mimic envisaged delivery of the CPR – in which women self-complete and self-score the CPR during a clinic visit using a paper-and-pencil format. Patient and Public Involvement (PPI) was conducted by consultation with women attending a GP, a Women's Centre and a Youth Forum (who comment on a range of health services and research) to decide this delivery method and finalise the following approach to data collection.

On arrival, women attending GPs were offered an envelope by research or Reception staff, except visibly-distressed women, those known to have insufficient English language skills, or those who were clearly outside of the eligible age-range. Each envelope contained a pen, participant information sheet (PIS) and a brief questionnaire, comprising potential CPR items and the outcomes of interest. The questionnaire was designed to take 5 minutes to complete, while awaiting an appointment. This was deemed feasible as a 2016 study of General Practices in the study location found a mean waiting time of 15.6 minutes (K. Maskell, personal communication, November 9, 2017). Participants were instructed to complete the questionnaire anonymously, sitting alone in the waiting area if possible. Consent was implied by completion of the questionnaire. Three questions on the front of the questionnaire were used to screen out ineligible patients (those who had completed the questionnaire previously and/or did not identify as female and/or were outside the eligible age-range were instructed by questionnaire text to seal and return their questionnaire without completing further). The questionnaire and PIS instructed participants to seal their questionnaire in the envelope before returning to staff to be securely returned to the researchers.

2.4. Data storage and management

Questionnaires were stored at Brighton & Sussex Medical School for the duration of the study. Data were double-entered by an external company and analysed in a statistical software dataset - Stata SE v13 (StataCorp, 2013). Accuracy checks were performed on a random 10% sample and anomalies checked and addressed individually by researchers.

2.5. Statistical analysis

Weighting techniques were not applied as pseudo-inclusion probabilities (Elliott and Valliant, 2017) (which approximate the likelihood of study inclusion) could not be calculated. Available case analysis was used because item non-response was low. Bivariate analyses were conducted between psychosocial questions and each outcome. To avoid over-fitting the models, these analyses were only used to exclude from model entry: 1. Exposures not associated with any outcomes and 2.

Exposures reported by < 10% of respondents. We selected which psychosocial variables to enter into each model and then generated three models, one for each of the outcomes listed above. We used backwards-stepwise multivariable logistic regression as an established methodology for CPR derivation (Adams and Leveson, 2012; Falasinnu et al., 2014c; Gotz et al., 2005).

To assess model performance as a potential set of CPR questions, we used the C-statistic to quantify how well the model discriminated between those with, or without, the sexual risk of interest. The amount of variance explained by each model was assessed using McFadden's pseudo-R squared and calibration using Hosmer-Lemeshow goodness-of-fit test. Bayesian Information Criteria (BIC) were used to compare the parsimony of models. Exposures were not manually removed from the models if the corresponding *p*-value was > 0.05 as CPR derivation is a process of estimation rather than hypothesis testing. This is a recognised statistical approach to clinical prediction modelling (King et al., 2016). For each finalised model the coefficients were used to generate a scoring system for the CPR (Sullivan et al., 2004). A cumulative CPR score was then calculated for each participant based on their survey responses. For each model, participants' scores were cross-tabulated against their self-report of that outcome to assess how many participants would be classified as false positive or false negatives using the derived scoring system. Optimal cut-off values for sensitivity and specificity were then selected in a group consultation with five General Practitioners at one of the participating recruitment sites.

A sample size of 1500 was set in order to achieve an anticipated 150 reporting 2PP based on previous analyses (Edelman et al., 2017), sufficient to generate a 10–12 item CPR using the 10 events-per-variable approach (Peduzzi et al., 1996). 2PP was chosen as the primary outcome of interest as a known broad indicator of sexual risk (Sonnenberg et al., 2013). We ceased data collection at *n* = 1200 when *n* = 150 reporting 2PP had been exceeded.

3. Results

The final sample comprised *n* = 1238 women. 21.7% (*n* = 269) were aged 16–24 years, 41.8% (*n* = 518) were aged 25–34 years, and 36.4% (*n* = 451) were aged 35–44 years. 69.4% (*n* = 859) reported that they were currently renting or living rent-free and 29.7% (*n* = 368) owned their own home. Fig. 1 presents completion rates. We were unable to estimate response rates or investigate reasons for non-participation. Low rates of missing data and 'prefer not to answer' responses indicated that the questions were highly acceptable. Those exposures without the latter category still captured missing data *n* ≤ 40, comparable with those exposures that did offer a 'prefer not to answer' option.

The final model identified for '2+ sexual partners in the last year' comprised 11 items (Table 2). The Variance Inflation Factor of 1.21 indicated no multi-collinearity. Therefore no candidate predictors for this model needed to be removed to avoid inflated standard errors. This generated reasonable discriminatory power of *C* = 0.83 but low amount of variance explained (McFadden's Pseudo *R*² = 0.27). Good model calibration was indicated by H-L = 5.15 (*p* = 0.74). Although statistical significance is not the primary concern of modelling for estimation purposes (as in this case), the majority of exposures demonstrated an independent statistically-significant association with the outcome modelled. Through GP consultation a cut-off score of nine or above (range = 0–18) was chosen. This afforded a sensitivity of 72.8% and specificity of 76.7%.

The model identified for 'combined risk through multiple partnerships or most recent partner' comprised 11 items (Table 3), similar to those in the model for 2PP. The Variance Inflation Factor was 1.89, therefore no candidate predictors for this model needed to be removed to improve the accuracy of coefficients. It generated reasonable discriminatory power of *C* = 0.79 but explained only a small amount of the variance in outcome values (McFadden's Pseudo *R*² = 0.21). H-




Fig. 1. Completion rates.

L = 3.85 ($p = 0.87$) indicated good model calibration and moderate to good level of discrimination (Alba et al., 2017), only slightly lower than 2PP alone. However, the BIC value was higher for this model (BIC = 912) than for 2PP (BIC = 762). Through GP consultation a cut-off score of six or above (range = 0–17) was chosen. This afforded a sensitivity of 79.2% and specificity of 62.9%.

The model identified for ‘risk of unintended pregnancy risk in the last 6 months’ (RUIP) is presented in Table 4. The model had VIF = 1.21 so that no exposures needed to be removed for reasons of multi-collinearity. The model contained only five predictors, within the allowable number for the entered exposures and events, and accordingly had a lower BIC score (BIC = 402) than the other models. Although it showed good calibration (H-L = 1.0, $p = 0.96$), the RUIP model demonstrated a lower amount of variance explained (Pseudo- $R^2 = 0.12$) and lower discriminatory power (C-statistic = 0.70) compared to the other models. Nonetheless, this indicates moderate discrimination (on the basis that 0.5 denotes random chance and 1.0 perfect discrimination) (Alba et al., 2017). Although statistical significance is not the primary concern of modelling for estimation purposes (as in this case), it is noteworthy that only two of the exposures demonstrated an independent statistically-significant association ($p < 0.05$) with the outcome modelled. Using the 5 items retained in the UIP model, a score was developed with range 0–11. A cut-off score of three or above optimised sensitivity and specificity, giving a sensitivity of 69.0% and specificity of 56.8% and yielding $n = 315$ false positives and $n = 52$ false negatives.

The stakeholder consultation indicated that the sensitivity and specificity afforded by the CPR scores were too low for ubiquitous use in General Practices. However, the group felt that further research was warranted to investigate targeted use.

Table 2
Questions retained in the model predicting 2+ male sexual partners in the last year.

Psychosocial questions (yes v no)	Coefficient (log-odds)	Standard error	z-Statistic	p value	95% confidence limits (lower, upper)
Age 16–24	0.68	0.23	2.93	0.003	0.22, 1.13
Living in rented accommodation	0.52	0.25	2.11	0.035	0.04, 1.00
Ever had a partner who insulted or swore at you often	0.34	0.20	1.68	0.094	−0.06, 0.73
Little or no help with chores	0.27	0.21	1.29	0.197	−0.14, 0.67
Tested for pregnancy in last 6 months	0.37	0.22	1.72	0.086	−0.05, 0.79
Reports good or high self-esteem	0.90	0.23	3.95	< 0.001	0.45, 1.34
Having a partner at all times is not important to me	0.42	0.21	2.01	0.045	0.01, 0.83
Binge-drinks weekly or more often	0.70	0.20	3.46	0.001	0.30, 1.10
Smokes cigarettes at all	0.45	0.22	2.08	0.038	0.02, 0.87
Ever used illegal or illicit drugs incl. legal highs	0.38	0.23	1.67	0.096	−0.07, 0.83
Not in a steady relationship	2.06	0.22	9.35	< 0.001	1.63, 2.50
Intercept	−4.39	0.40	−11.07	< 0.001	−5.16, −3.61

4. Discussion

Our findings demonstrate that a variety of psychosocial variables are associated with sexual risk among GP-attending women of reproductive age. The results indicate that it is possible to discriminate women attending General Practices who report experiencing multiple male partners and/or risk through partners, to a greater degree than those experiencing RUIP. The findings also suggest that RUIP is predicted by a different profile of psychosocial factors. However, the findings indicate that our CPR does not warrant further validation and evaluation for routine use in GP settings using the particular psychosocial questions that we tested, because the level of discrimination achieved is unlikely to be practical for decision-making in GP settings.

Interestingly, various tools for related issues are already validated for use in General Practices, and may already be used in practice despite large differences in sensitivity and specificity. For example the HITS (Hurt, Insulted, Threatened with harm, Screamed at) domestic violence screening tool has demonstrated 96% sensitivity and 91% specificity in a General Practice setting (Sherin et al., 1998). The AUDIT-C (Alcohol Use Disorders Identification Test of Consumption) demonstrated sensitivity of 73% and specificity of 91% among women attending General Practice (Bradley et al., 2007). This sensitivity was therefore akin to this CPR. However, the specificity of our CPR was much lower, raising concerns among our stakeholders about its routine use in GP environments. Nonetheless, it is important to note that the specificity of our CPR may be lower in the study than in practice, as women who have been sexually inactive in the last year are likely to decline its use.

Unlike many sexual health tools that are developed without preliminary investigation (Falasinnu et al., 2014b), our CPR included candidate predictor variables selected on the basis of two separate

Table 3

Questions retained in the model predicting 2+ male sexual partners in the last year OR most recent partner has not used condoms with previous partners in last year.

Psychosocial questions (yes v no)	Coefficient (log-odds)	Standard error	z-Statistic	p value	95% confidence limits (lower, upper)
Age 16–24	0.25	0.22	1.14	0.256	−0.18, 0.67
Living in rented accommodation	0.43	0.20	2.16	0.031	0.040, 0.82
Sexual debut (including unwanted) < 16 years of age	0.53	0.18	2.87	0.004	0.17, 0.89
Tested for pregnancy in last 6 months	0.48	0.19	2.60	0.009	0.12, 0.84
Ever used illegal or illicit drugs incl. legal highs	0.48	0.19	2.55	0.011	0.11, 0.85
Having a partner at all times is not important to me	0.51	0.18	2.89	0.004	0.16, 0.85
Little or no help to prepare meals	0.42	0.18	2.32	0.020	0.07, 0.77
Smokes cigarettes at all	0.62	0.20	3.03	0.002	0.22, 1.02
Has not had treatment for depression in the last year	0.30	0.20	1.47	0.143	−0.10, 0.69
Most recent partner 5+ years older	0.40	0.20	2.05	0.040	0.018, 0.79
Not in a steady relationship	2.01	0.23	8.82	< 0.001	1.56, 2.46
Intercept	−2.77	0.33	−8.53	< 0.001	−3.41, −2.14

preliminary studies of random probability surveys (Edelman et al., 2015; Edelman et al., 2017). This survey was conducted on the population for whom the CPR is intended, and in the same setting, using a similar paper-and-pencil self-completion approach. This improves the validity of the dataset and findings, although it is important to note that while the models demonstrate proof-of-concept, they are not validated or evaluated for use by this study.

This survey indicates that age group and housing tenure were socio-demographic factors independently associated with sexual risk, rather than being fully explained by psychosocial items. This mirrors previous studies (Edelman et al., 2017; Sheringham et al., 2013; Woodhall et al., 2015). Our findings concurred with previous research regarding a positive association between sexual risk/morbidity and early sexual debut (Edelman et al., 2017; Wellings et al., 2013b), binge-drinking (Wellings et al., 2013b; Xaverius et al., 2009), social support (Gao and Chen, 2011), drug use (Edelman et al., 2017; Wellings et al., 2013b; Cavazos-Rehg et al., 2011) and emotional dissatisfaction (Testa et al., 2005). However, we found opposite directions of association to previous studies examining self-esteem (Sterk et al., 2004) partner importance (Raiford et al., 2009) and treatment for depression (Edelman et al., 2017). The former two were studies of African-American women, which may account for the differences in direction of association. However the latter finding was from female participants in the National Survey of Sexual Attitudes and Lifestyles-3 (Natsal-3) who were aged 16–44 and who reported sexual activity in the last year. ‘Treatment for depression’ is a composite of both experiencing depression and accessing treatment for it, which may explain the different direction of association. Alternatively, our contradictory findings may reflect the use of a convenience sample from within an urban clinical setting in South-East England as described below.

The CPR was developed to meet patient delivery preferences and answered a recognised need for brief sexual health assessment using simple self-scoring (Duke et al., 2008). Low proportions of participants with missing data and reporting ‘prefer not to answer’ suggests that the questions asked were acceptable. However, if the CPR had been developed for digital delivery, e.g. as a computer-assisted self-interview, this may have resulted in even lower rates of missing data and potentially more accurate reporting of both outcomes and exposures (Richens et al., 2010). This may have enabled the CPR to achieve higher

sensitivity and specificity. Similarly, electronic delivery would have allowed for the development of a more complex and potentially more sensitive scoring system, using a larger number of response options.

In order to tailor the CPR towards primary prevention, the outcomes of interest in this study concerned recent sexual behaviour rather than morbidity. Thus, we could not verify self-report using more objective measures. Recall bias may also have occurred in the self-report of both exposures and outcomes, particularly as the study was cross-sectional and several items in the survey were unvalidated.

The study benefited from having a large sample, ensuring sufficient statistical power to undertake multivariable regression. However, using convenience sampling may have resulted in an under-representation of those reporting sexual risk behaviour and adjunct psychosocial issues, such that sample bias cannot be ruled out. Because brevity and self-completion were key concerns, we were unable to record immigrant or other minority status, and the findings may be biased towards those with stronger English language skills and towards those not experiencing current distress. Nonetheless, it is possible that those who were unable or chose not to participate in the study would be equally unlikely to self-complete the CPR in practice so our findings still anticipate the CPR's performance in practice.

The generalisability of the findings may also be limited by the particular demographics within the city where the study was undertaken. Compared with other geographical areas in the United Kingdom, the population is young, urban, wealthy, University-educated and ethnically homogenous (Brighton and Hove City Council, 2012). A 2016 survey of the city's General Practices attenders (K. Maskell, personal communication, November 9, 2017) found that 91.6% were White, 87.5% had English as their First Language, 42.4% were educated to degree level, and the mean age was 49.3 years (s.d. = 18.9).

Although low sensitivity and specificity suggest that wide-spread delivery of the CPR to all women of reproductive age is not advisable, the CPR may be worthy of further validation and evaluation for targeted use in GP settings to those where poor sexual health is suspected, particularly as a means of generating discussion where adjunct issues such as binge-drinking and intimate partner violence are also a potential cause for concern. In these scenarios the CPR may provide an efficient alternative to clinical decision-making based on either use of socio-demographic data (as a population health approach) or

Table 4

Questions retained in the model predicting risk of unintended pregnancy in the last six months.

Psychosocial questions (yes v no)	Coefficient (log-odds)	Standard error	z-Statistic	p value	95% confidence limits (lower, upper)
Sexual debut (including unwanted) < 16 years of age	0.38	0.30	1.29	0.196	−0.20, 0.96
Used emergency contraception in last 6 months	2.31	0.36	6.36	< 0.001	1.60, 3.02
Emotionally dissatisfied with current or most recent relationship	0.42	0.28	1.49	0.137	−0.13, 0.97
Has not had treatment for depression in the last year	0.58	0.40	1.45	0.147	−0.02, 1.36
Partner had other partners during relationship	0.72	0.49	1.49	0.137	−0.23, 1.68
Intercept	−2.78	0.42	−6.62	< 0.001	−3.60, −1.96

opportunistic, detailed sexual history (as an individual health approach).

5. Conclusion

Our findings contribute to the broader study of associations between social factors and sexual health. In particular, they indicate that a different set of issues are predictive of RUIP. Further research should take into account how differences in which populations are offered which interventions may affect predictors of sexual risk (e.g. young age may have dropped out of the RUIP model because young women are more likely to be fitted with long acting reversible contraception). Together these findings suggest that further research should be carried out to investigate the role of psychosocial factors both as causal factors in sexual risk and as a means of identifying and differentiating between those experiencing different types of sexual risk.

Funding

This is a summary of independent research funded by the National Institute for Health Research (NIHR)'s Doctoral Research Fellowship programme DRF-2013-06-004. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Ethical approval

Ethical approval for this study was granted by University of Sussex and by the National Research Ethics Service Ref 16/LO/0206.

Conflict of interests

There are no conflicts of interests to declare.

Acknowledgements

The authors wish to acknowledge all those who have helped with and contributed to the study. This includes attenders at Brighton Women's Centre and at Newhaven Youth Forum, the NIHR Clinical Research Network Kent, Surrey and Sussex, the site Principal Investigators: Dr. Jennifer Whetham, Dr. Abbey Gersten and Dr. Paul Deffley; and the staff and patients of all the participating sites: Pavilion Surgery, St Peter's Medical Centre, Brighton Health and Wellbeing Centre, Charter Medical Centre, Beaconsfield Medical Practice and Trinity Medical Centre (formerly Sackville).

References

- Adams, S.T., Leveson, S.H., 2012. Clinical prediction rules. *Br. Med. J.* 344. <http://dx.doi.org/10.1136/bmj.d8312>.
- Alba, A.C., Agoritsas, T., Walsh, M., Hanna, S., Iorio, A., Devereaux, P.J., et al., 2017. Discrimination and calibration of clinical prediction models: users' guides to the medical literature. *JAMA* 318 (14), 1377–1384. Epub 2017/10/20. <https://doi.org/10.1001/jama.2017.12126> (PubMed PMID: 29049590).
- Armstrong, N., Eborall, H., 2012. The sociology of medical screening: past, present and future: introduction: the sociology of medical screening. *Sociol. Health Illn.* 34 (2), 161–176. <http://dx.doi.org/10.1111/j.1467-9566.2011.01441.x>.
- Bradley, K.A., DeBenedetti, A.F., Volk, R.J., Williams, E.C., Frank, D., Kivlahan, D.R., 2007. AUDIT-C as a brief screen for alcohol misuse in primary care. *Alcohol. Clin. Exp. Res.* 31 (7), 1208–1217. Epub 2007/04/25. <https://doi.org/10.1111/j.1530-0277.2007.00403.x> (PubMed PMID: 17451397).
- Brighton & Hove City Council, 2012. Brighton & Hove 2001 Census Briefing. Brighton & Hove City Council, Brighton & Hove, UK.
- Cassell, J.A., Mercer, C.H., Fenton, K.A., Copas, A.J., Erens, B., Wellings, K., et al., 2006. A comparison of the population diagnosed with chlamydia in primary care with that diagnosed in sexual health clinics: implications for a national screening programme. *Public Health* 120 (10), 984–988. <http://dx.doi.org/10.1016/j.puhe.2006.05.025>.
- Cassell, J.A., Dodds, J., Lanza, S., Low, N., Roberts, T., Smith, H.E., et al., 2013. P3.385 Comparative experience and outcomes of clinic staff versus intensive researcher led recruitment to a sexual health intervention in UK primary care. *Sex. Transm. Infect.* 89 (Suppl. 1), A269-A70. <http://dx.doi.org/10.1136/sextrans-2013-051184.0838>.
- Cavazos-Rehg, P.A., Krauss, M.J., Spitznagel, E.L., Schootman, M., Cottler, L.B., Bierut, L.J., 2011. Number of sexual partners and associations with initiation and intensity of substance use. *AIDS Behav.* 15 (4), 869–874. <http://dx.doi.org/10.1007/s10461-010-9669-0>. PubMed PMID: 2011-08875-021. PMID. (First Author & Affiliation: Cavazos-Rehg, Patricia A).
- Define, 2008. Define Chlamydia Screening and Sexual Health Marketing - Young People Final Report. Define Research & Insight Ltd for Department of Health, London (Contract No.: Define Job Number: 1649/COI Reference: 285937).
- DH, 2001. The National Strategy for Sexual Health and HIV. Department of Health, London.
- DH, 2013. A Framework for Sexual Health Improvement in England. Department of Health, London.
- DiClemente, R.J., 2016. Validity of self-reported sexual behavior among adolescents: where do we go from Here? *AIDS Behav.* 20, S215-S7. <http://dx.doi.org/10.1007/s10461-015-1217-5>.
- Duke, N.N., Sieving, R.E., Pettingell, S.L., Skay, C.L., 2008. Associations between health screening questions and sexual risk behaviors in adolescent female clinic patients: identifying a brief question format to yield critical information. *Clin. Pediatr. (Phila.)* 47 (6), 564–572. Epub 2008/05/10. <https://doi.org/10.1177/0009922808314904> (PubMed PMID: 18467672).
- Edelman, N., Patel, H., Glasper, A., Bogen-Johnston, L., 2013. Understanding barriers to sexual health service access among substance-misusing women on the South East coast of England. *J. Fam. Plan. Reprod. Health Care* 39 (4), 258–263.
- Edelman, N.L., de Visser, R.O., Mercer, C.H., McCabe, L., Cassell, J.A., 2015. Targeting sexual health services in primary care: a systematic review of the psychosocial correlates of adverse sexual health outcomes reported in probability surveys of women of reproductive age. *Prev. Med.* 81, 345–356. <http://dx.doi.org/10.1016/j.pymed.2015.09.019>.
- Edelman, N., Cassell, J.A., de Visser, R., Prah, P., Mercer, C.H., 2017. Can psychosocial and socio-demographic questions help identify sexual risk among heterosexually-active women of reproductive age? Evidence from Britain's third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *BMC Public Health* 17 (1), 5. <http://dx.doi.org/10.1186/s12889-016-3918-8>.
- Elliott, M.R., Valliant, R., 2017. Inference for nonprobability samples. *Stat. Sci.* 32 (2), 249–264. <http://dx.doi.org/10.1214/16-STS598>.
- Everett, B.G., 2013. Sexual orientation disparities in sexually transmitted infections: examining the intersection between sexual identity and sexual behavior. *Arch. Sex. Behav.* 42 (2), 225–236. <http://dx.doi.org/10.1007/s10508-012-9902-1>. (PubMed PMID: 2011897472. Language: English. Entry Date: 20130705. Revision Date: 20130705. Publication Type: journal article).
- Fairley, C.K., 2016. STI Services From Three Continents: Past and Future. *International Union against Sexually Transmitted Infections, Marrakech, Morocco*.
- Falasinu, T., Gilbert, M., Gustafson, P., Shoveller, J., 2014a. Deriving and validating a risk estimation tool for screening asymptomatic chlamydia and gonorrhoea. *Sex. Transm. Dis.* 41 (12), 706–712. Epub 2015/01/13. <https://doi.org/10.1097/olq.0000000000000205> (PubMed PMID: 25581805).
- Falasinu, T., Gustafson, P., Hottes, T.S., Gilbert, M., Ogilvie, G., Shoveller, J., 2014b. A critical appraisal of risk models for predicting sexually transmitted infections. *Sex. Transm. Dis.* 41 (5), 321–330. Epub 2014/04/12. <https://doi.org/10.1097/olq.0000000000000120> (PubMed PMID: 24722388).
- Falasinu, T., Gilbert, M., Gustafson, P., Shoveller, J., 2014c. Deriving and validating a risk estimation tool for screening asymptomatic chlamydia and gonorrhoea. *Sex. Transm. Dis.* 41. <http://dx.doi.org/10.1097/olq.0000000000000205>.
- Gao, Y., Chen, Y., 2011. Social support associated with a reduced risk of sexually transmitted infection in Canadians. *J. Public Health* 19 (1), 49–56. <http://dx.doi.org/10.1007/s10389-010-0351-9>.
- Gaydos, C.A., Jett-Goheen, M., Barnes, M., Dize, L., Barnes, P., Hsieh, Y.-H., 2015. Use of a risk quiz to predict infection for sexually transmitted infections: a retrospective analysis of acceptability and positivity. *Sex. Transm. Infect.* <http://dx.doi.org/10.1136/sextrans-2015-052058>.
- Gotz, H.M., Bergen, J.E., Veldhuijzen, I.K., Broer, J., Hoebe, C.J., Steyerberg, E.W., et al., 2005. A prediction rule for selective screening of *Chlamydia trachomatis* infection. *Sex. Transm. Infect.* 81. <http://dx.doi.org/10.1136/sti.2004.010181>.
- Haukoos, J.S., Lyons, M.S., Lindsell, C.J., Hopkins, E., Bender, B., Rothman, R.E., et al., 2012. Derivation and validation of the Denver Human Immunodeficiency Virus (HIV) risk score for targeted HIV screening. *Am. J. Epidemiol.* 175 (8), 838–846. Epub 2012/03/21. <https://doi.org/10.1093/aje/kwr389> (PubMed PMID: 22431561; PubMed Central PMCID: PMC3390011).
- King, C., Mercer, C., Furegato, M., Mohammed, H., Copas, A., Shahmanesh, M., et al., 2016. O037 Predicting STI risk among people attending sexual health services: development of a triage tool targeting behavioural interventions among young people. *Sex. Transm. Infect.* 92 (Suppl. 1), A14-A. <http://dx.doi.org/10.1136/sextrans-2016-052718.36>.
- Macdowall, W., Parker, R., Nanchahal, K., Ford, C., Lowbury, R., Robinson, A., et al., 2010. 'Talking of sex': developing and piloting a sexual health communication tool for use in primary care. *Patient Educ. Couns.* 81 (3), 332–337. <http://dx.doi.org/10.1016/j.pec.2010.10.027>.
- Miller, W.B., Barber, J.S., Gatny, H.H., 2013. The effects of ambivalent fertility desires on pregnancy risk in young women in the USA. *Popul. Stud. J. Demogr.* 67 (1), 25–38. <http://dx.doi.org/10.1080/00324728.2012.738823>.
- Mittal, M., S, T.E., Carey, M.P., 2012. Intimate partner violence and condom use among women: does the information-motivation-behaviour skills model explain sexual risk behaviour? *AIDS. Behaviour* 16, 1011–1019.
- NHS Digital, 2018. Patients Registered at a GP Practice Online. NHS Digital cited 2018/22/02/2018. Available from. <https://digital.nhs.uk/patients-registered-at-a-GP-practice-GP-data-hub>.

- Peduzzi, P., Concato, J., Kemper, E., Holford, T.R., Feinstein, A.R., 1996. A simulation study of the number of events per variable in logistic regression analysis. *J. Clin. Epidemiol.* 49 (12), 1373–1379. [http://dx.doi.org/10.1016/S0895-4356\(96\)00236-3](http://dx.doi.org/10.1016/S0895-4356(96)00236-3).
- Raiford, J.L., DiClemente, R.J., Wingood, G.M., 2009. Effects of fear of abuse and possible STI acquisition on the sexual behavior of young African American women. *Am. J. Public Health* 99 (6), 1067–1071. <http://dx.doi.org/10.2105/AJPH.2007.131482>.
- Richens, J., Copas, A., Sadiq, S.T., Kingori, P., McCarthy, O., Jones, V., et al., 2010. A randomised controlled trial of computer-assisted interviewing in sexual health clinics. *Sex. Transm. Infect.* 86 (4), 310–314. <http://dx.doi.org/10.1136/sti.2010.043422>.
- Robins, R.W., Hendin, H.M., Trzesniewski, K.H., 2001. Measuring global self-esteem: construct validation of a single-item measure and the Rosenberg self-esteem scale. *Personal. Soc. Psychol. Bull.* 27 (2), 151–161. <http://dx.doi.org/10.1177/0146167201272002>.
- Sherin, K.M., Sinacore, J.M., Li, X.Q., Zitter, R.E., Shakil, A., 1998. HITS: a short domestic violence screening tool for use in a family practice setting. *Fam. Med.* 30 (7), 508–512 (Epub 1998/07/21. PubMed PMID: 9669164).
- Sheringham, J., Mann, S., Simms, I., Stafford, M., Hart, G.J., Raine, R., 2013. It matters what you measure: a systematic literature review examining whether young people in poorer socioeconomic circumstances are more at risk of chlamydia. *Sex. Transm. Infect.* 89, 175–180.
- Sonnenberg, P., Clifton, S., Beddows, S., Field, N., Soldan, K., Tanton, C., et al., 2013. Prevalence, risk factors, and uptake of interventions for sexually transmitted infections in Britain: findings from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). *Lancet* 382 (9907), 1795–1806.
- StataCorp., 2013. *Stata Statistical Software: Release. 13* StataCorp LP, College Station, TX.
- Sterk, C.E., Klein, H., Elifson, K.W., 2004. Self-esteem and “at risk” women: determinants and relevance to sexual and HIV-related risk behaviors. *Women Health* 40 (4), 75.
- Sullivan, L.M., Massaro, J.M., D’Agostino, R.B., 2004. Presentation of multivariate data for clinical use: the Framingham study risk score functions. *Stat. Med.* 23 (10), 1631–1660. <http://dx.doi.org/10.1002/sim.1742>.
- Testa, M., VanZile-Tamsen, C., Livingston, J.A., 2005. Childhood sexual abuse, relationship satisfaction, and sexual risk taking in a community sample of women. *J. Consult. Clin. Psychol.* 73 (6), 1116–1124. <http://dx.doi.org/10.1037/0022-006X.73.6.1116>.
- Town, K., Ricketts, E.J., Hartney, T., Dunbar, J.K., Nardone, A., Folkard, K.A., et al., 2015. Supporting general practices to provide sexual and reproductive health services: protocol for the 3Cs & HIV programme. *Public Health* 129 (9), 1244–1250. Epub 2015/08/19. <https://doi.org/10.1016/j.puhe.2015.07.016> (PubMed PMID: 26278476).
- Wakley, G., 2005. “Opening a can of worms”: GP and practice nurse barriers to talking about sexual health in primary care. *J. Fam. Plan. Reprod. Health Care* 31 (1), 70. <http://dx.doi.org/10.1783/0000000052973013>.
- Walker, J., Tabrizi, S.N., Fairley, C.K., Chen, M.Y., Bradshaw, C.S., Twin, J., et al., 2012. Chlamydia trachomatis incidence and re-infection among young women - behavioural and microbiological characteristics. *PLoS One* 7 (5), e37778. <http://dx.doi.org/10.1371/journal.pone.0037778>.
- Wellings, K., Johnson, A.M., 2013. Framing sexual health research: adopting a broader perspective. *Lancet* 382 (9907), 1759–1762.
- Wellings, K., Jones, K.G., Mercer, C.H., Tanton, C., Clifton, S., Datta, J., et al., 2013a. The prevalence of unplanned pregnancy and associated factors in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Lancet* 382. [http://dx.doi.org/10.1016/s0140-6736\(13\)62071-1](http://dx.doi.org/10.1016/s0140-6736(13)62071-1).
- Wellings, K., Jones, K.G., Mercer, C.H., Tanton, C., Clifton, S., Datta, J., et al., 2013b. The prevalence of unplanned pregnancy and associated factors in Britain: findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Lancet* 382 (9907), 1807–1816. [http://dx.doi.org/10.1016/S0140-6736\(13\)62071-1](http://dx.doi.org/10.1016/S0140-6736(13)62071-1). PubMed PMID: 2013-42963-029. PMID: 24286786. (First Author & Affiliation: Wellings, Kaye).
- Wellings, K., Brima, N., Sadler, K., Copas, A.J., McDaid, L., Mercer, C.H., et al., 2015. Stopping and switching contraceptive methods: findings from Contessa, a prospective longitudinal study of women of reproductive age in England. *Contraception* 91 (1), 57–66. <http://dx.doi.org/10.1016/j.contraception.2014.09.008>. (PubMed PMID: WOS:000347017900010).
- Woodhall, S.C., Soldan, K., Sonnenberg, P., Mercer, C.H., Clifton, S., Saunders, P., et al., 2015. Is chlamydia screening and testing in Britain reaching young adults at risk of infection? Findings from the third National Survey of Sexual Attitudes and Lifestyles (Natsal-3). *Sex. Transm. Infect.* <http://dx.doi.org/10.1136/sextrans-2015-052013>.
- Xaverius, P.K., Tenkku, L.E., Salas, J., 2009. Differences between women at higher and lower risk for an unintended pregnancy. *Womens Health Issues* 19 (5), 306–312. <http://dx.doi.org/10.1016/j.whi.2009.06.002>.