

Original Article

\*Equal contributors

†Corresponding author for statistical aspects

**Cite this article:** Kuipers J, Moffa G, Kuipers E, Freeman D, Bebbington P (2018). Links between psychotic and neurotic symptoms in the general population: an analysis of longitudinal British National Survey data using Directed Acyclic Graphs. *Psychological Medicine* **49**, 388–395. <https://doi.org/10.1017/S0033291718000879>

Received: 27 October 2017

Revised: 17 January 2018

Accepted: 20 March 2018

First published online: 29 May 2018

**Key words:**

Affective symptoms; alcohol; cannabis; directed acyclic graph analysis; hallucinations; longitudinal community survey; paranoia; psychosis; worry

**Author for correspondence:**

Paul Bebbington, E-mail: [p.bebbington@ucl.ac.uk](mailto:p.bebbington@ucl.ac.uk)

# Links between psychotic and neurotic symptoms in the general population: an analysis of longitudinal British National Survey data using Directed Acyclic Graphs

J. Kuipers<sup>1,\*</sup>, G. Moffa<sup>2,3,\*†</sup>, E. Kuipers<sup>4,5</sup>, D. Freeman<sup>6,7</sup> and P. Bebbington<sup>3</sup>

<sup>1</sup>D-BSSE, ETH Zurich, Basel, Switzerland; <sup>2</sup>Institute for Clinical Epidemiology and Biostatistics, University Hospital Basel and University of Basel, Basel, Switzerland; <sup>3</sup>Division of Psychiatry, University College London, London, UK; <sup>4</sup>Department of Psychology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, London, UK; <sup>5</sup>NIHR Biomedical Research Centre, South London and Maudsley NHS Foundation Trust, Beckenham, UK; <sup>6</sup>Department of Psychiatry, Warneford Hospital, University of Oxford, Oxford, UK and <sup>7</sup>Oxford Health NHS foundation Trust, Warneford Hospital, Oxford, UK

## Abstract

**Background.** Non-psychotic affective symptoms are important components of psychotic syndromes. They are frequent and are now thought to influence the emergence of paranoia and hallucinations. Evidence supporting this model of psychosis comes from recent cross-fertilising epidemiological and intervention studies. Epidemiological studies identify plausible targets for intervention but must be interpreted cautiously. Nevertheless, causal inference can be strengthened substantially using modern statistical methods.

**Methods.** Directed Acyclic Graphs were used in a dynamic Bayesian network approach to learn the overall dependence structure of chosen variables. DAG-based inference identifies the most likely directional links between multiple variables, thereby locating them in a putative causal cascade. We used initial and 18-month follow-up data from the 2000 British National Psychiatric Morbidity survey ( $N = 8580$  and  $N = 2406$ ).

**Results.** We analysed persecutory ideation, hallucinations, a range of affective symptoms and the effects of cannabis and problematic alcohol use. Worry was central to the links between symptoms, with plausible direct effects on insomnia, depressed mood and generalised anxiety, and recent cannabis use. Worry linked the other affective phenomena with paranoia. Hallucinations were connected only to worry and persecutory ideation. General anxiety, worry, sleep problems, and persecutory ideation were strongly self-predicting. Worry and persecutory ideation were connected over the 18-month interval in an apparent feedback loop.

**Conclusions.** These results have implications for understanding dynamic processes in psychosis and for targeting psychological interventions. The reciprocal influence of worry and paranoia implies that treating either symptom is likely to ameliorate the other.

## Introduction

The concept of causality is based on temporal relationships: causes must precede consequences. Some consequences may exist only as long as the cause applies, but in other cases, they persist after it stops, presumably because mediating processes have been recruited or modified. Sometimes the emergence of a consequence may modify its cause, either enhancing or reducing its effect. Such positive and negative feedback loops also have temporal attributes, and may be abrupt or leisurely.

The medical approach to disease defines disorders on the basis of concatenations of symptoms (syndromes), and then infers an underlying pathology (a diathesis): this, in turn, becomes the basis of causal explanation and the target of research. This approach is applied equally to psychiatric disorders, including psychosis. Current accounts of the origins of the psychotic diathesis acknowledge contributions from biological, social and psychological domains, which are seen as likely to act synergistically (Broome *et al.* 2005; Howes & Murray, 2014).

However, our understanding of psychiatric disorders has been amplified by the emergence of *interactional models*. These treat psychiatric symptoms as a network of individual interacting variables, exerting mutual causal influences without requiring the formulation of a consistent underlying pathology (Borsboom & Cramer, 2013; McNally *et al.* 2015; Isvoranu *et al.* 2016, 2017; McNally, 2016; Klippel *et al.* 2017; Fried *et al.* 2017; Murphy *et al.* 2017). The network then becomes the locus of causality. This allows the possibility that different causes may operate through different points of entry in the network, information that may have useful implications for treatment.

© Cambridge University Press 2018. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

We have long argued against the rigid distinction between psychotic and non-psychotic problems in psychiatric classifications and that it is important to model the relationships between psychotic, non-psychotic, and environmental variables. Non-psychotic experiences such as low self-esteem, worry, sleep disturbance, anxiety, and depression may be contributory causes of particular psychotic symptoms, especially paranoia (e.g. Kuipers *et al.* 2006; Bebbington *et al.* 2013; Freeman, 2016). Most researched has been the cross-sectional and longitudinal association of anxiety and depression with psychotic experiences (e.g. Krabbendam *et al.* 2004; Scott *et al.* 2009; Ben-Zeev *et al.* 2011; Thewissen *et al.* 2011), with a smaller number of studies indicating similar findings for low self-esteem, worry, and sleep disturbance (e.g. Freeman *et al.* 2012; Hartley *et al.* 2014; Sheaves *et al.* 2016). Damaging social experiences also increase the likelihood of a range of psychological symptoms, including psychotic experiences (e.g. Bebbington *et al.* 2011; Kelleher *et al.* 2013; McGrath *et al.* 2017). In particular, non-psychotic symptoms may modulate the relationship between adversity and psychotic experiences, as moderators and mediators (e.g. Gracie *et al.* 2007; Marwaha & Bebbington, 2015; Gibson *et al.* 2016). If so, this has clinical relevance, as such symptoms may then serve as significant candidate targets for psychological or pharmacological treatments. Moreover, while psychoactive substances such as cannabis may elicit paranoia by inducing anomalous perceptual experiences such as hallucinations directly, they may also operate through these common affective processes (Freeman *et al.* 2015a).

The gold standard for causal inference in putative causal systems is direct intervention, itself an attempt to capture the temporal element required for causal inference. Although they may have valuable practical implications, intervention studies can be expensive. They may also be difficult to implement, though there are new opportunities with digital interventions (Freeman *et al.* 2017). As a result, attempts have been made to optimise causal inference in datasets that do not involve interventions. Analysis of cross-sectional datasets using variants of logistic regression has been used to establish mediation (e.g. Marwaha *et al.* 2014, 2015; Catone *et al.* 2015). These approaches narrow the field of plausible candidate mediators, essentially by excluding variables that do not meet the criteria for mediation.

Nonetheless, there remains a due wariness about making positive causal inferences from cross-sectional data. However, we have argued elsewhere that such data capture participants at different stages of causal progressions; the time dependence remains to some extent encoded in the data, and is thus not completely lost. Different causal hypotheses would predict different patterns of relationship, each with different levels of support from the observed data. The task is then to decode the data, and this can be done to a significant extent through the use of advanced statistical methods. In particular, we have argued for the introduction in psychiatry of probabilistic graphical models in the form of Directed Acyclic Graphs (DAGs) to ascertain which causal patterns best fit cross-sectional datasets (Moffa *et al.* 2017).

DAGs are the representation of Bayesian networks and model the overall dependence structure of multiple variables. 'Edges' are directed links (lines with an arrow head) that connect 'nodes', the specific variables being analysed (these conventions are illustrated in the figure summarising our results). Nodes are described as the *parents* of those immediately below them, which are conversely referred to as *children*. Nodes that can be reached from another node by following a directed path are called its *descendants*. The connections between the nodes of a Bayesian network

represent conditional independence relationships between the represented variables. Each variable in a Bayesian network is independent of its non-descendants conditional on its parents. As a consequence, the joint probability distribution of the variables in the network decomposes as a product of the conditional distribution of each node, given its parent nodes in the graph.

Alongside the probabilistic interpretation of DAGs, a causal interpretation is also widespread. The logic behind using DAGs to depict hypothetical causal structures is set out in detail by Pearl (2009). In this interpretation, a direct edge in the graph is held to imply a causal relationship. A directed *pathway* from one variable to another also implies a causal relationship, though one operating through intermediary variables on the path.

Causal discovery through DAG remains controversial (Dawid, 2010). However, the probabilistic DAG-based inference is capable of pointing to the most likely directional links between multiple variables, thereby locating them in a putative causal cascade, in which upstream variables are potential causes of those downstream. Sometimes a given dataset is incapable of allowing the full specification of the direction of the edges of a DAG, thereby limiting insights about potential causal links between pairs of variables. These variables then need to be placed at the same level of the graph (i.e. neither upstream nor downstream). We recently applied this method to investigate mechanisms linking bullying with persecutory ideation and hallucinations, obtaining consistent findings with data from two British national surveys of psychiatric morbidity (Moffa *et al.* 2017).

While DAGs may be able to capture some of the time dependence between individual variables in cross-sectional data, they cannot provide information about the length of time over which putative causes have their effects. However, it may be possible to illuminate this using longitudinal studies, although these are constrained by the interval between assessments: if the interval is too short for a given cause to operate, or too long for it to be maintained, there may appear to be no relationship even where one exists. Nevertheless, applying DAG discovery to longitudinal data may illuminate the characteristics of specific causal relationships, in particular because it may allow the identification of the feedback loops likely in social/psychological systems. Bayesian networks are incapable of modelling feedback loops in single cross-sectional datasets; however such loops may be identifiable by subjecting time course data to analysis in terms of *dynamic Bayesian networks* (Friedman *et al.* 1998).

In the current study, we applied a dynamic Bayesian network approach for learning graphical structures to data from the 2000 British National Psychiatric Morbidity survey. This provided two equivalent waves, as it included follow-up interviews after 18 months with a stratified subsample. We took hallucinations and persecutory ideation as key symptoms of psychosis, and identified potential causal links between them and non-psychotic psychiatric symptoms both cross-sectionally and longitudinally.

## Survey methods

### Setting and design

The second National Psychiatric Morbidity Survey of adults living in private households in England, Scotland, and Wales was carried out in 2000 (Singleton *et al.* 2001). The initial assessment was based on a sample of 8580 participants, while a subsample of 2406 participants was reassessed 18 months later (Singleton and Lewis, 2003). The follow-up survey focused on participants

with disorders and those likely to develop them. Thus all people identified as having a mental disorder, and all others with six or more common mental disorder symptoms were re-interviewed, along with 20% of the remainder selected at random.

### Measures

The assessment of persecutory ideation was based on question PSQ3a in the Psychosis Screening Questionnaire (PSQ; Bebbington & Nayani, 1995): 'Have there been times when you felt that people were deliberately acting to harm you or your interests?' Auditory hallucinations were identified from question PSQ5a: 'Have you at any time heard voices saying quite a few words or sentences when there was no-one around that might account for it?' Both these items were assessed at baseline in relation to the past year, and over the intervening 18 months at follow up. At both time-points, the Clinical Interview Schedule-Revised (CIS-R; Lewis *et al.* 1992) was used to establish the presence of insomnia, generalised anxiety, situational anxiety, social anxiety, worry, and depressed mood over a consistent period lasting at least 2 weeks in the past month. Cannabis use, cannabis dependence, and hazardous alcohol use were also assessed at both interviews. Cannabis use was assessed in relation to the past month, and current cannabis dependence was identified on the basis of a positive answer to any of five questions characterising dependence in the Diagnostic Interview Schedule (Robins *et al.* 1981). Current hazardous alcohol use (a pattern of consumption carrying a risk of physical and psychological harm) was defined as a score of 8 or more on the Alcohol Use Disorders Identification Test (AUDIT; Saunders *et al.* 1993).

All variables were analysed in binary form, suitable for the DAG procedures employed. Data were weighted back to the first stage sample, using a weighting variable that took account both of the sampling strategy and of refusals.

### Statistical methods

Given the longitudinal nature of the data, we employed dynamic Bayesian networks to capture the time component and to describe the dynamic evolution of the variables (Friedman *et al.* 1998). Such networks model the overall dependence structure of multiple variables, measured over different time points, and visualised in DAGs. *Nodes* representing the specific variables being analysed are connected by directed *edges* to form a network (see Moffa *et al.* 2017 for a more detailed description). In dynamic Bayesian networks, different waves of data collection are referred to as *time slices*: we then have a node for each variable in each time slice. Nodes may be connected within a time slice, but may also be linked to variables in the follow-up time slice, thereby forming a larger Bayesian network incorporating measurements of all variable over time. The connections are assumed to be stationary over time and identical across time slices, implying that the way symptoms relate to each other does not change with time. Under this assumption, the problem of inferring the network that describes the observed data is essentially reduced to learning a network of size equal to the number of variables considered at each time point. However, for each variable in the follow-up time slice, we also need to learn the set of parents from the previous time slice (i.e. the variables from the previous time point with a direct edge into the variable under consideration, for which no conditional independence is found to hold). Analysing the same connections at different time points enables

us to characterise the networks describing both time slices, together with the relationships between the time slices. A feedback loop may be tentatively inferred if a variable A at time 1 affects a variable B at time 2, and B at time 1 affects A at time 2.

In the current analysis, the discovery of the DAG relied on an appropriately modified version of our partition MCMC method (Kuipers & Moffa, 2017), a novel Bayesian method for graphical structure learning. The range of causal effects was also predicted in a Bayesian fashion, as implemented by Moffa *et al.* (2017) for binary variables. The main underlying assumptions are that the variables are causally linked, and that the causal relationship can be represented by a causal DAG. The method has two special features. First, it quantifies both the *strength* and the *direction* of causal effects. Second, since several different graphical arrangements of variables may each be capable of explaining the data reasonably well, the uncertainty in the inference is captured by sampling from the range of possible DAGs in proportion to their posterior distribution. In total, we sampled 10 000 DAGs, such that better fitting graphs were chosen more often.

Each of the DAGs sampled from the posterior distribution provides a model of the data, and from this, we can derive an estimate of the probability of each variable taking the value 0 or 1 depending on the state of its parents in the network. The individual DAGs allow us to calculate the effect of one variable on another. The value of one variable is set to 0, and the probability that each remaining variable is 0 or 1 can then be calculated. The selected variable is then set to 1 and the process is repeated. In Pearl's terms, the difference between the two sets of values constitutes the causal effect of the selected variable on the others (Pearl, 2009). This procedure is followed in turn for each variable in the DAG. By averaging the estimated effects over all the DAGs in the sample we can then obtain the overall effect of changing each variable on all of the others. This enables us to identify the most plausible causal mechanisms. Further details of the method in action are given by Moffa *et al.* (2017). In addition, we improved the characterization of the distribution of intervention effects by taking into account the uncertainty in the parameter estimation of each DAG to provide the full posterior distribution.

### Results

The correlations between variables assessed initially and at follow up are displayed in Table 1, while the results of the DAG discovery are shown in Figs 1 and 2.

In Fig. 1 we show the distribution of causal effects of the Time 1 variables (the row variables) on the Time 2 variables (the column variables). In cases where there was no effect between two variables, we entered a zero in the relevant box. In all boxes, most of the probability mass of the effect distribution is within the range  $-0.1$  to  $0.75$  and so we truncated the graphs accordingly. The red vertical line in each box indicates zero causal effect. The whole box is coloured to highlight cases where the 95% credible interval (the Bayesian counterpart of confidence limits) does not cover the point corresponding to zero causal effect. The number reported across the boxes quantifies the relevant *average causal effect*.

In Fig. 2 we present a consensus graph generated by this analysis. In order to focus on the stronger effects, edges are only shown in the figures if they are present in at least 10% of the sampled DAGs. Note that the graphical structure within each time slice is identical, as this is imposed by the dynamic Bayesian network framework. This restriction then enabled us

Table 1. Correlation of variables at each time point

Time 1	Time 2										
	Persecutory ideation	Worry	Hallucination	Sleep	Depression	General anxiety	Situational anxiety	Social anxiety	Cannabis last month	Cannabis dependence	Alcohol problems
<b>Persecutory ideation</b>	0.31	0.23	0.05	0.14	0.15	0.19	0.14	0.11	0.10	0.10	0.18
<b>Worry</b>	0.19	0.35	0.08	0.21	0.22	0.26	0.12	0.12	0.06	0.02	0.06
<b>Hallucination</b>	0.10	0.07	0.29	0.05	0.08	0.06	0.09	0.11	0.003	-0.003	0.04
<b>Sleep</b>	0.10	0.24	0.07	0.36	0.16	0.19	0.09	0.08	0.04	0.04	0.03
<b>Depression</b>	0.14	0.22	0.08	0.18	0.21	0.21	0.13	0.10	0.07	0.08	0.08
<b>General anxiety</b>	0.15	0.23	0.05	0.16	0.20	0.26	0.12	0.14	0.05	0.02	0.06
<b>Situational anxiety</b>	-0.004	-0.04	-0.01	-0.01	0.00	-0.03	-0.01	-0.01	0.04	0.001	0.02
<b>Social anxiety</b>	0.02	-0.03	-0.01	0.003	-0.01	-0.004	-0.01	-0.01	0.01	0.02	-0.01
<b>Cannabis last month</b>	0.07	0.07	-0.02	0.04	0.03	0.05	0.06	0.04	0.65	0.53	0.26
<b>Cannabis dependence</b>	0.09	0.07	-0.01	0.01	0.03	0.06	0.06	0.07	0.53	0.58	0.17
<b>Alcohol problems</b>	0.07	0.04	0.04	-0.001	0.001	0.06	0.02	0.03	0.22	0.17	0.52

to quantify the links between the initial and follow up symptoms. The density of the arrows joining the variables represents the strength of the links. The variables are either joined by single arrows, indicating a plausible causal link, or by arrows in both directions, which implies that causal influences in either direction are compatible with the data (this corresponds to the presence of bimodal peaks in the plots in Fig. 1).

We will first consider the cross-sectional relationships apparent *within* the two waves of data (time slices). Worry appears to have a central role in the links between symptoms. It has plausible direct effects on insomnia, depressed mood and generalised anxiety, and on past-month cannabis use. Via cannabis use, it appears to be linked both to cannabis dependence and to alcohol problems. The relationship between persecutory ideation and worry is indeterminate, consistent with the analyses of Moffa *et al.* (2017). Finally, worry is weakly determined by social anxiety and hallucinations.

The relationship between social anxiety and situational anxiety is also indeterminate. While generalised anxiety is clearly upstream of situational anxiety, its link with depressed mood is undetermined. Insomnia is downstream of depressed mood and also of worry. The relationship between persecutory ideation and auditory hallucinations is indeterminate. Hallucinations have no direct link with any other variable, apart from a weak effect on worry (it should be noted the identification of hallucinations did not require that they should be distressing).


In relation to follow up, the variables fell into two groups, those in which a given variable at time 1 was clearly and strongly predictive of its equivalent at time 2, and those in which such relationships were weak or absent. Somewhat unexpectedly, the second group was wholly made up of affective variables: depression, social anxiety, and situational anxiety. The implication is that these variables fluctuate significantly over the 18 months of follow up and are consequently less likely to be endorsed at a given time-point. They may vary in response to changes in other psychological variables, both those included and those absent from the current analysis. However, general anxiety, worry, sleep problems, and persecutory ideation are strongly self-predicting, suggesting they tend to persist over the follow-up period. To a lesser extent, this is also true of auditory hallucination. Finally, the substance use variables relating to alcohol and cannabis are strongly persistent.

The relationship over the 18-month follow-up period between persecutory ideation and worry is of particular theoretical interest, as each variable has an effect on the other. There is no other instance of a putative feedback loop in these data.

## Discussion

The identification of causal effects is always hostage to the temporal attributes of data. In this investigation, this includes the time frame forming part of the definition of the individual variables. Thus the affective variables relate to the 2-week period immediately before assessment, while others are recorded in relation to the previous year. In consequence, the former may not have been apparent at an earlier stage in the interval between assessments, and the latter may not be present in the later part of the interval. This will tend to attenuate the apparent causal links between these two sorts of variable.

The analysis presented here distinguishes two groups of plausible causal effects. The first group involves analysis *within* each time point: this identifies potential causal relationships but cannot




**Fig. 1.** Distributions of downstream causal effects. The row variables are those at Time 1, the column variables those at Time 2. Where there was no effect between variables we entered a zero in the relevant box. For clarity, the graphs are truncated to cover the range  $-0.1$  to  $0.75$ . The red vertical line in each box indicates zero causal effect. Where the 95% credible interval (the Bayesian counterpart of confidence limits) does not cover the point corresponding to zero causal effect, the whole box is coloured. The numbers in the boxes quantify the relevant *average causal effect*.

provide information about the interval over which the causes operate. The second element of the analysis links the time-points, and relates to potential causes that operate over the 18-month interval. Edges *between* time points will only appear if the timescale of their effects is similar to the interval between the two assessments. Very slow effects would not be picked up, and shorter-term effects will not be identified unless they are cumulative. The structure of the network *within* a time point is likely (though not certain) to capture correlations in the data associated with mechanisms acting on a timescale different from the interval between the repeated measurements. Finally, we observe that in some cases nodes of the same variable do not show edges between time points. One plausible explanation is that the causal effects impinging on such nodes are represented entirely within the structure of the single time point, or flow from the previous time point only by transmission through other variables.

It is notable that most of the strongest links over the 18-month follow-up period were *within* given variables. Worry, persecutory

ideation, general anxiety, and sleep disturbance were self-perpetuating. So were hallucinations and depression, though less strongly. This raises the issues of the inception and maintenance of these psychological features. While many mental experiences appear short-lived, either with an inherent tendency to resolve, or dissipating swiftly when their cause abates, we know that other phenomena are self-perpetuating. Once triggered, the resulting emotional and cognitive reactions are self-fulfilling and so maintain the anxiety, worry or depression. Situational and social anxiety may be maintained because they involve repeated episodes of proximity to a feared context followed by retreat. In many cases the triggers are avoided, so that anxiety can never be resolved by the discovery that there is nothing to fear or that the threat has disappeared. Thus people with claustrophobia may avoid all small spaces and never find out that there is no real danger of death or suffocation. Other phenomena reflect propensities that are constantly reinforced by interaction with the social world. So, persecutory ideation may involve a cognitive



**Fig. 2.** Directed Acyclic Graph of relationships between variables at the two time-points. Arrows within time points are blue, those linking variables across the two time points are red. The density of the arrows represents the strength of the links. Single arrows indicate a plausible causal link, arrows in both directions imply that the data are compatible with causal influences in either direction.

predisposition that is constantly reinforced by inevitable contact with, and over-interpretation of, anodyne or ambiguous social experiences (e.g. Freeman *et al.* 2008). Worry similarly reveals a continuing propensity for handling cognitive content in a particular way. Worry brings implausible ideas to mind, keeps them there, and elaborates the content in negative directions.

Three of the persistent variables concerned the use of cannabis and alcohol. This may be definitional: habitude is written into the definition of alcohol problems and cannabis dependence. Reported last-month cannabis use seems more likely to identify sporadic use, but even so it may often reflect much more habitual use than is implied by the specific question. Our results suggest strongly that problems with substance use do not lead to persecutory ideation in this population sample, though persecutory ideation has reasonably strong effects on later alcohol problems. Our data also indicate that sleep disturbance is a persistent problem, consistent with other research (e.g. Morphy *et al.* 2007).

The relationship between affective variables and persecutory ideation is of particular interest, given that previous research suggests that affective responses induce paranoia, and may mediate the link between traumatic events and paranoia (e.g. Gracie *et al.* 2007; Gibson *et al.* 2016). In particular, Experience Sampling Methodology studies have found that clear affective responses to minor stress were paralleled by moment-to-moment variation in psychotic phenomena (Myin-Germeys *et al.* 2001, 2005; Delespaul *et al.* 2002; Myin-Germeys & van Os, 2007).

In the current analysis, the affective symptom with a clear direct relationship with paranoia is worry, which provides the only link between paranoia and the other affective phenomena analysed here.

The worry/persecutory ideation link can be interpreted in terms of *reciprocal* causality, as manifested over an interval of

18 months. This is consistent with the indeterminate relationship within each time point, but cannot be inferred from it. The relationship implies that targeting either symptom in treatment is likely to ameliorate the other, and will influence the other variables upon which they have an effect. The therapeutic benefit of targeting worry has received strong support in a direct intervention clinical trial involving 150 patients with persistent persecutory delusions (Freeman *et al.* 2015b). The intervention focused solely on reducing the cognitive style of worry and led to reductions in delusions; the mechanism of action was substantiated by mediation analysis.

In contrast, new research showing that reducing insomnia through sleep management techniques leads to reductions in both paranoia and hallucinations (Freeman *et al.* 2017) appears to run counter to the results of our DAGs analysis. Though the benefits of improved sleep might work via a reduction in worry, as indicated in the DAGs model, the picture seems less straightforward for hallucinations. There is also a discrepancy with work associating cannabis with paranoia and hallucinations (Freeman *et al.* 2015a; Murray *et al.* 2017), although these links may operate through shared environmental causes (Shakoor *et al.* 2015). Our results suggest that targeting more persistent features such as worry is likely to be productive. Targeting anxiety and depression may improve well-being, but whether this would ameliorate more established persecutory ideas remains to be tested.

## Conclusion

Our novel statistical analysis, based on dynamic Bayesian networks, enabled us to re-examine the relationships between interacting affective and psychotic variables and to evaluate these

relationships over time. This, in turn, identifies key variables for interventions aimed at improving outcomes for these distressing conditions.

**Acknowledgements.** The authors affirm that they have no financial involvement in or affiliation with any organisation whose financial interests may be affected by material in the manuscript or which might potentially bias it. The work was not directly funded: the dataset used in the analyses is publicly accessible.

## References

- Bebbington P, Jonas S, Kuipers E, King M, Cooper C, Brugha T, Meltzer H, McManus S and Jenkins R (2011) Childhood sexual abuse and psychosis: data from a cross-sectional national psychiatric survey in England. *British Journal of Psychiatry* **199**, 29–37.
- Bebbington P, McBride O, Steel C, Kuipers E, Radovanovic M, Brugha T, Jenkins R, Meltzer H and Freeman D (2013) The structure of paranoia in the general population. *British Journal of Psychiatry* **202**, 419–427.
- Bebbington P and Nayani T (1995) The psychosis screening questionnaire. *International Journal of Methods in Psychiatric Research* **5**, 11–19.
- Ben-Zeev D, Ellington K, Swendsen J and Granholm E (2011) Examining a cognitive model of persecutory ideation in the daily life of people with schizophrenia: a computerized experience sampling study. *Schizophrenia Bulletin* **37**, 1248–1256.
- Borsboom D and Cramer AO (2013) Network analysis: an integrative approach to the structure of psychopathology. *Annual Review of Clinical Psychology* **9**, 91–121.
- Broome MR, Woolley JB, Tabraham P, Johns LC, Bramon E, Murray GK, Pariante C, McGuire PK and Murray RM (2005) What causes the onset of psychosis? *Schizophrenia Research* **79**, 23–34.
- Catone G, Marwaha S, Kuipers E, Lennox B, Freeman D, Bebbington P and Broome M (2015) Bullying victimisation and risk of psychotic phenomena: analyses of British national survey data. *The Lancet. Psychiatry* **2**, 618–624.
- Dawid AP (2010) Beware of the DAG!. *Journal of Machine Learning Research Workshop and Conference Proceedings* **26**, 99–157.
- Delespaul P, deVries M and van Os J (2002) Determinants of occurrence and recovery from hallucinations in daily life. *Social Psychiatry and Psychiatric Epidemiology* **37**, 97–104.
- Freeman D (2016) Persecutory delusions: a cognitive perspective on understanding and treatment. *The Lancet Psychiatry* **3**, 685–692.
- Freeman D, Dunn G, Murray R, Evans N, Lister R, Antley A, Slater M, Godlewska B, Cornish R, Williams J, Di Simplicio M, Igoumenou A, Brenneisen R, Tunbridge E, Harrison P, Harmer C, Cowen P and Morrison P (2015a) How cannabis causes paranoia: using the intravenous administration of  $\Delta 9$ -tetrahydrocannabinol (THC) to identify key cognitive mechanisms leading to paranoia. *Schizophrenia Bulletin* **41**, 391–399.
- Freeman D, Dunn G, Startup H, Pugh K, Cordwell J, Mander H, Cernis E, Wingham G, Shirvell K and Kingdon D (2015b) Effects of cognitive behaviour therapy for worry on persecutory delusions in patients with psychosis (WIT): a parallel, single-blind, randomised controlled trial with a mediation analysis. *The Lancet Psychiatry* **2**, 305–313.
- Freeman D, Pugh K, Antley A, Slater M, Bebbington P, Gittins M, Dunn G, Kuipers E, Fowler D and Garety PA (2008) A virtual reality study of paranoid thinking in the general population. *British Journal of Psychiatry* **192**, 258–263.
- Freeman D, Sheaves B, Goodwin GM, Yu L.-M, Nickless A, Harrison PJ, Emsley R, Luik AI, Foster RG, Wadekar V, Hinds C, Gumley A, Jones R, Lightman S, Jones S, Bentall R, Kinderman P, Rowse G, Brugha T, Blagrove M, Gregory AM, Fleming L, Walklet E, Glazebrook C, Davies EB, Hollis C, Haddock G, John B, Coulson M, Fowler D, Pugh K, Cape J, Moseley P, Brown G, Hughes C, Obonsawin M, Coker S, Watkins E, Schwannauer M, MacMahon K, Siriwardena AN and Espie CA (2017) The effects of improving sleep on mental health (OASIS): a randomised controlled trial with mediation analysis. *The Lancet Psychiatry* **4**, 749–758.
- Freeman D, Stahl D, McManus S, Meltzer H, Brugha T, Wiles N and Bebbington P (2012) Insomnia, worry, anxiety and depression as predictors of the occurrence and persistence of paranoid thinking. *Social Psychiatry and Psychiatric Epidemiology* **47**, 1195–1203.
- Fried EI, van Borkulo CD, Cramer AOJ, Boschloo L, Schoevers RA and Borsboom D (2017) Mental disorders as networks of problems: a review of recent insights. *Social Psychiatry and Psychiatric Epidemiology* **52**, 1–10.
- Friedman N, Murphy K and Russell S (1998) Learning the structure of dynamic probabilistic networks. In *Proceedings of the Fourteenth Conference on Uncertainty in Artificial Intelligence*. San Francisco, CA: Morgan Kaufmann Publishers Inc, 139–147.
- Gibson LE, Alloy LB and Ellman LM (2016) Trauma and the psychosis spectrum: a review of symptom specificity and explanatory mechanisms. *Clinical Psychology Review* **49**, 92–105.
- Gracie A, Freeman D, Green S, Garety PA, Kuipers E, Hardy A, Ray K, Dunn G, Bebbington P and Fowler D (2007) The association between traumatic experience, paranoia and hallucinations: a test of the predictions of psychological models. *Acta Psychiatrica Scandinavica* **116**, 280–289.
- Hartley S, Haddock G, Vasconcelos E Sa D, Emsley R and Barrowclough C (2014) An experience sampling study of worry and rumination in psychosis. *Psychological Medicine* **44**, 1605–1614.
- Howes OD and Murray RM (2014) Schizophrenia: an integrated sociodevelopmental-cognitive model. *Lancet* **10**, 383(9929), 1677–1687.
- Isvoranu AM, Borsboom D, van Os J and Guloksuz S (2016) A network approach to environmental impact in psychotic disorder: brief theoretical framework. *Schizophrenia Bulletin* **42**, 870–873.
- Isvoranu AM, van Borkulo CD, Boyette LL, Wigman JT, Vinkers CH, Borsboom D and Group Investigators (2017) A network approach to psychosis: pathways between childhood trauma and psychotic symptoms. *Schizophrenia Bulletin* **43**, 187–196.
- Kelleher I, Keeley H, Corcoran P, Ramsay H, Wasserman C, Carli V, Sarchiapone M, Hoven C, Wasserman D and Cannon M (2013) Childhood trauma and psychosis in a prospective cohort study: cause, effect, and directionality. *American Journal of Psychiatry* **170**, 734–741.
- Klippel A, Viechtbauer W, Reininghaus U, Wigman J, van Borkulo C; MERGE, Myin-Germeys I and Wichers M (2017) The cascade of stress: a network approach to explore differential dynamics in populations varying in risk for psychosis. *Schizophrenia Bulletin* **43**, 302–315.
- Krabbedam L, Myin-Germeys I, De Graaf R, Vollebergh W, Nolen WA, Iedema J and Van Os J (2004) Dimensions of depression, mania and psychosis in the general population. *Psychological Medicine* **34**, 1177–1186.
- Kuipers E, Garety P, Fowler D, Freeman D, Dunn G and Bebbington P (2006) Cognitive, emotional and social processes in psychosis: refining cognitive behavioural therapy for persistent positive symptoms. *Schizophrenia Bulletin* **32**, s24–s31.
- Kuipers J and Moffa G (2017) Partition MCMC for inference on acyclic digraphs. *Journal of the American Statistical Association* **112**, 282–299.
- Lewis G, Pelosi AJ, Araya R and Dunn G (1992) Measuring psychiatric disorder in the community: a standardised assessment for use by lay interviewers. *Psychological Medicine* **22**, 465–486.
- Marwaha S and Bebbington P (2015) Mood as a mediator of the link between child sexual abuse and psychosis. *Social Psychiatry and Psychiatric Epidemiology* **50**, 661–663.
- Marwaha S, Broome MR, Bebbington PE, Kuipers E and Freeman D (2014) Mood instability and psychosis: analyses of British national survey data. *Schizophrenia Bulletin* **40**, 269–277.
- Marwaha S, Thompson A, Bebbington P, Singh SP, Freeman D, Winsper C and Broome M (2015) Adult attention deficit hyperactivity symptoms and psychosis: epidemiological evidence from a population survey in England. *Psychiatry Research* **229**, 49–56.
- McGrath JJ, Saha S, Lim CCW, Aguilar-Gaxiola S, Alonso J, Andrade LH, Bromet EJ, Bruffaerts R, Caldas de Almeida JM, Cardoso G, de Girolamo G, Fayyad J, Florescu S, Gureje O, Haro JM, Kawakami N, Koenen KC, Kovess-Masfety V, Lee S, Lepine JP, McLaughlin KA, Medina-Mora ME, Navarro-Mateu F, Ojagbemi A, Posada-Villa J, Sampson N, Scott KM, Tachimori H, Ten Have M, Kessler KS, Kessler RC and WHO World Mental Health Survey Collaborators (2017) Trauma and psychotic experiences: transnational data from the world mental health survey. *British Journal of Psychiatry* **211**, 373–380.

- McNally RJ** (2016) Can network analysis transform psychopathology? *Behaviour Research and Therapy*, **86**, 95–104.
- McNally RJ, Robinaugh DJ, Wu GWY, Wang L, Deserno M and Borsboom D** (2015) Mental disorders as causal systems: a network approach to post-traumatic stress disorder. *Clinical Psychological Science*, **3**, 836–849.
- Moffa G, Catone G, Kuipers J, Kuipers E, Freeman D, Marwaha S, Lennox BR, Broome MR and Bebbington P** (2017) Using directed acyclic graphs in epidemiological research in psychosis: an analysis of the role of bullying in psychosis. *Schizophrenia Bulletin* **43**, 1273–1279.
- Morphy H, Dunn KM, Lewis M, Boardman HF and Croft PR** (2007) Epidemiology of insomnia: a longitudinal study in a UK population. *Sleep* **30**, 274–280.
- Murphy J, McBride O, Fried E and Shevlin M** (2017) Distress, impairment and the extended psychosis phenotype: a network analysis of psychotic experiences in an US general population sample. *Schizophrenia Bulletin* 2017 Sep 23. doi: 10.1093/schbul/sbx134. [Epub ahead of print].
- Murray RM, Englund A, Abi-Dargham A, Lewis DA, Di Forti M, Davies C, Sherif M, McGuire P and D'Souza DC** (2017) Cannabis-associated psychosis: neural substrate and clinical impact. *Neuropharmacology* **124**, 89–104.
- Myin-Germeyns I, Delespaul P and van Os J** (2005) Behavioural sensitization to daily life stress in psychosis. *Psychological Medicine* **35**, 733–741.
- Myin-Germeyns I and van Os J** (2007) Stress-reactivity in psychosis: evidence for an affective pathway to psychosis. *Clinical Psychology Review* **27**, 409–424.
- Myin-Germeyns I, van Os J, Schwartz JE, Stone AA and Delespaul PA** (2001) Emotional reactivity to daily life stress in psychosis. *Archives of General Psychiatry* **58**, 1137–1144.
- Pearl J** (2009) *Causality: Models, Reasoning, and Inference*, 2nd edn., Cambridge, UK: Cambridge University Press.
- Robins LN, Helzer JE, Croughan J and Ratcliff KS** (1981) National institute of mental health diagnostic interview schedule. Its history, characteristics, and validity. *Archives of General Psychiatry* **38**, 381–389.
- Saunders JB, et al.** (1993) Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption—II. *Addiction* **88**, 791–804.
- Scott J, Martin G, Bor W, Sawyer M, Clark J and McGrath J** (2009) The prevalence and correlates of hallucinations in Australian adolescents: results from a national survey. *Schizophrenia Research* **107**, 179–185.
- Shakoor S, Zavos HMS, McGuire P, Cardno AG, Freeman D and Ronald A** (2015) Psychotic experiences are linked to cannabis use in adolescents in the community because of common underlying environmental risk factors. *Psychiatry Research* **227**, 144–151.
- Sheaves B, Bebbington PE, Goodwin GM, Harrison PJ, Espie CA, Foster RG and Freeman D** (2016) Insomnia and hallucinations in the general population: finding from the 2000 and 2007 British psychiatric morbidity surveys. *Psychiatry Research* **241**, 141–146.
- Singleton N, Bumpstead R, O'Brien M, Lee A and Meltzer H** (2001) *Psychiatric Morbidity among Adults Living in Private Households*. London: TSO.
- Singleton N and Lewis G** (2003) *Better or Worse: A Longitudinal Study of the Mental Health of Adults Living in Private Households in Great Britain*. London: TSO.
- Thewissen V, Bentall RP, Oorschot M, A Campo J, van Lierop T, van Os J and Myin-Germeyns I** (2011) Emotions, self-esteem, and paranoid episodes: an experience sampling study. *British Journal of Clinical Psychology* **50**, 178–195.