

1
2
3
4
5
6
7
8
9

Protocol for Direct Reporting of Awareness in Maternity patients (DREAMY): a prospective, multicentre cohort study of accidental awareness during general anaesthesia in obstetric patients

10
11
12
13
14
15
16
17
18
19

P.M. Odor, ^{a} S. Bampoe,^b D.N. Lucas,^c S.R. Moonesinghe,^b J. Andrade,^d J.J. Pandit^e for the DREAMY Investigators Group*

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

^a Department of Anaesthesia, St. George's University Hospital, London, UK

^b Centre for Anaesthesia and Perioperative Medicine, University College London Hospital, London, UK

^c Department of Anaesthesia, Northwick Park Hospital, London, UK

^d School of Psychology, University of Plymouth, Plymouth, UK

^e Nuffield Department of Anaesthetics, John Radcliffe Hospital, Oxford, UK

*Corresponding author. E-mail: peter.odor@nhs.net; 07746 399221

Abstract

Background

Accidental awareness during general anaesthesia (AAGA) is a complex and rare outcome to investigate in surgical patient populations, particularly obstetric patients. We report the protocol of the Direct Reporting of Awareness in Maternity patients (DREAMY) study, to illustrate how the research was designed to address practical and methodological challenges for studying AAGA in an obstetric cohort.

Methods

This is the trial protocol of a prospective, multicentre cohort study of patients undergoing obstetric surgery under general anaesthesia. AAGA will be detected using three repetitions of standardised direct questioning over thirty days, with responses indicating memories during general anaesthesia verified using structured interviews. Reports will be adjudicated then classified in accordance with pre-defined and pre-validated structures, including the Michigan Awareness Classification tool. Quantitative data will be collected on general anaesthesia conduct for all participants. This descriptive study is being conducted in England, aiming to recruit a minimum of 2015 patients.

Results

The DREAMY study was prospectively registered (ClinicalTrials.gov Identifier: NCT03100396) and ethical approval granted. Participant recruitment began in May 2017 and 1 year follow up concluded in August 2019. Publication of the results is anticipated in 2020.

Conclusions

The DREAMY study will provide data on incidence, experience and implications of AAGA for obstetric patients using a robust methodology that will reliably detect and translate subjective AAGA reports into objective outcomes. In addition, the study is expected to improve vigilance for AAGA in participating hospitals and encourage adoption of recommendations for support of patients suffering AAGA.

Keywords

Awareness, consciousness; obstetric surgery; perioperative complications; research protocol

Background

The 5th National Audit Project in the UK and Ireland (NAP5)¹ was the largest ever study of accidental awareness during general anaesthesia (AAGA). The NAP5 methodology combined national surveillance for spontaneous patient reports of AAGA and a parallel snapshot survey of general anaesthetic activity.² Obstetric patients undergoing caesarean delivery were markedly over-represented, constituting almost 10% of all NAP5 AAGA reports, but only 0.8% of general anaesthetic activity. Importantly, NAP5 adopted a different approach to study design to most previous investigations, relying solely upon descriptions *spontaneously* reported by patients to clinicians, rather than asking patients directly about their memories. Whilst it may be argued that NAP5 was successful in detecting the most clinically relevant AAGA experiences³, it is unknown how many cases of AAGA were missed, either due to lack of opportunity, motivation or anxiety in patients to declare their experiences.

A more common methodological approach to detect AAGA is to screen patients using a standardised interview format, typically based upon a set of questions termed the Brice interview.⁴ The Brice interview questions provoke patients to recall their memories of the induction and emergence stages of a general anaesthetic, including anything that may have occurred in between. Studies using such direct questioning consistently estimate the incidence of AAGA at 1-2:1000.⁵⁻¹¹ In contrast, NAP5 estimated the incidence of AAGA to be far lower, ~1:19,000 anaesthetics, albeit with considerable context-dependent variation: ~1:8,000 when neuromuscular blocking drugs (NMBDs) were used, compared with ~1:136,000 without NMBDs; and ~1:670 for caesarean delivery.

Obstetric general anaesthesia has multiple risk factors for AAGA: the universal use of NMBDs, rapid sequence induction, a high incidence of difficult airway management¹² and emergency surgery. Previous attempts to quantify AAGA in obstetric patients using the Brice interview have indicated a disproportionately high risk, ranging from an estimated incidence of 1:110 during the 1980s¹³, to 1:382 in the mid 2000s.¹⁴ These studies involved results with wide confidence intervals, a lack of consistent AAGA classification and are difficult to translate to current anaesthetic practice, given recent changes in anaesthetic management for obstetric general anaesthesia.¹⁵ Brice interviews were used by Paech et al, however the findings were limited by a relatively small sample size (n = 763) and a lack of verification for AAGA reports beyond one or two brief interviews.¹⁴

Aims and objectives

The aim of the Direct Reporting of Awareness in Maternity patients (DREAMY) study is to describe the epidemiology of AAGA in a cohort of adult patient undergoing obstetric surgery, describing the incidence, nature of experiences, risk factors and implications of AAGA. The study aims to fulfil research implications from NAP5, in relation to AAGA in the obstetric population. These implications are to “define the incidence of AAGA as identified by the Brice questionnaire” (Research Implication 16.6) and “explore whether

factors make obstetric patients more likely to report episodes of AAGA than the non-obstetric population” (Research Implication 16.7).¹⁶

The primary objective is to:

1. Describe the incidence of different classifications of AAGA identified using direct questioning following obstetric surgery.

Secondary objectives are:

1. Describe the characteristics of the patient population undergoing obstetric surgery under general anaesthesia.
2. Investigate the conduct of general anaesthesia for obstetric surgery, including pharmacological and airway management, workforce considerations, and the risk of any complications.
3. Describe the psychological implications of AAGA using a post-traumatic stress disorder symptom checklist follow up over 12 months and any prognostic factors for this.
4. Investigate the association between risk of AAGA and patient or anaesthetic characteristics.
5. Investigate the association between dreaming during general anaesthetic and patient or anaesthetic characteristics.

Methods

Ethical approval was granted by the UK National Research Ethics Service (London Fulham Committee; REC reference 17/LO/0071) for this consenting study. Trial registration was obtained prospectively (ClinicalTrials.gov Identifier: NCT03100396). This protocol is reported in accordance with the STROBE statement for reporting epidemiological observational studies.¹⁷

Study design

DREAMY is a prospective, non-randomised, descriptive, multi-centre cohort study. Following informed consent, patients undergoing general anaesthesia for obstetric surgery will be questioned about their experience of general anaesthesia, either in person or via telephone, using the Brice interview on three separate occasions within 30 days of anaesthesia. Patients who indicate possible memories during general anaesthesia will undergo structured follow up for up to 12 months. All patient reports will be analysed and adjudicated by a mixed-background expert panel, using pre-defined classification for AAGA outcomes.

Setting

Patients have been recruited from participating hospitals in England during a 15 month recruitment window, beginning in May 2017. Between 40 and 60 hospitals are expected to participate.

Participants

Table 1 shows patient eligibility criteria. All urgencies of surgery (classified according to a model proposed by Lucas et al¹⁸ and adopted by the Royal College of Obstetricians and Gynaecologist in the United Kingdom) are eligible, including emergency surgery.

Variables and Measurement

AAGA will be identified using a four-stage approach:

1. Detection

Screening for AAGA will be performed with a structured Brice interview, adapted from the originally published form⁴ (Appendix 1) to incorporate categorical and open patient responses, maintaining consistency with use in several comparable studies.^{19,20} Since detection of AAGA may be increased by multiple repetitions of structured post-operative questioning,^{11, 21, 22} the Brice interview will be provided on three separate occasions over thirty days; 0-24 hours post-operatively, 24-48 hours and at 30 days. This approach is termed “Thrice Brice” and maximises the opportunity for participants to declare experiences recalled immediately following surgery (when the prospect for recall of detail is highest) and later emerging memories.

The first two questionnaires will be performed by local hospital investigators, offering the questionnaire but clarifying responses and answering queries in person. The third Brice interview and all subsequent research follow up will be performed via telephone by the coordinating research team. Patients will be included for analysis even if fewer than three Brice responses are collected.

Any abnormal report indicating memories that a patient attributes to the period between "going to sleep" and "waking up" will trigger a second, verification process to improve AAGA detection specificity.

2. Verification

To assure validity, data on AAGA reports will be collected using multiple methods. Patients reporting possible awareness will be independently interviewed by a study author, using semi-structured questioning to verify the detail, characteristics, plausibility and psychological response to the experience. Suspected AAGA structured interview schedules (Appendix 2) are adapted from the BAG-RECALL trial.¹⁹ Any specific description of events made in verification interviews will be checked with staff involved with clinical care to provide corroboration or refutation. This includes investigating the timing and specific nature of reported sensory perceptions.

All suspected AAGA patients will also be asked to complete a PCL-5 post-traumatic stress disorder (PTSD) checklist²³ by telephone at day 30, then 3, 6, 9 and 12 months following the

1 anaesthetic episode. During first administration of the PCL-5, additional information will be
2 obtained using a structured and multi-dimensional questionnaire on the patient's self-reported
3 mental health history, infant's health status and psychological experiences in the gravidic-
4 puerperal cycle. Question domains have been derived from recognised risk factors for
5 developing PTSD during the post-natal period.²⁴ PCL-5 symptom checklists will also be
6 asked to a minimum comparator sample of 300 patients with no indication of AAGA on
7 "Thrice Brice" questioning.
8
9

10 An extended AAGA follow up will occur at 3 months post-operatively, to investigate the
11 nature of local clinical follow up received by patients and how the AAGA episode has
12 impacted on attitudes towards future anaesthetics and post-natal experience.
13
14

15 *3. Adjudication*

16 Adjudication of patients' reports according to the likelihood of each representing an AAGA
17 experience will be performed by a panel of experts independent to study data collection and
18 blinded to hospital site and any personal identifiers.
19
20

21 Panel members will be drawn from mixed anaesthetic and psychologist backgrounds, with
22 prior understanding of AAGA, including individuals who adjudicated on NAP5 panels. A
23 minimum of five panel members will review all reports.
24
25

26 Each panel member will first review the patient reports and anaesthetic episode data
27 separately. Cases will be discussed collectively by the panel, considering detail, plausibility
28 and consistency of reported experiences with intraoperative process. Panel members will be
29 reminded to be mindful of "outcome bias" (where knowledge of the poor outcome can lead to
30 a retrospective harsh judgement) and "groupthink" bias (where groups make irrational
31 decisions given a subconscious desire to agree with others).
32
33

34 Panel members will declare separate adjudication decisions and kappa statistic will be
35 reported to measure agreement. Final allocations will be decided based upon the majority
36 adjudication decision by the panel.
37
38

39 AAGA reports will be graded according to a confidence scale described in Table 2 (adapted
40 from NAP5).²⁵
41
42

43 *4. Classification*

44 AAGA events will be graded by consensus opinion by the review panel in accordance with
45 the Michigan Awareness Classification Instrument²⁶ (Table 3) and a modified version of the
46 National Patient Safety Agency tool²⁷ adapted for NAP5 to be suitable for predominantly
47 psychological harm related to AAGA (Table 4).¹
48
49

50 Where possible, events will also be classified according to the phase of anaesthesia (Table 5).
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Statistical analysis

The minimum sample size was based upon confidence to detect an incidence of AAGA at least three times higher than described following Brice interviews in non-obstetric surgical patients. The assumption for the comparator proportion was based upon data from the most recent study of AAGA in obstetric surgical patients.¹⁴ The baseline AAGA proportion was taken to be 0.15% (~1:666).^{5,6} An exact binomial test with a one-sided alpha of 0.05 showed that 2015 patients would be needed to give a power of 80% to detect a comparator proportion more than 0.45%.

Commonly applied limits on alpha and beta error may yield poor precision estimates for the binomial proportion when rare events are expected. Likewise, disparity of AAGA detection and reporting may lead to under-or over-estimates of sample size power. On the basis that AAGA was expected to be rare outcome in the sampled population and the study had a descriptive objective (i.e. to describe the incidence of AAGA), no maximum sample size was pre-defined. Instead the study will use a combination of a minimum recruitment threshold with a descriptive approach, aiming to recruit a maximal number of eligible patients during a 15-month recruitment period. The recruitment duration was pragmatically selected, and feasibility was supported by an activity survey of obstetric general anaesthesia in hospitals expected to be involved in the study.

The primary outcome of proportion of obstetric patients reporting a composite of certain/probable and possible AAGA will be expressed using binomial confidence intervals and compared with established values using Fisher's exact or chi-squared testing. Analysis will be stratified for certain/probable and outcomes for other classification groups to provide optimistic or pessimistic confidence intervals for the estimated incidence of AAGA. Associations between anaesthetic, surgical and patient factors will be tested using univariate analysis. If sufficient AAGA events are identified, then a multivariate regression model will be used to evaluate the independent association of specific anaesthetic episode variables with AAGA, with results expressed as odds ratios. For secondary outcomes, data will be presented as a number or percentage; mean and standard deviation for normally distributed data and median and interquartile range for non-parametric data, with 95% confidence intervals.

PCL-5 results will be expressed as a total symptom severity score (range = 0-80), by summing scores from the 20 checklist items, and differentiated according to symptom criteria domains as re-living, avoidance behaviour, emotional blunting and hyper-excitability.²⁸ PTSD prevalence rates will be estimated for the entire sample and subgroups, including AAGA and non-AAGA patients. Results will be expressed with the respective 95% confidence intervals.

Case descriptions of the experiences of AAGA undergone by patients with AAGA will be presented in anonymised format. The phase of anaesthesia/surgery when the AAGA event occurred, alongside possibility contributory, causal or mitigating factors, will be described where possible.

1 Anaesthetic episode data will be collected by local investigation from patient notes and
2 anaesthetic records, before transferring to a central electronic database. Local clinical teams
3 will maintain responsibility for provision of support to patients with AAGA. All sites will be
4 encouraged to provide care in accordance with the NAP5 Anaesthesia Awareness Support
5 Pack guidelines.²⁹
6

7
8 The DREAMY study is coordinated by a chief investigator, supported by a study steering
9 group and an affiliated anaesthetic trainee network, the Pan-London Perioperative Audit and
10 Research Network (PLAN). PLAN is led by anaesthetic trainees to benefit trainee learning
11 and supports the development of resources to encourage large scale participation in multi-
12 centre studies of anaesthetic practice. Collaborations with regional anaesthetic trainee
13 networks were invited.
14
15

16 **Discussion**

17
18 We present the DREAMY study protocol to demonstrate how the study has been designed to
19 address prior challenges with the methodology and interpretation of AAGA research. The
20 study combines approaches tested by NAP5 with direct questioning.
21
22

23 The first challenge relates to the nature of the outcome to be reported. AAGA is a complex
24 phenomenon, occurring with a wide spectrum of experiences, and descriptions of events are
25 subject to variable interpretation by clinicians and researchers alike.³⁰ It is expected that only
26 a minority of patients reporting AAGA will present clear evidence that can be corroborated,
27 such as memory of specific intraoperative events or conversations between operating theatre
28 staff. Since AAGA detection relies upon subjective reporting of a spectrum of experiences
29 occurring in an unfamiliar hospital environment, a report indicating AAGA likely will be
30 unordered, confused, lacking in points of reference and potentially psychologically traumatic.
31 The DREAMY protocol therefore includes varied approaches to enable detailed capture of
32 patient reports. In addition, the 30-day post-operative interview is conducted by a trained
33 interviewer independent to each hospital, presenting an alternative route for disclosure.
34 Training, guidance documents and advice regarding local communication and management of
35 patients reporting distressing memories support patient wellbeing,²⁹ which may also improve
36 conditions for reporting by patients.³¹
37
38

39 The second challenge is that AAGA can only be readily detected if memory of the experience
40 is present. The relationship between memory and unintended consciousness during general
41 anaesthesia is neither simple nor mutual. It is unknown how long a period of AAGA must last
42 to produce a memory that can be recalled after waking from general anaesthesia.
43 Respondents in isolated forearm testing rarely have explicit recall afterwards.³² A minority of
44 patients experiencing AAGA have no declarative memory of the event until several weeks
45 post-operatively.^{5, 33}
46
47

48 General anaesthesia affects perceptual and episodic memory formation at lower effect site
49 concentrations than for hypnotic effects, inhibiting the communication pathways needed to
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 translate memory from a short-term buffer (“working” memory) into explicit recall.³⁴ Hence,
2 suppression of memory formation at anaesthetic concentrations that enable perceptive AAGA
3 may interfere with the process that is fundamental to detection of AAGA.
4

5 Alternatively, memories of intraoperative events can be subject to recall bias, or disruption by
6 procedural factors. The timing and format of questions about memories is important. Best
7 practices to identify AAGA reports are unknown, lacking validation of optimal phrasing,
8 timing and frequency of questions asked. Recall is a process of reconstructing rather than
9 replaying a past event. Memories can distort, or false memories be inserted by encouraging
10 patients to generate that information themselves.³⁵ Such phenomenon tends to occur when
11 patients are presented with repeated, leading questions; a process that may potentially occur
12 with the Brice interview.
13
14
15
16

17 Ultimately credibility for AAGA outcomes relies upon having the consistent, unbiased,
18 detailed and contemporaneous records of patient memories following surgery. Whilst the
19 Brice interview has face validity, construct and content validity are unquantified. Different
20 variations of Brice question delivery or phrasing have not been investigated. Comparisons
21 between a single repetition of the Brice interview at post-operative day 28-30 versus open-
22 ended quality assurance interviews at post-operative day 1 have indicated higher detection of
23 AAGA events using the Brice approach.²¹ Thus Brice has become a de facto standard, based
24 upon commonality rather than any formal testing of validity.
25
26
27
28
29

30 The approach in the DREAMY protocol of three Brice interview repetitions over thirty days
31 was designed to balance pragmatism, sensitivity for identifying transient memories and
32 minimise memory implantation. Although we acknowledge that there are limitations to Brice
33 interviews, this methodology remains the only option to enable comparisons with previous
34 studies. We consider Brice to be a sensitive tool for detecting AAGA, but lacking specificity;
35 hence the additional verification processes in DREAMY.
36
37
38
39

40 The third challenge relates to translating subjective descriptions of experiences into objective
41 outcomes. The process of adjudication is often poorly reported with wide variation in blinded
42 panel outcomes;³⁰ a finding that may reflect inconsistency in study methodologies. The
43 DREAMY protocol therefore employs a range of tools that have been tested by other studies,
44 including the Michigan Awareness Classification Scale, the National Patient Safety Agency
45 tool and PCL-5. We are also report intra-panel agreement to ensure transparency for how
46 AAGA reports are considered and outcomes structured.
47
48
49
50

51 The study design retains several limitations. The small number of anticipated AAGA events
52 means that confidence intervals may still be wide. Patients with severely traumatic AAGA
53 experiences may decline to consent for research participation, potentially introducing bias.
54 However, this effect has not been described for previous (non-obstetric) patients. A minimum
55 proficiency in English language potentially restricts participation in the study, but avoids the
56 introduction of unknown bias from translational difficulties in patient reporting.
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

In conclusion, DREAMY will provide vital descriptive data on AAGA incidence, experience and implications for obstetric patients. In addition, the study is expected to improve vigilance for AAGA in participating hospitals, to encourage adoption of recommendations on patient support from NAP5³⁶ and raise the profile for an important patient-facing complication of anaesthetic practice. We urge others to improve and debate the methodological strategy used in DREAMY to establish and support standardised best practice for AAGA investigation.

Declaration of interests

Funding is via a National Institute of Academic Anaesthesia (NIAA) and Obstetric Anaesthetists' Association (OAA) research grant. Peter Odor was Chair of the Pan-London Perioperative Audit and Research Network (PLAN) from 2015-2018. Jaideep Pandit was the clinical lead for the 5th National Audit Project (NAP5). Nuala Lucas is an editor of IJOA and Chair of the OAA Education subcommittee, but was not involved in the editorial process for this manuscript. There are no other conflicts of interest to declare.

Author Contributions

Study conception: PO, SB

Study design: PO, SB, JA, JP, RM, NL

Data collection: all investigators

Data analysis: PO, SB, JA, JP, NL

Manuscript preparation: PO, SB

Manuscript editing and review: PO, SB, JA, JP, NL, RM

References

1. Pandit JJ, Andrade J, Bogod DG, et al. 5th National Audit Project (NAP5) on accidental awareness during general anaesthesia: protocol, methods, and analysis of data. *Br J Anaesth* 2014; **113**: 540-548.
2. Sury MR, Palmer JH, Cook TM, Pandit JJ. The state of UK anaesthesia: a survey of National Health Service activity in 2013. *Br J Anaesth* 2014; **113**: 575-584.
3. Cook TM, Pandit JJ. Pitfalls of comparing incidences of awareness from NAP5 and from Brice studies. *Br J Anaesth* 2015; **115**: 471-472.
4. Brice DD, Hetherington RR, Utting JE. A simple study of awareness and dreaming during anaesthesia. *Br J Anaesth* 1970; **42**: 535-542.
5. Sandin RH, Enlund G, Samuelsson P, Lennmarken C. Awareness during anaesthesia: a prospective case study. *Lancet* 2000; **355**: 707-711.
6. Sebel PS, Bowdle TA, Ghoneim MM, et al. The incidence of awareness during anesthesia: a multicenter United States study. *Anesth Analg* 2004; **99**: 833-839.
7. Ranta SO, Laurila R, Saario J, Ali-Melkkila T, Hynynen M. Awareness with recall during general anesthesia: incidence and risk factors. *Anesth Analg* 1998; **86**: 1084-1089.
8. Walker EMK, Bell M, Cook TM, Grocott MPW, Moonesinghe SR. Patient reported outcome of adult perioperative anaesthesia in the United Kingdom: a cross-sectional observational study. *Br J Anaesth* 2016; **117**: 758-766.

9. Myles PS, Leslie K, McNeil J, Forbes A, Chan MT. Bispectral index monitoring to prevent awareness during anaesthesia: the B-Aware randomised controlled trial. *Lancet* 2004; **363**: 1757-1763.
10. Avidan MS, Zhang L, Burnside BA, et al. Anesthesia awareness and the bispectral index. *N Engl J Med* 2008; **358**: 1097-1108.
11. Avidan MS, Jacobsohn E, Glick D, et al. Prevention of intraoperative awareness in a high-risk surgical population. *N Engl J Med* 2011; **365**: 591-600.
12. Kinsella SM, Winton AL, Mushambi MC, et al. Failed tracheal intubation during obstetric general anaesthesia: a literature review. *Int J Obstet Anesth* 2015; **24**: 356-374.
13. Lyons G, Macdonald R. Awareness during caesarean section. *Anaesthesia* 1991; **46**: 62-64.
14. Paech M. J. SKL, Clavisi O., Chua S., McDonnell N.,. A prospective study of awareness and recall associated with general anaesthesia for caesarean section. *Int J Obstet Anesth* 2009; **17**: 298-303.
15. Desai N, Wicker J, Sajayan A, Mendonca C. A survey of practice of rapid sequence induction for caesarean section in England. *Int J Obstet Anesth* 2018; **36**: 3-10.
16. Pandit JJ, Andrade J, Bogod DG, et al. 5th National Audit Project (NAP5) of the Royal College of Anaesthetist and Association of Anaesthetists of Great Britain and Ireland. Accidental Awareness during General Anaesthesia in the United Kingdom and Ireland <http://nap5.org.uk/NAP5report2014>. Accessed June 15th 2019.
17. von Elm E, Altman DG, Egger M, Pocock SJ, Gotsche PC, Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008; **61**: 344-349.
18. Lucas DN, Yentis SM, Kinsella SM, et al. Urgency of caesarean section: a new classification. *J R Soc Med* 2000; **93**: 346-350.
19. Avidan MS, Palanca BJ, Glick D, et al. Protocol for the BAG-RECALL clinical trial: a prospective, multi-center, randomized, controlled trial to determine whether a bispectral index-guided protocol is superior to an anesthesia gas-guided protocol in reducing intraoperative awareness with explicit recall in high risk surgical patients. *BMC Anesthesiol* 2009; **9**: 8.
20. Moonesinghe SR, Walker EMK, Bell M, Group S-I. Design and methodology of SNAP-1: a Sprint National Anaesthesia Project to measure patient reported outcome after anaesthesia. *Periop Med* 2015; **4**: 4-4.
21. Mashour GA, Kent C, Picton P, et al. Assessment of intraoperative awareness with explicit recall: a comparison of 2 methods. *Anesth Analg* 2013; **116**: 889-891.
22. Pollard RJ, Coyle JP, Gilbert RL, Beck JE. Intraoperative awareness in a regional medical system: a review of 3 years' data. *Anesthesiology* 2007; **106**: 269-274.
23. Weathers FW, Litz, B.T., Keane, T.M., Palmieri, P.A., Marx, B.P., & Schnurr, P.P. The PTSD Checklist for DSM-5 (PCL-5). Scale available from the National Center for PTSD 2013. Available from: www.ptsd.va.gov. Accessed June 15th 2019.
24. Andersen LB, Melvaer LB, Videbech P, Lamont RF, Joergensen JS. Risk factors for developing post-traumatic stress disorder following childbirth: a systematic review. *Acta Obstet Gynecol Scand* 2012; **91**: 1261-1272.
25. Pandit JJ, Andrade J, Bogod DG, et al. The 5th National Audit Project (NAP5) on accidental awareness during general anaesthesia: Protocol, methods and analysis of data. *Anaesthesia* 2014; **69**: 1078-1088.
26. Mashour GA, Esaki RK, Tremper KK, Glick DB, O'Connor M, Avidan MS. A novel classification instrument for intraoperative awareness events. *Anesth Analg* 2010; **110**: 813-815.
27. Agency NPS. A Risk Matrix for Risk Managers. NPSA: London, 2008.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
28. Blevins CA, Weathers FW, Davis MT, Witte TK, Domino JL. The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): Development and Initial Psychometric Evaluation. *J Trauma Stress* 2015; **28**: 489-498.
 29. Cook T PJ, Andrade J, Wang M. NAP5 Anaesthesia Awareness Support Pack 2014. Available from: <http://www.nationalauditprojects.org.uk/NAP5-Anaesthesia-Awareness-Pathway>. Accessed June 15th 2019.
 30. Messina AG, Wang M, Vezina DP, Pace NL. Adjudication of awareness events. *Br J Anaesth* 2018; **121**: 329-330.
 31. Dickson-Swift V, James EL, Kippen S, Liamputtong P. Doing sensitive research: What challenges do qualitative researchers face? *Qual Res* 2007; **7**: 327-353.
 32. Linassi F, Zanatta P, Tellaroli P, Ori C, Carron M. Isolated forearm technique: a meta-analysis of connected consciousness during different general anaesthesia regimens. *Br J Anaesth* 2018; **121**: 198-209.
 33. Ghoneim MM, Block RI, Haffarnan M, Mathews MJ. Awareness during anesthesia: risk factors, causes and sequelae: a review of reported cases in the literature. *Anesth Analg* 2009; **108**: 527-535.
 34. Veselis RA. Memory formation during anaesthesia: plausibility of a neurophysiological basis. *Br J Anaesth* 2015; **115 Suppl 1**: i13-i19.
 35. Loftus EF, Pickrell JE. The formation of false memories. *Psychiatr Ann* 1995; **25**: 720-725.
 36. Pandit JJ CT, Shinde S, Ferguson K, Hitchman J, Jonker W, Odor PM, Meek T. The 'NAP5 Handbook': Concise practice guidance on the prevention and management of accidental awareness during general anaesthesia. Association of Anaesthetists of Great Britain and Ireland and the Royal College of Anaesthetists, 2019. Available from: <https://www.nationalauditprojects.org.uk/downloads/AccidentalAwareness2019.pdf>. Accessed June 15th 2019.

Figure Legends

Table 1. Eligibility criteria.

Table 2. Likelihood of a patient report representing AAGA.

Table 3. Michigan Awareness Classification Instrument. An additional designation of ‘D’ is applied where the report described distress during the experience (e.g. fear, suffocation, sense of impending death, etc.)

Table 4. Modified NPSA classification including psychological impact on the patient devised for use in NAP5

Table 5. Definitions used for phase of anaesthesia.

Figure 1. Study design flow chart.

List of Contributors

Alicja A’Court, Dina Abdel-Gadir, Ayman Abdu, Caroline Abisogun, Zainab Aboud, Judith Abrams, Andrea Ackerman, Christine Adamso, Rachel Addison, Adedayo Adeyeye, Rachel Adler, Maame Aduse-Poku, Siddharth Adyanthaya, Najmiah Ahmad, Dabeeruddeen Ahmed, Atif Ahmed, Bucky Akindele, Olubukola Akindele, Sonia Akrimi, Samar Al-Rawi, Yousif Ali, Jo Allam, Alisha Allana, Kellie Allen, Orlanda Allen, Natasha Amaradasa, Lareina Amarasekara, Freda Amoakwa-adu, Padmanabhan Anandageetha, Suresh Anandkrishnan, Rathai Anandanadesan, Michelle Anderson, Simon Apps, Audrey Aquilina, Gill Arbane, Alice Arch, Sarah Armstrong, Rita Arya, Gloria Ashiru, Keri Ashpole, Claire Atkinson, Francesca Atkinson, Elizabeth Auer, Ben Avery, Maite Babio-Galan, Hareth Bader, George Badham, Subha Bagchi, Sarah Bailey, Yolanda Baird, Cristina Balaka, Melissa Baldwin, Paul Balfour, Shreya Bali, Samantha Banks, Philip Barclay, Lucy Barnes, Thomas Barnes, Niraj Barot, Stephen Barrett, Venetia Barrett, Kate Barrett, Louise Bates, Katherine Batte, Beki Baytug, Mona Behraves, Suzanne Bell, Raul Benloch, Richard Bentley, Jennifer Berg, Charlotte Berwick, Richard Berwick, Rashmi Bhadange, Sanjoy Bhattacharyya, Egidija Bielskute, Sian Birch, Scott Bird, Ruth Bird, William Birts, Becky Black, Tatyana Blagova, Holly Blake, Oliver Blightman, Susara Blunden, Richard Bolton, Caroline Borkett-Jones, John Boselli, Manuella Bowen, Ruth Bowen, James Bowye, Henry Boyle, Zameen Brar, Jo Bray, Samantha Brayshaw, Catherine Bressington, Alison Brewer, Naomi Brice, Laura Bridge, James Briscoe, Sonia Brocklesby, Harry Brown, Sarah Brown, Derek Brunnen, Keshavareddy Burijintichenna, Sara Burnard, Agnieszka Burt, Victoria Buswell, Hannah Bykar, Matthew Cairney, Clare Calvert, Laura Camarasa, Natasha Campbell, Frederick Campbell-Jones, Jane Cantliffe, William Carrol, Joanna Carvalho, Catherine Cashell, Sarah Cassie, Kadu Cassim, Marika Chandler, Rachel Chapman, Robert Charles, Ping Chen, Deanne Cheyne, Kiran Chima, Floju Chin, Radu Chirvasuta, Martin Shao Chong, Sadia Choudhury, Priyakam Chowdhury, Tracey Christmas, Shilpa Chughwani, Sarah Ciechanowicz, Emma Clarey, Rob Coe, Jolyon Cohen, Nicholas Coker, Karen Collins, Lauren Collis, Jasna Comar, Marianne Conroy, Kathy Constantin, Jeremy Corfe, Elaine Coulborn, Vanessa Cowie, Rachel Crone, James Cronin, Jennifer Crooks, Nicola Crowther, Elizabeth Crowther, Carina Cruz, Alexa Curtis, Sarah Curtis, Sam Curtis, Adrian Dabrowicz,

1 Nick Daines, Viral Dalal, Peter Dannatt, Dinesh Das, Jennifer Dash, Katherine Davidson,
2 Sarah Davies, Yvette Davis, James Dawson, Jonathan Dean, Caroline Dean, Jane Denman,
3 Neel Desai, Preeti Dewan, Sarah Dimont, Clare Donovan, Mano Doraiswami, Kate Doughty,
4 Jamie Douglass, Michelle Dower, Samantha Downing, Wendy Duberry, Emily Duckham,
5 Lucy Dudgeon, Stephanie Dukes, Leigh Dunn, Venkat Duraiswamy, Anthony O' Dwyer,
6 Kristyn Dyer, Suresh Eapen, Mark Earl, Sally Eason, Kylie-Ellen Edwards, Zara Edwards,
7 Osinachi Egole, John Ekpa, Olivia El-Amin, Kariem El-Boghdadly, Okba Elbasir, James
8 Eldridge, Laura Elgie, Matt Ellington, Katharine Elliott, Justine Elliott, Mohamed Elmi,
9 Ramy Elnoumeir, Eunice Emeakaraoha, Matt Evans, Mary Everett, Philippa Fabb, Hadia
10 Farooq, Rohan Farrimond, Fiona Faulds, Erik Fawcett, Andrew Feneley, Dinushi Fernando,
11 Janis Ferns, Claire Finlay, Simon Fitzgerald, Doireann O'Flaherty, Mark Fleet, Lydia
12 Fletcher, Vanessa Fludder, Terri Follet, Jane Forbes, Mark Forth, Greg Foster, John Francis,
13 Kate Fraser, Lauren Friedman, Luca Fruggeri, Laura Fulton, Sam Funnell, Amit Gadre, Ajay
14 Gandhi, Harriet Gardiner, Zoe Garner, Gráinne Garvey, Tom Gately, Richard George,
15 Siobhan Gillespie, Stephanie Glover, Jennifer Goddard, Ben Goodman, Tarikere Gopal,
16 George Graham, David Green, Denise Griffin, Jessica Griffith, Stephanie Grigsby, Julie
17 Grindey, Huw Griffiths, Jonathan Groome, Christine Grother, Gabrielle Grounds, Alice
18 Groves, Abhik Guha, Amila Gunawardhana, Aman Gupta, Rekha Gupta, James Gutsell,
19 Richard Haddon, Dina Hadi, Nicola Hadjipavlou, Henry Hammerbeck, Laura Hammon,
20 Sarah Hammond, Hampesh Hampanna, Holly Hancock, Hemantha Handapangoda, Yumna
21 Haroon-Mowahed, Dawn Harpham, Grant Harris, Abigail Harrison, Divya Harshan, Andrew
22 Hartopp, Eoin Harty, Nick Haslam, Gemma Hawkins, Edith Hawkins, Sarah Hawksey,
23 Carole Hays, Tracy Hazelton, Andrew Heavyside, Chibuzo Hemeson, Katrina Henderson,
24 Oliver Henry, Lara Herbert, Nikki Higgins, James Hilton, Claire Hindmoor, Rachael
25 Hitchcock, Laura Hobbs, Michele Homsy, Cassandra Honeywell, Nazima Hoque, Kirsty
26 House, Ryan Howle, Amanda Tiller, Matthew Huniak, Jeremy Hunte, Tauqeer Husain,
27 Coralie Huson, Catriona Hussain, Tawhida Hussain, Zainab Hussein, Jeremy Hyams, Eleanor
28 Hyde, Marina Iaverdino, Alex Ignacka, Erin Innes, Stefanos Ioannidis, Rehana Iqbal, Fahja
29 Ismail, Jenny Jackson, Michael Jackson, Guy Jackson, Robyn Jacobs, Poonam Jadhav, Aisha
30 Jalaly, Lorraine James, Monica James, Surekha Jani, Chandran Jeganathan, Claire Joannides,
31 Robert Johnson, Thosa Johnson, Carolyn Johnston, ReeAnne Jones, Tom Jones, Miran Kadr,
32 Rasmeet Kainth, Jenna Kane, Rafiq Kanji, Sunil Kannanparambil, Girish Kar, Theanalli
33 Kasianandan, Husam Kaskos, Lisa Kavanagh, Richard Kaye, Leith Kelliher, Serena Kelliot,
34 Jemma Kelly, Jenna Kelly, Carol Kenyon, Laura Kessack, Samuel Kestner, Mohammed
35 Khaku, Sophia Khaleeq, Pervaz Khan, Sabeen Khan, Usamah Kidwai, Christopher King,
36 Helen King, Elizabeth Kingston, Waisun Kok, Rebekka Konig, Zheyne Konstantinova, Priya
37 Krishnan, Justin Kua, Kavitha Kuntumalla, Emira Kursumovic, Krzysztof Kurzatkowski,
38 Hariprasad Kuttambakam, Katherine Lane, Sophie Lane, Andrew Langton, Hamad Latif,
39 Norlan Lau, Shyam Laxman, Helen Laycock, Rachel Lee, Silvia Leonardi, Karen Light,
40 Heidi Lightfoot, Shuang Liu, Sanduni Liyanage, Jessica Lowe, Nuala Lucas, Miyoba Lungu,
41 Marianne Lunn, Heather Lynes, Kishore Machavarapu, Matthew Mackenzie, Jonathan
42 Major, Victoria Male, Zain Malik, Katarina Manso, Myrna Maquinana, Katarzyna Marciniak,
43 Lizzie Maronge, Chris Marsh, Carmela Martella, Nicki Martin, Nicole Martins, Jaspreet
44 Marway, Louisa Mason, Lucy Mason, Nadia Masood, James Masters, Matthew Maton-
45 Howarth, Francesca Mazzola, Tamsin McAllister, Rachel McCarthy, Claire McCormick,
46 Sharon McCready, Steffan McDougall, Lorna Mcewan, Jon McGarry, Helen McKevitt,
47 Stephanie Mckinley, Anna Mckskeane, Eloise McMaster, Martina McMonagle, Helen
48 McNamara, Hannah McPhee, Laura McRae, Dee Mead, Emma Meadows, Madhur Mehta,
49 Joanne Meikle, Yavor Metodiev, Chloe Michael, Victoria Millar, Sarah Miller, Gregg Miller,
50 Sezanne Milne, Kostas Miltsios, Lucia Misquita, Simone Misquita, Manasi Mittal, Mohamed
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Mohamed, Katherine Powell Monaghan, Joanna Monk, Alexandra Monkhouse, David
2 Monks, Laura Montague, Angela Moon, Jennifer Moran, Annette Moreton, Endaf Morgan,
3 Oliver Morgan, Danny Morland, Monica Morosan, Kimberley Morris, Andrea Morris, Clare
4 Moser, Manulella Mount, Carol Muir, McDonald Mupudzi, Mayur Murali, Iona Murdoch,
5 Heather Murray, Theresa Murray, Karen Murrell, Girish Narasimha Murthy, Diana Neeley,
6 Hannah Nei, Kerry Neil, Tara Nejm, Mark Nel, Anne Nicholson, Angela Nicklin, Catherine
7 Nolan, Tracey Nolan, Eveliina Nurmi, Bridie O'Neill, Christie Oakes, Neil Oakes, Maria
8 Ochoa-Ferraro, Nike Odeleye, Katy Oliver, Mark Oliver, Julie Onslow, Desire Onwochei,
9 Thomas Oommen, Tim Orr, Osato Osagie, Hannah Osborn, Jennie Overend, Hazel Owston,
10 Emma Pack, Pulak Padhi, Prabvathy Palani, Rajesh Pandey, Dhruvi Pandya, Navjot Panesar,
11 Con Papageorgiou, Georgios Papanastasiou, Costas Papoutsos, Suji Pararajasingham, James
12 Parry, Hasita Patel, Jaishel Patel, Jiten Patel, Kaumudi Patel, Kiran Patel, Mitul Patel,
13 Reshma Patel, Ruchira Patel, Nimisha Patel, Sangeeta Pathak, Fiona Pearson, Viktorija
14 Peciulene, Beth Peers, Benjamin Peirce, Stacey Pepper, Jasmina Perinpanayagam , Hollie
15 Perry, Nadya Petrova, Trudie Phillips, Sioned Phillips, Leonidas Phylactides, Felicity
16 Pilkington, James Plumb, Evangelia Poimenidi, Anna Sau Kuk Poon, Thomas Potter, Una
17 Poultney, Lucy Powell, Andrew Prenter, Katie Preston, Anna Price, Naomi Pritchard, Jenny
18 Pullen, Manishi Purohit, Charlotte Quamina, Jibrán Qureshi, Zahra Rajput, Stephen Ramage,
19 Tamilselvi Ramanathan, Upeka Ranasinghe, Kalum Ranatunga, Abby Rand, Seema Randive,
20 Desikan Rangarajan, Chandana Rao, Sambasiva Rao Pelluri, Antony Ratnasingham, Jamil
21 Razzaque, Anuvidya Reddy, Katie Redington, Emma Reel, Peter Remeta, Francesca Ricco,
22 Anna Riccoboni, Polly Rice, Mel Rich, Nicole Richards, Joanne Riches, Simon Ripoll, Fleur
23 Roberts, Kitty Roberts, Kay Robins, Susie Robinson, Samantha Roche, Maria Rojo, Milena
24 Carmela Romano, Hilary Rosser, Lindsay Roughley, Catriona Routley, Christine Rowley,
25 Pallab Rudra, Robin Russell, Christine Ryan, Chloe Saad, Abtin Sadeghi, Armored Salberg,
26 Matt Samuel, Rebecca Samuels, Suresh Sanapala, Seliat Sanusi, Sarbpreet Sarao,
27 Sathyabhama Sathyabhama, Zoe Saunders, Bernadetta Sawarzynska-ryszka, Panagiota
28 Sceales, Penny Sceales, Natasha Schumacher, Nuala Schwartz, Claudia Sellers, Heather
29 Sellers, Jessica Sellick, Soumen Sen, Dhaneesha Senaratne, Sarah Senbeto, Dineth
30 Seneviratna, Thunga Setty, Raj Shah, Sonia Shah, James Shambly, Saju Sharafudeen. Imran
31 Sharieff, Laurence Sharifi, Lisa Sharpe, Michael Shaw, Ian Sheldrake, Priya Shinde, Adam
32 Shonfeld. Jonathon Short, Julian Siah, Sheena Sibug. Omar Siddique, Sara Siew, Matthew
33 Simpson, Georgina Singleton, Kathryn Sinha, Aneeta Sinha, Matthew Sinnott, Harry
34 Sivadhas, Sivanth Sivakumar, Boopathi Sivarajan, Sinduja Sivarajan, Chris Skeoch, Samuel
35 Slade, Paul Slater, Camilla Smith, Carys Smith, Christopher Smith, James Smith, Lorraine
36 Smith, Annika Smith, Edward Smith, Ruth Smith, Sue Smith, Tim Smith, Helen Smithers,
37 Sue Smolen, Claire Smyth, Toni Snel, Carol Snipe, Sam Soltanifar, Nilesh Sonawane, Andal
38 Soundararaja, Emily Spence, Mark Spiliopoulos, Chhavi Srivastava, Karen Stacey, Helena
39 Stafford, Nikki Staines, Richard Stead, Emma Stevens, Alex Stilwell, Gary Stocks, Aaron
40 Stokes, Christopher Stone, Ben Straughan, Vanitha Subbarathnam, Srinath Sudunagunta,
41 Pervez Sultan, Puvan Suppiah , Priyanka Surve , Angus Sutherland, Rob Swanton, Claire
42 Swarbrick, Amy Swinson, Eleni Syrrakou, Shahrzad Tadbiri, Preetam Tamhane, Perumal
43 Tamilselvan, Andrew Tan , Shamil Tanna, Hayley Tarft, Laura Tarry, Ian Taylor, Suzanne
44 Taylor, Julie Tebbot, Svetlana Theron, Megan Thomas , Sarah Todd, Hermione Tolliday,
45 Charlotte Topham, Nicholas Tovell, Martyn Traves, Dawn Trodd, Aseem Tufchi, Katie
46 Turley, Marc Turnbull, Chris Turnbull , Oliver Turner, William Turner, Sharon Turney,
47 Eleanor Tyagi, David Uncles , Vanessa Unsworth, Pradnya Vadnere, Rama Varadan, Vik
48 Vasishta, Andrew Veal, Lalitha Vedham, Jessica Venkaya, Miriam Verghese, Icel Veronica,
49 Dinesh Vidanagamage, Rachel Vincent, Vinodhan Vyapury, Harris Wain , Fiona Walbridge,
50 Elaine Walker, Pete Walsh, Eleanor Walshe, Michelle Walters, Yize Wan, Cherry Wang,
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Kavita Wankhade, Gareth Waters , Christopher Watts, Alex Webber, Tom Wedgwood,
2 Michael Wee, Susan Wellstead, Alison White, Michael Whitear, Lucy Whitefield, Sarah
3 Wilkinson, Lauren Williams, Rhys Williams, Dawn Wilson, Samantha Wilson, Katie
4 Wimble, Elaine Winkley, Luke Winslow, Paul Winwright, Karolina Wloch, Gideon Wong,
5 Hannah Wong, Jan Man Wong, Tim Wood, Sarah Wray, Ian Wrench, James Wu, Kynn
6 Wynn, Yeng Yap, Chia Kuan Yeow, Emily Young, Alex Yusaf, Saeed Uz Zafar, Darius
7 Zeinali, Sheldon Zhang, Sarvesh Zope, Liana Zucco. Pan-London Perioperative Audit and
8 Research Network (PLAN) Committee: Sibtain Anwar, Nadia Blunt, John Cronin, Kariem
9 El-Boghdadly, Vimal Grover, Kate Grailey, Martin Gray, David Highton, Phil Hopkins,
10 Carolyn Johnston, Harriet Kemp, Helen Laycock, Queenie Lo, Daniel Martin, Clare
11 Morkane, James O'Carroll, Charles Oliver, Mitul Patel, Sioned Phillips, Benjamin Post, Anil
12 Visram, Alex Wickham.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1





Figure 1 (B&W for print purposes)



Tables

Inclusion criteria	Exclusion criteria
Adults (≥ 18 years) of $\geq 24/40$ gestation.	Patients too unwell or confused to be able to complete the questionnaire.
Received general anaesthesia (de novo or regional anaesthesia converted to general anaesthesia) for surgery with an obstetric indication. Eligible obstetric indication surgical procedures include, but are not limited to: caesarean delivery, manual removal of placenta (MROP), exploration under anaesthesia (EUA) before or after delivery.	General anaesthesia for non-obstetric indication (e.g. colorectal or orthopaedic surgery in a pregnant patient).
Written informed consent obtained.	Surgery ≥ 48 hours post-partum.
	Unable to communicate verbally/in writing in English language.

Table 1. Eligibility criteria.

Likelihood that the report represents AAGA	Description
Certain/Probable	A report of AAGA in a “surgical setting” in which the detail of the patient story was judged consistent with AAGA, especially where report detail was verified independently or contained descriptions that would not otherwise be known to a patient.
Possible	A report of AAGA in a “surgical setting” in which details were judged to be consistent with AAGA or the circumstances might have reasonably led to AAGA, but where otherwise the report lacked a degree of verifiability or detail.
Unlikely	Details of the patient story were deemed unlikely, or judged to have occurred outside of the period of anaesthesia.
None	Evidence that the report was not AAGA or events occurred outside of the period of anaesthesia.

Table 2. Likelihood of a patient report representing AAGA.

MAC classification	Description
Class 0	No AAGA
Class 1	Isolated auditory perceptions
Class 2	Tactile perceptions (with or without auditory)
Class 3	Pain (with or without tactile or auditory)

Class 4	Paralysis (with or without tactile or auditory)
Class 5	Paralysis and pain (with or without tactile or auditory)

Table 3. Michigan Awareness Classification Instrument. An additional designation of ‘D’ is applied where the report described distress during the experience (e.g. fear, suffocation, sense of impending death, etc.)

Severity	Revised definitions used in NAP5
0	No harm occurred
1	Resolved (or likely to resolve) with no or minimal professional intervention. No consequences for daily living, minimal or no continuing anxiety about future healthcare
2	Moderate anxiety about future anaesthesia or related healthcare. Symptoms may have some impact on daily living. Patient has sought or would likely benefit from professional intervention
3	Striking or long-term psychological effects that have required, or might benefit from professional intervention or treatment: severe anxiety about future healthcare and/or impact on daily living. Recurrent nightmares or adverse thoughts or ideations about events. This may also result in formal complaint or legal action (but these alone may not be signs of severity)
4	Caused death

Table 4. Modified NPSA classification including psychological impact on the patient devised for use in NAP5

Phase of anaesthesia	Description
Pre-induction	Defined as drug error before intended anaesthesia.
Induction	defined as the start of administration of a hypnotic drug with the intention of producing general anaesthesia. Cases in which there is a drug administration error during induction, resulting in failure to deliver a hypnotic agent, will be included as induction phase, since the anaesthetist’s intention was to induce general anaesthesia irrespective of any drug error.
Intraoperative	Defined as the period between skin incision and cessation of the administration of maintenance anaesthesia agents.
Emergence	Defined as occurring after surgery has ended until a time when the patient feels awake and expects to be conscious. Emergence phase reports include residual neuromuscular blockade, but we will categorise uncomplicated “awake” emergence memories of extubation as non-AAGA, since delivery of anaesthesia has intentional ceased and the patient

	is expected to have sufficient return of consciousness and reversal of neuromuscular blockade for extubation.
Other	Uncertain time.

Table 4. Definitions used for phase of anaesthesia.