

$\Sigma$ -MATRIX IDEALS AND  $\Sigma$ -INVERTING HOMOMORPHISMS

by

NICHOLAS JOHN GREENWOOD

thesis for the degree of

M.Phil

UNIVERSITY COLLEGE LONDON

May 1993


ProQuest Number: 10044417

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10044417

Published by ProQuest LLC(2016). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code.  
Microform Edition © ProQuest LLC.

ProQuest LLC  
789 East Eisenhower Parkway  
P.O. Box 1346  
Ann Arbor, MI 48106-1346

## Abstract

In Free Rings and their Relations, P.M.Cohn constructed a skew field from a prime matrix ideal using admissible matrices. When  $\Sigma$  is a lower multiplicative set of matrices over any non-commutative ring this method can be generalised to construct any epic  $\Sigma$ -inverting homomorphism upto isomorphism. This depends on the introduction of the concept of a  $\Sigma$ -matrix ideal. Every  $\Sigma$ -inverting homomorphism gives rise to a  $\Sigma$ -matrix ideal and conversely our main theorem shows that, given a  $\Sigma$ -matrix ideal, an epic  $\Sigma$ -inverting homomorphism can be constructed and that the matrices which are admissible for zero are precisely those lying in the  $\Sigma$ -matrix ideal.

It is shown that the least  $\Sigma$ -matrix ideal induces the universal  $\Sigma$ -inverting homomorphism. A description of the least  $\Sigma$ -matrix ideal is then obtained yielding a new description of the kernel of the  $\Sigma$ -inverting homomorphism and a criterion for it to be an embedding; Malcolmson and Gerasimovs' respective descriptions of the kernel are also proved.

When  $\Sigma$  is taken to be the complement of a prime matrix ideal the construction reduces to that used by Cohn to construct a skew field.

It is further shown that the definition of a prime matrix ideal  $\mathcal{P}$  can be simplified by restricting the class of matrices necessarily lying in  $\mathcal{P}$  to be the hollow and degenerate matrices. The condition that  $\mathcal{P}$  be closed with respect to row determinantal sums can be dropped completely. As a consequence, Cohn's criterion for the existence of a homomorphism from a ring to a field and the criterion for the existence of a field of fractions for a ring can be refined somewhat. For completeness Dicks and Sontag's result

that Sylvester domains form the precise class of rings which have a universal field of fractions inverting all full matrices is also included.

## **Acknowledgement**

I wish to acknowledge the guidance received from my supervisor Professor P.M.Cohn. Without his constructive criticisms and suggestions this work would not have been possible. Also I would like to thank my joint supervisor Dr.W.Stephenson for his friendly help and advice.

This research was supported by a studentship from the Science and Engineering Research Council.

## Contents

Abstract	2
Acknowledgement	4
Chapter 1 $\Sigma$ -Inverting Homomorphisms	6
Chapter 2 $\Sigma$ -Matrix Ideals	15
Chapter 3 Universal Localisation	32
Chapter 4 The Criteria of Malcolmson and Gerasimov	40
Chapter 5 Prime Matrix Ideals	50
Chapter 6 Fields of Fractions	62
References	67

We shall be concerned with inverting certain sets of matrices over a non-commutative ring or more generally homomorphisms which invert certain matrices. In the commutative case this reduces to inverting certain sets of elements of the ring since we can invert any square matrix  $A$  by adjoining an inverse of  $\det A$ . However for a non-commutative ring we shall have to adjoin all the entries in the matrix's inverse. In this chapter we consider general  $\Sigma$ -inverting homomorphisms and in particular the universal localisation at a set of matrices. We conclude with a description of the universal localisation in terms of the universal right localisation.

Definitions If  $R$  is any ring and  $\Sigma$  a set of matrices over  $R$  then a ring homomorphism  $f:R \rightarrow S$  is said to be  $\Sigma$ -inverting if every matrix in  $\Sigma$  is mapped to an invertible matrix over  $S$ .

Similar notions of  $f$  being  $\Sigma$ -right (left) inverting are clear.

If  $f:R \rightarrow S$  is a ring homomorphism and  $\Sigma$  a set of matrices over  $R$  then we define  $R_{\Sigma, f}(S)$  as the set of all final components to solutions  $u$  over  $S$  of equations,

$$Au = A_0 \text{ where } A \in \Sigma f \text{ and } A_0 \text{ is a column over } \text{im} f. \quad (1)$$

When the context is clear we write  $R_{\Sigma}(S)$  for  $R_{\Sigma, f}(S)$ .

We will have to impose some restrictions on  $\Sigma$  so we also make the following definition:

A set of matrices  $\Sigma$  over a ring is said to be **lower multiplicative** if  $A, B \in \Sigma$  implies that  $\begin{bmatrix} A & 0 \\ C & B \end{bmatrix} \in \Sigma$  where  $C$  is any matrix of the appropriate size and  $1 \in \Sigma$ .

Lemma 1.1 Let  $f:R \rightarrow S$  be a ring homomorphism and  $\Sigma$  a lower multiplicative set of matrices over  $R$ . Then  $R_{\Sigma}(S)$  is a subring of  $S$  containing  $\text{im}f$ .

Proof. If  $c \in \text{im}f$  then  $c=af$  for some  $a \in R$  and  $c$  satisfies the equation  $1.u=af$ . Hence  $\text{im}f \subseteq R_{\Sigma}(S)$ , so  $R_{\Sigma}(S)$  is non-empty containing in particular  $0$  and  $1$ .

Let  $p$  be the final component of the solution  $u=(u_* p)^T$  of (1) where  $A=(A_* A_{\infty})$  and  $A_{\infty}$  is the final column of  $A$  and also let  $q$  be the final component of the solution  $v$  of  $Bv=B_0$  where  $B \in \Sigma f$  etc. Then  $p-q$  is the final component of a solution  $w$  of the equation,

$$\begin{bmatrix} B_* & B_{\infty} & 0 & 0 \\ 0 & A_{\infty} & A_* & A_{\infty} \end{bmatrix} w = \begin{bmatrix} B_0 \\ A_0 \end{bmatrix}$$

Namely  $w=(v^T u_* p-q)^T$  is a solution. Similarly  $pq$  is the final component of a solution  $w$  of the equation,

$$\begin{bmatrix} B_* & B_{\infty} & 0 & 0 \\ 0 & -A_0 & A_* & A_{\infty} \end{bmatrix} w = \begin{bmatrix} B_0 \\ 0 \end{bmatrix}$$

That is to say  $w=(v^T u_* q pq)^T$  is a solution. This shows that  $R_{\Sigma}(S)$  is closed under subtraction and multiplication and since we have already shown that  $1 \in R_{\Sigma}(S)$  we have that  $R_{\Sigma}(S)$  is a subring of  $S$  as claimed. ■

Now any given equation (1) need not necessarily have any solution in  $S$ . However if  $f$  is  $\Sigma$ -right inverting then every equation has a (not necessarily unique) solution. In this case we have the

following characterisation of  $R_\Sigma(S)$ .

**Theorem 1.2** *If  $R$  is any ring,  $\Sigma$  a lower multiplicative set of matrices over  $R$  and let  $f:R \rightarrow S$  be a  $\Sigma$ -right inverting ring homomorphism. Then  $x \in R_\Sigma(S)$  if and only if  $x$  is an entry in the right inverse of the image of a matrix in  $\Sigma$ .*

**Proof.** ( $\Leftarrow$ )  $x$  is (say) the  $i$ th entry in a column  $u = (u_* u_n)^T$  which satisfies the equation,

$$Au = e_j \quad A \in \Sigma_n f; \quad n \geq j \in \mathbb{N} .$$

[ $e_j$  is the column which is zero except in the  $j$ th entry which is one and likewise  $e_\infty$  is zero except in the final entry.] If  $i=n$  then the conclusion follows by the definition of  $R_\Sigma(S)$ . Otherwise we note that the following holds;

$$\begin{bmatrix} A_* & A_\infty & 0 & 0 \\ C & A_\infty & A_* & A_\infty \end{bmatrix} \begin{bmatrix} u_*^T \\ u_n \\ u_*^T \\ x \end{bmatrix} = \begin{bmatrix} e_j \\ e_j \end{bmatrix} ,$$

where  $C$  has as  $i$ th column  $-A_\infty$  and the rest zero. Since the matrix on the L.H.S. lies in  $\Sigma f$  and  $e_j$  has entries in  $\text{im} f$  the conclusion follows.

( $\Rightarrow$ )  $x$  is the final component of the solution to (1). Now  $A$  has a right inverse over  $S$ , say  $B$ . Then

$$\begin{bmatrix} 1 & 0 \\ -A_\infty & A \end{bmatrix} \begin{bmatrix} 1 & 0 \\ u & B \end{bmatrix} = I .$$

Hence  $x$  is an entry in the right inverse of an element of  $\Sigma f$ . ■

This result highlights the one-sided nature of the definition of

$R_\Sigma(S)$ . We shall return to this matter at the end of Chapter 2. When  $R_\Sigma(S)=S$  for a  $\Sigma$ -inverting homomorphism  $f:R \rightarrow S$  then we shall call  $f$  **epic  $\Sigma$ -inverting**.

If we have an equation (1) then we have an important analogue of Cramer's Rule;

**Proposition 1.3** *Let  $R$  be any ring,  $\Sigma$  a lower multiplicative set of matrices over  $R$  and  $f:R \rightarrow S$  a  $\Sigma$ -inverting homomorphism. If we have an equation  $Au=A_0$  over  $S$  where  $u=(u_* p)^T$  has final component  $p$ ,  $A=(A_* A_\infty) \in \Sigma f$  and  $A_0$  is a column over  $imf$  then  $p$  is stably associated to  $(A_0 A_*)$  over  $S$ .*

Proof. If so we have

$$(A_0 A_*) = (A_* A_\infty) \begin{bmatrix} u_*^T & I \\ p & 0 \end{bmatrix}, \text{ and } (A_* A_\infty) \text{ is invertible over } S. \quad \blacksquare$$

**Definition** For a set  $\Sigma$  of matrices over a ring  $R$  a universal  $\Sigma$ -inverting homomorphism is a homomorphism  $\lambda_\Sigma:R \rightarrow R_\Sigma$  such that any  $\Sigma$ -inverting homomorphism  $f:R \rightarrow S$  factors uniquely through  $\lambda$  via a homomorphism  $f':R_\Sigma \rightarrow S$ . The ring  $R_\Sigma$  is determined up to isomorphism by these conditions and is called the universal  $\Sigma$ -inverting ring or the universal localisation of  $R$ . This was first considered by P.M.Cohn [1] and below is his proof of its existence. Similarly there is a universal  $\Sigma$ -right inverting homomorphism  $\lambda_{(\Sigma)}:R \rightarrow R_{(\Sigma)}$ .

**Theorem 1.4** (Cohn) *Let  $R$  be any ring and  $\Sigma$  any set of matrices over  $R$ . Then there is a universal [right] localisation  $R_\Sigma$  [ $R_{(\Sigma)}$ ], unique up to isomorphism, with a universal  $\Sigma$ -[right] inverting homomorphism*

$$\lambda_\Sigma:R \rightarrow R_\Sigma \quad [ \lambda_{(\Sigma)}:R \rightarrow R_{(\Sigma)} ] .$$

Moreover,  $\lambda_\Sigma [\lambda_{(\Sigma)}]$  is injective if and only if  $R$  can be embedded in a ring over which all the matrices of  $\Sigma$  have [right] inverses.

Proof. For each  $m \times n$  matrix  $A=(a_{ij})$  in  $\Sigma$  we take a set of  $mn$  symbols, arranged as an  $n \times m$  matrix  $A'=(a'_{ji})$  and take a ring presentation of  $R_\Sigma [R_{(\Sigma)}]$  consisting of all the elements of  $R$ , as well as all the  $a'_{ji}$  as generators, and as defining relations take all the relations holding in  $R$ , together with the relations, in matrix form,

$$AA'=A'A=I \text{ for each } A \in \Sigma. \quad [AA'=I \text{ each } A \in \Sigma]$$

The mapping taking each element of  $R$  to the corresponding element of  $R_\Sigma [R_{(\Sigma)}]$  is clearly a homomorphism  $\lambda_\Sigma: R \rightarrow R_\Sigma [\lambda_{(\Sigma)}: R \rightarrow R_{(\Sigma)}]$ , which is  $\Sigma$ -[right] inverting by construction. If  $f: R \rightarrow S$  is any  $\Sigma$ -[right] inverting homomorphism, we define a homomorphism  $f': R_\Sigma \rightarrow S [f': R_{(\Sigma)} \rightarrow S]$  by putting  $xf'=xf$  for all  $x \in R$  and for any matrix  $A \in \Sigma$  defining  $f'$  on  $(A\lambda)^{-1}$  by putting  $(A\lambda)^{-1}f'=(Af)^{-1}$ . This gives a well defined homomorphism  $f'$ , because any relation in  $R_\Sigma [R_{(\Sigma)}]$  is a consequence of the defining relations in  $R$  and the relations above and all these relations also hold in  $S$ . If there is a  $\Sigma$ -[right] inverting homomorphism  $f$  which is injective then it must factor through  $\lambda_\Sigma [\lambda_{(\Sigma)}]$  which then must also be injective. ■

It is now possible to show how  $R_\Sigma$  is related to  $R_{(\Sigma)}$  when  $\Sigma$  is a lower multiplicative set of matrices over  $R$ .

Lemma 1.5 If  $f: R \rightarrow S$  is a ring homomorphism and  $\Sigma$  a lower multiplicative set of matrices over  $R$  then the set  $J_{\Sigma, f}(S)$  of final components of solutions  $u$  to equations,

$$Au = 0 \quad \text{where} \quad A \in \Sigma f \quad (2)$$

is a two sided ideal of  $R_{\Sigma, f}(S)$ .

Proof. If  $p, q$  are the final components of solutions  $u, v$  respectively to equations  $Au=0$  and  $Bv=0$  where  $A, B \in \Sigma f$  then  $p-q$  is the final component to the solution  $w$  of

$$\begin{bmatrix} B_* & B_\infty & 0 & 0 \\ 0 & A_\infty & A_* & A_\infty \end{bmatrix} w = 0 \quad .$$

$J_\Sigma(S)$  is non-empty since zero is easily seen to be a member. If  $r \in R_\Sigma(S)$  and (2) holds then  $Aur=0$ . It remains to show closure under multiplication by elements of  $R_\Sigma(S)$  on the left. By the definition of  $R_\Sigma(S)$   $r$  is the final component of the solution  $v$  to say  $Cv=C_0$  where  $C \in \Sigma f$  and  $C_0$  has entries in  $\text{im}f$ . Then  $rp$  is the final component of a solution  $w$  of,

$$\begin{bmatrix} A_* & A_\infty & 0 & 0 \\ 0 & -C_0 & C_* & C_\infty \end{bmatrix} w = 0.$$

Hence  $rp \in J_\Sigma(S)$  and  $J_\Sigma(S)$  is a two sided ideal of  $R_\Sigma(S)$ . ■

When  $\Sigma$  is lower multiplicative then by Lemma 1.1 we have that  $R_{\Sigma, \lambda(\Sigma)}(R_\Sigma)$  is a subring of  $R_\Sigma$  and from the ring presentation construction of the universal  $\Sigma$ -right inverting homomorphism and Theorem 1.2 it is clear that  $R_\Sigma(R_\Sigma)$  contains all of the generators of  $R_\Sigma$  hence  $R_\Sigma(R_\Sigma) = R_\Sigma$ . Consequently  $R_\Sigma$  has a two sided ideal  $J$  where every element of  $J$  is the final component of the solution  $u$  to an equation  $Au=0$  where  $A \in \Sigma \lambda(\Sigma)$ .

Theorem 1.6 If  $R$  is any ring,  $\Sigma$  a lower multiplicative set of matrices over  $R$  then the universal  $\Sigma$ -inverting homomorphism  $\lambda_{\Sigma}: R \rightarrow R_{\Sigma}$  factors through the universal  $\Sigma$ -right inverting homomorphism  $\lambda_{(\Sigma)}: R \rightarrow R_{(\Sigma)}$  as follows;

$$R \xrightarrow{\lambda_{(\Sigma)}} R_{(\Sigma)} \xrightarrow{\nu} R_{(\Sigma)/J} \cong R_{\Sigma}$$

where  $J$  is the two sided ideal of  $R_{(\Sigma)}$  consisting of final components of solutions to equations  $Au=0$ ,  $A \in \Sigma \lambda_{(\Sigma)}$  and  $\nu$  is the natural homomorphism. Also  $\ker \lambda_{\Sigma} = \lambda_{(\Sigma)}^{-1}(J)$ .

Proof. Write  $\lambda = \lambda_{(\Sigma)} \cdot \nu$  then  $\lambda$  is  $\Sigma$ -right inverting since  $\lambda_{(\Sigma)}$  is  $\Sigma$ -right inverting.

Now we shall denote by  $\bar{r}$  the equivalence class of  $R_{(\Sigma)/J}$  which has  $r$  as a representative. If  $\bar{A}\bar{u}=0$  in  $R_{(\Sigma)/J}$  where  $A \in \Sigma \lambda_{(\Sigma)}$  then we have an equation  $(A+U)(u+u_2)=u_3$  in  $R_{(\Sigma)}$  where  $U, u_2$  and  $u_3$  all have entries in  $J$ . Expanding and rearranging we get

$$Au = u_3 - Au_2 - Uu - Uu_2 .$$

All the terms on the right hand side are over  $J$  since  $J$  is a two sided ideal of  $R_{(\Sigma)}$ . Say  $Au=u_4 \in J$ . Since  $A$  lies in  $\Sigma \lambda_{(\Sigma)}$  it has a right inverse,  $B$  say. Then  $Au=ABu_4$  and so  $A(u-Bu_4)=0$  in  $R_{(\Sigma)}$ . Hence by the definition of  $J$  the final component of  $u-Bu_4$  lies in  $J$ . But again  $Bu_4$  has entries in  $J$  since  $u_4$  is over  $J$ , so the final component of  $u$  lies in  $J$  also. Therefore the final component of  $\bar{u}$  is zero. It is straightforward to show that the other entries in  $\bar{u}$  are zero since we can use the trick employed in Theorem 1.2 to write an equation in which they occur as final components. We have shown therefore that  $\bar{A}\bar{u}=0$  implies that  $\bar{u}=0$ . Since

$$\bar{A}(\bar{B}\bar{A} - I) = (\bar{A}\bar{B})\bar{A} - \bar{A} = I\bar{A} - \bar{A} = 0,$$

we must have  $\bar{B}\bar{A}-I=0$  and we see that  $\lambda$  is  $\Sigma$ -invertible.

Any  $\Sigma$ -invertible homomorphism  $f:R\rightarrow S$  factors through the universal  $\Sigma$ -right invertible homomorphism via a unique homomorphism  $f'$  say.

$$\begin{array}{ccc}
 & R(\Sigma) & \\
 \lambda(\Sigma) \nearrow & & \searrow f' \\
 R & \xrightarrow{f} & S
 \end{array}$$

Now  $J \subseteq \ker f'$  since otherwise there is an equation in which a member of  $\Sigma f$  is a left zero divisor. Hence  $f'$  factors through  $\nu$  and the following diagram commutes.

$$\begin{array}{ccc}
 & \nu & \\
 R(\Sigma) & \xrightarrow{\quad} & R(\Sigma/J) \\
 \lambda(\Sigma) \uparrow & \searrow f' & \downarrow f'' \\
 R & \xrightarrow{f} & S
 \end{array}$$

So  $f$  factors through  $\lambda$  via  $f''$  and this is unique by the uniqueness of  $f'$  so since  $\lambda$  is  $\Sigma$ -invertible we see that  $\lambda$  is the universal  $\Sigma$ -invertible homomorphism. Also it is immediate that  $\ker \lambda_{\Sigma} = \lambda_{\Sigma}^{-1}(J)$ . ■

In fact P.M.Cohn has shown in [2] that any ring  $R$  can be embedded in a fully right invertible ring  $S$  such that every left regular matrix over  $R$  is mapped to a right invertible matrix over  $S$ . What this implies is the following:

**Theorem 1.7** (Cohn [2]) *If  $R$  is any ring and  $\Sigma$  any set of matrices over  $R$  then the universal  $\Sigma$ -right invertible homomorphism is an*

embedding if and only if  $\Sigma$  does not contain any right zero  
divisors. ■

This chapter contains our main construction. From the definition of a  $\Sigma$ -matrix ideal given later we generalise Cohn's construction using admissible matrices (Chap.7 [1]) to show that every  $\Sigma$ -matrix ideal gives rise to an epic  $\Sigma$ -inverting ring homomorphism and conversely every  $\Sigma$ -inverting homomorphism factors through such a map.

From Theorem 1.2 we see that when  $\Sigma$  is lower multiplicative then formally attaching to a ring  $R$  the entries in right inverses of elements of  $\Sigma$  is equivalent to attaching the final components of solutions  $u$  to equations  $Au=A_0$ , where  $A \in \Sigma$ . This latter system is more conveniently written  $(A_0 \ A_* \ A_\infty)$  where  $A=(A_*A_\infty)$ ,  $A_\infty$  is the final column of  $A$  and  $A_*$  the remaining columns of  $A$ . The matrix  $(A_*A_\infty) \in \Sigma$  which we aim to invert will be called the **denominator** of the **admissible matrix**  $(A_0 \ A_* \ A_\infty)$ ; the matrix  $(A_*)$  will be called the **core**; and noting Cramer's rule (Proposition 1.3), the **numerator** of the admissible matrix is the matrix  $(A_0 \ A_*)$ .

Definition If  $R$  is any ring and  $\Sigma$  a lower multiplicative set of matrices over  $R$  then we define  $M_\Sigma$  to be the set of all admissible matrices, i.e. all matrices  $A$  such that  $A=(A_0 \ A_* \ A_\infty)$  with  $(A_*A_\infty) \in \Sigma$  and  $A_0$  a column over  $R$ . For any two admissible matrices  $A, B$  we will write  $A \rightarrow B$  if they are obtainable from each other by a series of the following invertible **permissible operations**;

(a) row operations, i.e multiplication on the left by an invertible matrix over  $R$ .

(b) column operations within the core  $A_*$  over the characteristic ring and we also permit the addition of a right multiple (over the characteristic ring of  $R$ ) of a column of the

core to  $A_\infty$  i.e. this amounts to multiplication on the right by a half bordered matrix over the characteristic ring of R where a half bordered matrix is an invertible matrix of the form  $1 \oplus Q$  where the bottom row of Q is  $e_\infty^T$ .

The above relation is easily shown to be an equivalence and by  $M_\Sigma$  we shall understand  $M_\Sigma / \sim$ . Any application of the above operations will be denoted by an arrow.

Definition. If  $f: R \rightarrow S$  is a  $\Sigma$ -inverting homomorphism then  $(A_\circ A_* A_\infty) \in M_\Sigma$  is said to be **admissible for zero** under  $f$  if  $e_\infty^T (A_* f A_\circ f)^{-1} A_\circ f = 0$ . Similarly a subset of  $M_\Sigma$  is said to be **admissible for zero** under  $f$  if every member of the set has that property.

We now proceed by defining a commutative semigroup structure on  $M_\Sigma$  as follows;

$$A \square B = \begin{bmatrix} B_\circ & B_* & B_\infty & 0 & 0 \\ A_\circ & 0 & -A_\infty & A_* & A_\infty \end{bmatrix} \quad (1)$$

Firstly this is compatible with the permissible operations (a) and (b), i.e. if  $A \rightarrow \hat{A}$  then  $A \square B \rightarrow \hat{A} \square B$  etc., and hence is well defined.

Also  $A \square B \in M_\Sigma$  since  $\Sigma$  is lower multiplicative and A and B are admissible. Now we verify that  $\square$  is an associative operation:

$$(A \square B) \square C = \begin{bmatrix} C_\circ & C_* & C_\infty & 0 & 0 & 0 & 0 \\ B_\circ & 0 & 0 & B_* & B_\infty & 0 & 0 \\ A_\circ & 0 & -A_\infty & 0 & -A_\infty & A_* & A_\infty \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} C_0 & C_* & C_\infty & 0 & 0 & 0 & 0 \\ B_0 & 0 & -B_\infty & B_* & B_\infty & 0 & 0 \\ A_0 & 0 & 0 & 0 & -A_\infty & A_* & A_\infty \end{bmatrix} = A \square (B \square C)$$

So  $M_\Sigma$  forms a semigroup with respect to this operation.

Commutativity follows as easily;

$$A \square B = \begin{bmatrix} B_0 & B_* & B_\infty & 0 & 0 \\ A_0 & 0 & -A_\infty & A_* & A_\infty \end{bmatrix} \rightarrow \begin{bmatrix} A_0 & 0 & -A_\infty & A_* & A_\infty \\ B_0 & B_* & B_\infty & 0 & 0 \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} A_0 & A_* & A_\infty & 0 & 0 \\ B_0 & 0 & -B_\infty & B_* & B_\infty \end{bmatrix} = B \square A .$$

It is also possible to define another binary operation on  $M_\Sigma$ , multiplication, which is also associative.

$$A.B = \begin{bmatrix} B_0 & B_* & B_\infty & 0 & 0 \\ 0 & 0 & -A_0 & A_* & A_\infty \end{bmatrix} \quad (2)$$

Again this is compatible with the permissible operations and so is a well defined binary operation. Also since  $\Sigma$  is lower multiplicative  $A.B$  lies in  $M_\Sigma$  if  $A$  and  $B$  lie in  $M_\Sigma$ . The operation is immediately seen to be associative. Hence we have shown;

Proposition 2.1 *If  $R$  is any ring and  $\Sigma$  a lower multiplicative set of matrices over  $R$  then the set of admissible matrices  $M_\Sigma$  is a*

commutative semigroup with respect to addition ' $\square$ ' defined by (1) and is a semigroup with respect to multiplication '.' defined by (2). ■

We now investigate when  $M_{\Sigma}$  can have a ring structure imposed on it. First we need a lemma on commutative semigroups. If  $H$  is a subset of a commutative semigroup then  $H$  is said to be **unitary** if  $a \in H$  and  $a+b \in H$  imply that  $b \in H$ . The following is a commutative version of a result from Clifford and Preston [3].

Lemma 2.2 *Let  $S$  be a non-empty commutative semigroup,  $U$  a unitary subsemigroup of  $S$  and let there exist a unary operation  $^{\circ}: S \rightarrow S$  such that  $m+m^{\circ} \in U$  for any  $m \in S$ . Then the relation ' $\sim$ ' defined by*

$$a \sim b \quad \text{iff } \exists x \in S; a+x, b+x \in U \quad (3)$$

*is a congruence on  $S$ , the quotient  $S/\sim$  is an abelian group and moreover if  $e$  is a representative of the equivalence class which is the identity in  $S/\sim$  then  $a \sim e$  if and only if  $a \in U$ .*

Proof. It is clear that ' $\sim$ ' is symmetric by definition and  $a+a^{\circ} \in U$  for any  $a \in S$  hence  $a \sim a$ . If  $a \sim b, b \sim c$  with  $a+x, b+x, b+y, c+y \in U$  then  $a+x+b+y = (a+y)+(b+x) \in U$ . Since  $U$  is unitary and  $b+x \in U$  then  $a+y \in U$  and  $a \sim c$ . Also for any  $d \in S$  we have  $(a+x)+(d+d^{\circ}), (b+x)+(d+d^{\circ}) \in U$  hence  $a+d \sim b+d$  and we have established that (3) defines a congruence on  $S$ .

Let  $S/U$  denote  $S/\sim$  the quotient semigroup which is commutative since  $S$  is commutative. Since  $U$  is closed under addition it is clear that  $U$  is contained in a single equivalence class. Conversely if  $b \sim u \in U$  then  $b+x, u+x \in U$  so  $u \in U$  implies that  $x \in U$  which in turn implies that  $b \in U$ . We denote the equivalence class of  $U$  by  $0$ . Now  $b+0=b$  in  $S/U$  since  $e, b+e+x \in U$  imply that  $b+x \in U$  and  $x=(b+e)^{\circ}$  gives

$b+e+x \in U$ . So 0 is the identity element in  $S/U$ . That  $S/U$  is a group follows since  $b+b^0 \in U$  for any  $b \in S$ . ■

If  $\Sigma$  is a lower multiplicative set of matrices over a ring  $R$  then we consider the case when  $S$  is taken to be  $M_\Sigma$ ; the commutative semigroup of admissible matrices with respect to addition '□'. In this case  $U$  will be the unitary subsemigroup of admissible matrices which are going to represent zero in the quotient semigroup. We know that the unary operation  $^0: M_\Sigma \rightarrow M_\Sigma$  corresponds to finding an additive inverse for each  $m \in M_\Sigma/U$ . Within our construction we define such an operation by

$$(A_0 \ A_* \ A_\infty)^0 = (-A_0 \ A_* \ A_\infty) . \quad (4)$$

Since if  $(A_* \ A_\infty)$  is invertible (under some homomorphism to a ring) then,  $e_\infty^T (A_* \ A_\infty)^{-1} (-A_0) = -e_\infty^T (A_* \ A_\infty)^{-1} A_0$ . Taking (4) as the definition of the unary operation we see that  $U$  must contain all matrices of the form  $(A_0 \ A_* \ A_\infty) \square (-A_0 \ A_* \ A_\infty)$ . This sum can be written as follows;

$$\begin{bmatrix} -A_0 & A_* & A_\infty & 0 & 0 \\ A_0 & 0 & -A_\infty & A_* & A_\infty \end{bmatrix} \rightarrow \begin{bmatrix} -A_0 & A_* & A_\infty & 0 & 0 \\ 0 & 0 & 0 & A_* & A_\infty \end{bmatrix}$$

In fact we shall impose a stronger condition on  $U$ . Define  $\mathcal{L}$  to be the set of admissible matrices which after a sequence of permissible operations can be written;

$$\begin{bmatrix} K_0 & K & 0 \\ 0 & 0 & L \end{bmatrix} \text{ where } K, L \in \Sigma \text{ and } K_0 \text{ is a column.}$$

Certainly it is easy to see that any matrix in  $\mathcal{L}$  must be admissible for zero under any  $\Sigma$ -inverting homomorphism. For this reason  $\mathcal{L}$  will

be called the minimal set of matrices admissible for zero. It is important to realise that  $\mathcal{L}$  need not contain all the matrices admissible for zero. We are now in a position to define a  $\Sigma$ -matrix ideal.

Definition Let  $R$  be any ring and  $\Sigma$  a lower multiplicative set of matrices over  $R$ . Then a  $\Sigma$ -matrix ideal  $\mathfrak{U}$  of  $R$  is a well defined subset of  $M_\Sigma$  with the following properties:

- (i)  $\mathfrak{U}$  contains  $\mathcal{L}$ , the minimal set of matrices admissible for zero.
- (ii)  $\mathfrak{U}$  is an additive unitary subsemigroup of  $M_\Sigma$ .
- (iii)  $\mathfrak{U}$  is a multiplicative semigroup ideal of  $M_\Sigma$ .

We note that the intersection of any family of  $\Sigma$ -matrix ideals is also a  $\Sigma$ -matrix ideal and hence we can talk of the least  $\Sigma$ -matrix ideal which we shall write  $\mathfrak{U}_\Sigma$ . This  $\Sigma$ -matrix ideal has important properties which are taken up in the next chapter. Now we investigate the properties of a general  $\Sigma$ -matrix ideal. Firstly we note that it is straightforward to prove that a  $\Sigma$ -inverting homomorphism gives rise to a  $\Sigma$ -matrix ideal.

Lemma 2.3 Let  $R$  be a ring,  $\Sigma$  a lower multiplicative set of matrices over  $R$  and  $f:R \rightarrow S$  a  $\Sigma$ -inverting homomorphism then  $\Sigma\text{-Ker}f = \{(A_{\infty} A_{\infty} A_{\infty}) \in M_\Sigma \mid e_{\infty}^T (A_{\infty} f A_{\infty} f)^{-1} A_{\infty} f = 0\}$  is a  $\Sigma$ -matrix ideal. ■

Given a  $\Sigma$ -matrix ideal for a ring  $R$  we can construct an  $R$ -ring as follows;

Theorem 2.4 Let  $R$  be any ring and let  $\Sigma$  be a set of lower multiplicative matrices over  $R$ . If  $\mathfrak{U}$  is a  $\Sigma$ -matrix ideal of  $R$  then  $M_\Sigma/\mathfrak{U}$  has a ring structure and the map  $\lambda_{\mathfrak{U}}:R \rightarrow M_\Sigma/\mathfrak{U}$  given by  $\lambda_{\mathfrak{U}}(r) =$

(r 1) is a ring homomorphism from  $R$  to  $M_\Sigma/\mathfrak{U}$ . Moreover  $M_\Sigma/\mathfrak{U}$  is the zero ring if and only if  $\mathfrak{U}=M_\Sigma$ .

Proof. From the above remarks  $\mathfrak{U}$  satisfies the conditions of Lemma 2.2 with respect to the unary operation defined in (4). Applying Lemma 2.2,  $M_\Sigma/\mathfrak{U}$  is an abelian group under ' $\square$ ' and we have  $A \sim B$  if and only if  $A \square B^0 \in \mathfrak{U}$ . Since  $M_\Sigma$  has a multiplication defined on it we define multiplication on  $M_\Sigma/\mathfrak{U}$  in the obvious way. It remains to be shown that this operation is well defined. However we can prove the following identities in  $M_\Sigma/\mathfrak{U}$  for any  $(A_0, A_*, A_\infty), P \in \Sigma$  by noting that  $\mathfrak{U}$  contains  $\mathcal{L}$ .

$$\begin{bmatrix} 0 & P & 0 & 0 \\ A_0 & Q & A_* & A_\infty \end{bmatrix} \sim (A_0 \ A_* \ A_\infty) \quad \dots\dots(5)$$

$$\begin{bmatrix} B_0 & P & B_* & B_\infty \\ A_0 & 0 & A_* & A_\infty \end{bmatrix} \sim (A_0 \ A_* \ A_\infty) \quad \dots\dots(6)$$

whenever the denominator of the l.h.s of (6) lies in  $\Sigma$ . We prove these by showing  $(L.H.S.) \square (R.H.S.)^0 \in \mathcal{L}$ , so we have for (5);

$$\begin{bmatrix} -A_0 & A_* & A_\infty & 0 & 0 & 0 \\ 0 & 0 & 0 & P & 0 & 0 \\ A_0 & 0 & -A_\infty & Q & A_* & A_\infty \end{bmatrix} \rightarrow \begin{bmatrix} -A_0 & A_* & A_\infty & 0 & 0 & 0 \\ 0 & 0 & 0 & P & 0 & 0 \\ 0 & 0 & 0 & Q & A_* & A_\infty \end{bmatrix} \in \mathcal{L}$$

and similarly for (6);

$$\begin{bmatrix} -A_0 & A_* & A_\infty & 0 & 0 & 0 \\ B_0 & 0 & -B_\infty & P & B_* & B_\infty \\ A_0 & 0 & -A_\infty & 0 & A_* & A_\infty \end{bmatrix} \rightarrow \begin{bmatrix} -A_0 & A_* & A_\infty & 0 & 0 & 0 \\ B_0 & -B_* & -B_\infty & P & B_* & B_\infty \\ 0 & 0 & 0 & 0 & A_* & A_\infty \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} -A_0 & A_* & A_\infty & 0 & A_* & A_\infty \\ B_0 & -B_* & -B_\infty & P & 0 & 0 \\ 0 & 0 & 0 & 0 & A_* & A_\infty \end{bmatrix} \rightarrow \begin{bmatrix} -A_0 & A_* & A_\infty & 0 & 0 & 0 \\ B_0 & -B_* & -B_\infty & P & 0 & 0 \\ 0 & 0 & 0 & 0 & A_* & A_\infty \end{bmatrix} \in \mathcal{L}$$

These identities can in turn be used to show that the distributive identities hold in  $M_\Sigma/\mathcal{U}$ .

$$A.C \square B.C \sim (A \square B).C \quad \dots\dots\dots(7)$$

$$C.A \square C.B \sim C.(A \square B) \quad \dots\dots\dots(8)$$

[We note for future use that we actually show here that  $A.C \square B.C \square \{(A \square B).C\} \in \mathcal{L}$  and similarly for (8).] The l.h.s. of (7) is, (with a dot representing a zero);

$$\begin{bmatrix} C_0 & C_* & C_\infty & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & -B_0 & B_* & B_\infty & \cdot & \cdot & \cdot \\ C_0 & \cdot & \cdot & \cdot & \cdot & C_* & C_\infty & \cdot \\ \cdot & \cdot & \cdot & \cdot & -A_\infty & \cdot & -A_0 & A_* & A_\infty \end{bmatrix} \rightarrow \begin{bmatrix} C_0 & C_* & C_\infty & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & -B_0 & B_* & B_\infty & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & C_* & C_\infty & \cdot \\ \cdot & \cdot & -A_0 & \cdot & -A_\infty & \cdot & -A_0 & A_* & A_\infty \end{bmatrix}$$

which by (5),

$$\sim \begin{bmatrix} C_0 & C_* & C_\infty & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & -B_0 & B_* & B_\infty & \cdot & \cdot \\ \cdot & \cdot & -A_0 & \cdot & -A_\infty & A_* & A_\infty \end{bmatrix} = (A \square B).C$$

And again the l.h.s. of (8) is,

$$\begin{bmatrix} B_0 & B_* & B_\infty & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & -C_0 & C_* & C_\infty & \cdot & \cdot & \cdot & \cdot \\ A_0 & \cdot & \cdot & \cdot & \cdot & A_* & A_\infty & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & -C_\infty & \cdot & -C_0 & C_* & C_\infty \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} B_0 & B_* & B_\infty & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & -C_0 & C_* & C_\infty & \cdot & \cdot & \cdot & \cdot \\ A_0 & \cdot & -A_\infty & \cdot & \cdot & A_* & A_\infty & \cdot & \cdot \\ \cdot & \cdot & C_0 & -C_* & -C_\infty & \cdot & -C_0 & C_* & C_\infty \end{bmatrix}$$

After adding the second row block to the fourth row block we can apply (6) to get;

$$\sim \begin{bmatrix} B_0 & B_* & B_\infty & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ A_0 & \cdot & -A_\infty & A_* & A_\infty & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & -C_0 & C_* & C_\infty & \cdot & \cdot \end{bmatrix} = C. (A \square B)$$

We can also use (5) and (6) to establish the existence of a multiplicative identity element (1 1) since we have;

$$(A_0 A_* A_\infty). (1 \ 1) = \begin{bmatrix} 1 & 1 & \cdot & \cdot \\ \cdot & -A_0 & A_* & A_\infty \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & \cdot & \cdot \\ A_0 & \cdot & A_* & A_\infty \end{bmatrix} \sim (A_0 A_* A_\infty)$$

Also we note that similarly;

$$(1 \ 1). (A_0 A_* A_\infty) = \begin{bmatrix} A_0 & A_* & A_\infty & \cdot \\ \cdot & \cdot & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} A_0 & A_* & \cdot & A_\infty \\ \cdot & \cdot & -1 & 1 \end{bmatrix} \sim (A_0 A_* A_\infty)$$

We can now show that multiplication is well defined. Suppose that  $A \sim B$  i.e. that  $A \sqcup B^{\circ} \in \mathfrak{U}$ . Then we require that  $A.C \sim B.C$  and  $C.A \sim C.B$  for any  $C \in M_{\Sigma}$ . Now  $A.C \sim B.C$  if and only if  $A.C \sqcup (B.C)^{\circ} \sim 0$ . It is an easy check that  $(B.C)^{\circ} = B.C^{\circ} = B^{\circ}.C$  in  $M_{\Sigma}$ . Hence  $A.C \sqcup (B.C)^{\circ} \sim A.C \sqcup B^{\circ}.C$  and using the distributive identity (7) we get  $A.C \sqcup (B.C)^{\circ} \sim (A \sqcup B^{\circ}).C$  and  $A \sqcup B^{\circ} \sim 0$  so it suffices to show that if  $A \sim 0$  then  $A.C \sim 0$ . Similarly on the l.h.s. we require that  $C.A \sim 0$ . But this is precisely the condition that  $\mathfrak{U}$  is an ideal w.r.t. the multiplicative semigroup structure of  $M_{\Sigma}$ . Hence  $M_{\Sigma}/\mathfrak{U}$  has a ring structure.

Now define a map  $\lambda_{\mathfrak{U}}: R \rightarrow M_{\Sigma}/\mathfrak{U}$  by  $\lambda_{\mathfrak{U}}(r) = (r \ 1)$ . We observe immediately that  $\lambda_{\mathfrak{U}}$  preserves 1 and the following identities show  $\lambda_{\mathfrak{U}}$  to be a ring homomorphism.

$$(r \ 1).(s \ 1) = \begin{bmatrix} s & 1 & . \\ . & -r & 1 \end{bmatrix} \rightarrow \begin{bmatrix} s & 1 & . \\ rs & . & 1 \end{bmatrix} \sim (rs \ 1)$$

$$(r \ 1) \sqcup (s \ 1) = \begin{bmatrix} s & 1 & . \\ r & -1 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} s & 1 & . \\ r+s & . & 1 \end{bmatrix} \sim (r+s \ 1)$$

That  $M_{\Sigma}/\mathfrak{U}$  is trivial precisely when  $\mathfrak{U} = M_{\Sigma}$  follows from Lemma 2.2. ■

From now on we will denote  $M_{\Sigma}/\mathfrak{U}$  by  $R_{\mathfrak{U}}$ . In fact the above homomorphism is  $\Sigma$ -inverting but first we require a lemma.

**Lemma 2.5** *Under the conditions of Theorem 2.4 we have the following identities in  $R_{\mathfrak{U}}$ ;*

(i)  $(A \begin{smallmatrix} o \\ o \end{smallmatrix} A_* A_{\infty}) \sqcup (B \begin{smallmatrix} o \\ o \end{smallmatrix} A_* A_{\infty}) = (A + B \begin{smallmatrix} o \\ o \end{smallmatrix} A_* A_{\infty})$

(ii) If  $p \in R_{\mathfrak{U}}$  has as an admissible matrix  $(A \begin{smallmatrix} o \\ o \end{smallmatrix} A_* A_{\infty})$  then  $p=1$  in  $R_{\mathfrak{U}}$ .

(iii)  $(A \begin{smallmatrix} o \\ o \end{smallmatrix} A_* A_{\infty}) = (A + A_* q \begin{smallmatrix} o \\ o \end{smallmatrix} A_* A_{\infty})$  when  $q$  is a column of the appropriate size over the characteristic ring of  $R$ .

Proof. (i) Writing out the l.h.s. we have

$$\begin{bmatrix} B_{\circ} & A_{*} & A_{\infty} & 0 & 0 \\ A_{\circ} & 0 & -A_{\infty} & A_{*} & A_{\infty} \end{bmatrix} \rightarrow \begin{bmatrix} B_{\circ} & A_{*} & A_{\infty} & 0 & 0 \\ A_{\circ}+B_{\circ} & 0 & 0 & A_{*} & A_{\infty} \end{bmatrix}$$

which by (6) is equivalent to  $(A_{\circ}+B_{\circ} \ A_{*} \ A_{\infty})$ .

$$(ii) \quad (A_{\infty} A_{*} A_{\infty}) \square (1 \ 1)^{\circ} = \begin{bmatrix} -1 & 1 & 0 & 0 \\ A_{\infty} & -A_{\infty} & A_{*} & A_{\infty} \end{bmatrix} \rightarrow \begin{bmatrix} -1 & 1 & 0 & 0 \\ 0 & 0 & A_{*} & A_{\infty} \end{bmatrix}$$

The final matrix on the r.h.s. lies in  $\mathcal{L}$  and hence is equivalent to zero.

$$(iii) \quad (A_{\circ} A_{*} A_{\infty}) \sim \begin{bmatrix} 1 & 1 & 0 & 0 \\ A_{\circ} & 0 & A_{*} & A_{\infty} \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 1 & 0 & 0 \\ A_{\circ}+A_{*}q & A_{*}q & A_{*} & A_{\infty} \end{bmatrix}$$

(The second operation here is achieved by row operations.)

$$\rightarrow \begin{bmatrix} 1 & 1 & 0 & 0 \\ A_{\circ}+A_{*}q & 0 & A_{*} & A_{\infty} \end{bmatrix} \sim (A_{\circ}+A_{*}q \ A_{*} \ A_{\infty})$$

(The first operation is permitted by column operations within the core.) ■

[ n.b. when two matrices differ in one column only e.g.  $A=(A_{\circ} A_{*})$ ,  $B=(B_{\circ} A_{*})$  then the **determinantal sum** of the two matrices is defined w.r.t. that column and is written  $A \nabla B=(A_{\circ}+B_{\circ} \ A_{*})$ . This is developed later in Chapter 5. In (i) above we have shown that in  $R_{\mathbb{U}}$  the determinantal sum w.r.t. the initial column is admissible for the sum of the two elements represented by the original two matrices.]

Theorem 2.6     *Under the same hypotheses as Theorem 2.4 the homomorphism  $\lambda_{\mathbb{U}}:R \rightarrow R_{\mathbb{U}}$  is epic  $\Sigma$ -inverting.*

Proof. Let  $A=(A_*A_\infty)\in\Sigma_n$ . We shall denote the  $(i,j)$ th element of  $A$  by  $a_{ij}$  and the  $k$ th column of  $A$  by  $A_k$ . Also if  $i<n$  we will write  $A_\infty^{(1)}$  for the square matrix whose columns are all zero except the  $i$ th which is  $-A_\infty$  and the  $n$ th which is  $A_\infty$ ; for  $i=n$  define  $A_\infty^{(1)}=0$ .

Let  $B$  be the square matrix over  $R_U$  whose  $(i,j)$ th element,  $b_{ij}$  has as an admissible matrix

$$\begin{bmatrix} e_j & A & 0 \\ e_j & A_\infty^{(1)} & A \end{bmatrix} .$$

Then we can show that  $B$  is a left inverse for  $\lambda_U(A)$  over  $R_U$ . If  $\Sigma$  admitted column permutations then we could (as in Cohn [1] for his skew field construction) define a simpler admissible matrix for  $b_{ij}$  but we shall use the above. Then we have

$$(B\lambda_U(A))_{ik} = \sum_j b_{ij}\lambda_U(A)_{jk} ,$$

where the r.h.s. has an admissible matrix;

$$\boxed{\Sigma_j} \begin{bmatrix} a_{jk} & 1 & . & . \\ . & -e_j & A & . \\ . & -e_j & A_\infty^{(1)} & A \end{bmatrix}$$

Considering the  $j$ th individual summand we see that after adding the top row to the  $j+1$  th and  $n+j+1$  th row we have;

$$\begin{bmatrix} a_{jk} & 1 & . & . \\ a_{jk}e_j & . & A & . \\ a_{jk}e_j & . & A_\infty^{(1)} & A \end{bmatrix}$$

which simplifies by (6) hence,

$$(B\lambda_{\mathbb{U}}(A))_{ik} = \boxed{\sum_j} \begin{bmatrix} a_{jk}e_j & A & \cdot \\ a_{jk}e_j & A_{\infty}^{(1)} & A \end{bmatrix} .$$

We notice that each summand is only dependent on  $j$  in the first column so we can apply Lemma 2.5(i) inductively to obtain;

$$(B\lambda_{\mathbb{U}}(A))_{ik} = \begin{bmatrix} A_k & A & \cdot \\ A_k & A_{\infty}^{(1)} & A \end{bmatrix}$$

If  $i \neq k$  then by simply subtracting the relevant two columns ( $k \neq n$ ) or relevant column ( $k = n$ ) from the first column we get a zero initial column hence;

$$(B\lambda_{\mathbb{U}}(A))_{ik} = 0 \quad \text{for } i \neq k.$$

For  $i = k$  there are two cases. Firstly if  $i = k = n$  then  $A_{\infty}^{(1)} = 0$  and subtracting the  $(n+1)$ th column from the first we see that the initial and final columns agree since  $A_n = A_{\infty}$ . By Lemma 2.5 (ii) we have an admissible matrix for 1. Secondly if  $i = k \neq n$  we have;

$$\begin{bmatrix} A_i & A_1 & A_2 & \dots & A_i & \dots & A_{\infty} & 0 \\ A_i & 0 & 0 & \dots & -A_{\infty} & 0 & \dots & A_{\infty} & A \end{bmatrix} \rightarrow \begin{bmatrix} 0 & A & 0 \\ A_i + A_{\infty} & A_{\infty}^{(1)} & A \end{bmatrix}$$

and subtracting the  $(n+i+1)$ th column from the first we can again apply Lemma 2.5 (ii) to obtain a matrix that is admissible for 1. Hence we have proved that  $(B\lambda_{\mathbb{U}}(A))_{ik} = \delta_{ik}$  and  $B$  is a left inverse for the image of  $A$  under  $\lambda_{\mathbb{U}}$ .

If  $v_n \in R_{\mathbb{U}}$  has as admissible matrix  $(A_{\infty} A_* A_{\infty})$  where  $(A_* A_{\infty}) \in \Sigma_n$  then we can show that there is an equation in  $R_{\mathbb{U}}$ ;

$$\lambda_{\mathbb{1}}(A_{\infty} * A_{\infty})(v_1 \ v_2 \ \dots \ v_n)^T - \lambda_{\mathbb{1}}(A_0) = 0 \quad (9)$$

where each  $v_i$  has as an admissible matrix

$$\begin{bmatrix} A_0 & A & 0 \\ A_0 & A_{\infty}^{(i)} & A \end{bmatrix}$$

To prove this we shall take  $n=2$  for simplicity. The proof for larger  $n$  is entirely analogous. Then the  $i$ th row of the l.h.s. of (9) has as an admissible matrix;

$$(-a_{i0} \ 1) \square (a_{i1} \ 1) \cdot \begin{bmatrix} A_0 & A & 0 \\ A_0 & A_{\infty}^{(1)} & A \end{bmatrix} \square (a_{i2} \ 1) \cdot \begin{bmatrix} A_0 & A & 0 \\ A_0 & A_{\infty}^{(2)} & A \end{bmatrix}$$

However since  $n=2$   $A_{\infty}^{(2)}=0$  and  $\begin{bmatrix} A_0 & A & 0 \\ A_0 & 0 & A \end{bmatrix} \sim (A_0 A)$  by (6).

So we have,

$$(-a_{i0} \ 1) \square \begin{bmatrix} A_0 & A_1 & A_2 & 0 & 0 & 0 \\ A_0 & -A_2 & A_2 & A_1 & A_2 & 0 \\ 0 & 0 & 0 & 0 & -a_{i1} & 1 \end{bmatrix} \square \begin{bmatrix} A_0 & A_1 & A_2 & 0 \\ 0 & 0 & -a_{i2} & 1 \end{bmatrix} =$$

$$\begin{bmatrix} A_0 & A_1 & A_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -a_{i2} & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ A_0 & 0 & 0 & 0 & A_1 & A_2 & 0 & 0 & 0 & 0 \\ A_0 & 0 & 0 & 0 & -A_2 & A_2 & A_1 & A_2 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 & 0 & 0 & -a_{i1} & 1 & 0 \\ -a_{i0} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 \end{bmatrix} \cdot$$

After adding the seventh and the eight columns to the fifth and then subtracting the third row block from the fourth we get;

$$\begin{bmatrix} A_0 & A_1 & A_2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -a_{i2} & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ A_0 & 0 & 0 & 0 & A_1 & A_2 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & A_1 & A_2 & 0 & 0 \\ 0 & 0 & 0 & -1 & -a_{i1} & 0 & 0 & -a_{i1} & 1 & 0 \\ -a_{i0} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$

which by (5) is equivalent to,

$$\begin{bmatrix} A_0 & A_1 & A_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -a_{i2} & 1 & 0 & 0 & 0 & 0 \\ A_0 & 0 & 0 & 0 & A_1 & A_2 & 0 & 0 \\ 0 & 0 & 0 & -1 & -a_{i1} & 0 & 1 & 0 \\ -a_{i0} & 0 & 0 & 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$

Similarly by repeating this process we get

$$\begin{bmatrix} A_0 & A_1 & A_2 & 0 & 0 & 0 \\ 0 & 0 & -a_{i2} & 1 & 0 & 0 \\ 0 & -a_{i1} & 0 & -1 & 1 & 0 \\ -a_{i0} & 0 & 0 & 0 & -1 & 1 \end{bmatrix}$$

After several row operations we can reduce this to

$$\begin{bmatrix} A_0 & A_1 & A_2 & 0 & 0 & 0 \\ 0 & 0 & -a_{i2} & 1 & 0 & 0 \\ 0 & -a_{i1} & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

which we see lies in  $\mathcal{L}$ . Hence we have shown that (9) holds and taking  $A_o = e_j$  ( $j=1$  to  $n$ ) we have that the image of  $A$  has a right inverse over  $R_{\mathcal{U}}$ . Hence by Theorem 1.2 every element of  $R_{\mathcal{U}}$  is an entry in the inverse of the image of an element of  $\Sigma$  and  $\lambda_{\mathcal{U}}:R \rightarrow R_{\mathcal{U}}$  is epic  $\Sigma$ -inverting. ■

Corollary If  $R$  is any ring,  $\Sigma$  a lower multiplicative set of matrices over  $R$  and  $\mathcal{U}$  a  $\Sigma$ -matrix ideal then  $\Sigma\text{-Ker}\lambda_{\mathcal{U}} = \mathcal{U}$ .

Proof. If we have an equation (9) in  $R_{\mathcal{U}}$  then  $v_n$  is unique since  $(A_*A_{\infty})\lambda_{\mathcal{U}}$  is invertible. Hence if  $v_n$  is zero then  $(A_o A_* A_{\infty}) \in \mathcal{U}$  or else it is admissible for a non-zero element  $v_n$  and we have the same equation by what precedes (9), which is a contradiction. ■

We have the following result which shows that a general  $\Sigma$ -inverting homomorphism will factor through certain homomorphisms  $\lambda_{\mathcal{U}}:R \rightarrow R_{\mathcal{U}}$ .

Lemma 2.7 If  $R$  is a ring,  $\Sigma$  a set of lower multiplicative matrices over  $R$ ,  $f:R \rightarrow S$  a  $\Sigma$ -inverting homomorphism and  $\mathcal{U}$  a  $\Sigma$  matrix ideal such that  $\mathcal{U} \subseteq \Sigma\text{-Ker}f$  then  $f$  factors uniquely through  $\lambda_{\mathcal{U}}:R \rightarrow R_{\mathcal{U}}$ .

Proof. Define a map  $f':R_{\mathcal{U}} \rightarrow S$  by  $(A_o A_* A_{\infty}) \mapsto e_{\infty}^T (A_* f A_{\infty} f)^{-1} A_o f$ . We know that this map is linear and multiplicative by construction (e.g. Lemma 1.1, 2.3). Hence to show  $f'$  is well defined it suffices to show that if  $(A_o A_* A_{\infty}) \sim 0$  in  $R_{\mathcal{U}}$  then  $(A_o A_* A_{\infty})f' = 0$ , but this is precisely the property that  $\mathcal{U}$  has since  $\mathcal{U}$  is admissible for zero under  $f$ . If  $r \in R$  then  $r\lambda_{\mathcal{U}}f' = (r \ 1)f' = 1.rf = rf$ . Hence  $f$  factors through  $\lambda_{\mathcal{U}}$ . Suppose  $f$  also factors through  $\lambda_{\mathcal{U}}$  via  $f''$ . If  $v_n \in R_{\mathcal{U}}$  with admissible matrix  $(A_o A_* A_{\infty})$  then we have an equation (9) as in Theorem 2.6 taking the image of this equation under  $f'$  and  $f''$  we

see that  $v_n f' = v_n f''$  since  $f = \lambda_u f' = \lambda_u f''$  is  $\Sigma$ -inverting. ■

There is an alternative way of doing this whole construction using rows as opposed to columns. The set of matrices  $\Sigma$  will then have to be upper multiplicative and our set of admissible matrices will be given by  $A = \begin{bmatrix} a_o \\ a_* \\ a_\infty \end{bmatrix}$  where  $a_o$  is a row and the remaining matrix  $A^\infty$  lies in  $\Sigma$  with final row  $a_\infty$ .

The two binary operations are then;

$$A \square B = \begin{bmatrix} b_o & a_o \\ b_* & . \\ b_\infty & -a_\infty \\ . & a_* \\ . & a_\infty \end{bmatrix} \quad A.B = \begin{bmatrix} a_o & . \\ a_* & . \\ a_\infty & -b_o \\ . & b_* \\ . & b_\infty \end{bmatrix}$$

The minimal set of admissible matrices for zero  $\mathcal{L}$  consists of all matrices which after a series of permissible operations (these differ also) can be written in the form below;

$$\begin{bmatrix} k_o & . \\ K & . \\ . & L \end{bmatrix}, \text{ where } K, L \in \Sigma \text{ and } k_o \text{ is a row.}$$

This we shall call the dual construction. We do not go into detail here but much is analogous to the proofs used above. In the next chapter we shall quote without proof the dual of our results on universal localisation.

We start by showing that the universal  $\Sigma$ -inverting homomorphism is the  $R$ -ring constructed from the least  $\Sigma$ -matrix ideal  $\mathfrak{U}_\Sigma$ . We then proceed to describe  $\mathfrak{U}_\Sigma$  in terms of  $\mathcal{L}$  the minimal set of matrices admissible for zero and once this is achieved we have a description of the kernel. Having determined  $\ker \lambda_\Sigma$  we give a necessary and sufficient condition for the universal  $\Sigma$ -inverting homomorphism to be an embedding.

**Theorem 3.1** *If  $R$  is any ring,  $\Sigma$  a lower multiplicative set of matrices over  $R$  and  $\mathfrak{U}_\Sigma$  the least  $\Sigma$ -matrix ideal then the universal  $\Sigma$ -inverting homomorphism is given by  $\lambda_{\mathfrak{U}_\Sigma}: R \rightarrow R_{\mathfrak{U}_\Sigma}$ .*

**Proof.** Let  $\sigma$  be the collection of all  $\Sigma$ -inverting homomorphisms then  $\mathfrak{U} = \bigcap_{\{f \in \sigma\}} \Sigma\text{-Ker} f$  is a  $\Sigma$ -matrix ideal which is admissible for zero under all  $\Sigma$ -inverting homomorphisms. Hence by Lemma 2.7 every  $f \in \sigma$  factors uniquely through  $\lambda_{\mathfrak{U}}: R \rightarrow R_{\mathfrak{U}}$  which is itself a  $\Sigma$ -inverting homomorphism by Theorem 2.6 i.e. this is the universal  $\Sigma$ -inverting homomorphism. Since  $\mathfrak{U}_\Sigma$  is the least  $\Sigma$ -matrix ideal we have  $\mathfrak{U} \supseteq \mathfrak{U}_\Sigma$  but since  $\Sigma\text{-Ker} \lambda_{\mathfrak{U}_\Sigma} = \mathfrak{U}_\Sigma$  by the corollary to Theorem 2.6 we have  $\mathfrak{U}_\Sigma \supseteq \mathfrak{U}$  and we have proved the result. ■

Now we proceed to describe  $\mathfrak{U}_\Sigma$ .

**Lemma 3.2** *Let  $R$  be a ring,  $\Sigma$  a lower multiplicative set of matrices and  $\mathcal{L}$  the minimal set of admissible matrices for zero. Then  $\mathcal{L}$  is an additive subsemigroup of  $M_\Sigma$  and a multiplicative ideal of  $M_\Sigma$ .*

Proof If  $L_1$  and  $L_2$  lie in  $\mathcal{L}$  then by definition we have;

$$L_1 \rightarrow \begin{bmatrix} A_0 & A & \cdot & \cdot \\ \cdot & \cdot & B_* & B_\infty \end{bmatrix} \quad L_2 \rightarrow \begin{bmatrix} C_0 & C & \cdot & \cdot \\ \cdot & \cdot & D_* & D_\infty \end{bmatrix}$$

where  $A$ ,  $(B_* B_\infty)$ ,  $C$  and  $(D_* D_\infty)$  lie in  $\Sigma$ ;  $A_0$  and  $C_0$  are columns over  $R$ . Then we have;

$$\begin{aligned} L_1 \square L_2 &\rightarrow \begin{bmatrix} C_0 & C & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & D_* & D_\infty & \cdot & \cdot & \cdot \\ A_0 & \cdot & \cdot & \cdot & A & \cdot & \cdot \\ \cdot & \cdot & \cdot & -B_\infty & \cdot & B_* & B_\infty \end{bmatrix} \\ &\rightarrow \begin{bmatrix} C_0 & C & \cdot & \cdot & \cdot & \cdot & \cdot \\ A_0 & \cdot & \cdot & \cdot & A & \cdot & \cdot \\ \cdot & \cdot & D_* & D_\infty & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & -B_\infty & \cdot & B_* & B_\infty \end{bmatrix} \\ &\rightarrow \begin{bmatrix} C_0 & C & \cdot & \cdot & \cdot & \cdot & \cdot \\ A_0 & \cdot & A & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & D_* & D_\infty & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & -B_\infty & B_* & B_\infty \end{bmatrix} \in \mathcal{L} \end{aligned}$$

Hence  $L_1 \square L_2 \in \mathcal{L}$  so  $\mathcal{L}$  is an additive subsemigroup of  $M_\Sigma$ .

If now  $(E_0 E_* E_\infty) = E$  is any admissible matrix then we have the following;

$$L_1 \cdot E \rightarrow \begin{bmatrix} E_0 & E_* & E_\infty & \cdot & \cdot & \cdot \\ \cdot & \cdot & -A_0 & A & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & B_* & B_\infty \end{bmatrix} \in \mathcal{L}$$

and also;

$$E.L_1 \rightarrow \begin{bmatrix} A_0 & A & . & . & . & . \\ . & . & B_* & B_\infty & . & . \\ . & . & . & -E_0 & E_* & E_\infty \end{bmatrix} \in \mathcal{L}$$

So  $\mathcal{L}$  is also a multiplicative ideal of  $M_\Sigma$ . ■

Now we require a lemma on commutative semigroups.

Lemma 3.3 *If  $H$  is a subsemigroup of a commutative semigroup  $S$  then  $H:H = \{a \in S \mid \exists b \in H; a+b \in H\}$  is the least unitary subsemigroup of  $S$  containing  $H$ .*

Proof Firstly  $H:H$  is a subsemigroup of  $S$ . For if  $a, c \in H:H$  then there are  $b, d \in H$  such that  $a+b, c+d \in H$ . Then  $(a+b)+(c+d) = (a+c)+(b+d) \in H$  and  $b+d \in H$ , hence  $a+c \in H:H$ .

$H:H$  is unitary since if  $a+b, b \in H:H$  then there exist  $c, d \in H$  with  $(a+b)+c, b+d \in H$ . So  $b+c+d \in H$  and  $a+b+c+d \in H$  hence  $a \in H:H$ .

If  $K$  is a unitary subsemigroup of  $S$  containing  $H$  then  $a \in H:H$  implies that there exists  $b \in H$  with  $a+b \in H$ . So  $b, a+b \in K$  and since  $K$  is unitary  $a \in K$ . Hence  $K \supseteq H:H$ . ■

Applying this to  $\mathcal{U}$ , an additive subsemigroup of  $M_\Sigma$  we see that  $\mathcal{U}:\mathcal{U}$  is the least unitary subsemigroup containing  $\mathcal{U}$ .

Lemma 3.4 *If  $R$  is any ring,  $\Sigma$  a lower multiplicative set of matrices and  $\mathcal{U} \supseteq \mathcal{L}$  is an additive subsemigroup and a multiplicative ideal of  $M_\Sigma$  then  $\mathcal{U}:\mathcal{U}$  is the least  $\Sigma$ -matrix ideal containing  $\mathcal{U}$ .*

Proof From the above remark the result will follow if we can show that  $\mathcal{U}:\mathcal{U}$  is a multiplicative ideal of  $M_\Sigma$ .

If  $A \in \mathcal{U}:\mathcal{U}$  then there exists  $B \in \mathcal{U}$  such that  $A \square B \in \mathcal{U}$ . Now let  $C$  be any admissible matrix in  $M_\Sigma$  then from the remark succeeding (7) and (8) in Theorem 2.4 we have;

$$\{C.(A \square B)\}^\circ \square C.A \square C.B \in \mathcal{L} \subseteq \mathcal{U}:\mathcal{U}$$

Since  $B \in \mathcal{U}$  and  $\mathcal{U}$  is an ideal we have that  $C.B \in \mathcal{U}$ . So since  $\mathcal{U}:\mathcal{U}$  is unitary the following holds;

$$\{C.(A \square B)\}^\circ \square C.A \in \mathcal{U}:\mathcal{U}$$

Now  $C^\circ.(A \square B) \in \mathcal{U}$  by the same argument that holds for  $C.B$ . And in  $M_\Sigma$  we have  $C^\circ.X = (C.X)^\circ$  so again since  $\mathcal{U}:\mathcal{U}$  is unitary we have that  $C.A \in \mathcal{U}:\mathcal{U}$ .

The argument showing  $A.C \in \mathcal{U}:\mathcal{U}$  follows in exactly the same manner using the distributive law on the other side. ■

By applying this to  $\mathcal{L}$  we have;

Theorem 3.5 *If  $R$  is any ring and  $\Sigma$  a lower multiplicative set of matrices over  $R$  then the least  $\Sigma$ -matrix ideal  $\mathcal{U}_\Sigma$  is  $\mathcal{L}:\mathcal{L}$  where  $\mathcal{L}$  is the minimal set of matrices admissible for zero. ■*

Corollary *If  $\mathcal{M}$  is an additive unitary subset of  $M_\Sigma$  which contains  $\mathcal{L}$ , the minimal set of admissible matrices for zero, and every admissible matrix in  $\mathcal{M}$  necessarily represents zero under any  $\Sigma$ -inverting homomorphism then  $\mathcal{M}$  is the least  $\Sigma$ -matrix ideal.*

Proof  $\mathcal{M}$  contains  $\mathcal{L}$  so  $\mathcal{M}:\mathcal{M}$  contains  $\mathcal{L}:\mathcal{L}$  which is the least  $\Sigma$ -matrix ideal from the above. However  $\mathcal{M}$  is unitary so  $\mathcal{M}:\mathcal{M}$  is just  $\mathcal{M}$  and hence we have that  $\mathcal{M}$  contains  $\mathcal{U}_\Sigma$ . Conversely since all admissible matrices in  $\mathcal{M}$  represent zero under any  $\Sigma$ -inverting homomorphism  $\mathcal{M}$  must be contained in the least  $\Sigma$ -matrix ideal. ■

Now we have our description of the kernel of the universal  $\Sigma$ -inverting homomorphism.

**Theorem 3.6** *If  $R$  is any ring and  $\Sigma$  a lower multiplicative set of matrices over  $R$  then  $r \in R$  lies in the kernel of the universal  $\Sigma$ -inverting homomorphism if and only if there exists an equation:*

$$A_*Q = (A_r)_* \Sigma \quad (1)$$

Where  $(A_r)_*$  and  $\Sigma$  lie in  $\Sigma$ ,  $Q$  has entries in the characteristic ring, and  $*$  is some matrix of the appropriate size.

**Proof** An element  $r$  of  $R$  lies in the kernel if and only if  $(r \ 1)$  lies in  $\mathcal{U}_\Sigma$ , which we have shown is  $\mathcal{L}:\mathcal{L}$ . Then by the definition of  $\mathcal{L}:\mathcal{L}$  we must have an equation;  $\{(U_0 U) \oplus V\} \square (r \ 1) \in \mathcal{L}$  where  $U, V \in \Sigma$ . Hence we get a matrix equation as below.

$$\begin{bmatrix} r & 1 & . & . & . \\ U_0 & . & U & . & . \\ . & -V_\infty & . & V_* & V_\infty \end{bmatrix} \begin{bmatrix} 1 & . & . \\ . & Q_{11} & Q_{12} \\ . & Q_{21} & Q_{22} \\ . & Q_{31} & Q_{32} \\ . & . & e_\infty^T \end{bmatrix} = \begin{bmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \\ P_{31} & P_{32} \end{bmatrix} \begin{bmatrix} L_0 & L & . \\ . & . & M \end{bmatrix}$$

Where  $U_0, L_0$  are columns;  $U, (V_* V_\infty), L, M \in \Sigma$ . The half bordered matrix together with the other invertible matrix (with entries in the  $P_{ij}$ ) have been blocked appropriately. From this we can obtain directly the following matrix equation;

$$\begin{bmatrix} r & 1 & . \\ . & -V_\infty & V_* \end{bmatrix} \begin{bmatrix} 1 & . \\ . & Q_{11} \\ . & Q_{31} \end{bmatrix} = \begin{bmatrix} P_{11} \\ P_{31} \end{bmatrix} [L_0 \ L] \quad (2)$$

After eliminating  $V_\infty$  by row operations this gives us the equations,

$V_{\infty} r = P_{31}' L_0$  and  $V_* Q_{31} = P_{31}' L$ , where  $P_{31}'$  is the transformed  $P_{31}$  (after eliminating  $V_{\infty}$  by row operations). Rewriting these in the following form we have proved the necessity of the result.

$$V_*(0 \ Q_{31}) = (V_{\infty} r \ P_{31}') \begin{bmatrix} 1 & 0 \\ -L_0 & L \end{bmatrix}$$

It is a straightforward calculation to show that the condition is also sufficient and this is discussed below. ■

We also have the dual result:

**Theorem 3.6\*** *If  $R$  is any ring and  $\Sigma$  an upper multiplicative set of matrices over  $R$  then  $r \in R$  lies in the kernel of the universal  $\Sigma$ -inverting homomorphism if and only if there exists an equation:*

$$Ql_* = \Sigma \begin{bmatrix} r l_{\infty} \\ * \end{bmatrix} \quad (3)$$

where  $\Sigma$  and  $\begin{bmatrix} l_* \\ l_{\infty} \end{bmatrix} \in \Sigma$ ,  $Q$  has entries in the characteristic ring and  $*$  is some matrix of the appropriate size. ■

From this description of the kernel we can state a necessary and sufficient condition on a ring for the universal  $\Sigma$ -inverting homomorphism to be an embedding. If we have an equation (1) for  $(A_* A_{\infty}) \in \Sigma$  where  $r$  is non-zero we shall say that  $(A_* A_{\infty})$  is a **column  $\Sigma$ -zero divisor** (and for an equation (3) we say **row  $\Sigma$ -zero divisor**) since if  $\Sigma$  is right-inverted  $(A_* A_{\infty})$  becomes a left zero divisor via the first column in the expression below;

$$A_* Q \Sigma^{-1} = (A_{\infty} r \ *).$$

Hence we have the following.

Theorem 3.7 If  $R$  is any ring,  $\Sigma$  a lower multiplicative set of matrices over  $R$  then the universal  $\Sigma$ -inverting homomorphism is an embedding if and only if  $\Sigma$  contains no column  $\Sigma$ -zero divisors. ■

And the dual result:

Theorem 3.7\* If  $R$  is any ring,  $\Sigma$  an upper multiplicative set of matrices over  $R$  then the universal  $\Sigma$ -inverting homomorphism is an embedding if and only if  $\Sigma$  contains no row  $\Sigma$ -zero divisors. ■

There is a special case where this condition simplifies.

Lemma 3.8 If  $R$  is any ring,  $\Sigma$  a lower multiplicative set of matrices and if  $\mathcal{L}$  the minimal set of admissible matrices for zero is closed w.r.t. first column determinantal sums then the universal  $\Sigma$ -inverting homomorphism is an embedding if and only if  $\Sigma$  does not contain a left zero divisor.

Proof If  $\sum_1 u = 0$  in  $R$  where  $\Sigma_1 \in \Sigma$  and  $u$  is a non-zero column over  $R$  then in  $R_\Sigma$  we can cancel the image of  $\Sigma_1$  to show that the components of  $u$  must lie in the kernel. Conversely we have;

$$\begin{bmatrix} r & 1 & . & . & . \\ U_0 & . & U & . & . \\ . & -V_\infty & . & V_* & V_\infty \end{bmatrix} \nabla \begin{bmatrix} . & 1 & . & . & . \\ -U_0 & . & U & . & . \\ . & -V_\infty & . & V_* & V_\infty \end{bmatrix} = \begin{bmatrix} r & 1 & . & . & . \\ . & . & U & . & . \\ . & -V_\infty & . & V_* & V_\infty \end{bmatrix}$$

If  $r$  lies in the kernel then by Theorem 3.5 there exist  $U, (V_* V_\infty) \in \Sigma$  and  $U_0$  a column such that the first matrix on the l.h.s. lies in  $\mathcal{L}$ .

After permuting some rows and columns of the second matrix on the l.h.s. we see that it lies in  $\mathcal{L}$ . Hence by hypothesis the r.h.s. must lie in  $\mathcal{L}$  also. After adding left multiples of the top row to

rows within the bottom row block and then a column operation we see that there exist invertible matrices P and Q where Q is half bordered such that there is a factorisation,

$$P \begin{bmatrix} r & 1 & . & . & 1 \\ . & . & U & . & . \\ V_{\infty} r & . & . & V_* & V_{\infty} \end{bmatrix} = \begin{bmatrix} L_o & L & . & . \\ . & . & M_* & M_{\infty} \end{bmatrix} Q$$

Equating first and last columns on both sides and noting that Q is half bordered we see that,

$$P \begin{bmatrix} 1 \\ . \\ V_{\infty} \end{bmatrix} r = \begin{bmatrix} L_o \\ . \end{bmatrix} \quad \text{and} \quad P \begin{bmatrix} 1 \\ . \\ V_{\infty} \end{bmatrix} = \begin{bmatrix} * \\ M_* q + M_{\infty} \end{bmatrix}$$

where an asterisk denotes a non-specific column and q is also a column. Hence if r is a non-zero member of the kernel then we have a non-trivial column u such that  $(M_* M_{\infty})u=0$  as required. ■

Chapter 4      The Criteria of Malcolmsen and Gerasimov

The results of the previous chapter now enable us to prove the results of Malcolmsen and Gerasimov respectively on the kernel of the universal  $\Sigma$ -inverting homomorphism. Of course this can be achieved directly by using the equations in (2) of Theorem 3.6 in the previous chapter but we shall verify what amounts to their description of the least  $\Sigma$ -matrix ideal by using the corollary to Theorem 3.5.

Malcolmsen's construction, using a so called "zigzag" method, constructs  $R_\Sigma$  as a set of equivalence classes of triples  $(b,A,c)$  where  $b$  is a row,  $c$  a column and  $A$  lies in  $\Sigma$ . The equivalence class of  $(b,A,c)$  is to be interpreted as the element  $bA^{-1}c$  of  $R_\Sigma$ . In the present construction we are considering elements  $e_\omega^T A^{-1} A_0$ . Hence by substituting  $b=e_\omega^T$  and  $c=A_0$  we can determine Malcolmsen's description of the elements  $(e_\omega^T, A, A_0)$  which lie in the equivalence class of  $(1,1,0)$  i.e. those admissible matrices which represent zero and therefore lie in  $\mathbb{U}_\Sigma$ .

In Malcolmsen [4] we see that  $(e_\omega^T, A, A_0) \sim (1,1,0)$  if there exist  $L, M, P, Q \in \Sigma$ , rows  $j$  and  $u$  the sizes of  $L$  and  $P$ , respectively, and columns  $w$  and  $v$  the sizes of  $M$  and  $Q$ , respectively, such that

$$\begin{bmatrix} A & . & . & . & A_0 \\ . & 1 & . & . & . \\ . & . & L & . & . \\ . & . & . & M & w \\ e_\omega^T & 1 & j & . & . \end{bmatrix} = \begin{bmatrix} P \\ u \end{bmatrix} \begin{bmatrix} Q & v \end{bmatrix}$$

In fact we can simplify this slightly. We shall prove the following theorem which is essentially Malcolmsen's description of  $\mathbb{U}_\Sigma$  the least  $\Sigma$ -matrix ideal. Then we will gain a description of the kernel

of the universal  $\Sigma$ -inverting homomorphism which is Malcolmson's criterion without having used his "zigzag" construction. However our result will have to be slightly less general.

Theorem 4.1 *If  $R$  is a ring and  $\Sigma$  a lower multiplicative set of matrices closed w.r.t. multiplication on the left and right by elements of  $GL(R)$  then  $U_\Sigma$  the least  $\Sigma$ -matrix ideal consists precisely of those admissible matrices  $(A_{\infty} A_* A_0)$  for which there exist matrices  $L, M, P, Q \in \Sigma$ ; rows  $j$  and  $u$  the sizes of  $L$  and  $P$  respectively; and columns  $w$  and  $v$  the sizes of  $M$  and  $Q$  respectively, such that there exists a factorisation (1) below.*

$$\begin{bmatrix} A & \cdot & \cdot & A_0 \\ \cdot & L & \cdot & \cdot \\ \cdot & \cdot & M & w \\ e_\infty^T & j & \cdot & \cdot \end{bmatrix} = \begin{bmatrix} P \\ u \end{bmatrix} \begin{bmatrix} Q & v \end{bmatrix} \quad (1)$$

Where  $A = (A_* A_\infty)$ .

Proof Such a factorisation will be called an allowable factorisation for  $(A_\infty A_* A_0)$ .

Given the R.H.S. of (1) we note that we can perform certain elementary operations viz. if  $S, T \in GL(R)$  and  $t$  is any column of the appropriate size then,

$$\begin{bmatrix} S & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} P \\ u \end{bmatrix} \begin{bmatrix} Q & v \end{bmatrix} \begin{bmatrix} T & t \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} SP \\ u \end{bmatrix} \begin{bmatrix} QT & Qt+v \end{bmatrix}.$$

Consequently we can apply these restricted operations to both sides of (1) and maintain the form of the factorisation on the R.H.S.. Any application of these restricted operations will be denoted by

an arrow. We note that given a factorisation (1) for an admissible matrix A then by using these restricted operations we can obtain an allowable factorisation for any matrix obtainable from A by a series of permissible operations. Hence  $\mathcal{U}$  is a well defined subset of  $M_\Sigma$ . We can now proceed to show that  $\mathcal{U}$  is the least  $\Sigma$ -matrix ideal by using the corollary to Theorem 3.5. Observing the following factorisation;

$$\begin{bmatrix} K & . & . & K_0 \\ . & L & . & . \\ . & . & I & . \\ . & e_\infty^T & . & . \end{bmatrix} = \begin{bmatrix} I & . & . \\ . & L & . \\ . & . & I \\ . & e_\infty^T & . \end{bmatrix} \begin{bmatrix} K & . & . & K_0 \\ . & I & . & . \\ . & . & I & . \end{bmatrix}$$

where  $K, L \in \Sigma$  and  $K_0$  is any column we see that  $\mathcal{U}$  contains  $\mathcal{L}$ . Now suppose that  $(A_0 A_* A_\infty) \square (B_0 B_* B_\infty) \in \mathcal{U}$  with an allowable factorisation (1) and also  $(B_0 B_* B_\infty) \in \mathcal{U}$  with a similar factorisation in which P is replaced by  $P_1$  etc. then,

$$\begin{bmatrix} P & . \\ . & P_1 \\ u & -u_1 \end{bmatrix} \begin{bmatrix} Q & . & v \\ . & Q_1 & v_1 \end{bmatrix} = \begin{bmatrix} A_* & A_\infty & . & . & . & . & . & . & A_0 \\ . & -B_\infty & B_* & B_\infty & . & . & . & . & B_0 \\ . & . & . & . & L & . & . & . & . \\ . & . & . & . & . & M & . & . & w \\ . & . & . & . & . & . & B_* & B_\infty & . & B_0 \\ . & . & . & . & . & . & . & . & L_1 & . \\ . & . & . & . & . & . & . & . & . & M_1 & w_1 \\ . & . & . & 1 & j & . & . & -1 & -j_1 & . & . \end{bmatrix}$$

After adding the fourth column block to the second and to the eighth column blocks, and the third to the seventh we have;

$$\rightarrow \begin{bmatrix} A_* & A_\infty & . & . & . & . & . & . & . & . & A_0 \\ . & . & B_* & B_\infty & . & . & B_* & B_\infty & . & . & B_0 \\ . & . & . & . & L & . & . & . & . & . & . \\ . & . & . & . & . & M & . & . & . & . & w \\ . & . & . & . & . & . & B_* & B_\infty & . & . & B_0 \\ . & . & . & . & . & . & . & . & L_1 & . & . \\ . & . & . & . & . & . & . & . & . & M_1 & w_1 \\ . & 1 & . & 1 & j & . & . & . & -j_1 & . & . \end{bmatrix}$$

Then subtracting the fifth row block from the second and permuting allowable rows and columns we get an allowable factorisation for  $(A_0 A_* A_\infty)$  so that  $U$  is unitary.

It is clear that any admissible matrix with an allowable factorisation (1) represents zero under any  $\Sigma$ -inverting homomorphism for then we have;

$$0 = uv = uQ(PQ)^{-1}Pv = e_\infty^T A^{-1} A_0 + jL^{-1}0 + 0M^{-1}w = e_\infty^T A^{-1} A_0$$

Indeed this was Malcolmson's initial motivation for defining his equivalence. So consequently by the corollary to Theorem 3.5 we deduce that  $U$  is the least  $\Sigma$ -matrix ideal. ■

### Malcolmson's Description of the Kernel

Now  $r \in \ker(\lambda)$  if and only if  $-r \in \ker(\lambda)$  which is if and only if  $(-r \ 1) \in U_\Sigma$  and this is if and only if there is an allowable factorisation for  $(-r \ 1)$ . Suppose so then;

$$\begin{bmatrix} P \\ u \end{bmatrix} \begin{bmatrix} Q & v \end{bmatrix} = \begin{bmatrix} 1 & . & . & -r \\ . & L & . & . \\ . & . & M & w \\ 1 & j & . & . \end{bmatrix} \rightarrow \begin{bmatrix} 1 & . & . & . \\ . & L & . & . \\ . & . & M & w \\ 1 & j & . & r \end{bmatrix}$$

This is an instance of a factorisation (2) below.

$$\begin{bmatrix} P \\ u \end{bmatrix} \begin{bmatrix} Q & v \end{bmatrix} = \begin{bmatrix} L & . & . \\ . & M & w \\ j & . & r \end{bmatrix} \quad (2)$$

where  $P, Q, \dots$  etc. satisfy the conditions of (1). If we have such a factorisation then we have  $\text{reker}(\lambda_\Sigma)$  by the following allowable factorisation for  $(-r \ 1)$ .

$$\begin{bmatrix} 1 & . \\ . & P \\ 1 & u \end{bmatrix} \begin{bmatrix} 1 & . & -r \\ . & Q & v \end{bmatrix} = \begin{bmatrix} 1 & . & -r \\ . & PQ & . \\ 1 & uQ & . \end{bmatrix}$$

We have proved the following description of the kernel due to Malcolmson [4].

Theorem 4.2 *If  $R$  is any ring,  $\Sigma$  a set of lower multiplicative matrices closed w.r.t. multiplication on the left and right by elements of  $GL(R)$  then the kernel of the universal  $\Sigma$ -inverting homomorphism consists precisely of those elements  $r \in R$  for which there exist matrices  $L, M, P, Q \in \Sigma$ ; rows  $j$  and  $u$  the sizes of  $L$  and  $P$  respectively; and columns  $w$  and  $v$  the sizes of  $M$  and  $Q$  respectively, such that*

$$\begin{bmatrix} L & 0 & 0 \\ 0 & M & w \\ j & 0 & r \end{bmatrix} = \begin{bmatrix} P \\ u \end{bmatrix} \begin{bmatrix} Q & v \end{bmatrix} .$$

■

### Gerasimov's Criterion

Gerasimov constructed the universal  $\Sigma$ -inverting ring by considering not elements of  $R_\Sigma$  but all matrices over  $R_\Sigma$ . There is an account of the construction in Chapter 7 [1]. As before we will take the description of those admissible matrices over  $R$  which represent zero and prove that they form the least  $\Sigma$ -matrix ideal.

Gerasimov constructed all matrices over  $R_\Sigma$  by considering 4-block matrices over  $R$  of the form

$$A = \left[ \begin{array}{c|c} A' & \tilde{A} \\ \hline A^o & \backslash A \end{array} \right]$$

where  $A^o \in \Sigma$ . Then  $A$  was to be interpreted as the matrix over  $R_\Sigma$  given by  $\tilde{A} - A'(A^o)^{-1} \backslash A$ . Hence in this construction an admissible matrix  $(A_o A_* A_\infty)$  is represented by the 4-block matrix,

$$A = \left[ \begin{array}{c|c} \cdot & -1 & | & \cdot \\ \hline A_* & A_\infty & | & A_o \end{array} \right] \cdot$$

From the account of Gerasimov's construction in 7.11 [1] we see that this matrix represents zero in  $R_\Sigma$  if and only if there exists a factorisation

$$\left[ \begin{array}{cccc} \cdot & -1 & * & * \\ A_* & A_\infty & * & * \\ \cdot & \cdot & \Sigma & * \end{array} \right] \left[ \begin{array}{cc} * & * \\ * & * \\ * & * \\ \Sigma & * \end{array} \right] = \left[ \begin{array}{cc} \cdot & \cdot \\ \cdot & A_o \\ \cdot & \cdot \end{array} \right]$$

where  $\Sigma$  denotes any matrix lying in  $\Sigma$  and an asterisk denotes any matrix of the appropriate size. In fact we shall simplify this

slightly and prove the following result.

Theorem 4.3 If  $R$  is any ring and  $\Sigma$  a lower multiplicative set of matrices over  $R$  then the least  $\Sigma$ -matrix ideal consists precisely of those admissible matrices  $(A_o A_* A_\omega)$  for which there exists a factorisation (3) below.

$$\begin{bmatrix} A_* & * & * \\ 0 & \Sigma & * \end{bmatrix} \begin{bmatrix} * & * \\ * & * \\ \Sigma & * \end{bmatrix} = \begin{bmatrix} 0 & A_o \\ 0 & 0 \end{bmatrix} \quad (3)$$

Where an asterisk denotes any matrix over  $R$  of the appropriate size and  $\Sigma$  denotes any matrix lying in  $\Sigma$ .

Proof Let  $\mathcal{U}$  be all those admissible matrices for which there exists such a factorisation. Firstly we show that  $\mathcal{U}$  is indeed a well defined subset of  $M_\Sigma$ . If  $P$  is any matrix over  $R$  such that  $PA_*$  is defined then multiplying both sides of (3) on the left by  $P \circ I$  gives us,

$$\begin{bmatrix} PA_* & P* & P* \\ 0 & \Sigma & * \end{bmatrix} \begin{bmatrix} * & * \\ * & * \\ \Sigma & * \end{bmatrix} = \begin{bmatrix} 0 & PA_o \\ 0 & 0 \end{bmatrix}$$

If  $Q$  is an invertible matrix and  $A_*Q$  is defined then we note;

$$\begin{bmatrix} A_*Q & * & * \\ 0 & \Sigma & * \end{bmatrix} \begin{bmatrix} Q^{-1}* & Q^{-1}* \\ * & * \\ \Sigma & * \end{bmatrix} = \begin{bmatrix} 0 & A_o \\ 0 & 0 \end{bmatrix}$$

We have shown that if  $(A_o A_* A_\infty) \rightarrow (B_o B_* B_\infty)$  then  $(A_o A_* A_\infty) \in \mathcal{U}$  implies that  $(B_o B_* B_\infty) \in \mathcal{U}$ . Hence  $\mathcal{U}$  is a well defined subset of  $M_\Sigma$ . We can now show that  $\mathcal{U}$  is the least  $\Sigma$ -matrix ideal of  $R$  by using the corollary to Theorem 3.5.

Firstly  $\mathcal{U}$  contains  $\mathcal{L}$  since if  $K, L \in \Sigma$  and  $K_o$  is any column over  $R$  then we have a factorisation for  $(K_o K) \circ L_*$ ;

$$\begin{bmatrix} K & 0 & 0 & I \\ 0 & L_* & 0 & 0 \\ 0 & 0 & I & 0 \end{bmatrix} \begin{bmatrix} -I & 0 \\ 0 & 0 \\ 0 & 0 \\ K & K_o \end{bmatrix} = \begin{bmatrix} 0 & K_o \\ 0 & 0 \\ 0 & 0 \end{bmatrix} .$$

To show that  $\mathcal{U}$  is a unitary subset of the additive semigroup  $M_\Sigma$  we suppose that  $(A_o A_* A_\infty) \square (B_o B_* B_\infty)$  has an allowable factorisation below;

$$\begin{bmatrix} A_* & A_\infty & \cdot & U & V \\ \cdot & -B_\infty & B_* & \bar{U} & \bar{V} \\ \cdot & \cdot & \cdot & \Sigma_1 & W \end{bmatrix} \begin{bmatrix} M & m \\ T & t \\ Q & q \\ N & n \\ \Sigma_2 & s \end{bmatrix} = \begin{bmatrix} \cdot & A_o \\ \cdot & B_o \\ \cdot & \cdot \end{bmatrix}$$

;and also that  $(B_o B_* B_\infty)$  lies in  $\mathcal{U}$  with a factorisation given below;

$$\begin{bmatrix} B_* & U^+ & V^+ \\ \cdot & \Sigma_1^+ & W^+ \end{bmatrix} \begin{bmatrix} M^+ & m^+ \\ N^+ & n^+ \\ \Sigma^+ & s^+ \end{bmatrix} = \begin{bmatrix} \cdot & B_o \\ \cdot & \cdot \end{bmatrix}$$

Then we see from the following factorisation that  $(A_o A_* A_\infty) \in \mathcal{U}$ .

$$\begin{bmatrix} A_* & . & -A_\infty & U & . & V & . \\ . & \Sigma_1^+ & . & . & . & . & W^+ \\ . & . & \Sigma_1 & . & . & W & . \\ . & U^+ & \bar{U} & B_* & B_\infty & \bar{V} & V^+ \end{bmatrix} \begin{bmatrix} M & . & m \\ . & N^+ & -n^+ \\ N & . & n \\ Q & M^+ & q-m^+ \\ -T & . & -t \\ \Sigma_2 & . & s \\ . & \Sigma_2^+ & -s^+ \end{bmatrix} = \begin{bmatrix} . & . & A_o \\ . & . & . \\ . & . & . \end{bmatrix}$$

By the corollary to Theorem 3.5 it is now possible to show that  $\mathfrak{U}$  is the least  $\Sigma$ -matrix ideal by showing that every admissible matrix in  $\mathfrak{U}$  represents zero under any  $\Sigma$ -inverting homomorphism.

Suppose that we have a factorisation below for an admissible matrix  $(A_o A_* A_\infty) \in \mathfrak{U}$ .

$$\begin{bmatrix} A_* & U & V \\ . & \Sigma_1 & W \end{bmatrix} \begin{bmatrix} M & m \\ N & n \\ \Sigma_2 & s \end{bmatrix} = \begin{bmatrix} . & A_o \\ . & . \end{bmatrix} \quad (4)$$

Then upon expansion we get;

$$\Sigma_1 N + W \Sigma_2 = 0 \quad (5)$$

$$\Sigma_1 n + W s = 0 \quad (6)$$

$$A_* M + U N + V \Sigma_2 = 0 \quad (7)$$

$$A_* m + U n + V s = A_o \quad (8)$$

From (5) and (6) we get;

$$U N = -U \Sigma_1^{-1} W \Sigma_2 \quad \text{and} \quad U n = -U \Sigma_1^{-1} W s .$$

Substituting into (7) and (8) respectively we get;

$$A_* M + (V - U \Sigma_1^{-1} W) \Sigma_2 = 0$$

$$A_* m + (V - U \Sigma_1^{-1} W) s = A_o$$

Thus we see that  $A_* (m - M \Sigma_2^{-1} s) = A_o$ . After multiplying both sides on the left by  $(A_* A_\infty)^{-1}$  we conclude that the last component of

$(A_* A_\infty)^{-1} A_0$  is zero. Hence  $\mathfrak{U}$  is the least  $\Sigma$ -matrix ideal. ■

Now the following theorem is Gerasimov's description of the kernel of the universal  $\Sigma$ -inverting homomorphism.

**Theorem 4.4** *If  $R$  is any ring and  $\Sigma$  a lower multiplicative set of matrices over  $R$  then the kernel of the universal  $\Sigma$ -inverting homomorphism consists precisely of those elements  $r$  of  $R$  for which there exists a factorisation below.*

$$\begin{bmatrix} \cdot & r \\ \cdot & \cdot \end{bmatrix} = \begin{bmatrix} * & * \\ \Sigma & * \end{bmatrix} \begin{bmatrix} * & * \\ \Sigma & * \end{bmatrix}$$

Where  $\Sigma$  represents any matrix lying in  $\Sigma$  and  $*$  denotes any matrix over  $R$  of the appropriate size.

**Proof** For any  $r \in R$ ,  $r$  lies in the kernel if and only if  $(r \ 1) \in \mathfrak{U}$ . Then we have a factorisation (4) for  $(r \ 1)$ ;

$$\begin{bmatrix} u & v \\ \Sigma_1 & W \end{bmatrix} \begin{bmatrix} N & n \\ \Sigma_2 & s \end{bmatrix} = \begin{bmatrix} \cdot & r \\ \cdot & \cdot \end{bmatrix}$$

Since the core of  $(r \ 1)$  is null  $u$  and  $v$  are rows,  $M$  and  $m$  are null. ■

By comparison it is straightforward to show the equations in (2) Theorem 3.6 give the following:

**Theorem 4.5** *Under the same conditions as Theorem 4.4 then  $r \in \ker \lambda_\Sigma$  if and only if there exists a factorisation below.*

$$\begin{bmatrix} \cdot & r \\ \cdot & \cdot \end{bmatrix} = \begin{bmatrix} e_\infty^T & * \\ \Sigma & * \end{bmatrix} \begin{bmatrix} Q & 0 \\ \Sigma & * \end{bmatrix} \quad \Sigma \in \Sigma \text{ etc.} \quad \blacksquare$$

The concept of a prime matrix ideal  $\mathcal{P}$  was introduced by P.M.Cohn to construct a skew field using admissible matrices whose numerators lie in  $\mathcal{P}$ . This construction was the basis of the more general construction in Chapter 2. Here we define, slightly more simply than Cohn, a prime matrix ideal going on to determine conditions for a ring to have a prime matrix ideal. This criterion will be shown to refine Cohn's result for a ring  $R$  to have a homomorphism to a skew field.

Prime matrix ideals are collections of square matrices over a ring  $R$  with certain defined properties. We will show that given a prime matrix ideal  $\mathcal{P}$  the set of admissible matrices whose numerators lie in  $\mathcal{P}$  is a  $\Sigma$ -matrix ideal where  $\Sigma$  is the complement of  $\mathcal{P}$ . This can then be shown to lead directly to the construction of a skew field using Theorem 2.6.

Definitions. The inner rank of an  $r \times s$  matrix  $A$  over a ring  $R$  is defined as the least  $n$  such that  $A=PQ$  where  $P$  is  $r \times n$  and  $Q$  is  $n \times s$  over  $R$ . We shall write  $\rho(A)=n$ . A square matrix is called **full** if its inner rank equals its order and **non-full** otherwise.

Let  $R$  be any ring,  $f:R \rightarrow F$  a homomorphism from  $R$  to a field  $F$ ; then we can consider  $\mathcal{P}$ , the set of all square matrices over  $R$  which are mapped to singular matrices over  $F$ . This set is called the **singular kernel** of  $f$  and is denoted by  $\text{Ker}f$ . We make the following observations;

N1. All non-full matrices lie in  $\mathcal{P}$ . This is simply because  $f$  preserves the property of being non-full and since a field has UGN only full matrices can be invertible.

N2. If the determinantal sum of  $A, B \in \mathcal{P}$  is defined w.r.t. some column then  $A \vee B \in \mathcal{P}$ . This is since  $f(A \vee B) = f(A) \vee f(B)$  and the expression on

the right is the determinantal sum of two singular matrices over a field which must be singular.

N3. If  $A \in \mathcal{P}$  then for any square matrix  $B$  over  $R$ ,  $A \oplus B \in \mathcal{P}$ .

N4. If  $1 \oplus A \in \mathcal{P}$  then  $A \in \mathcal{P}$ .

N5.  $1 \notin \mathcal{P}$  and if  $A, B$  are square matrices with  $A \oplus B \in \mathcal{P}$  then at least one of  $A$  or  $B$  must lie in  $\mathcal{P}$ .

A square matrix is said to be **hollow** if it has an  $r \times s$  block of zeros (maybe after column permutations) and  $r+s > n$  where  $n$  is the order of the matrix. A square matrix is said to be **degenerate** if one column is a right multiple of another (distinct) column. Both hollow and degenerate matrices can easily be shown to be non-full. With the above definitions and observations in mind we now introduce the notions of matrix pre-ideal, matrix ideal and prime matrix ideal and aim to derive a condition for the existence of the latter.

If  $\mathcal{P}$  is a set of square matrices over a ring  $R$  then  $\mathcal{P}$  is said to be a **matrix pre-ideal** if the following conditions hold.

M1. All square matrices which are hollow or degenerate lie in  $\mathcal{P}$ .

(Such matrices will be said to have property M1.)

M2.  $\mathcal{P}$  is closed w.r.t. column determinantal sums (when defined).

M3. If  $A \in \mathcal{P}$  and  $B$  is any square matrix over  $R$  then  $A \oplus B \in \mathcal{P}$ .

A **matrix ideal** is a matrix pre-ideal which also satisfies,

M4. If  $1 \oplus A \in \mathcal{P}$  then  $A \in \mathcal{P}$ .

A matrix ideal is said to be **proper** if  $1 \notin \mathcal{P}$ . A **prime matrix ideal** is a proper matrix ideal which additionally satisfies,

M5. If  $A \oplus B \in \mathcal{P}$  where both A and B are square then at least one of A or B lies in  $\mathcal{P}$ .

It is clear from what has been said above that,

Proposition 5.1 *If  $f:R \rightarrow F$  is a homomorphism from a ring R to a field F then the singular kernel of f is a prime matrix ideal of R.* ■

The following hold for a matrix pre-ideal.

(a) If  $C=AVB$  and B has property M1 then  $C \in \mathcal{P}$  if and only if  $A \in \mathcal{P}$ .

For if  $A \in \mathcal{P}$  then by M2  $C \in \mathcal{P}$ . Conversely changing the sign of a column of B preserves property M1, so taking B to the other side of the equation the result follows.

(b) If  $A \in \mathcal{P}$  then the result of adding a right multiple of one column of A to another also lies in  $\mathcal{P}$ .

Since if  $A=(A_1, A_2, \dots, A_n)$  where  $A_i$  is the  $i$ th column of A then  $(A_1 + A_2^c, A_2, A_3, \dots, A_n) = A \nabla (A_2^c, A_2, A_3, \dots, A_n)$  and the second matrix on the r.h.s. is in  $\mathcal{P}$  by M1. So the result follows by M2.

(c) If A and B are square and C,D are matrices of the appropriate size then the following hold.

$$(i) \quad \begin{bmatrix} A & C \\ 0 & B \end{bmatrix} \in \mathcal{P} \stackrel{(*)}{\Leftrightarrow} \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} A & 0 \\ D & B \end{bmatrix} \in \mathcal{P} .$$

$$(ii) \quad \begin{bmatrix} 0 & B \\ A & C \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} 0 & B \\ A & 0 \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} D & B \\ A & 0 \end{bmatrix} \in \mathcal{P} .$$

We shall prove (i)(\*). The other results follow in an entirely analogous way.

If  $B = (B_1, B')$  where  $B_1$  is the first column of  $B$  and similarly  $C = (C_1, C')$  then

$$\begin{bmatrix} A & C \\ 0 & B \end{bmatrix} = \begin{bmatrix} A & C_1 & C' \\ 0 & 0 & B' \end{bmatrix} \quad \nabla \quad \begin{bmatrix} A & 0 & C' \\ 0 & B & B' \end{bmatrix}$$

The first matrix on the R.H.S. is hollow and hence has property M1.

Thus by (a) we have,

$$\begin{bmatrix} A & C \\ 0 & B \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} A & 0 & C' \\ 0 & B_1 & B' \end{bmatrix} \in \mathcal{P} .$$

In a similar way the columns of  $C'$  can be varied and the assertion proved.

(d) If  $A$  and  $B$  are square and of the same size (although not necessarily in  $\mathcal{P}$ ) then

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} 0 & -A \\ B & 0 \end{bmatrix} \in \mathcal{P} .$$

Since by column operations as in (b) we get

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} A & 0 \\ B & B \end{bmatrix} \in \mathcal{P} \Leftrightarrow$$

$$\begin{bmatrix} A & -A \\ B & 0 \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} 0 & -A \\ B & 0 \end{bmatrix} \in \mathcal{P} .$$

Now if further  $\mathcal{P}$  is a matrix ideal then we have

(e) If  $A$  and  $B$  are square and of the same size and if at least one of them lies in  $\mathcal{P}$  then  $AB$  lies in  $\mathcal{P}$ .

Firstly let  $A \in \mathcal{P}$  then by M3  $A \circ B \in \mathcal{P}$  and so by (c) we obtain,

$$\begin{bmatrix} A & -I \\ 0 & B \end{bmatrix} \in \mathcal{P}.$$

By successive column operations as in (b) we can get,

$$\begin{bmatrix} 0 & -I \\ BA & B \end{bmatrix} \in \mathcal{P}.$$

So by (c)  $\begin{bmatrix} 0 & -I \\ BA & 0 \end{bmatrix} \in \mathcal{P}$ , and by (d) this gives us  $\begin{bmatrix} I & 0 \\ 0 & BA \end{bmatrix} \in \mathcal{P}$ .

Now M4 implies that  $BA \in \mathcal{P}$ . Hence  $\mathcal{P}$  admits left multiplication by square matrices. In particular  $\mathcal{P}$  allows the corresponding row operations to (b). As a consequence

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} 0 & -A \\ B & 0 \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} B & 0 \\ 0 & -A \end{bmatrix} \in \mathcal{P} \Leftrightarrow \begin{bmatrix} B & 0 \\ 0 & A \end{bmatrix} \in \mathcal{P}.$$

By an identical argument to above we then get  $AB \in \mathcal{P}$ .

(f) If  $A$  belongs to a matrix ideal  $\mathcal{P}$  then the result of permuting the rows or columns of  $A$  in any way belongs to  $\mathcal{P}$ .

For we can achieve any permutation by multiplying by an appropriate permutation matrix on the left or right.

(g) A matrix ideal  $\mathcal{P}$  is proper if and only if  $\mathcal{P} \neq \mathfrak{M}(R)$ .

If  $\mathcal{P}$  is proper then  $1 \notin \mathcal{P}$  by definition. If  $\mathcal{P}$  is not proper then  $I \in \mathcal{P}$  and hence by (e)  $A = A \cdot I \in \mathcal{P}$  for any  $A \in \mathfrak{M}(R)$ .

We now proceed to establish a necessary and sufficient condition on a ring  $R$  for a proper matrix ideal to exist.

Let  $(\mathcal{P}_\lambda)$  be any family of matrix ideals; then it is clear that

$\mathcal{P} = \bigcap \mathcal{P}_\lambda$  is again a matrix ideal. We can therefore speak of the 'least' matrix ideal containing a given subset  $\mathcal{X}$  of  $\mathfrak{M}(R)$ . This least matrix ideal is also called the matrix ideal generated by  $\mathcal{X}$ . Similarly we can define the matrix pre-ideal generated by  $\mathcal{X}$ .

Let  $\mathfrak{L}$  be the matrix pre-ideal generated by the empty set. Clearly this is the least matrix pre-ideal and it consists precisely of all column determinantal sums of square matrices which are hollow or degenerate.

We note that  $\mathfrak{L}$  has an additional property apart from M1-M3. If  $B \in \mathfrak{M}(R)$  and  $A \in \mathfrak{L}$  then  $B \circ A \in \mathfrak{L}$ . This is similar to but not equivalent to M3 (although it is in the presence of M1, M2 and M4). In particular we shall use  $I \circ A \in \mathfrak{L}$  for all  $A \in \mathfrak{L}$ , where  $I$  is any unit matrix. We shall denote by  $\ell$  the set of all unit matrices.

Now define  $\mathfrak{L}/\ell = \{A \in \mathfrak{M}(R) \mid I \circ A \in \mathfrak{L}\}$ . From the above remark we see that  $\mathfrak{L}/\ell \supseteq \mathfrak{L}$  and hence  $\mathfrak{L}/\ell$  satisfies M1. If  $A_i \in \mathfrak{L}/\ell$  ( $i=1,2$ ), then  $I_{n_1} \circ A_1 \in \mathfrak{L}$ . By the above remark  $I_{n_1} \circ I_{n_2} \circ A_1 \in \mathfrak{L}$  and if  $A_1 \nabla A_2$  is defined then,

$$(I_{n_1} \circ I_{n_2} \circ A_1) \nabla (I_{n_1} \circ I_{n_2} \circ A_2) = I_{n_1} \circ I_{n_2} \circ (A_1 \nabla A_2) .$$

The L.H.S. lies in  $\mathfrak{L}$  hence  $A_1 \nabla A_2 \in \mathfrak{L}/\ell$  and so we have M2.

If  $B$  is square over  $R$  and  $A \in \mathfrak{L}/\ell$  then  $I \circ A \in \mathfrak{L}$  so  $I \circ A \circ B \in \mathfrak{L}$  and hence  $A \circ B \in \mathfrak{L}/\ell$ . Therefore M3 holds. Also if  $1 \circ A \in \mathfrak{L}/\ell$  then  $I_n \circ 1 \circ A \in \mathfrak{L}$ . But  $I_n \circ 1 = I_{n+1}$  so  $A \in \mathfrak{L}/\ell$ . Consequently M4 holds and we have shown that  $\mathfrak{L}/\ell$  is a matrix ideal. If  $\mathcal{P}$  is a matrix ideal containing  $\mathfrak{L}$  then;

$$A \in \mathfrak{L}/\ell \Rightarrow I \circ A \in \mathfrak{L} \Rightarrow I \circ A \in \mathcal{P} \Rightarrow A \in \mathcal{P} \Rightarrow \mathcal{P} \supseteq \mathfrak{L}/\ell .$$

Hence  $\mathfrak{L}/\ell$  is the least matrix ideal containing  $\mathfrak{L}$ . Now  $\mathfrak{L}/\ell$  is proper if and only if  $\mathfrak{L}/\ell \neq \emptyset$ . When  $\mathcal{P}$  is a matrix ideal and  $\Sigma$  is either a

singleton set or a set of square matrices closed under diagonal sums and containing 1 then in a similar fashion we can show that  $\mathcal{P}/\Sigma = \{B \in \mathfrak{M}(R) \mid B \oplus A \in \mathcal{P}, A \in \Sigma\}$  is also a matrix ideal containing  $\mathcal{P}$ . Summarizing these results we see,

Proposition 5.2  $\mathfrak{L}/\mathfrak{l}$  is the least matrix ideal containing  $\mathfrak{L}$  and  $\mathfrak{L}/\mathfrak{l}$  is proper if and only if  $\mathfrak{L} \cap \mathfrak{l} = \emptyset$ . ■

Since every matrix ideal contains  $\mathfrak{L}$  we see  $\mathfrak{L}/\mathfrak{l}$  is the least matrix ideal (written as  $\mathcal{P}_0$ ) and we have,

Proposition 5.3 Let  $R$  be any ring. Then  $R$  has a proper matrix ideal if and only if no unit matrix can be written as a column determinantal sum of matrices which are hollow or degenerate. ■

It is now possible to consider when a ring  $R$  has a prime matrix ideal. The following is taken directly from Cohn [1].

Definition. Given two matrix ideals  $\mathcal{P}_1, \mathcal{P}_2$  in a ring  $R$ , their product, denoted by  $\mathcal{P}_1 \mathcal{P}_2$ , is defined as the matrix ideal generated by all  $A_1 \oplus A_2$  with  $A_i \in \mathcal{P}_i$  ( $i=1,2$ ).

The product so defined is easily seen to be associative and from property (f) it follows that the product is commutative.

Lemma 5.4 In any ring  $R$ , let  $\mathcal{X}_i \subseteq \mathfrak{M}(R)$  ( $i=1,2$ ),  $\mathcal{X}$  the set of matrices  $A_1 \oplus A_2$  ( $A_i \in \mathcal{X}_i$ ) and  $\mathcal{P}_1, \mathcal{P}_2, \mathcal{P}$  the matrix ideals generated by  $\mathcal{X}_1, \mathcal{X}_2, \mathcal{X}$ , respectively. Then  $\mathcal{P} = \mathcal{P}_1 \mathcal{P}_2$ .

Proof. Clearly  $\mathcal{X} \subseteq \mathcal{P}_1 \mathcal{P}_2$ , hence  $\mathcal{P} \subseteq \mathcal{P}_1 \mathcal{P}_2$ . To establish equality, let  $A_i \in \mathcal{X}_i$ , then  $A_1 \oplus A_2 \in \mathcal{P}$  by definition, hence  $\mathcal{X}_1 \subseteq \mathcal{P}/A_2$  and so  $\mathcal{P}_1 \subseteq \mathcal{P}/A_2$ . It

follows that  $B_1 \oplus A_2 \in \mathcal{P}$  for all  $B_1 \in \mathcal{P}_1$ , so fixing  $B_1$  we have  $\mathcal{X}_2 \subseteq \mathcal{P}/B_1$ , hence  $\mathcal{P}_2 \subseteq \mathcal{P}/B_1$  and so  $B_1 \oplus B_2 \in \mathcal{P}$  for all  $B_1 \in \mathcal{P}_1$ , and it follows that  $\mathcal{P}_1 \mathcal{P}_2 \subseteq \mathcal{P}$ . ■

**Proposition 5.5** *for any matrix ideal  $\mathcal{P}$  in a ring  $R$  the following three properties are equivalent.*

- (a)  $\mathcal{P}$  is prime,
- (b)  $\mathcal{P}$  is proper and for any matrix ideals  $\mathcal{P}_1, \mathcal{P}_2$  we have  $\mathcal{P}_1 \mathcal{P}_2 \subseteq \mathcal{P} \Rightarrow \mathcal{P}_1 \subseteq \mathcal{P}$  or  $\mathcal{P}_2 \subseteq \mathcal{P}$ ,
- (c)  $\mathcal{P}$  is proper and for any matrix ideals  $\mathcal{P}_1, \mathcal{P}_2$  such that  $\mathcal{P}_1 \supseteq \mathcal{P}$  we have  $\mathcal{P}_1 \mathcal{P}_2 \subseteq \mathcal{P} \Rightarrow \mathcal{P}_1 = \mathcal{P}$  or  $\mathcal{P}_2 = \mathcal{P}$ .

**Proof.** (a) $\Rightarrow$ (b). Let  $\mathcal{P}$  be prime and  $\mathcal{P}_1 \mathcal{P}_2 \subseteq \mathcal{P}$  but  $\mathcal{P}_i \not\subseteq \mathcal{P}$  ( $i=1,2$ ). Then there exist  $A_i \in \mathcal{P}_i$  but  $A_i \notin \mathcal{P}$ . Since  $\mathcal{P}$  is prime,  $A_1 \oplus A_2 \notin \mathcal{P}$ , but  $A_1 \oplus A_2 \in \mathcal{P}_1 \mathcal{P}_2 \subseteq \mathcal{P}$ , a contradiction. Clearly  $\mathcal{P}$  is also proper.

(b) $\Rightarrow$ (c) is clear; to prove (c) $\Rightarrow$ (a), suppose that  $A_1 \oplus A_2 \in \mathcal{P}$ . Consider the matrix ideal  $\mathcal{P}_i$  generated by  $\mathcal{P}$  and  $A_i$ ; for any  $B_i \in \mathcal{P} \cup \{A_i\}$ ,  $B_1 \oplus B_2 \in \mathcal{P}$ , hence  $\mathcal{P}_1 \mathcal{P}_2 \subseteq \mathcal{P}$ . By hypothesis,  $\mathcal{P}_1$  or  $\mathcal{P}_2$  must equal  $\mathcal{P}$ , so  $A_1$  or  $A_2$  lies in  $\mathcal{P}$ , i.e. (a). ■

The usual method of constructing prime ideals also works for prime matrix ideals:

**Theorem 5.6** *Let  $R$  be any ring,  $\Sigma$  a non-empty subset of  $\mathfrak{M}(R)$  closed under diagonal sums and  $\mathcal{A}$  any matrix ideal such that  $\mathcal{A} \cap \Sigma = \emptyset$ . Then there exists a matrix ideal  $\mathcal{P}$  which is maximal subject to the conditions  $\mathcal{P} \supseteq \mathcal{A}$ ,  $\mathcal{P} \cap \Sigma = \emptyset$ , and any such matrix ideal is prime.*

**Proof.** The collection  $\mathcal{C}$  of all matrix ideals containing  $\mathcal{A}$  and disjoint from  $\Sigma$  is clearly inductive, so by Zorn's lemma it has a

maximal member  $\mathcal{P}$ , and this satisfies the conditions of the theorem. Any such  $\mathcal{P}$  is proper, because  $\Sigma$  is non-empty; now let  $\mathcal{P}_i \supseteq \mathcal{P}$  be matrix ideals such that  $\mathcal{P}_1, \mathcal{P}_2 \subseteq \mathcal{P}$ . If  $\mathcal{P}_i \neq \mathcal{P}$  ( $i=1,2$ ), then  $\mathcal{P}_i \supset \mathcal{P}$ , so by maximality of  $\mathcal{P}$ ,  $\mathcal{P}_i \cap \Sigma \neq \emptyset$ . Take  $A_i \in \mathcal{P}_i \cap \Sigma$ , then  $A_1 \oplus A_2 \in \mathcal{P} \cap \Sigma$ , which is a contradiction. Hence  $\mathcal{P}_1$  or  $\mathcal{P}_2$  equals  $\mathcal{P}$ , and so  $\mathcal{P}$  is prime by the previous proposition. ■

This theorem shows for example that every maximal proper matrix ideal is prime; we need only take  $\Sigma = \ell$ .

Corollary. *Any ring  $R$  has a prime matrix ideal if and only if  $R$  has a proper matrix ideal.* ■

Hence by Proposition 5.3 we have;

Theorem 5.7 *Any ring  $R$  has a prime matrix ideal if and only if no unit matrix can be written as a column determinantal sum of matrices which are either hollow or degenerate.* ■

#### Localisation at a Prime Pair.

We can now proceed to show that when a ring  $R$  does indeed have a prime matrix ideal then there exists a homomorphism from  $R$  to a field. We shall first consider the concept of localisation at a prime pair introduced by Malcolmson [4]. If  $\Sigma$  is a lower multiplicative set of matrices over a ring  $R$  and  $\mathcal{P}$  is a matrix ideal of  $R$  then a pair  $Q=(\Sigma, \mathcal{P})$  is said to be a **prime pair** if  $A \oplus B \in \mathcal{P}$  and  $A \in \Sigma$  implies that  $B \in \mathcal{P}$ . The idea is to form a localisation where the image of  $\Sigma$  becomes invertible while somehow setting the elements of  $\mathcal{P}$  to become 'singular'. If  $(\Sigma, \mathcal{P})$  is a pair Malcolmson showed that  $(\Sigma, \mathcal{P}/\Sigma)$  is in fact prime and  $\mathcal{P}/\Sigma$  is proper iff  $\mathcal{P} \cap \Sigma = \emptyset$ .

Given a prime pair  $Q=(\Sigma, \mathcal{I})$  we can form the well defined **associated subset** of  $M_\Sigma$  as  $\mathcal{Q}=\{(A_\circ A_* A_\infty) \in M_\Sigma \mid (A_\circ A_*) \in \mathcal{I}\}$ . Below we show that  $\mathcal{Q}$  is a  $\Sigma$ -matrix ideal and then we call the homomorphism  $\lambda_{\mathcal{Q}}:R \rightarrow R_{\mathcal{Q}}$  the localisation at  $\mathcal{Q}$ .

**Proposition 5.8** *If  $Q=(\Sigma, \mathcal{I})$  is a prime pair then the associated subset of  $M_\Sigma$ ,  $\mathcal{Q}$ , is a  $\Sigma$ -matrix ideal, and the localisation at  $\mathcal{Q}$  is  $\Sigma$ -invertible.*

**Proof.** Since  $\mathcal{I}$  is a matrix ideal it contains all hollow matrices and so  $\mathcal{Q} \supseteq \mathcal{L}$ . If  $A, B \in \mathcal{Q}$  then the numerator of  $A \square B$  can be written;

$$\begin{bmatrix} B_\circ & B_* & B_\infty & 0 \\ 0 & 0 & -A_\infty & A_* \end{bmatrix} \quad \nabla \quad \begin{bmatrix} 0 & B_* & B_\infty & 0 \\ A_\circ & 0 & -A_\infty & A_* \end{bmatrix}$$

and the matrix on the left is in  $\mathcal{I}$  since  $(B_\circ B_*) \in \mathcal{I}$  and  $\mathcal{I}$  is a matrix ideal. After some row and column permutations the matrix on the right is similarly seen to lie in  $\mathcal{I}$  since  $(A_\circ A_*) \in \mathcal{I}$ . Hence since  $\mathcal{I}$  is closed w.r.t. determinantal sums we see that  $A \square B \in \mathcal{Q}$  and  $\mathcal{Q}$  is an additive subsemigroup of  $M_\Sigma$ . We now show that  $\mathcal{Q}$  is unitary. Assume that  $A \square B, B \in \mathcal{Q}$ . Then the numerator of  $A \square B$ ,

$$\begin{bmatrix} B_\circ & B_* & B_\infty & 0 \\ A_\circ & 0 & -A_\infty & A_* \end{bmatrix} \text{ lies in } \mathcal{I}.$$

If  $(B_\circ B_*) \in \mathcal{I}$  then by the properties of a matrix ideal,

$$\begin{bmatrix} -B_\circ & B_* & B_\infty & 0 \\ 0 & 0 & -A_\infty & A_* \end{bmatrix} \in \mathcal{I}.$$

The determinantal sum of the above two matrices is defined w.r.t. the first column and hence,

$$\begin{bmatrix} 0 & B_* & B_\infty & 0 \\ A_\circ & 0 & -A_\infty & A_* \end{bmatrix} \in \mathcal{P} \Rightarrow \begin{bmatrix} B_* & B_\infty & 0 & 0 \\ 0 & 0 & A_\circ & A_* \end{bmatrix} \in \mathcal{P}.$$

Since  $Q$  is a prime pair we get  $(A_\circ A_*) \in \mathcal{P}$  and  $A \in \Omega$ . So  $\Omega$  is unitary and it just remains to show that  $\Omega$  is an ideal w.r.t. the multiplicative semigroup structure of  $M_\Sigma$ . If  $A \in \Omega$  and  $B \in M_\Sigma$  then the numerator of both  $A.B$  and  $B.A$  is immediately seen to lie in  $\mathcal{P}$  by the properties of a matrix ideal. Consequently by Theorem 2.6  $\lambda_\Omega: R \rightarrow R_\Omega$  is  $\Sigma$ -invertible. ■

Malcolmson used this result to get a bound on the kernel of the universal  $\Sigma$ -invertible homomorphism by taking a 'minimal' prime pair  $(\Sigma, \mathcal{P}_\circ / \Sigma)$  where  $\mathcal{P}_\circ$  is the least matrix ideal. This is no longer necessary as we have a complete description of the kernel.

When a ring  $R$  has a prime matrix ideal  $\mathcal{P}$  then the complement of  $\mathcal{P}$ ,  $\setminus \mathcal{P}$ , is easily seen to be a lower multiplicative set of square matrices over  $R$  and  $(\setminus \mathcal{P}, \mathcal{P})$  is a prime pair. With the present notation the localisation at this prime pair is written  $\lambda_{\mathcal{P}}: R \rightarrow R_{\mathcal{P}}$ .

Theorem 5.9 (Cohn) *If  $R$  is any ring with a prime matrix ideal  $\mathcal{P}$  then the localisation at  $\mathcal{P}$  (the associated  $\setminus \mathcal{P}$ -matrix ideal) is a homomorphism from  $R$  to a skew field whose singular kernel is precisely  $\mathcal{P}$ .*

Proof  $\mathcal{P}$  is the associated  $\setminus \mathcal{P}$ -matrix ideal and consists of all those admissible matrices with numerators lying in  $\mathcal{P}$  so  $(A_\circ A_* A_\infty) \notin \mathcal{P}$  implies that  $(A_\circ A_*) \notin \mathcal{P}$  and  $(A_\infty A_* A_\circ)$  is an admissible matrix. We observe the following;

$$(A_{\infty} \ A_* \ A_0) \cdot (A_0 \ A_* \ A_{\infty}) = \begin{bmatrix} A_0 & A_* & A_{\infty} & 0 & 0 \\ 0 & 0 & -A_{\infty} & A_* & A_0 \end{bmatrix} \rightarrow$$

$$\begin{bmatrix} A_0 & A_* & A_{\infty} & 0 & 0 \\ 0 & -A_* & -A_{\infty} & A_* & A_0 \end{bmatrix} \rightarrow \begin{bmatrix} A_0 & A_* & A_{\infty} & 0 & 0 \\ 0 & 0 & -A_{\infty} & A_* & A_0 \end{bmatrix} \sim (A_0 \ A_* \ A_0) \sim 1$$

Similarly we can get  $(A_0 \ A_* \ A_{\infty}) \cdot (A_{\infty} \ A_* \ A_0) \sim 1$ , and hence all non-zero elements in  $R_{\mathfrak{p}}$  have multiplicative inverses and  $R_{\mathfrak{p}}$  is seen to be a field.

We already have that  $\text{Ker} \lambda_{\mathfrak{p}} \subseteq \mathcal{P}$  since  $\lambda_{\mathfrak{p}}$  inverts the complement of  $\mathcal{P}$ . Now we proceed to show that  $\mathcal{P} \subseteq \text{Ker} \lambda_{\mathfrak{p}}$ . Let  $P \in \mathcal{P}$  and denote by  $P^{(1)}$  the matrix obtained from  $P$  by omitting the first column and the  $i$ th row. If  $P^{(1)} \in \mathcal{P}$  for all  $i$ , then by induction and the use of column determinantal sums w.r.t. the initial column we get that  $P \in \text{Ker} \lambda_{\mathfrak{p}}$ . So w.l.o.g. we can assume that  $P^{(1)} \notin \mathcal{P}$ . Then since  $\setminus \mathcal{P}$  is lower multiplicative and admits row permutations we have that  $(Pe_1)$  is admissible, and moreover it is admissible for zero since the numerator lies in  $\mathcal{P}$ . Hence from Theorem 2.6 there is an equation in  $R_{\mathfrak{p}}$ ;  $P_0 = (P_* e_1)(u_* 0)^T$  i.e  $P$  is a zero divisor in  $R_{\mathfrak{p}}$  and so as required we have  $P \in \text{Ker} \lambda_{\mathfrak{p}}$ . ■

(Cohn actually showed that  $R_{\mathfrak{p}}$  the universal  $\setminus \mathcal{P}$ -inverting ring is local with residue class field  $R_{\mathfrak{p}}$ .)

Corollary For any ring  $R$  there exists a homomorphism from  $R$  to a field if and only if no unit matrix can be written as a column determinantal sum of hollow or degenerate matrices. ■

With the results of the previous chapter we are now in a position to determine a criterion for any ring to have a field of fractions. This is a refinement of Cohn's criterion (a reference is given after the proof). For completeness we then go on to prove Dicks and Sontag's result classifying all rings having a universal field of fractions inverting all full matrices over that ring.

Theorem 6.1 *Any non-zero ring  $R$  has a field of fractions if and only if no diagonal matrix over  $R$  with non-zero diagonal entries can be expressed as the column determinantal sum of matrices that are either hollow or degenerate.*

Proof If  $R$  has a field of fractions then  $R$  is embeddable in a field over which every diagonal matrix with non-zero diagonal entries is invertible. However over a field the column determinantal sum of non-full matrices is singular, so we have proved the necessity of the condition.

To prove sufficiency we note that  $\Sigma$ , the set of diagonal matrices over  $R$  with non-zero diagonal entries is closed w.r.t. diagonal sums and non-empty. Then the condition states that  $\Sigma$  is disjoint from the least matrix pre-ideal. From the description of the least matrix ideal in Proposition 5.2 we can see that  $\Sigma$  is also disjoint from the least matrix ideal. Hence by applying Theorem 5.6 we have a prime matrix ideal  $\mathcal{P}$  disjoint from  $\Sigma$  which by Theorem 5.9 gives rise to a homomorphism from  $R$  to a field  $F$  with singular kernel precisely  $\mathcal{P}$  i.e.  $\Sigma$  is inverted which is sufficient to prove that we have an embedding in a field of fractions. ■

(cf. Cohn [1], Th. 7.5.6.). We note that this condition ensures that  $R$  is an integral domain for if  $ab=0$ , then

$$\begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} = \begin{bmatrix} a & 0 \\ 1 & b \end{bmatrix} \nabla \begin{bmatrix} 0 & 0 \\ -1 & b \end{bmatrix}$$

where the second matrix on the right is clearly hollow, and the first matrix on the right is degenerate since the first column multiplied by  $b$  on the right is just the second column.

If there is an homomorphism from a ring  $R$  to a field then no non-full matrix over  $R$  can be inverted since a non-full matrix over a field is singular. Hence the most that can be hoped for is that all full matrices over  $R$  be inverted and this is only possible if the singular kernel of the homomorphism consists precisely of the non-full matrices over  $R$ . In this case we must have an embedding since any non-zero  $1 \times 1$  matrix is full and so maps to an invertible element. A ring homomorphism which keeps all full matrices full is called **honest** so by the above we have that there is an honest homomorphism from a ring to a field if and only if there is a fully inverting homomorphism from the ring to a field. Let  $\Phi = \Phi(R)$  denote the set of all full matrices over  $R$ , if this set is lower multiplicative then we can consider  $R_\Phi$  the universal localisation at  $\Phi$ .

Proposition 6.2 (Cohn) *Let  $R$  be any ring. If  $\Phi$ , the set of full matrices over  $R$ , is lower multiplicative and  $R_\Phi \neq 0$  then  $R_\Phi$  is a field.*

Proof Let  $p$  be a non-unit in  $R_\Phi$  with an admissible matrix

$A = (A_0 A_* A_\infty)$ . By Cramer's rule the image of the numerator over  $R_\Phi$  is stably associated to  $p$  and hence must be non-full over  $R$  or else  $p$  is invertible. So  $(A_0 A_* A_\infty) = PQ$  where  $P \in {}^n R^{n-1}$  and  $Q \in {}^{n-1} R^n$  and then we can write  $A$  as follows;

$$A = (A_0 A_* A_\infty) = (PQ A_\infty) = (P A_\infty) \begin{bmatrix} Q & 0 \\ 0 & 1 \end{bmatrix} .$$

Now  $(P A_\infty)$  must be full for it is a left factor of the denominator of  $A$  which is full by definition and so the image of  $(P A_\infty)$  is invertible over  $R_\Phi$ . Since  $A$  is an admissible matrix for  $p$  then we have an equation  $Au=0$  over  $R_\Phi$  where the last component of  $u$  is  $p$ , but from above we can cancel the left factor  $(P A_\infty)$  in the equation to deduce that  $p=0$  as required. ■

We are now in a position to determine exactly when  $R_\Phi$  is a field.

Theorem 6.3 (Cohn) *If  $R$  is any ring then the two following conditions are equivalent:*

- (i) *The set of non-full matrices over  $R$  is a prime matrix ideal.*
- (ii)  *$R$  has a universal field of fractions inverting all full matrices over  $R$ .*

Proof If (i) holds then the set  $\Phi$  of full matrices is lower multiplicative and so by Proposition 6.2  $R_\Phi$  is either a field or the zero ring. If  $R_\Phi=0$  then the admissible matrix  $(1 \ 1)$  represents zero. However from Theorem 5.7 we know that the least  $\Phi$ -matrix ideal has numerators which are non-full because (i) ensures that those admissible matrices with full denominators and non-full numerators form a  $\Phi$ -matrix ideal which necessarily contains the

least  $\Phi$ -matrix ideal. Hence if  $R_\Phi = 0$  then 1 is non-full; a contradiction, so  $R_\Phi$  is a field.

Conversely if  $R$  is embeddable in such a field then we have a fully inverting homomorphism from  $R$  to a field. Hence the singular kernel is a prime matrix ideal and from an earlier comment we know that the singular kernel must consist precisely of the non-full matrices over  $R$ . ■

We are now in a position to classify all rings with a universal field of fractions inverting all full matrices but first we shall have to state (without proof) a couple of results from Cohn [1] Chapter 5.

Theorem 6.4 *Every honest homomorphism of a ring preserves the inner rank.* ■

Let  $R$  be a ring. If  $PQ=0$  where  $P \in R^{r \times n}$ ,  $Q \in R^{n \times s}$  and  $r+s > n$  implies that,

$$\rho(P) + \rho(Q) \leq n$$

then  $R$  is said to be a **Sylvester domain**.

Lemma 6.5 *Let  $R$  be a Sylvester domain. Then*

(i) *if  $A, B$  are full matrices over  $R$  then  $A \otimes B$  is also full.*

(ii) *if  $A, B, C$  are any matrices over  $R$  with the same number of rows and if  $\rho(A, B) = \rho(A, C) = \rho(A)$  and  $A$  is right full, then*

$$\rho(A \begin{pmatrix} B \\ C \end{pmatrix}) = \rho(A) .$$

■

With these two results we can prove the main theorem of this chapter.

Theorem 6.6 (Dicks and Sontag [6]) *A ring  $R$  has a universal field of fractions inverting all full matrices over  $R$  if and only if  $R$  is a Sylvester domain.*

Proof If  $R_\phi$  is a field then  $\lambda:R \rightarrow R_\phi$  is an honest homomorphism from  $R$  to a field, hence by Theorem 6.4 we have an inner rank preserving homomorphism from  $R$  to a field. Any field necessarily satisfies Sylvester's law of nullity and so is seen to be a Sylvester domain. Therefore  $R$  must also be a Sylvester domain as the inner rank is preserved by the homomorphism.

Conversely by Theorem 6.3 it suffices to show that the set of non-full matrices over a Sylvester domain form a prime matrix ideal. Using the basic properties of non-full matrices this amounts to showing that the set of non-full matrices is closed under column determinantal sums and that the set of full matrices is closed under diagonal sums. Lemma 6.5 (i) shows that the set of full matrices over a Sylvester domain is closed w.r.t. diagonal sums. To prove that the set of non-full matrices is closed w.r.t. column determinantal sums suppose that  $A=(A_*,a)$ ,  $B=(A_*,b)$ ,  $C=(A_*,a+b)$  where  $A_* \in R^{n \times n-1}$ ;  $a,b$  are columns over  $R$  and  $A,B$  are not full. If  $A_*$  is right full then  $n-1 = \rho(A_*) = \rho(A_*,a) = \rho(A_*,b)$ , so by Lemma 6.5 (ii)  $n-1 = \rho(A_*,a,b) = \rho(A_*,a+b,b) \geq \rho(A_*,a+b) = \rho(C)$ . If  $A_*$  is not right full then  $\rho(A_*) < n-1$  and then  $\rho(C) < n$ . In either case  $C$  is non-full as required. ■

In particular, since a semifir is a Sylvester domain we have the following corollary.

Corollary 6.7 (Cohn) *Any semifir  $R$  has a universal field of fractions  $K$  and any full matrix over  $R$  is invertible over  $K$ . ■*

## References

- [1] P.M.Cohn, Free Rings and their Relations, 2nd Ed. LMS Monographs No.19, Academic Press, London and New York 1985.
  
- [2] P.M.Cohn, One-sided Localization in Rings, to appear
  
- [3] Clifford and Preston, Algebraic Theory of Semigroups Vol.2, AMS 1961.
  
- [4] P.Malcolmson, Construction of Universal Matrix Localizations, Advances in Non-Comm. Ring Theory, Proc. 12th G.H.Hudson Symp., Plattsburgh, 1981. Lecture Notes in Math. No.951, Springer-Verlag, Berlin, pp.117-131.
  
- [5] P.Malcolmson, A Prime Matrix Ideal Yields a Skew Field, J. London Math. Soc. (2), 18 (1978), 221-233.
  
- [6] W.Dicks and E.D.Sontag, Sylvester Domains, J. Pure Appl. Algebra 13 (1978), 243-275.