

Journal Pre-proofs

Psychosocial factors and hospitalisations for COVID-19: Prospective cohort study based on a community sample

G.D. Batty, I.J. Deary, M. Luciano, D.M. Altschul, M. Kivimäki, C.R. Gale

PII: S0889-1591(20)31309-X
DOI: <https://doi.org/10.1016/j.bbi.2020.06.021>
Reference: YBRBI 4198


To appear in: *Brain, Behavior, and Immunity*

Received Date: 9 June 2020
Accepted Date: 14 June 2020

Please cite this article as: Batty, G.D., Deary, I.J., Luciano, M., Altschul, D.M., Kivimäki, M., Gale, C.R., Psychosocial factors and hospitalisations for COVID-19: Prospective cohort study based on a community sample, *Brain, Behavior, and Immunity* (2020), doi: <https://doi.org/10.1016/j.bbi.2020.06.021>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc.



Psychosocial factors and hospitalisations for COVID-19: Prospective cohort study based on a community sample

G.D. Batty^a (Email: david.batty@ucl.ac.uk; ORCID: 0000-0003-1822-5753)
Professor of Epidemiology

I.J. Deary^b (ian.deary@ed.ac.uk; 0000-0002-1733-263X)
Professor of Differential Psychology

M. Luciano^c (michelle.luciano@ed.ac.uk; 0000-0003-0935-7682)
Reader in Psychology

D.M. Altschul^c (drew.altschul@ed.ac.uk; 0000-0001-7053-4209)
Postdoctoral Researcher in Psychology

M. Kivimäki^a (m.kivimaki@ucl.ac.uk; 0000-0002-4699-5627)
Professor of Social Epidemiology

C.R. Gale^{b,d} (crg@mrc.soton.ac.uk; 0000-0002-3361-8638)
Emeritus Professor of Cognitive Epidemiology

^aDepartment of Epidemiology and Public Health, University College London, UK

^bLothian Birth Cohorts, Department of Psychology, University of Edinburgh, UK

^cDepartment of Psychology, The University of Edinburgh, Edinburgh, UK

^dMRC Lifecourse Epidemiology Unit, University of Southampton, UK

Manuscript statistics: 4331 words (including 263 abstract), 3 tables, 2 figures, 5 supplemental tables

Correspondence: David Batty, Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London, UK, WC1E 6BT. E. david.batty@ucl.ac.uk

Funding: GDB is supported by the UK Medical Research Council (MR/P023444/1) and the US National Institute on Aging (1R56AG052519-01; 1R01AG052519-01A1); IJD by the UK Medical Research Council (MR/R024065/1), UK Economic and Social Research Council (ES/S015604/1), and US National Institute on Aging (NIH), US (1R01AG054628-01A1); DMA by the UK Medical Research Council (MRC_MC_PC_17209); and MK by the UK Medical Research Council (MR/R024227), US National Institute on Aging (NIH), US (R01AG056477), NordForsk, and Academy of Finland (311492). These funders, who provided no direct financial or material support for the work, had no role in study design, data collection, data analysis, data interpretation, or report preparation.

Acknowledgement: We thank UK Biobank study members for their generosity in participating.

Access to data: Data from UK Biobank (<http://www.ukbiobank.ac.uk/>) are available to *bona fide* researchers upon application. Part of this research has been conducted using the UK Biobank Resource under Application 10279.

Competing of interest: Ian Deary was responsible for the design of some of the cognitive function tests in the revised battery used in the imaging sessions in UK Biobank; he is also a study participant. There are no other potential competing interests to report.

Transparency: GDB affirms that the manuscript is an honest, accurate, and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned have been explained.

Contributions: The authors collectively generated the idea for the present paper and formulated an analytical plan; ML built the dataset; CRG carried out the data analyses; DMA prepared the figures; and GDB wrote the first draft of manuscript. All authors commented on an earlier version of the manuscript. GDB, CRG, ML and IJD will act as guarantors for this work. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. CRG and ML had full access to UK Biobank data. GDB takes responsibility for the decision to submit the manuscript for publication.

Dissemination to participants and related patient and public communities: Findings will be disseminated via the media departments of the authors' institutes. Results from UK Biobank are routinely disseminated to study participants via the study website and Twitter feed.

Ethical approval: In UK Biobank, ethical approval for data collection was received from the North-West Multi-centre Research Ethics Committee and the research was carried out in accordance with the Declaration of Helsinki of the World Medical Association.

Open access statement: University College London will cover open access charges (subject to confirmation).

Abstract

Background: While certain infectious diseases have been linked to socioeconomic disadvantage, mental health problems, and lower cognitive function, relationships with COVID-19 are either uncertain or untested. Our objective was to examine the association of a range of psychosocial factors with hospitalisation for COVID-19.

Methods: UK Biobank, a prospective cohort study, comprises around half a million people who were aged 40 to 69 years at study induction between 2006 and 2010 when information on psychosocial factors and covariates were captured. Hospitalisation for COVID-19 were ascertained between 16th March and 26th April 2020.

Results: There were 908 hospitalisations for COVID-19 in an analytical sample of 431,051 England-based study members. In age- and sex-adjusted analyses, an elevated risk of COVID-19 was related to disadvantaged levels of education (odds ratio; 95% confidence interval: 2.05; 1.70, 2.47), income (2.00; 1.63, 2.47), area deprivation (2.20; 1.86, 2.59), occupation (1.39; 1.14, 1.69), psychological distress (1.58; 1.32, 1.89), mental health (1.50; 1.25, 1.79), neuroticism (1.19; 1.00, 1.42), and performance on two tests of cognitive function – verbal and numerical reasoning (2.66; 2.06, 3.34) and reaction speed (1.27; 1.08, 1.51). These associations were graded (p -value for trend ≤ 0.038) such that effects were apparent across the full psychosocial continua. After mutual adjustment for these characteristics plus ethnicity, comorbidity, and lifestyle factors, only the relationship between lower cognitive function as measured using the reasoning test and a doubling in the risk of the infection remained (1.98; 1.38, 2.85).

Conclusions: A range of psychosocial factors revealed associations with hospitalisations for COVID-19 of which the relation with cognitive function, a marker of health literacy, was most robust.

Key words: risk factors, COVID-19, hospitalisation, cohort study, UK Biobank

1. Introduction

With outbreaks reported across 114 countries, the novel coronavirus referred to as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was declared pandemic by the World Health Organization on 11th March 2020.(World Health Organization Director-General, 2020) By 11th May, in the absence of widespread testing in most countries, there was global notification of 4 million confirmed cases of coronavirus disease 2019 (COVID-19) – the disease caused by SARS-CoV-2 – with it being implicated in more than 270,000 deaths.(World Health Organization, 2020) Equivalent data releases for the UK indicated 223,060 cases and 32,065 fatalities.(UK Government, 2020)

Prior pandemics – Spanish influenza in 1918 and Swine influenza in 2009 – were notable for marked inequalities in their occurrence, whereby more socioeconomically disadvantaged countries,(Simonsen et al., 2013; Wiemken et al., 2020) cities,(Rutter et al., 2012) neighbourhoods,(Grantz et al., 2016; Mamelund, 2006) and individuals(Bengtsson et al., 2018) experienced the highest mortality rates from the infection. Recent findings from analyses of data for COVID-19 hospitalisations across the five boroughs of New York City(Wadhera et al., 2020) and deaths involving the infection in the UK(Office for National Statistics, 2020) reveal higher rates in more deprived areas. The mechanisms that underlie these disparities are likely to be numerous and might involve overcrowded living and working conditions, comorbidity, poor access to healthcare, and a relative lack of understanding of prevention advice among socially disadvantaged individuals.(La et al., 2018) Indirect pathways might include the higher prevalence of unfavourable health behaviours (Hamer et al., 2020) – cigarette smoking, alcohol intake, and suboptimal nutrition – in lower social groups which in themselves have been linked to selected lower respiratory tract infections.(Yap et al., 2000)

Although correlated with socioeconomic status,(Neisser et al., 1996; Russ et al., 2012) mental health and cognitive function might have independent utility in understanding the burden of respiratory disease.

Poor mental health has been hypothesised to be a potential consequence of COVID-19 based on the findings of studies of survivors of the severe acute respiratory syndrome (SARS) pandemic,(Lam et al., 2009; Lee et al., 2007) but it may also influence the risk of contracting the infection, at least in part by impairing innate or adaptive immunity(Cohen et al., 1991) and diminishing the precautions taken to minimise risk. In a cross-sectional study, mental health problems were correlated with a higher likelihood of reporting the common cold,(Adam et al., 2013) a species of coronavirus. In cohort studies generated using linked electronic registries, people with a history of depression,(Andersson et al., 2016) psychosis,(Seminog and Goldacre, 2013) and stress disorders(Jiang et al., 2019) serious enough to warrant treatment in a psychiatric care facility subsequently experienced elevated rates of an array of respiratory infections. Additionally, in the general population, even moderate levels of self-reported symptoms of psychological distress (depression and anxiety) have been prospectively linked to an elevated risk of death from pneumonia despite adjustment for confounding factors which include socioeconomic position.(Hamer et al., 2019)

In the COVID-19 pandemic, the public has been offered much preventative advice and guidelines which span the simple and practical to the complex, contradictory and false.(Cuan-Baltazar et al., 2020; Orso et al., 2020; Zarocostas, 2020) In order to diminish their risk of the infection, the population has to acquire, synthesise, and deploy this information but the ability to do so seems to vary by levels of health literacy(Wolf et al., 2020) just as it may for its close correlate, cognitive function. Although traditionally studied in the context of non-communicable disease,(Batty et al., 2010; Batty et al., 2008b; Deary et al., 2011) higher levels of cognitive ability – a psychological trait that involves the storage, selection, manipulation, and organisation of information – appear to be related to markedly lower rates of mortality from infectious disease after taking into account social circumstances.(Calvin et al., 2017; Gale et al., 2019)

With this evidence base giving us reason to anticipate a relation of these socioeconomic and psychological characteristics with incident COVID-19 infection, we explored them using data from UK Biobank, a large prospective cohort study. While we have reported on the influence of ethnicity (Lassale et al., 2020) and lifestyle factors (Hamer et al., 2020) on COVID-19 hospitalisation, to the best of our knowledge, this is the first examination of the role of selected socioeconomic characteristics (income, occupation) and any psychological factors (cognition, mental health, personality type) in the primary prevention of the infection.

2. Methods

We used data from both UK Biobank, a prospective cohort study, the sampling and procedures of which have been well described. (Sudlow et al., 2015) In brief, baseline data collection took place between 2006 and 2010 in twenty-two research assessment centres across the UK, resulting in a sample of 502,655 people aged 40 to 69 years (response rate 5.5%). (Sudlow et al., 2015) In UK Biobank, ethical approval was received from the North-West Multi-centre Research Ethics Committee, and the research was carried out in accordance with the Declaration of Helsinki of the World Medical Association, and participants gave informed consent. No additional ethical approval was required for present analyses of anonymised data.

Assessment of socioeconomic factors

We used four indicators of socioeconomic status. Total annual household income before tax was self-reported and classified into three groups (<18,000, -30,999, -51,999, \geq £52,000 GBP). For educational qualifications, we used a three category variable (degree, other qualifications, no qualifications). Using Standard Occupational Classifications of current job, or most recent if participants were not working or data on current job were missing, we produced three categories with managerial positions having the highest prestige: managers & senior officials, professional, associate professional & technical;

administrative & secretarial, & skilled; and personal service, sales & customer service, process, plant & machine operatives, elementary. Lastly, we used the Townsend deprivation index as our indicator of neighbourhood socioeconomic circumstances. Based on a composite of four characteristics (home and car ownership, employment, and number of household resident), participants' postcodes at recruitment were matched to areas from the most recent national census. A continuously scored variable, higher values denote greater deprivation.

Assessment of psychological factors

We used five psychological factors. Study members were asked if they had ever been under the care of a psychiatrist for any mental health problem; in the UK, such a referral would ordinarily have been triaged via a general practitioner. Symptoms of psychological distress – anxiety, worrying, anhedonia, and depression – were measured using the four item version of the Patient Health Questionnaire (PHQ-4)⁶ in which individual items are rated on a 4 point Likert scale from 0 (“not at all”) to 3 (“nearly every day”) such that total scores range from 0 to 12 (higher scores denote greater distress). Scores on the PHQ-4 show good agreement with longer scales, and reveal known correlations with demographic risk factors for depression and anxiety.⁷ Neuroticism was measured with the 12-item Eysenck Personality Questionnaire-Revised Short Form;⁸ higher scores denote higher levels.

Scores from two tests of cognitive functioning were used. Verbal and numerical reasoning was measured using a computerized 13-item multiple-choice test with a two-minute time limit. The score was the number of correct answers.(Davies et al., 2016) This test was introduced after the beginning of the baseline assessment period so data are available for a subset of study members (N=180,914). Reaction time was measured using a computerized Go/No-Go “Snap” game. Participants were presented with electronic images of two cards. If symbols on the cards were identical, participants were instructed to immediately push the button-box using their dominant hand. The first five pairs were used as a practice

with the remaining seven pairs, containing four identical cards, forming the assessment. Reaction time score was the mean time to press the button when each of these four pairs was presented. Choice reaction time correlates strongly with single mental tests that involve complex reasoning and knowledge.(Deary et al., 2001)

2.1. Assessment of confounding factors

Ethnicity was self-reported and categorised as White, Asian, Black, Chinese, Mixed, or other ethnic group. A social isolation scale was derived from enquiries concerning number of people in household, visiting friends/family, and social activities.(Elovainio et al., 2017) One point was allocated for living alone, one for friends/family visits less than once/month, and one for no weekly participation in social activities. A dichotomous variable was derived with social isolation denoted by a score of 3. Self-reported physician diagnosis was collected for vascular or heart problems, diabetes, chronic lung disease, asthma, and cancer. Cigarette smoking, physical activity, and alcohol consumption were measured using standard enquiries. Height and weight were measured directly during a medical examination from which body mass index was calculated using the usual formula (weight, kg/height,² m²). Forced expiratory volume in one second, a measure of pulmonary function, was quantified using spirometry with the best of three technically satisfactory exhalations used in our analyses. Handgrip strength was measured using a hydraulic hand dynamometer (Jamar J00105) with the participant maximally squeezing the handle of the dynamometer while seated for 3 seconds; an average of the readings from the right and left hand was used. Seated systolic and diastolic blood pressure measurements were made twice using the Omron HEM-7015IT digital blood pressure monitor (Omron Healthcare)²⁰ or, exceptionally, a manual sphygmomanometer; an average of the two readings was used herein. We defined hypertension according to existing guidelines as systolic/diastolic blood pressure $\geq 140/90$ mmHg and/or self-reported use of antihypertensive medication.(Chobanian et al., 2003) Non-fasting venous blood, available in a subsample, was drawn with assaying conducted at dedicated central laboratory for C-reactive protein,

glycated haemoglobin A1c, and high-density lipoprotein cholesterol.(Elliott and Peakman, 2008)

2.2. Ascertainment of hospitalisation for COVID-19

Provided by Public Health England, data on COVID-19 status downloaded on 1st May 2020 covered the period 16th March 2020 until 26th April 2020.(UK Biobank, 2020) Nose and/or throat swabs were taken from hospitalised patients and detection of SARS-CoV-2 can be regarded as an indication of severe disease.(UK Biobank, 2020) With coverage being for England only, study members from Scotland and Wales were omitted from our analytical sample.

In preliminary analyses, we used three different COVID-19 case definitions based on these data: all apparent cases of the disease (N=908); cases based on samples from in-patients only (N=751); and cases based on two or more samples from in-patients (N=445) – the notion being that these patients were amongst the most severe cases. Evidence from prognostic studies of hospitalised patients in the USA(Richardson et al., 2020) and China(Wu et al., 2020) suggest that men, older individuals, ethnic minorities, and those with existing disease experience greater rates of progression to intensive care and death. Preliminary analyses of the present data on incidence of severe disease revealed similar associations irrespective of case definition (supplemental table 1). On the basis of the similarity of this predictive validity, we proceeded with our main analyses in which we used all COVID-19 cases (N=908).

3. Statistical analyses

We omitted from our analyses men and women who had died before 5th March 2020 – the latest date to which vital status data were available – as they could not contribute to the risk set for COVID-19. Odds ratios and accompanying 95% confidence intervals were computed using logistic regression models to summarise the relationship between psychosocial factors and COVID-19 hospitalisations. In the main analyses, we initially adjusted odds ratios for age and sex, followed by ethnicity, then covariates

organised into comorbidities (vascular disease, diabetes etc.), lifestyle factors (cigarette smoking etc.), and, depending on the psychosocial exposures of interest, socioeconomic or psychological factors. In preliminary analyses, the addition of biomarkers to the final model had no appreciable impact on the effects estimates relative to the final model in which they did not feature (supplemental tables 4 and 5 versus tables 2 and 3); these covariates were therefore not included in the main analyses. Analyses were conducted using Stata version 13.

4. Results

In 431,051 study members (236,725 women) there were 908 hospitalisations for COVID-19 between 16 March 2020 and 26th April in England (402 in women). Of the 28 baseline characteristics featured in table 1, only four – extant cancer, grip strength, neuroticism, and social isolation – did not reveal relationships with COVID-19 at conventional levels of statistical significance in unadjusted analyses. These were therefore excluded as covariates from subsequent multiple regression analyses.

In table 2 and figure 1 we depict the association between various socio-economic characteristics and the risk of hospitalisation for COVID-19 infection. After adjustment for age and sex, those study members who were most disadvantaged educationally, financially, and geographically experienced around a doubling in the risk of infection. Effects in these analyses were apparent across the full socioeconomic continuum (p for trend <0.0001). Whereas controlling for ethnicity had little impact on these gradients, partial attenuation was apparent after taking into account comorbidities and lifestyle factors.

However, adjusting for psychological characteristics had the largest attenuating effect relative to the minimally-adjusted (age, sex, and ethnicity) odds ratios. Although the risk of hospitalisation remained somewhat elevated at both lower levels of education and income, statistical significance at conventional levels was lost. Given the known correlation between education and cognitive ability (herein, $r=-0.40$, $p-$

value <0.0001), in sensitivity analyses we removed verbal and numerical reasoning test scores from the model containing the 5 psychological factors. This resulted in the magnitude of the low education–COVID-19 relationship being restored (odds ratio; 95% confidence interval for no qualifications: 2.08; 1.69, 2.56) and suggested most of the marked attention seen for this relationship after taking into account psychological factors could be ascribed to individual differences in cognitive ability rather than education. The association between area deprivation and risk of infection was more robust to these various statistical adjustments.

Of the socioeconomic variables, occupational classification of the study members revealed the weakest association with hospitalisation for COVID-19 and, in all analyses, study members in the administrative/secretarial occupations in fact experienced some protection against the infection. Lastly, after including up to seventeen covariates in the most complex multivariable models, there was evidence of some weak residual associations for income and deprivation but not for education.

In table 2 and figure 2 we illustrate the associations between psychological traits and the risk of COVID-19. In minimally-adjusted (age, sex, and ethnicity) analyses, all five psychological factors were related to the risk of hospitalisation with the infection. Effects for neuroticism and reaction time – weak initially – were essentially eliminated after control for comorbidities and any subsequent group of covariates. Adjustment for comorbidities also had a partial impact on the relation of distress, psychiatric consultation, and verbal and numerical reasoning with the infection, but associations largely remained, most obviously for reasoning score. After multiple control for all covariates, however, the only relationship that remained with COVID-19 was that for verbal and numerical reasoning such that the most disadvantaged group experienced around a doubling of hospitalisation risk.

We also carried out some planned sensitivity analyses. With the verbal and numerical reasoning test having been introduced part way through baseline data collection, as indicated, analyses featuring this variable were based on a subgroup of study members. To ensure direct comparison across statistical models, for each exposure we therefore recomputed our analyses based on a non-missing dataset (supplemental table 2 for socioeconomic characteristics, and supplemental table 3 for psychological characteristics). The same patterns of association was apparent in these sensitivity analyses.

5. Discussion

5.1. Principal findings

Our main findings were that disadvantaged levels of a series of psychosocial characteristics – education, income, area deprivation, mental health, and cognitive function – were related to an elevated risk of hospitalisations with COVID-19 in most of the analyses conducted. Net of mutual control for these factors, and after taking into account several other potential confounders, including lifestyle factors (Hamer et al., 2020) and ethnicity, (Lassale et al., 2020) however, only the association of lower cognitive function based on a test of verbal and numerical reasoning with a higher risk of this infection remained. That we were able to replicate findings for apparently known risk factors for COVID-19 from prognostic studies – being male, having an ethnic minority background, carrying a comorbidity – provides some support for the more novel findings for these psychosocial factors.

Our finding that the intermediate occupational group experienced a lower risk of hospitalisation was unexpected. A *post hoc* explanation is that this apparent ‘J’-shaped relation could in part be driven by the higher prestige category containing some medical professionals and, at the opposite end of the continuum, the personal services group being partially composed of carers, both of whom would be more likely to be exposed to the virus. Cardiovascular disease death is an exemplar of socioeconomic inequalities in disease risk, (Batty et al., 2009) and analyses of this endpoint in relation to these occupational groups

revealed a graded effect such that a doubling of risk was apparent in the most disadvantage group (age- and sex-adjusted hazard ratio; 95% confidence interval: 1.90; 1.61, 2.25) with intermediate rates evident in the administrative class (1.32; 1.11, 1.58). As such, these occupational classifications have some predictive validity, so lending some support to the apparently surprising result for COVID-19.

5.2. Comparison with results from existing studies

While this manuscript was under editorial evaluation, other analyses were published which used UK Biobank data to explore the relation of area-based deprivation and educational achievement with hospitalisation for COVID-19.(Niedzwiedz et al., 2020) While that study found that people who were disadvantaged experienced a higher risk of the infection, no account was made for cognitive function – seemingly an important explanatory factor in our analyses – and the role of other indices of socioeconomic position – occupational status and income – were not examined. Prognostic studies using area-based statistics have also recently appeared. In New York City, Manhattan, the most socioeconomically advantaged borough based on routinely collected education and poverty statistics, had the lowest rates of hospitalisations for COVID-19 relative to the four remaining areas.(Wadhwa et al., 2020) While, by contrast, the Bronx, the least favourable socioeconomically, had the highest level of hospitalisations, rates were graded across the boroughs for education but not poverty. In a recent report from the Office for National statistics in the UK, rates of death in which COVID-19 was implicated were directly related to neighbourhood deprivation in a step-wise manner.(Office for National Statistics, 2020) Outside the eras of pandemics, other respiratory diseases such as tuberculosis,(Cantwell et al., 1998; Spence et al., 1993) pneumonia,(Burton et al., 2010) influenza,(Tam et al., 2014) and, importantly, the common cold (Stone et al., 2010) – also appear to be similarly socioeconomically patterned, although these are not universal observations.(Charland et al., 2011; Vrbova et al., 2005) We are unaware of any studies exploring the relation of indicators of cognitive function and mental health with COVID-19, though up to a doubling in rates of death from respiratory disease has been reported in people with lower

cognition test scores,(Calvin et al., 2017; Gale et al., 2019) individuals with a serious mental illness,(Seminog and Goldacre, 2013) and those with higher levels of psychological distress.(Hamer et al., 2019)

5.3. *Mechanisms of effect*

Specific and non-specific mechanisms may link these psychosocial variables to the risk of COVID-19. A plausible explanation for the association between cognition and respiratory infection is that people with higher ability, and indeed the educationally advantaged,(La et al., 2018) may be more likely to take-up influenza and pneumococcal inoculation; however, in absence of any effective vaccination for COVID-19 this is implausible. In our analyses we took into account unfavourable health behaviours which are more common in lower cognition scoring groups(Batty et al., 2006, 2007a; Batty et al., 2008a; Batty et al., 2007b, c) and have also been implicated in the occurrence of pneumonia,(Baik et al., 2000) but the effect for cognition remained. It may be that the deluge of health advice in the current pandemic during a period when news outlets and social media platforms have never been more ubiquitous, has highlighted that lower cognition and therefore poor health literacy in the population is a public health concern.(Mottus et al., 2014; Reeve and Basalik, 2014) In a small-scale cross-sectional study, people with low health literacy also reported being less concerned about the current pandemic and to believe they were at lower risk.(Wolf et al., 2020)

Mental health problems may influence the risk of acquiring a respiratory infection by negatively impacting cognitive function,(Pedrelli et al., 2004) potentially compromising the ability to effectively take precautions to minimise the risk, adequately recognise a deterioration in health, actively seek medical attention, and communicate effectively with health care professionals. An unhealthy lifestyle and sub-optimal circumstances including poor housing and lower income are also more common in people with mental health problems(Phillips et al., 2009; Russ et al., 2012) but we were careful to covary on these

factors in our analyses. It could also be the case that people experiencing higher levels of psychological distress have diminished learned resistance to infection owing to fewer social interactions, although a pre-pandemic measure of social isolation in our analyses did not confer the expected protection against the infection.

More speculative explanations for some of the effects found herein are that our outcome, hospitalisation for COVID-19 infection, represents not only the occurrence of the infection itself but also a sub-optimal viral-response. People with a higher burden of psychological distress – which includes worry about physical health – might be more concerned about becoming seriously unwell and therefore have a lower symptom severity threshold for visiting hospital. Similarly, individuals with lower cognition may have less confidence in their own decision-making, self-care, and UK government messages to remain at home when unwell, instead resorting to hospital-based advice.

5.4. Study strengths and limitations

The strengths of our study include it being well characterised for exposures and covariates despite its scale, allowing us to attempt to identify independent effects. That the study is prospective means assessment of these baseline data preceded that of disease onset; as such, reverse causality is not a concern such that the infection could not, for instance, influence mental health and job loss leading to downward social mobility. Our work has its weaknesses. Samples were taken from hospitalised patients but it is unclear if all cases had been exclusively hospitalised because of COVID-19-type symptoms, or, as seem likely given mass testing within hospitals, some patients were found to be positive for the infection while an inpatient for other reasons. Our outcome also represents an unfavourable response to a viral challenge as opposed to disease incidence across the full population; the latter could only be ascertained with comprehensive testing of our study sample or indeed the population of England as a whole. We excluded study members who had died prior to 5th March 2020 because they could not

contribute to the risk set, however, ascertainment of hospitalisations for COVID-19 did not reliably begin until 16th March. It is unlikely, however, that the absence of vital status data for this 11 day period would have substantially biased our effect estimates in this large dataset.

The UK Biobank study sample comprises only the 5.5% of the target population who agreed to participate.(Sudlow et al., 2015) As has been demonstrated,(Batty et al., 2020; Fry et al., 2017) the data material is therefore inappropriate for estimation of risk factor or disease prevalence and incidence of COVID-19 infection, and any data simulations of its dissemination. These observations do not, however, seem to influence reproducibility of the association of established risk factors for non-communicable disease such as vascular disease and selected cancers, and other health endpoints such as suicide.(Batty et al., 2020) We think the same reasoning can be applied to associations with communicable diseases.

As with all studies, the characteristics collected at baseline are, with very few exceptions (sex, ethnicity), time-varying. Repeat assessment of selected subgroups gives us the opportunity to explore the stability of the exposure variables in the present analyses. Over a median of 8.18 years between baseline and follow-up in study members participating in an imaging sub-study, Pearson correlation coefficients were high for those socioeconomic factors reassessed (annual income 0.66, $p < 0.001$, $N = 26,322$; educational qualifications 0.86, $p < 0.001$, $N = 30,350$) and somewhat lower for indices of cognitive function (reasoning 0.63, $p < 0.001$, $N = 9689$; reaction time 0.49, $p < 0.001$, $N = 28,810$) and mental health (seen psychiatrist 0.64, $p < 0.001$, $N = 47,291$; psychological distress score 0.52, $p < 0.001$, $N = 42782$). Any instability in test-retest correlations is likely to lead to an underestimated of risk factor–disease associations.(Clarke et al., 1999)

6.0. *Conclusions*

In conclusion, in aetiological-orientated analyses of data from this prospective cohort study, a range of psychosocial factors showed associations with subsequent hospitalisations for COVID-19, among which cognitive function – a potential marker of health literacy – was most robustly related. These findings have important implications for public health messaging, but replication is required before policy recommendations can be advanced.

References

- Adam, Y., Meinlschmidt, G., Lieb, R., 2013. Associations between mental disorders and the common cold in adults: a population-based cross-sectional study. *Journal of psychosomatic research* 74, 69-73.
- Andersson, N.W., Goodwin, R.D., Okkels, N., Gustafsson, L.N., Taha, F., Cole, S.W., Munk-Jorgensen, P., 2016. Depression and the risk of severe infections: prospective analyses on a nationwide representative sample. *International journal of epidemiology* 45, 131-139.
- Baik, I., Curhan, G.C., Rimm, E.B., Bendich, A., Willett, W.C., Fawzi, W.W., 2000. A prospective study of age and lifestyle factors in relation to community-acquired pneumonia in US men and women. *Archives of internal medicine* 160, 3082-3088.
- Batty, G.D., Deary, I.J., Benzeval, M., Der, G., 2010. Does IQ predict cardiovascular disease mortality as strongly as established risk factors? Comparison of effect estimates using the West of Scotland Twenty-07 cohort study. *Eur J Cardiovasc Prev Rehabil* 17, 24-27.
- Batty, G.D., Deary, I.J., MacIntyre, S., 2006. Childhood IQ and life course socioeconomic position in relation to alcohol induced hangovers in adulthood: the Aberdeen children of the 1950s study. *Journal of epidemiology and community health* 60, 872-874.
- Batty, G.D., Deary, I.J., MacIntyre, S., 2007a. Childhood IQ in relation to risk factors for premature mortality in middle-aged persons: the Aberdeen Children of the 1950s study. *Journal of epidemiology and community health* 61, 241-247.
- Batty, G.D., Deary, I.J., Schoon, I., Emslie, C., Hunt, K., Gale, C.R., 2008a. Childhood mental ability and adult alcohol intake and alcohol problems: the 1970 British cohort study. *American journal of public health* 98, 2237-2243.
- Batty, G.D., Deary, I.J., Schoon, I., Gale, C.R., 2007b. Childhood mental ability in relation to food intake and physical activity in adulthood: the 1970 British Cohort Study. *Pediatrics* 119, e38-e45.
- Batty, G.D., Deary, I.J., Schoon, I., Gale, C.R., 2007c. Mental ability across childhood in relation to risk factors for premature mortality in adult life: the 1970 British Cohort Study. *Journal of epidemiology and community health* 61, 997-1003.
- Batty, G.D., Gale, C.R., Kivimäki, M., Deary, I.J., Bell, S., 2020. Comparison of risk factor associations in UK Biobank against representative, general population based studies with conventional response rates: prospective cohort study and individual participant meta-analysis. *BMJ (Clinical research ed.)* 368.
- Batty, G.D., Shipley, M.J., Dundas, R., MacIntyre, S., Der, G., Mortensen, L.H., Deary, I.J., 2009. Does IQ explain socio-economic differentials in total and cardiovascular disease mortality? Comparison with the explanatory power of traditional cardiovascular disease risk factors in the Vietnam Experience Study. *Eur Heart J* 30, 1903-1909.
- Batty, G.D., Shipley, M.J., Mortensen, L.H., Gale, C.R., Deary, I.J., 2008b. IQ in late adolescence/early adulthood, risk factors in middle-age and later coronary heart disease mortality in men: the Vietnam Experience Study. *Eur J Cardiovasc Prev Rehabil* 15, 359-361.
- Bengtsson, T., Dribe, M., Eriksson, B., 2018. Social Class and Excess Mortality in Sweden During the 1918 Influenza Pandemic. *Am J Epidemiol* 187, 2568-2576.

- Burton, D.C., Flannery, B., Bennett, N.M., Farley, M.M., Gershman, K., Harrison, L.H., Lynfield, R., Petit, S., Reingold, A.L., Schaffner, W., Thomas, A., Plikaytis, B.D., Rose, C.E., Jr., Whitney, C.G., Schuchat, A., 2010. Socioeconomic and racial/ethnic disparities in the incidence of bacteremic pneumonia among US adults. *American journal of public health* 100, 1904-1911.
- Calvin, C.M., Batty, G.D., Der, G., Brett, C.E., Taylor, A., Pattie, A., Cukic, I., Deary, I.J., 2017. Childhood intelligence in relation to major causes of death in 68 year follow-up: prospective population study. *BMJ (Clinical research ed.)* 357, j2708.
- Cantwell, M.F., McKenna, M.T., McCray, E., Onorato, I.M., 1998. Tuberculosis and race/ethnicity in the United States: impact of socioeconomic status. *American journal of respiratory and critical care medicine* 157, 1016-1020.
- Charland, K.M., Brownstein, J.S., Verma, A., Brien, S., Buckeridge, D.L., 2011. Socio-economic disparities in the burden of seasonal influenza: the effect of social and material deprivation on rates of influenza infection. *PloS one* 6, e17207.
- Chobanian, A.V., Bakris, G.L., Black, H.R., Cushman, W.C., Green, L.A., Izzo, J.L., Jr., Jones, D.W., Materson, B.J., Oparil, S., Wright, J.T., Jr., Roccella, E.J., 2003. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *Jama* 289, 2560-2572.
- Clarke, R., Shipley, M., Lewington, S., Youngman, L., Collins, R., Marmot, M., Peto, R., 1999. Underestimation of risk associations due to regression dilution in long-term follow-up of prospective studies. *Am. J. Epidemiol* 150, 341-353.
- Cohen, S., Tyrrell, D.A., Smith, A.P., 1991. Psychological stress and susceptibility to the common cold. *The New England journal of medicine* 325, 606-612.
- Cuan-Baltazar, J.Y., Munoz-Perez, M.J., Robledo-Vega, C., Perez-Zepeda, M.F., Soto-Vega, E., 2020. Misinformation of COVID-19 on the Internet: Infodemiology Study. *JMIR public health and surveillance* 6, e18444.
- Davies, G., Marioni, R.E., Liewald, D.C., Hill, W.D., Hagenaars, S.P., Harris, S.E., Ritchie, S.J., Luciano, M., Fawns-Ritchie, C., Lyall, D., Cullen, B., Cox, S.R., Hayward, C., Porteous, D.J., Evans, J., McIntosh, A.M., Gallacher, J., Craddock, N., Pell, J.P., Smith, D.J., Gale, C.R., Deary, I.J., 2016. Genome-wide association study of cognitive functions and educational attainment in UK Biobank (N=112 151). *Molecular psychiatry* 21, 758-767.
- Deary, I., Weiss, A., Batty, G., 2011. Intelligence and personality as predictors of illness and death: How researchers in differential psychology and chronic disease epidemiology are collaborating to understand and address health inequalities. *Psychological Science in the Public Interest* 11, 53-79.
- Deary, I.J., Der, G., Ford, G., 2001. Reaction times and intelligence differences: a population-based cohort study. *Intelligence* 29, 389-399.
- Elliott, P., Peakman, T.C., 2008. The UK Biobank sample handling and storage protocol for the collection, processing and archiving of human blood and urine. *International journal of epidemiology* 37, 234-244.

- Elovainio, M., Hakulinen, C., Pulkki-Raback, L., Virtanen, M., Josefsson, K., Jokela, M., Vahtera, J., Kivimäki, M., 2017. Contribution of risk factors to excess mortality in isolated and lonely individuals: an analysis of data from the UK Biobank cohort study. *Lancet Public Health* 2, e260-e266.
- Fry, A., Littlejohns, T.J., Sudlow, C., Doherty, N., Adamska, L., Sprosen, T., Collins, R., Allen, N.E., 2017. Comparison of Sociodemographic and Health-Related Characteristics of UK Biobank Participants With Those of the General Population. *Am J Epidemiol* 186, 1026-1034.
- Gale, C.R., Deary, I.J., Batty, G.D., 2019. Cognitive ability and risk of death from lower respiratory tract infection: findings from UK Biobank. *Scientific reports* 9, 1342.
- Grantz, K.H., Rane, M.S., Salje, H., Glass, G.E., Schachterle, S.E., Cummings, D.A., 2016. Disparities in influenza mortality and transmission related to sociodemographic factors within Chicago in the pandemic of 1918. *Proceedings of the National Academy of Sciences of the United States of America* 113, 13839-13844.
- Hamer, M., Kivimäki, M., Gale, C.R., David Batty, G., 2020. Lifestyle risk factors, inflammatory mechanisms, and COVID-19 hospitalization: A community-based cohort study of 387,109 adults in UK. *Brain, behavior, and immunity*.
- Hamer, M., Kivimäki, M., Stamatakis, E., Batty, G.D., 2019. Psychological distress and infectious disease mortality in the general population. *Brain, behavior, and immunity* 76, 280-283.
- Jiang, T., Farkas, D.K., Ahern, T.P., Lash, T.L., Sorensen, H.T., Gradus, J.L., 2019. Posttraumatic Stress Disorder and Incident Infections: A Nationwide Cohort Study. *Epidemiology (Cambridge, Mass.)* 30, 911-917.
- La, E.M., Trantham, L., Kurosky, S.K., Odom, D., Aris, E., Hoge, C., 2018. An analysis of factors associated with influenza, pneumococcal, Tdap, and herpes zoster vaccine uptake in the US adult population and corresponding inter-state variability. *Human vaccines & immunotherapeutics* 14, 430-441.
- Lam, M.H., Wing, Y.K., Yu, M.W., Leung, C.M., Ma, R.C., Kong, A.P., So, W.Y., Fong, S.Y., Lam, S.P., 2009. Mental morbidities and chronic fatigue in severe acute respiratory syndrome survivors: long-term follow-up. *Archives of internal medicine* 169, 2142-2147.
- Lassale, C., Gaye, B., Hamer, M., Gale, C.R., David Batty, G., 2020. Ethnic Disparities in Hospitalisation for COVID-19 in England: The Role of Socioeconomic Factors, Mental Health, and Inflammatory and Pro-inflammatory Factors in a Community-based Cohort Study. *Brain, behavior, and immunity*.
- Lee, A.M., Wong, J.G., McAlonan, G.M., Cheung, V., Cheung, C., Sham, P.C., Chu, C.M., Wong, P.C., Tsang, K.W., Chua, S.E., 2007. Stress and psychological distress among SARS survivors 1 year after the outbreak. *Canadian journal of psychiatry. Revue canadienne de psychiatrie* 52, 233-240.
- Mamelund, S.E., 2006. A socially neutral disease? Individual social class, household wealth and mortality from Spanish influenza in two socially contrasting parishes in Kristiania 1918-19. *Social science & medicine* (1982) 62, 923-940.
- Mottus, R., Johnson, W., Murray, C., Wolf, M.S., Starr, J.M., Deary, I.J., 2014. Towards understanding the links between health literacy and physical health. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association* 33, 164-173.

- Neisser, U., Boodoo, G., Bouchard Jr, T., Boykin, A.W., Brody, N., Ceci, S.J., Halpern, D.F., Loehlin, J.C., Perloff, R., Sternberg, R.J., Urbina, S., 1996. Intelligence: knowns and unknowns. *Am Psychol* 51, 77-101.
- Niedzwiedz, C.L., O'Donnell, C.A., Jani, B.D., Demou, E., Ho, F.K., Celis-Morales, C., Nicholl, B.I., Mair, F.S., Welsh, P., Sattar, N., Pell, J.P., Katikireddi, S.V., 2020. Ethnic and socioeconomic differences in SARS-CoV-2 infection: prospective cohort study using UK Biobank. *BMC medicine* 18, 160.
- Office for National Statistics, 2020. Deaths involving COVID-19 by local area and socioeconomic deprivation: deaths occurring between 1 March and 17 April 2020. <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/bulletins/deathsinvolvingcovid19bylocalareasanddeprivation/deathsoccurringbetween1marchand17april> (access 12 May 2020).
- Orso, D., Federici, N., Copetti, R., Vetrugno, L., Bove, T., 2020. Infodemic and the spread of fake news in the COVID-19-era. *European journal of emergency medicine : official journal of the European Society for Emergency Medicine*.
- Pedrelli, P., McQuaid, J.R., Granholm, E., Patterson, T.L., McClure, F., Beck, A.T., Jeste, D.V., 2004. Measuring cognitive insight in middle-aged and older patients with psychotic disorders. *Schizophrenia research* 71, 297-305.
- Phillips, A.C., Batty, G.D., Gale, C.R., Deary, I.J., Osborn, D., MacIntyre, K., Carroll, D., 2009. Generalized anxiety disorder, major depressive disorder, and their comorbidity as predictors of all-cause and cardiovascular mortality: the Vietnam experience study. *Psychosomatic medicine* 71, 395-403.
- Reeve, C.L., Basalik, D., 2014. Is health literacy an example of construct proliferation? A conceptual and empirical evaluation of its redundancy with general cognitive ability. *Intelligence* 44, 93-102.
- Richardson, S., Hirsch, J.S., Narasimhan, M., Crawford, J.M., McGinn, T., Davidson, K.W., Barnaby, D.P., Becker, L.B., Chelico, J.D., Cohen, S.L., Cookingham, J., Coppa, K., Diefenbach, M.A., Dominello, A.J., Duer-Hefele, J., Falzon, L., Gitlin, J., Hajizadeh, N., Harvin, T.G., Hirschwerk, D.A., Kim, E.J., Kozel, Z.M., Marrast, L.M., Mogavero, J.N., Osorio, G.A., Qiu, M., Zanos, T.P., 2020. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *Jama*.
- Russ, T.C., Stamatakis, E., Hamer, M., Starr, J.M., Kivimaki, M., Batty, G.D., 2012. Association between psychological distress and mortality: individual participant pooled analysis of 10 prospective cohort studies. *BMJ (Clinical research ed.)* 345, e4933.
- Rutter, P.D., Mytton, O.T., Mak, M., Donaldson, L.J., 2012. Socio-economic disparities in mortality due to pandemic influenza in England. *International journal of public health* 57, 745-750.
- Seminog, O.O., Goldacre, M.J., 2013. Risk of pneumonia and pneumococcal disease in people with severe mental illness: English record linkage studies. *Thorax* 68, 171-176.
- Simonsen, L., Spreeuwenberg, P., Lustig, R., Taylor, R.J., Fleming, D.M., Kroneman, M., Van Kerkhove, M.D., Mounts, A.W., Paget, W.J., 2013. Global mortality estimates for the 2009 Influenza Pandemic from the GLaMOR project: a modeling study. *PLoS medicine* 10, e1001558.

- Spence, D.P., Hotchkiss, J., Williams, C.S., Davies, P.D., 1993. Tuberculosis and poverty. *BMJ (Clinical research ed.)* 307, 759-761.
- Stone, A.A., Krueger, A.B., Steptoe, A., Harter, J.K., 2010. The socioeconomic gradient in daily colds and influenza, headaches, and pain. *Archives of internal medicine* 170, 570-572.
- Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., Landray, M., Liu, B., Matthews, P., Ong, G., Pell, J., Silman, A., Young, A., Sprosen, T., Peakman, T., Collins, R., 2015. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS medicine* 12, e1001779.
- Tam, K., Yousey-Hindes, K., Hadler, J.L., 2014. Influenza-related hospitalization of adults associated with low census tract socioeconomic status and female sex in New Haven County, Connecticut, 2007-2011. *Influenza and other respiratory viruses* 8, 274-281.
- UK Biobank, 2020. Records of COVID-19 test results in UK Biobank. <http://biobank.ndph.ox.ac.uk/showcase/field.cgi?id=40100> (access 12 May 2020).
- UK Government, 2020. Number of coronavirus (COVID-19) cases and risk in the UK. <https://web.archive.org/web/20200512082155/https://www.gov.uk/guidance/coronavirus-covid-19-information-for-the-public> (accessed 12 May 2020).
- Vrbova, L., Mamdani, M., Moineddin, R., Jaakimainen, L., Upshur, R.E., 2005. Does socioeconomic status affect mortality subsequent to hospital admission for community acquired pneumonia among older persons? *Journal of negative results in biomedicine* 4, 4.
- Wadhwa, R.K., Wadhwa, P., Gaba, P., Figueroa, J.F., Joynt Maddox, K.E., Yeh, R.W., Shen, C., 2020. Variation in COVID-19 Hospitalizations and Deaths Across New York City Boroughs. *Jama*.
- Wiemken, T.L., Carrico, R.M., Furmanek, S.P., Guinn, B.E., Mattingly, W.A., Peyrani, P., Ramirez, J.A., 2020. Socioeconomic Position and the Incidence, Severity, and Clinical Outcomes of Hospitalized Patients With Community-Acquired Pneumonia. *Public health reports (Washington, D.C. : 1974)* 135, 364-371.
- Wolf, M.S., Serper, M., Opsasnick, L., O'Connor, R.M., Curtis, L.M., Benavente, J.Y., Wismer, G., Batio, S., Eifler, M., Zheng, P., Russell, A., Arvanitis, M., Ladner, D., Kwasny, M., Persell, S.D., Rowe, T., Linder, J.A., Bailey, S.C., 2020. Awareness, Attitudes, and Actions Related to COVID-19 Among Adults With Chronic Conditions at the Onset of the U.S. Outbreak: A Cross-sectional Survey. *Annals of internal medicine*.
- World Health Organization, 2020. Coronavirus disease (COVID-19) Situation Report – 112. Data as received by WHO from national authorities by 10:00 CEST, 11 May 2020 (accessed 12 May 2020). https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200511-covid-19-sitrep-112.pdf?sfvrsn=813f2669_2.
- World Health Organization Director-General, 2020. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. <https://web.archive.org/web/20200502133342/https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020> (accessed 12 May 2020).

Wu, C., Chen, X., Cai, Y., Xia, J., Zhou, X., Xu, S., Huang, H., Zhang, L., Zhou, X., Du, C., Zhang, Y., Song, J., Wang, S., Chao, Y., Yang, Z., Xu, J., Zhou, X., Chen, D., Xiong, W., Xu, L., Zhou, F., Jiang, J., Bai, C., Zheng, J., Song, Y., 2020. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA internal medicine*.

Yap, A.G., Baladi, N., Allman, G., Avenmarg, J., Yap, S., Shaw, R.E., 2000. Coronary artery bypass surgery on small patients. *J. Invasive. Cardiol* 12, 242-246.

Zarocostas, J., 2020. How to fight an infodemic. *Lancet (London, England)* 395, 676.

Table 1. Psychosocial factors and covariates at baseline according to hospitalisations for COVID-19

	COVID-19 hospitalisation at follow-up		P value	Unadjusted OR (95% CI) ^a
	Yes (n=908)	No (n=430,143)		
<i>Demographic factors</i>				
Age, yr, mean (SD)	57.27 (8.99)	56.36 (8.10)	0.0007	1.12 (1.05, 1.20)
Female, N (%)	402 (44.27)	236,323 (54.94)	<0.0001	0.65 (0.57, 0.74)
Non-white ethnicity	128 (14.22)	25,170 (5.89)	<0.0001	2.65 (2.20, 3.20)
Socially isolated, N (%)	90 (9.91)	38,353 (8.92)	0.293	1.12 (0.90, 1.40)
<i>Comorbidities</i>				
Vascular or heart disease, N (%)	383 (42.7)	124,306 (29.03)	<0.0001	1.82 (1.60, 2.08)
Hypertension, N (%)	566 (64.61)	238,111 (56.35)	<0.0001	1.44 (1.22, 1.69)
Diabetes, N (%)	90 (10.03)	21,316 (4.98)	<0.0001	2.13 (1.71, 2.64)
Chronic bronchitis or emphysema, N (%)	28 (3.08)	6,311 (1.47)	<0.0001	2.14 (1.47, 3.11)
Asthma, N (%)	126 (13.88)	49,600 (11.53)	0.027	1.24 (1.02, 1.49)
Cancer, N (%)	75 (8.42)	31,051 (7.26)	0.183	1.17 (0.93, 1.49)
<i>Lifestyle factors</i>				
Current smoker, N (%)	102 (11.37)	42,636 (9.97)	<0.0001	1.35 (1.09, 1.68)
No physical activity, N (%)	117 (13.31)	26,096 (6.16)	<0.0001	2.34 (1.93, 2.84)
Drinks alcohol daily/almost daily, N (%)	148 (16.41)	87,754 (20.46)	0.003	0.76 (0.64, 0.91)
Body mass index, kg/m ² , mean (SD)	29.2 (5.45)	27.4 (4.77)	<0.0001	1.37 (1.30, 1.44)
<i>Biomarkers</i>				
Lung function, L, mean (SD)	2.69 (0.82)	2.82 (0.80)	<0.0001	0.84 (0.78, 0.90)
Hand grip strength, kg, mean (SD)	32.5 (11.0)	32.5 (11.3)	0.970	1.00 (0.94, 1.07)
C-reactive protein, mg/L, median (IQR)	1.63 (0.84-3.04)	1.24 (0.63-2.44)	0.0001	1.29 (1.20, 1.39)
High-density lipoprotein, mmol/L, median (IQR)	1.33 (1.11-1.56)	1.43 (1.20-1.71)	0.0009	0.69 (0.63, 0.76)
HbA1C, mmol/mol, median (IQR)	35.6 (33.3-38.2)	35.0 (32.6-37.4)	0.0001	1.29 (1.20, 1.40)
<i>Psychological factors</i>				
Psychological distress score ≥ 3 , N (%)	224 (28.64)	90,981 (23.69)	0.001	1.29 (1.11, 1.51)
Psychiatric consultation, N (%)	140 (15.71)	48,599 (11.38)	<0.0001	1.45 (1.21, 1.74)
Neuroticism, mean (SD)	4.37 (3.39)	4.27 (3.28)	0.371	1.03 (0.96, 1.10)
Reasoning, mean (SD)	5.24 (2.08)	6.03 (2.16)	<0.0001	1.47 (1.32, 1.63)
Reaction time, msec, mean (SD)	574.91 (134.88)	558.83 (117.73)	<0.0001	1.13 (1.06, 1.20)
<i>Socioeconomic factors</i>				
No university education, N (%)	647 (73.86)	283,693 (67.36)	<0.0001	1.37 (1.18, 1.59)
Annual household income <£18,000, N (%)	241 (33.1)	80,966 (22.27)	<0.0001	1.73 (1.48, 2.01)
Neighbourhood deprivation score	-0.08 (3.53)	-1.32 (3.06)	<0.0001	1.43 (1.35, 1.51)
Personal service, sales occupations etc, N (%)	149 (26.28)	58,766 (19.07)	<0.0001	1.51 (1.25, 1.82)

^aOdds ratios are expressed per category, or per SD, increase for continuous variables except for reasoning which is expressed per SD decrease. The maximum analytical sample of 431051 people was lower in selected analyses owing to missing data.

Table 2. Odds ratios (95% CI) for the relation of socioeconomic factors with COVID-19 hospitalisation

	Case N /Risk N ¹	Adjustments					
		Age & sex	Age, sex & ethnicity	Age, sex, ethnicity & comorbidity ²	Age, sex, ethnicity & lifestyle factors ³	Age, sex, ethnicity & psychological factors ⁴	All covariates
Educational attainment		N=422057	N=420502	N=415945	N=415367	N=155244	N=152739
University degree	229/137717	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Other qualifications	406/214337	1.16 (0.98, 1.36)	1.19 (1.01, 1.41)	1.16 (0.98, 1.37)	1.03 (0.87, 1.22)	1.06 (0.81, 1.39)	0.94 (0.71, 1.24)
No qualifications	241/70003	2.05 (1.70, 2.47)	2.07 (1.71, 2.50)	1.85 (1.53, 2.25)	1.47 (1.20, 1.80)	1.35 (0.93, 1.95)	1.01 (0.68, 1.49)
P for trend		<0.0001	<0.0001	<0.0001	<0.0001	0.151	0.945
Annual household income		N=364219	N=363175	N=359853	N=359491	N=137808	N=135773
<£18,000	241/81207	2.00 (1.63, 2.47)	1.89 (1.51, 2.35)	1.74 (1.39, 2.17)	1.39 (1.10, 1.75)	1.34 (0.91, 1.97)	1.15 (0.77, 1.73)
18,000-30,999	179/92461	1.31 (1.05, 1.63)	1.27 (1.01, 1.60)	1.22 (0.97, 1.54)	1.05 (0.83, 1.32)	1.29 (0.90, 1.85)	1.15 (0.79, 1.68)
31,000-51,999	167/95454	1.18 (0.94, 1.48)	1.17 (0.94, 1.47)	1.16 (0.92, 1.45)	1.07 (0.85, 1.34)	1.03 (0.72, 1.49)	1.02 (0.70, 1.48)
≥52,000	141/95097	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
P for trend		<0.0001	<0.0001	<0.0001	0.006	0.077	0.401
Neighbourhood deprivation		N=430538	N=427986	N=419593	N=418942	N=156360	N=153384
1 (low)	205/143483	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
2	267/143548	1.32 (1.10, 1.58)	1.29 (1.07, 1.55)	1.25 (1.04, 1.50)	1.20 (1.00, 1.45)	1.32 (0.97, 1.79)	1.22 (0.89, 1.65)
3	436/143517	2.20 (1.86, 2.59)	1.97 (1.66, 2.34)	1.79 (1.51, 2.13)	1.57 (1.31, 1.88)	1.52 (1.12, 2.05)	1.20 (0.87, 1.63)
P for trend		<0.0001	<0.0001	<0.0001	<0.0001	0.007	0.297
Occupational classification		N=308689	N=307262	N=302239	N=302495	N=130238	N=128079
Managers, senior officials, etc	324/175637	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Administrative, secretarial, etc	94/74137	0.70 (0.55, 0.88)	0.69 (0.55, 0.87)	0.68 (0.54, 0.86)	0.63 (0.50, 0.80)	0.65 (0.46, 0.92)	0.59 (0.42, 0.84)
Personal service, sales, etc	149/58915	1.39 (1.14, 1.69)	1.30 (1.07, 1.59)	1.22 (1.00, 1.49)	1.06 (0.86, 1.30)	0.90 (0.65, 1.26)	0.75 (0.53, 1.06)
P for trend		0.024	0.091	0.314	0.780	0.242	0.027


¹ Numbers based on unadjusted model. ² Comorbidity includes diagnoses of vascular or heart disease, diabetes, chronic bronchitis or emphysema, asthma, and hypertension defined according to measured blood pressure and/or use of anti-hypertensive medication. ³ Lifestyle factors includes body mass index, smoking status, alcohol intake frequency & number of types of physical activity taken in last four weeks. ⁴ Psychological factors include psychological distress, psychiatric consultation, neuroticism, verbal and numerical reasoning, & reaction time. Categories of neighbourhood deprivation are based on tertiles.

Table 3. Odds ratios (95% CI) for the relation of psychological factors with COVID-19 hospitalisation

	Case N / Risk N ¹	Adjustments					
		Age & sex	Age, sex & ethnicity	Age, sex, ethnicity & comorbidity ²	Age, sex, ethnicity & lifestyle factors ³	Age, sex, ethnicity & socioeconomic factors ⁴	All covariates
Psychological distress		N=384909	N=383655	N=377290	N=376562	N=248162	N=245119
1 (low)	267/153504	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
2	291/140200	1.28 (1.08, 1.51)	1.29 (1.09, 1.53)	1.22 (1.03, 1.45)	1.16 (0.98, 1.38)	1.16 (0.93, 1.45)	1.07 (0.86, 1.35)
3	224/91205	1.58 (1.32, 1.89)	1.51 (1.26, 1.81)	1.37 (1.14, 1.65)	1.18 (0.98, 1.43)	1.26 (0.98, 1.61)	1.09 (0.84, 1.41)
P for trend		<0.0001	<0.0001	0.001	0.068	0.064	0.487
Per SD increase		1.22 (1.14, 1.29)	1.19 (1.12, 1.26)	1.15 (1.08, 1.23)	1.09 (1.02, 1.17)	1.12 (1.02, 1.22)	1.07 (0.97, 1.17)
Psychiatric consultation		N=427819	N=426823	N=418218	N=417481	N=269373	N=265566
No	751/379080	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
Yes	140/487739	1.50 (1.25, 1.79)	1.51 (1.26, 1.81)	1.45 (1.21, 1.75)	1.32 (1.09, 1.59)	1.23 (0.94, 1.62)	1.15 (0.87, 1.52)
Neuroticism		N=425707	N=424212	N=416378	N=415622	N=265538	N=264784
1 (low)	224/106910	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
2	345/174705	1.01 (0.85, 1.19)	1.03 (0.87, 1.22)	0.99 (0.83, 1.17)	0.99 (0.83, 1.18)	1.03 (0.82, 1.30)	1.01 (0.80, 1.27)
3	319/144092	1.19 (1.00, 1.42)	1.21 (1.02, 1.44)	1.10 (0.92, 1.31)	1.08 (0.90, 1.29)	1.06 (0.84, 1.35)	1.00 (0.78, 1.28)
P for trend		0.038	0.023	0.277	0.382	0.621	0.985
Per SD increase		1.08 (1.01, 1.15)	1.08 (1.01, 1.16)	1.05 (0.98, 1.12)	1.03 (0.96, 1.10)	1.00 (0.92, 1.10)	0.99 (0.90, 1.09)
Verbal numerical reasoning		N=175267	N=174581	N=172530	N=415777	N=126721	N=124890
1 (low)	152/43988	2.66 (2.06, 3.34)	2.31 (1.77, 3.02)	2.17 (1.65, 2.86)	1.92 (1.45, 2.53)	2.14 (1.50, 3.05)	1.98 (1.38, 2.85)
2	115/58446	1.52 (1.16, 1.99)	1.45 (1.10, 1.90)	1.46 (1.10, 1.92)	1.36 (1.03, 1.80)	1.57 (1.14, 2.17)	1.58 (1.14, 2.18)
3	96/72833	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
P for trend		<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001
Per SD decrease		1.47 (1.32, 1.64)	1.37 (1.23, 1.53)	1.33 (1.19, 1.49)	1.27 (1.13, 1.42)	1.35 (1.17, 1.57)	1.31 (1.13, 1.52)
Reaction time		N=426147	N=424432	N=416366	N=415777	N=268826	N=265002
1 (low)	262/140934	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)	1.0 (ref)
2	274/141575	1.04 (0.87, 1.23)	1.00 (0.84, 1.19)	0.97 (0.82, 1.16)	0.95 (0.79, 1.13)	1.02 (0.82, 1.27)	1.02 (0.81, 1.27)
3	345/143368	1.27 (1.08, 1.51)	1.16 (0.98, 1.37)	1.11 (0.93, 1.32)	1.04 (0.88, 1.24)	1.06 (0.84, 1.34)	1.02 (0.80, 1.29)
P for trend		0.004	0.078	0.205	0.572	0.608	0.876
Per SD increase		1.12 (1.06, 1.19)	1.07(1.01, 1.14)	1.06 (0.99, 1.13)	1.03 (0.97, 1.10)	1.08 (0.98, 1.18)	1.07 (0.97, 1.17)


¹Numbers based on age & sex adjusted model. ² Comorbidity includes diagnoses of vascular or heart disease, diabetes, chronic bronchitis or emphysema, asthma, and hypertension defined according to measured blood pressure and/or use of anti-hypertensive drugs. ³Lifestyle factors included body mass index, smoking status, alcohol intake frequency & number of types of physical activity taken in last four weeks. ⁴Socioeconomic factors included occupational classification, educational attainment, Townsend deprivation index, & household income before tax. With the exception of psychiatric consultation, categories are based on tertiles.

Figure 1. Odds ratios (95% CI) for the relation of socioeconomic factors with COVID-19 hospitalisation



Adjustments are as per table 2.

Figure 2. Odds ratios (95% CI) for the relation of psychological factors with COVID-19 hospitalisation



Adjustments are as per table 2

Highlights

- Socioeconomic differentials in COVID-19 have been reported but their origin is uncertain.
- Psychological factors have an unknown association with this infection.
- In a community-based cohort – UK Biobank – socioeconomic disadvantage and indices of mental health, lower cognitive function, and personality type (neuroticism) were associated an elevated risk of hospitalisation for COVID-19.
- The most robust relationship was that for cognitive function, a marker of health literacy.