

TITLE PAGE

Title

Combining evidence based healthcare with environmental sustainability: using the toothbrush as a model.

Abstract word count: 199

Word count (excl abstract, acknowledgements, references, tables): 2,652

Total number of figures: 3

Total number of tables: 2

Total number of references: 23

ABSTRACT

Introduction Healthcare professionals should consider environmental sustainability when recommending medical devices to patients, although there is currently little quantitative data available. The toothbrush is a widely recommended healthcare device worldwide. The aim of this study was to compare the sustainability of different types of toothbrush.

Materials and Methods Four types of toothbrush were studied: a traditional plastic and electric toothbrush, as well as a plastic manual toothbrush with replaceable heads, and a bamboo manual toothbrush. Life cycle assessment (LCA) methodology was applied to quantify the environmental impact of these toothbrushes over 5 years.

Results The electric toothbrush performed consistently poorly compared to the 3 manual toothbrush types, and had the greatest impact in 15 out of 16 environmental categories. The bamboo and replaceable head plastic toothbrushes had the lowest impact in all categories. The climate change potential of the electric toothbrush was 11 times greater than the bamboo toothbrush.

Discussion Switching toothbrush, from the traditional toothbrushes to bamboo or replaceable head plastic, is more environmentally sustainable. These results could be used to inform individual consumer choice, oral health recommendations, procurement of toothbrushes for public health programmes, and toothbrush manufacturers. LCA methodology can be used to make healthcare more sustainable.

INTRODUCTION

Environmental sustainability is a worldwide public health issue.¹ The planet and its global population face a range of challenges, including climate change, reduction in biodiversity, air and water pollution, and ozone depletion. Global healthcare is a significant contributor to national carbon dioxide emissions – and on average is responsible for 5% of national carbon dioxide emissions.² In England, the National Health Service (NHS) is responsible for 25% of England's public sector carbon footprint.³ Established to improve population health, healthcare systems are harming the planet.

As an 'anchor organisation' in the UK, the NHS aims to be a sustainable healthcare system.⁴ The NHS long term plan commits to 3 environmental aims: to reduce air pollution, waste, and greenhouse gas emissions.⁵ In England, this is driven by the Sustainable Development Unit (SDU), which was established to embed sustainable development at all levels of healthcare.⁶ One measure of success is that '*professionals are encouraged to consider sustainability principles when deciding what is right for patients and clients*'. For NHS professionals to understand the environmental impact of a product or service, different methods can be used. These range from carbon footprint analysis, to more detailed methods which look at wider environmental impacts.

Life cycle assessment (LCA) is used to measure the environmental impact of different services or products.^{7,8} Also referred to as a cradle-to-grave analysis, LCA considers all aspects of a product along its life cycle, including raw materials, manufacture, use, transport, and disposal. The NHS, along with other healthcare companies, established the Coalition of Sustainable Pharmaceuticals and Medical Devices (CSPM), which recommends LCA to compare services and enable policy makers to make informed recommendations.⁹ More recently, the European Union adopted the Product Environment Framework (PEF) to provide a consistent, standardised, comparable approach to undertaking LCA.¹⁰

There is currently little evidence or guidance regarding the sustainability of specific healthcare interventions, services or devices. This includes both evidence-based interventions within a healthcare setting (such as a hospital), and those carried out within the home setting. In this paper, we have selected a commonly prescribed

intervention (the toothbrush) to explore the impact of this preventative device on the environment. Its efficacy as an intervention to prevent oral disease is well established.¹¹ There are several different types of toothbrush available in the UK, with different sustainability 'credentials'. Although there is evidence that electric toothbrushes are associated with a greater level of plaque and gingivitis reduction compared to manual toothbrushes, there is no evidence that any type of toothbrush is more clinically effective for the prevention of dental caries and periodontal disease.¹² Therefore, potentially the environmental impact could be the prime consideration for NHS providers when selecting or recommending a product. The national Scottish oral health programme, Childsmile, has pledged to include sustainability as part of the product specification when procuring toothbrushes.¹³

This paper uses LCA methodology to quantify the environmental impact of perhaps the most used healthcare device worldwide: the toothbrush. The aim was to compare the sustainability of different types of toothbrush, and identify which aspects of the life cycle have the greatest environmental impact.

MATERIALS AND METHODS

A comparative life cycle assessment of four different types of toothbrushes was undertaken at the Eastman Dental Hospital, London, in partnership with the Dublin Dental University Hospital (Trinity College Dublin, Ireland).

The software OpenLCA v1.8 was used for the life cycle assessment, alongside the reference database Ecoinvent v3.5. The LCA methodology was applied in line with ISO standards and PEF guidelines.^{8,9,10}

The four types of toothbrush were:

1. Plastic manual: plastic handle with fixed head.
2. Bamboo manual: bamboo handle with fixed head.
3. Plastic manual replaceable head: reusable plastic handle (made from a bio-plastic) with replaceable heads.
4. Electric: handle and charging unit, with replaceable heads.

Four individual products, available in the UK, were selected to represent each type of toothbrush. The specific brands and manufacturers have been anonymised.

An attributional LCA was conducted from cradle to grave, using physical allocation by mass. The functional unit was defined as individual toothbrush use over 5 years. The time period of 5 years was chosen as this is the average life span of the battery in an electric toothbrush.¹⁴

The system boundaries are shown in Figure 1. The entire product system, including geographical location, was compared, as only the bamboo toothbrush was manufactured outside of Europe (bamboo was cultivated and manufactured in China).

Environmental data for raw bamboo, used to form the handle of the bamboo manual toothbrush, was not available. Therefore, inputs from cultivation of raw bamboo in China were estimated by the consultancy firm GreenDelta (GmbH, 2019) and authors (SS and MC). All assumptions made in this study are listed in Table 1.

A life cycle inventory was created for each type of toothbrush. A sample of each product was dismantled to identify and weigh the component materials. Manufacturers were contacted to clarify any materials, manufacturing and packaging process, and the transport route. The number of products needed over 5 years was calculated. For machinery not available in the database, the energy consumption (kWh) of the machine was used.

Data from the life cycle inventory was modelled in OpenLCA v1.8 for the life cycle impact assessment (LCIA). The impact categories and LCIA methods are shown in Table 2. The LCIA method for each category was selected based on the PEF Category Rules Guidance.¹⁰

RESULTS

The life cycle inventory for each product is available as an appendix.

The results of the LCIA is shown in Figure 2. The electric toothbrush had the greatest environmental impact in all categories, except water scarcity. The plastic manual replaceable head and bamboo manual toothbrushes had the lowest environment impact in 11 and 5 of the impact categories respectively.

Figure 3 demonstrates how each life cycle stage contributed to the impact assessment. For the bamboo manual toothbrush, the biggest contributing factor was consumer use (the tap water used during brushing). The materials contributed the most in both the manual plastic toothbrush and plastic manual replaceable head toothbrush. The material polypropylene, used to the make the plastic handle, was the single biggest contributing factor in both these toothbrushes (37% and 33% respectively). The electric toothbrush was the heaviest product at 1.42kg, had the greatest contributor to it's overall environmental impact was the transport (47%), followed closely by the materials (46%). All other aspects of the electric toothbrush had much less contribution to it's overall impact, including the consumer energy use from charging the handle (0.69%), and disposal (0.16%).

The disposal of the products had the smallest contribution to the environmental impact for all toothbrushes.

DISCUSSION

This study found that both the bamboo manual and plastic manual replaceable-head toothbrushes performed consistently better than the plastic manual and electric toothbrushes, in all impact categories. The sustainability of the electric toothbrush was poor, having the greatest environmental impact in all but one category (water scarcity). The climate change impact of the electric toothbrush was over 11 times greater than the bamboo toothbrush. When considering land use, and the consequential reduction in biodiversity and habitat, the negative impact of the electric toothbrush was over 36 times that of the bamboo toothbrush.

There is increasing public concern about the use of plastics and this alone may be the most important environmental consideration for individual consumers.^{19,20} All products in this study used plastic to make the toothbrush bristles (Nylon), and all except the bamboo product also used plastic for the toothbrush handles (polypropylene) and as part of the packaging (polyethylene). The bamboo toothbrush used just 11g of plastic over the 5 years, the lowest of all products (97% less plastic than the plastic manual toothbrush). The polypropylene in the handle of both the traditional plastic manual and the plastic manual replaceable head toothbrushes had the greatest contribution to the overall environmental impact. The replaceable heads did use a bio-plastic, with 30% of the polymer derived from starch, but the effect of this was unclear, and our results suggest that the lower weight of plastic, from only replacing the head and not the handle, had a greater impact. Further research to identify the 'ideal' sustainable toothbrush could investigate the exact impact of switching polypropylene for biopolymers. If the average life expectancy in the UK is 80 years, then an individual using plastic manual toothbrushes over their lifetime equates to 6.3kg of plastic. A decrease in public demand for plastic and electric toothbrushes may in turn encourage manufacturers to use LCA to improve the environmental impact of their products and use more sustainable materials and processes.

There are limitations to using LCA to compare different healthcare products. There is a range of impact categories and allocation methods that can be used, along with different methods for the LCIA, and this can make the results difficult to interpret. Although guidance by PEF aims to standardise the methodology, it advises that the toxicity-related results are interpreted with caution, as the corresponding three LCIA methods are still in development.¹⁰ However in this analysis, clear differences between the two manual toothbrushes and the electric toothbrush were seen.

In this study, four individual toothbrush products were selected to represent each type of toothbrush; however, the market is constantly changing. In particular, electric toothbrushes have a wide variability in design, and continually updating their features and composition. In this study, the simplest rechargeable electric toothbrush from a market leading brand was selected, but is not necessarily representative of all electric toothbrushes. Since this study was commenced, new toothbrush materials have come to market, including reusable handles made from aluminium. As the market evolves and manufacturers change their materials, LCA should be repeated and recommendations reviewed.

LCA is usually conducted 'in-house', by the manufacturer, which was not the case in this study. In order to correctly identify all the correct product materials and processes, the authors had to request the relevant data from the manufacturers. Where it was not possible to confirm an exact material or process, or the manufacturer was unwilling to supply the information, assumptions were made by the authors, based on industry knowledge. This would have affected the accuracy of the LCA inventory. Ideally, a sensitivity analysis of the most impactful processes and materials would have been carried out, to identify what changes in the material and manufacturing processes could be altered to improve the environmental impact of the product. However, this requires in-depth knowledge and data disclosure, and the authors feel that the responsibility to analyse this and make changes accordingly lies with the product manufacturer.

Including a bamboo product was challenging, as there was no available data for this raw material in the reference database. Therefore the background processes to cultivate bamboo had to be separately modelled by GreenDelta and authors (SS and MC). Several assumptions and exclusions (Table 1) had to be made in order to

produce the dataset, and the effect these assumptions had on the results would require further research and sensitivity analysis. However, bamboo cultivation practices can vary widely, altering the environmental impact of using bamboo as a product material. For example, fertilizers are used in less than 5% of industrial bamboo plantations as the fallen bamboo leaves provide sufficient nutrients for new shoots.¹⁷ As a conservative estimate, in this study we assumed yearly fertilizer application. Ideally, a robust dataset produced with the specific bamboo plantation used by the manufacturer would have been created, but this was beyond the scope of this study. Bamboo cultivation is currently assumed to be carbon neutral, as bamboo ecosystems are carbon sinks, but an increasing demand for this material may in future lead to modifications in the bamboo ecosystem and bamboo cultivation could even become a carbon source.²¹ It was confirmed that no glue is used in the manufacture of the bamboo toothbrush (the handle is made from shaping raw bamboo and heat treating the surface to sterilise), as concerns have been raised regarding the melamine resins used in the production of bamboo products such as reusable cups.²²

Some of the assumptions made about consumer use and disposal are likely to be unrealistic. Individual use of a toothbrush was based on clinical recommendations^{11,18} and informed the number of products needed in 5 years, as well as the waste water used during toothbrushing. There is no data available on public compliance with these oral health recommendations, although studies of health recommendations, such as physical activity, suggest that public compliance is moderate at best.²³ For the electric toothbrush, it was assumed that the handle is only used by one individual, and was disposed of after 5 years.¹⁴ However, families may share one electric toothbrush handle, and may upgrade the handle and charging unit more often or less often than every 5 years. There is currently no data available about the sharing habits of electric toothbrush users. This LCA further assumed that individuals would dispose of their products according to manufacturer's recommendations – for the bamboo toothbrush this includes removing the bristles and metal staples from the bamboo handle. This was undertaken by one of the authors (AL) as part of the product inventory; and as it took almost 30 minutes to remove all the bristles and metal staples using tweezers, it was considered that is unrealistic to expect from consumers. However, the authors felt

that using the manufacturers recommendations was reasonable, given that the disposal processes in the LCIA had little contribution to the overall impact for all the toothbrushes in this study. However, a sensitivity analysis using different disposal scenarios, including bamboo ending up in the ocean, could affect the results and is a topic for further research.

Furthermore, some toothbrush manufacturers have started offering recycling schemes for their products, which could reduce the impact of their materials. However, at the time of this study, the products in this trial were not offering a recycling scheme, and the exact procedure used in any recycling scheme should be specifically analysed by the individual manufacturer, to ensure that the impact created from the transport and recycling processes don't outweigh the benefits of reusing the materials.

Other oral health cleaning aids, such as interdental brushes and floss, will also have an environmental impact and would be subject to a separate life cycle analysis, given their different recommendations and disposal. Research into this is already underway at the author's institutions (UCL and Trinity College Dublin).

The NHS should recommend healthcare devices that are clinically effective, cost effective, and environmentally sustainable. There is currently no evidence that using an electric toothbrush reduces incidence of dental caries or periodontal disease, even if it is better for reducing plaque levels¹². For this reason, there is currently no evidence that individuals switching to the more sustainable manual toothbrushes from this study will develop more dental disease, which could in turn increase the environmental impact of providing dental care. However, should new evidence emerge, the clinical benefit of one type of toothbrush should be considered together with sustainability and cost. Electric toothbrushes are more expensive, and less environmentally sustainable, than the manual toothbrushes. This should be a strong consideration when recommending toothbrushes to the public. Similar principles should apply to toothbrushes procured by the NHS for public oral health programmes, and based on this study; either bamboo toothbrushes or replaceable head manual toothbrushes should be considered over traditional plastic and electric toothbrushes. Admittedly, the cost of bamboo and replaceable head manual toothbrushes, which are usually greater than simple plastic manual toothbrushes,

may present a barrier to their widespread use by consumers, and by public oral health improvement programmes.

This simple comparative LCA has shown that a plastic manual replaceable head toothbrush and bamboo manual toothbrush perform better than traditional plastic manual and electric toothbrushes in every environmental impact outcome measure used in this study. These results could be used to inform individual consumer choice, oral health recommendations, procurement of toothbrushes for public health programmes, and toothbrush manufacturers. Using LCA to inform healthcare policies and recommendations will help move the NHS towards a more environmentally sustainable system.

DECLARATION OF INTERESTS

This study was funded by the Eastman Dental Institute (University College London). The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

The authors would like to thank the manufacturers of the toothbrush products who helped clarify the processes and product materials. This study was funded by University College London (Eastman Dental Institute). All authors declare no conflict of interest. Author contributions: AL collected the data and drafted the paper; PA co-initiated the collaborative project, monitored data collection and revised the draft paper; SS and MC refined the bamboo cultivation dataset, monitored data analysis and revised the paper; BU revised the paper; BD co-initiated the collaborative project, carried out all data analysis and revised the paper. All authors give their final approval and agree to be accountable for all aspects of the work.

REFERENCES

1. Costello A, Abbas M, Allen A et al. Managing the health effects of climate change. *Lancet*, 2009;**373**:1693-1733.

2. Pichler PP, Jaccard IS, Weisz U, Weisz H. International comparison of health care carbon footprints. *Environ Res Lett* 2019;14:064004.
3. Sustainable Development Unit. Carbon footprint update for NHS in England 2015 [Internet]. Cambridge: SDU; 2016 [cited 2019 Nov 18]. 8 p. Available from: <https://www.sduhealth.org.uk/policy-strategy/reporting/nhs-carbon-footprint.aspx>
4. The Health Foundation. The NHS as an anchor institution [Internet]. London: The Health Foundation; 2019 [cited 2019 Nov 18]. Available from: <https://www.health.org.uk/news-and-comment/charts-and-infographics/the-nhs-as-an-anchor-institution>
5. National Health Service. The NHS Long Term Plan [Internet]. London: NHS; 2019 [cited 2019 Nov 18]. Available from: <https://www.longtermplan.nhs.uk/publication/nhs-long-term-plan/>
6. Sustainable Development Unit. What we do [Internet]. Cambridge: SDU; 2019 [cited 2019 Nov 18]. Available from: <https://www.sduhealth.org.uk/about-us/what-we-do.aspx>
7. Sustainable Development Unit. Sustainable clinical and care models [Internet]. London: SDU; 2014 [cited 2019 Nov 18]. Available from: <https://www.sduhealth.org.uk/areas-of-focus/clinical-and-care-models.aspx>
8. International Standards Organisation. ISO 14001 – Environmental management systems – Requirements with guidance for use [Internet]. 2015 [cited 2019 Nov 18]. Available from: <https://www.iso.org/standard/60857.html>
9. Sustainable Development Unit. Coalition for Sustainable Pharmaceuticals and Medical Devices [Internet]. London: SDU; 2019 [cited 2019 Nov 18]. Available from: <https://www.sduhealth.org.uk/areas-of-focus/carbon-hotspots/pharmaceuticals/cspm.aspx>
10. European Commission Joint Research Centre. Product environmental footprint category rules guidance v6.3 [Internet]. Brussels: European Commission; 2018 [cited 2019 Nov 18]. Available from: https://eplca.jrc.ec.europa.eu/permalink/PEFCR_guidance_v6.3-2.pdf
11. Department of Health. Delivering better oral health: an evidence-based toolkit for prevention [Internet]. London: Public Health England; 2017 [cited 2019 Nov 18]. Report no.: 2016224 Available from:

<https://www.gov.uk/government/publications/delivering-better-oral-health-an-evidence-based-toolkit-for-prevention>

12. Yaacob M, Worthington HV, Deacon SA et al. Powered versus manual toothbrushing for oral health. *Cochrane Database Syst Rev* 2014;6:No.:CD002281.
13. Scottish Dental. Sustainability and the Childsmile programme [Internet]. Dundee: NHS Scotland; 2019 [cited 2019 Nov 18]. Available from: <https://www.scottishdental.org/sustainability-and-the-childsmile-programme/>
14. Lee H. The best electric toothbrush [Internet]. New York: Reviews.com LLC 2019 [cited 2019 Nov 18]. Available from: <https://www.reviews.com/best-electric-toothbrush/>
15. Bamboo Grove. Bamboo agriculture [Internet]. Moline: BambooGrove 2019 [cited 2019 Nov 18]. Available from: <https://www.bamboogrove.com/bamboo-agriculture.html>
16. De Klein C, Novoa RSA, Ogle S et al. Guidelines for National Greenhouse Gas Inventories, Volume 4: Agriculture, forestry and other land use [Internet]. Geneva: Intergovernmental Panel on Climate Change (IPCC) [cited 2019 Dec 05] Available from: https://www.ipcc-nggip.iges.or.jp/public/2006gl/pdf/4_Volume4/V4_11_Ch11_N2O&CO2.pdf
17. Yang Y, Hui C. China's bamboo: culture, resources, cultivation, utilization. Beijing: International Network for Bamboo and Rattan (INBAR); 2010. Report no.: 33.
18. American Dental Association. Toothbrush care: cleaning, storing and replacement. *J Am Dent Assoc* 2006;3:415.
19. Buranyi S. The plastic backlash: what's behind our sudden rage – and will it make a difference [Internet]. London: The Guardian News & Media Limited 2018 [cited 2019 Nov 18]. Available from: <https://www.theguardian.com/environment/2018/nov/13/the-plastic-backlash-whats-behind-our-sudden-rage-and-will-it-make-a-difference>
20. Laville S. Plastic waste set to beat price as UK shoppers' top concern [Internet]. London: The Guardian News & Media Limited 2018 [cited 2019 Nov 18]. Available from: <https://www.theguardian.com/environment/2018/sep/10/plastic-waste-set-to-beat-price-as-uk-shoppers-top-concern-study>

21. Maoyi F. Sustainable management and utilization of sympodial bamboos. Hong Kong: China Forestry Publishing House; 2007.
22. Zheng Z. Bamboo reusable cups are not always biodegradable and can be harmful to health [Internet]. Singapore: Mothership 2019 [cited 2019 Nov 18]. Available from: <https://mothership.sg/2019/11/bamboo-reusable-cups-environment/>
23. Macek P, Terek-Derszniak M, Zak M et al. WHO recommendations on physical activity versus compliance rate within a specific urban population as assessed through IPAQ survey: a cross-sectional cohort study. *BMJ Open* 2019;9:e028334.

FIGURE LEGENDS

Figure 1. LCA boundaries.

Figure 2. LCIA results.

Figure 3. Contribution analysis for: a. plastic manual toothbrush, b. bamboo manual toothbrush, c. plastic manual replaceable head toothbrush, d. electric toothbrush.

TABLE LEGENDS

Table 1. Assumptions and exclusions

Table 2. Impact categories and LCIA methods used in this study.¹⁰