

Association of illicit drug use in adolescence with socioeconomic and criminal justice outcomes in adulthood: prospective findings from a UK national birth cohort

James White, Steven Bell, G. David Batty

James White, PhD. Centre for Trials Research, Cardiff University. 4th floor, Neuadd Meirionnydd, Heath Park, Cardiff, CF14 4YS, U.K. whitej11@cf.ac.uk; orcid.org/0000-0001-8371-8453

Steven Bell, PhD. Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK and The National Institute for Health Research Blood and Transplant Unit in Donor Health and Genomics at the University of Cambridge, Cambridge, UK. scb81@medschl.cam.ac.uk.

G. David Batty, DSc. Department of Epidemiology and Public Health, University College London, London, WC1E 7HB, UK. david.batty@ucl.ac.uk; School of Biological and Population Health Sciences, Oregon State University, Corvallis, Oregon, USA.

Manuscript statistics: 2021 words (excluding abstract of 227), 1 figure, 1 table, 28 references

Correspondence to: Dr James White

4th floor Neuadd Merinionydd,

Centre for Trials Research,

College of Biomedical and Life Sciences,

Cardiff University,

Cardiff.

CF14 4YS.

Wales, United Kingdom.

Email: whitej11@cf.ac.uk

Tel: +44 (0) 29 20 687054

Fax: +44 (0) 29 20 687236

What is already known on this subject

Illicit drug use in adolescence has previously been shown to be associated with an increased risk of drug use, delinquency, violence and poor mental health during adulthood.

Few studies have examined associations between adolescent illicit drug use with adult social, economic, and criminal justice outcomes.

What this study adds

We found new links between illicit drug use in adolescence and an increased risk of experiencing homelessness, being arrested and/or cautioned by the police, and being found guilty at court. These associations were not explained by parental socioeconomic status or psychological morbidity in adolescence.

Illicit drug use in adolescence may contribute to the development of health and economic inequalities in the transition to adulthood.

Abstract

Background: Illicit drug use in adolescence has been linked to drug use and poor mental health in adult life, but few studies have examined the relation with adult economic and criminal justice outcomes.

Methods: We analysed data from 14,082 participants (6,999 women) in the 1970 British Birth Cohort Study with data at 16 and 30 years of age. Illicit drug use over study members' lifetime and in the past year were self-reported at 16-years of age. Thirteen outcomes were self-reported at 30-years of age, including illicit drug use, smoking, problems with alcohol, mental and physical health, experience of socioeconomic disadvantage, and experiences with the criminal justice system.

Results: At 16 years of age, 20.3% of participants had ever tried an illicit drug and 7.2% had used in the past year. After adjustment for parental socioeconomic status and childhood mental health problems, and following correction for multiple testing, there was a dose-response association between illicit drug use at 16-years with illicit drug use in adult life the past year (odds ratio; 95% confidence interval: 1.83, 1.51 to 3.12), experiencing homelessness (1.74, 1.16 to 2.62), being arrested (1.57, 1.29 to 1.92) and cautioned (1.97, 1.50 to 2.57) by the police, and being found guilty at court (1.73, 1.34 to 2.23).

Conclusions: Illicit drug use in adolescence may contribute to the development of health and social inequalities in early adulthood.

Introduction

Illicit drugs use in adolescence has been well-examined in the context of later drug use^{1 2} and mental health.^{3 4 5} Studies of twins discordant for cannabis use (i.e. one used cannabis and the other did not) suggest a casual relation with later other illicit drugs,^{6 7} and systematic reviews report that regular cannabis use is associated with incident anxiety and mood disorders,^{2 4} and psychosis.^{8 9} Less well understood is the impact of adolescent drug use on socioeconomic adversity and criminal justice outcomes. In the few studies conducted, early-onset cannabis use has been linked to poor educational achievement,^{10 11} delinquency and violence.¹² In the Victoria Adolescent Health Cohort Study, amphetamine use by 18 years of age was associated with increased risks for leaving with no qualifications, but not with unemployment, or receiving government benefits at 24-years.⁵ The potential confounding effect of early socioeconomic adversity was not accounted for in these analyses and the role of psychological problems has also rarely been considered. Accordingly, we examined whether illicit drug use at 16-years of age is associated with an increased risk of an array of outcomes at 30 years, including poor health, socioeconomic disadvantage and experiences with the criminal justice system.

Methods

We used data from the 1970 British Birth Cohort Study, an ongoing longitudinal study of children born in Great Britain between the 5th and 11th April 1970. A total of 16, 571 babies born in England, Scotland and Wales were enrolled at birth and have been followed up on eight occasions across the life course. For the purposes of the present analyses, we used data from enquiries about illicit drug taking at 16 years of age, the first occasion these were collected.¹³ Study follow-up was at age 30 years. Parents of study participants gave written informed consent. This manuscript adheres to the guidelines for STrengthening the Reporting of OBservational studies in Epidemiology (STROBE).¹⁴

Study participants

At the age of 16 (1986), a national teachers strike and school examinations reduced study member participation.¹⁵ Of the 15,999 members traced and invited to participate, information

was obtained from 11,615 (72.6%). At the age of 30, 14,087 members were traced and invited to participate and 11,261 (68%) responded. Preliminary analyses showed there was very little difference in the prevalence of illicit drug use at 16-years among those who did and did not participate in the survey at 30-years (7.0% vs. 7.2%).

Assessment of illicit drug use at age 16 years

Exposures, assessed at 16-years of age, were derived from questions asking, “Have you ever tried taking cannabis?” (repeated for glue/solvents, amphetamines, barbiturates, cocaine, heroin and lysergic acid diethylamide (LSD)). Street names of drugs were also provided. Responses were combined across drugs and categorised into: never, taken but not in the past year, and taken in the past year.

Outcome assessment at age 30 years

Thirteen outcomes were chosen from the assessment at 30-years. These were chosen to test associations across a number of health and economic domains. Health behaviours comprised illicit drug use (ever and in the past year), including the use of cannabis, cocaine, crack, ecstasy, amphetamines, poppers, temazepam, ketamine, magic mushrooms, LSD, heroin, methadone, and other drugs. Street names of drugs were provided. All participants, apart from lifelong teetotallers, were asked to complete questions on possible problems with alcohol using the cutting down, being annoyed by criticism, feeling guilty, and eye-openers (CAGE) questionnaire.¹⁶ Smoking every day was defined as smoking at least one cigarette every day in the past week. Mental health problems were assessed using enquiries on: having seen a specialist medical provider since age 16 because of a problem with illicit drugs; a mental health problem; or a psychological morbidity based on a score of ≥ 7 on the Rutter Malaise Inventory.¹⁷ Physical health was ascertained from self-reports of accidents that occurred since 16-years that required treatment by a doctor. These included a sub-set that were due to a mugging or a violent or sexual assault. Socioeconomic disadvantage was based on employment status and whether participants had been homeless since 16-years. Finally, study members reported having been arrested, formally cautioned or found guilty in a court since 16-years of age.

Covariates

Covariates, identified *a priori*, were chosen based on previous studies showing an association with adolescent and/or adult illicit drug use. At 16-years of age these included the number of units of alcohol consumed in the past week, smoking in the past week, psychological morbidity (score of ≥ 7 on the Rutter Malaise inventory),¹⁷ and occupational social class based on mother's and father's occupation provided by parents at interview and coded using the Registrar General's classification system.¹⁸

Statistical methods

We took two approaches to dealing with missing data. The primary approach was to impute all missing exposure, outcome and covariate data (MI) using information from over 40 variables associated with our measures or missingness to make the assumption that data were 'missing at random' more plausible. We conducted a sensitivity analysis using multiple imputation then deletion (MID) whereby exposure, outcome and covariate data are imputed, but outcome is then deleted before analysis. MID has been found to produce more precise estimates than MI and can help minimize bias if the model for imputing outcome data is misspecified (i.e., because imputed outcome data are removed before analysis).¹⁹ For both MI and MID we generated 20 imputed datasets.

We compared the characteristics of participants with and without complete data. Logistic regression was used to calculate odds ratios (ORs) and 95% confidence intervals (CI) to summarize the association between illicit drug use at 16-years of age and later outcomes. In preliminary analyses, there was no difference in results in men and women so data were pooled and sex-adjusted. Illicit drug use at 16-years was modelled as a binary variable (never vs. use in the past year), or as a categorical variable (never vs. taken but not in the past year, used in the past year). The primary analysis was the testing of dose-response associations by modelling the categorical drug use exposure variable as a linear term. To reduce the risk of generating spurious findings due to multiple testing we applied the Bonferroni correction to the tests of trend across categories of drug use.²⁰ This meant that only p-values for a test of trend of <0.004 ($p=0.05/13$) were considered significant. Odds ratios were adjusted for sex, parental social class and the Rutter Malaise Inventory and at 16 year, plus the baseline status of the outcome variable if it was available at 16-years. All analyses were performed using Stata, version 15.1.

Results

There were 14,082 participants who provided data at either 16 or 30 years of age. The MID sample with complete outcome data was 9,651 and with complete exposure, outcomes and covariates was 1,858. Participants with complete data were more likely at 16 years to be female, have parents with a managerial or professional occupation, seen a doctor for a mental health problem, have a psychological morbidity, have had an accident and less likely to have used an illicit drug (ever, or in the past year), seen a specialist or been to a hospital because of a drug problems, been homeless, or had any police/criminal justice experiences, than those who did not have complete data. Our primary results are based on analyses with the MI sample as they offered greater precision (n = 14,082 participants; 6,999 women).

At 16-years, 20.3% of participants had ever tried an illicit drug, with 7.2% having used in the past year. Out of those who used in the past year, 87.0% only used cannabis. Of the 13 outcomes studied, after adjustment for confounding factors, adolescent illicit drug use was associated with all adult outcomes measured except for having possible problems with alcohol, getting treatment for an accident, or being unemployed (Figure). After correction for multiple testing, there was a significant linear trend across categories of illicit drug use for five outcomes at 30-years of age: illicit drug use in the past year, having been homeless, arrested, formally cautioned and found guilty in a court (Table 1). Sensitivity analysis using the MID sample (online supplementary tables 1 and 2) and the sample with no missing data were not materially different to those using the MI sample (online supplementary tables 3 and 4). As there were no participants in the sample with no missing data who had seen a specialist with a problem with drugs, were unemployed, or been homeless we could not generate estimates for these outcomes.

Discussion

In this population-wide birth cohort study, illicit drug use during adolescence was associated with a wide range of adverse outcomes in early adulthood. These included an association with illicit drug use in the past year, having been homeless, and criminal justice experiences. These associations were not explained by parental socioeconomic status or psychological morbidity in adolescence.

Comparison with existing studies

Our analyses examined illicit drug use but replicate those from smaller cohort studies that have investigated cannabis and amphetamine use in the United States, New Zealand and Australia. These studies have shown that cannabis use in adolescence is associated with negative outcomes across a number of domains, including drug use in early adulthood,^{1 2} daily smoking,¹² alcohol use disorders,³ mental health problems^{3 4 21} (in particular cannabis with psychosis²² and psychotic disorders²³) and involvement in criminal activity.^{12 24} They replicate the findings of the Victoria Adolescent Health Cohort Study showing amphetamine use in adolescence was associated with daily smoking, alcohol dependency, drug use in the past year, and psychological distress in early adulthood.²⁵ Our findings extend these by showing that after adjusting for early life socioeconomic disadvantage and multiple comparisons there was an association between adolescent drug use with being homeless and being convicted in a criminal court.

Potential mechanisms

The associations between drug use and adult outcomes we observed might be direct, indirect or due to confounding. In support of a direct effect, discordant twin studies have found the cannabis using twin was more likely to use other illicit drugs at a later date,^{6 7} and report psychotic like experiences,²⁶ than the twin who did not. In the Christchurch birth cohort, associations between cannabis use with juvenile offending were attenuated after adjusting for affiliation with substance using or delinquent peers.²⁴ In the present submission, criminal justice experiences might be acting as a mediator by increasing time with delinquent or older peers in whom drug use may be more prevalent. This may in-turn lead adolescents to taking drugs themselves. An alternative explanation is that these associations are brought about by confounding, whereby both adolescent drug use and later adulthood outcomes share common antecedents such as conduct/oppositional disorders,²⁴ delinquency,¹² and peer²⁷ or parental drug use.²⁸ This explanation suggests illicit drug use is better characterised as a marker rather than a cause of a life trajectory that is more likely to experience substance misuse, homelessness, the criminal justice system and poor mental health in later life.

Strengths and limitations

This study has several strengths, including the wide range of outcomes available, extended follow-up into early adulthood, and the wealth of potential confounding factors that we adjusted for in our analyses. The main limitation of this study is loss to follow-up and missing data. These may have introduced selection bias into the complete case sample. We therefore used multiple imputation to minimise this bias. Sensitivity analysis comparing the results from the complete case to the imputed datasets provided no evidence that missing data introduced bias. The sensitive nature of reporting illicit drug use and police and criminal justice experiences may have resulted in under-reporting. In 2004, the UK's 1971 Misuse of Drugs Act was amended introducing discretionary warnings for cannabis possession. As drug use was reported in the present cohort in 1986, it is possible that cases of cannabis possession now would not result in the criminal justice outcomes we observed.

Conclusions

We found illicit drug use in adolescence is associated with significant socioeconomic adversity and poorer health outcomes in adulthood. If causal, these associations would suggest that illicit drug use in adolescence contributes to the development of health and economic inequalities in the transition to adulthood.

Acknowledgements

Contributors: JW and GDB had the idea and designed the analyses. JW acquired the data and undertook the analyses. SB contributed to refining the analyses and drafted the figure. All authors drafted the manuscript and approved its publication.

Competing interest: None declared.

Ethics approval: Plewis I, Calderwood L, Hawkes D, Nathan G. National Child

Development Study and 1970 British Cohort Study technical report: Changes in the

NCDS and BCS70 populations and samples over time. London: Institute of

Education. 2004.

Provenance and peer review: Not commissioned; externally peer reviewed.

Data sharing statement: All data are openly available from the UK Data Archive (<https://discover.ukdataservice.ac.uk/>).

Grants and/or financial support: JW is funded by The Centre for the Development and Evaluation of Complex Interventions for Public Health Improvement (DECIPHer), a UKCRC Public Health Research Centre of Excellence. Joint funding (MR/KO232331/1) from the British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, the Welsh Government and the Wellcome Trust, under the auspices of the UK Clinical Research Collaboration, and is gratefully acknowledged. GDB is supported by the UK Medical Research Council (MR/P023444/1) and the US National Institute on Aging (1R56AG052519-01; 1R01AG052519-01A1). SB is supported by the National Institute for Health Research (NIHR) Blood and Transplant Research Unit in Donor Health and Genomics (NIHR BTRU-2014-10024), UK Medical Research Council (MR/L003120/1), British Heart Foundation (SP/09/002; RG/13/13/30194; RG/18/13/33946), and the NIHR Cambridge Biomedical Research Centre at the Cambridge University Hospitals NHS Foundation Trust.

Licence for Publication: The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd to permit this article (if accepted) to be published in JECH and any other BMJ PGL products and sublicences such use and exploit all subsidiary rights, as set out in our licence (<https://eur03.safelinks.protection.outlook.com/?url=http%3A%2F%2Fgroup.bmj.com%2Fproducts%2Fjournals%2Finstructions-for-authors%2Flicence-forms&data=01%7C01%7CWhiteJ11%40cardiff.ac.uk%7C0e49c5cbd2b84f01d89908d79233ed2a%7Cbdb74b3095684856bdbf06759778fcbc%7C1&data=oMgdVPdjCUmitMSnLtxtucVuBt2wC1C3E72SZ6Xnhc%3D&reserved=0>).

References

1. Swift W, Coffey C, Degenhardt L, Carlin JB, Romaniuk H, Patton GC. Cannabis and progression to other substance use in young adults: findings from a 13-year prospective population-based study. *J Epidemiol Community Health*. 2012 Jul;66(7):e26.

2. Hall W. What has research over the past two decades revealed about the adverse health effects of recreational cannabis use? *Addiction*. 2015;110(1):19–35.
3. Blanco C, Hasin DS, Wall MM, Flórez-Salamanca L, Hoertel N, Wang S, et al. Cannabis Use and Risk of Psychiatric Disorders: Prospective Evidence From a US National Longitudinal Study. *JAMA Psychiatry*. 2016 Apr 1;73(4):388–95.
4. Gobbi G, Atkin T, Zytynski T, Wang S, Askari S, Boruff J, et al. Association of Cannabis Use in Adolescence and Risk of Depression, Anxiety, and Suicidality in Young Adulthood: A Systematic Review and Meta-analysis. *JAMA Psychiatry* [Internet]. 2019 Feb 13 [cited 2019 Mar 22]; Available from: <https://jamanetwork.com/journals/jamapsychiatry/fullarticle/2723657>
5. Degenhardt L, Coffey C, Moran P, Carlin JB, Patton GC. The predictors and consequences of adolescent amphetamine use: findings from the Victoria Adolescent Health Cohort Study. *Addiction*. 2007;102(7):1076–84.
6. Lynskey MT, Heath AC, Bucholz KK, Slutske WS, Madden PAF, Nelson EC, et al. Escalation of drug use in early-onset cannabis users vs co-twin controls. *JAMA*. 2003 Jan 22;289(4):427–33.
7. Lynskey MT, Vink JM, Boomsma DI. Early onset cannabis use and progression to other drug use in a sample of Dutch twins. *Behav Genet*. 2006 Mar;36(2):195–200.
8. Marconi A, Di Forti M, Lewis CM, Murray RM, Vassos E. Meta-analysis of the Association Between the Level of Cannabis Use and Risk of Psychosis. *Schizophr Bull*. 2016;42(5):1262–9.
9. Large M, Sharma S, Compton MT, Slade T, Nielssen O. Cannabis Use and Earlier Onset of Psychosis: A Systematic Meta-analysis. *Arch Gen Psychiatry*. 2011 Jun 6;68(6):555–61.
10. Stiby AI, Hickman M, Munafò MR, Heron J, Yip VL, Macleod J. Adolescent cannabis and tobacco use and educational outcomes at age 16: birth cohort study. *Addict Abingdon Engl*. 2015 Apr;110(4):658–68.
11. Lynskey M, Hall W. The effects of adolescent cannabis use on educational attainment: a review. *Addict Abingdon Engl*. 2000 Nov;95(11):1621–30.
12. Brook JS, Balka EB, Whiteman M. The risks for late adolescence of early adolescent marijuana use. *Am J Public Health*. 1999 Oct;89(10):1549–54.
13. Elliott J, Shepherd P. Cohort Profile: 1970 British Birth Cohort (BCS70). *Int J Epidemiol*. 2006 Aug 1;35(4):836–43.
14. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ*. 2007 Oct 20;335(7624):806–8.
15. Goodman A, Butler N. BCS70—The 1970 British Cohort Study: The Sixteen-Year Follow-Up. *Inst Educ Lond*. 1987;

16. Ewing JA. Detecting alcoholism. The CAGE questionnaire. *JAMA*. 1984 Oct 12;252(14):1905–7.
17. Rodgers B, Pickles A, Power C, Collishaw S, Maughan B. Validity of the Malaise Inventory in general population samples. *Soc Psychiatry Psychiatr Epidemiol*. 1999 Jun 1;34(6):333–41.
18. General R. Classification of occupations. Lond HMSO. 1980;
19. Von Hippel PT. 4. Regression with missing Ys: An improved strategy for analyzing multiply imputed data. *Sociol Methodol*. 2007;37(1):83–117.
20. VanderWeele TJ, Mathur MB. Some desirable properties of the Bonferroni correction: Is the Bonferroni correction really so bad? *Am J Epidemiol*. 2019 Mar 1;188(3):617–8.
21. Patton GC, Coffey C, Carlin JB, Degenhardt L, Lynskey M, Hall W. Cannabis use and mental health in young people: cohort study. *BMJ*. 2002 Nov 23;325(7374):1195–8.
22. McGrath J, Welham J, Scott J, Varghese D, Degenhardt L, Hayatbakhsh MR, et al. Association Between Cannabis Use and Psychosis-Related Outcomes Using Sibling Pair Analysis in a Cohort of Young Adults. *Arch Gen Psychiatry*. 2010 May 1;67(5):440–7.
23. Forti MD, Quattrone D, Freeman TP, Tripoli G, Gayer-Anderson C, Quigley H, et al. The contribution of cannabis use to variation in the incidence of psychotic disorder across Europe (EU-GEI): a multicentre case-control study. *Lancet Psychiatry* [Internet]. 2019 Mar 19 [cited 2019 Mar 22];0(0). Available from: [https://www.thelancet.com/journals/lanpsy/article/PIIS2215-0366\(19\)30048-3/abstract](https://www.thelancet.com/journals/lanpsy/article/PIIS2215-0366(19)30048-3/abstract)
24. Fergusson DM, Lynskey MT, Horwood LJ. The short-term consequences of early onset cannabis use. *J Abnorm Child Psychol*. 1996 Aug 1;24(4):499–512.
25. Degenhardt L, Coffey C, Moran P, Carlin JB, Patton GC. The predictors and consequences of adolescent amphetamine use: findings from the Victoria Adolescent Health Cohort Study. *Addiction*. 2007;102(7):1076–84.
26. Karcher NR, Barch DM, Demers CH, Baranger DAA, Heath AC, Lynskey MT, et al. Genetic Predisposition vs Individual-Specific Processes in the Association Between Psychotic-like Experiences and Cannabis Use. *JAMA Psychiatry*. 2019 Jan 1;76(1):87–94.
27. Van Ryzin MJ, Fosco GM, Dishion TJ. Family and peer predictors of substance use from early adolescence to early adulthood: an 11-year prospective analysis. *Addict Behav*. 2012 Dec;37(12):1314–24.
28. Bahr SJ, Hoffmann JP, Yang X. Parental and Peer Influences on the Risk of Adolescent Drug Use. *J Prim Prev*. 2005 Nov 1;26(6):529–51.




Figure. Odds ratios (95% confidence intervals) for adult outcomes of lifetime illicit drug use in adolescence (n = 14, 082)

Reference category was never used illicit drugs. Odds ratios (ORs) are adjusted for sex, parental social class and the Rutter Malaise Inventory and at 16 years. When available, age 16-year (baseline) assessments of outcomes were added to multivariable models. This included adjusting for screening positive on the Rutter Malaise Inventory at 16 years for the outcome at 30-years of having seen a specialist for a psychiatric

the number of cigarettes smoked per week at 16 years for the outcome at 30-years of smoking every day.

Table 1. Odds ratios (95% confidence intervals) for adult outcomes of illicit drug use in adolescence (n = 14, 082)

Outcome at 30 years ^a	Illicit drug use at 16 years			P-value for trend
	Never	Taken but not in past year	Taken in past year	
Health behavior				
Illicit drug use in the past year	1 (Reference)	1.57 (1.27, 1.94)	2.36 (1.80, 3.09)	<0.001
Smoke every day	1	1.20 (0.93, 1.54)	1.52 (1.10, 2.12)	0.02
Alcohol problem (CAGE score ≥ 2)	1	1.17 (0.86, 1.58)	1.59 (1.10, 2.26)	0.02
Mental health				
Seen specialist for problem with illicit drugs since 16	1	1.63 (0.86, 3.10)	2.71 (1.22, 6.01)	0.01
Seen specialist for psychiatric problem since 16	1	1.19 (1.01, 1.40)	1.22 (0.94, 1.60)	0.03
Psychiatric morbidity (Malaise score ≥ 7)	1	1.35 (1.09, 1.67)	1.31 (1.00, 1.72)	0.01
Physical health				
Seen doctor since 16 years of age to get treatment for an accident	1	1.03 (0.88, 1.20)	1.00 (0.83, 1.22)	0.88
Seen doctor since 16 years of age to get treatment for a violent assault, mugging or sexual assault	1	1.27 (0.99, 1.63)	1.27 (0.90, 1.80)	0.03
Socioeconomic disadvantage				
Unemployed	1	1.24 (0.74, 2.08)	1.27 (0.68, 2.37)	0.36
Homeless since 16 years of age	1	1.54 (1.04, 2.29)	2.11 (1.32, 3.37)	0.002
Police/ criminal justice experiences				
Arrested by police since 16 years	1	1.55 (1.23, 1.94)	1.63 (1.23, 2.17)	<0.001
Formally cautioned by police since 16 years	1	1.74 (1.34, 2.25)	2.41 (1.59, 3.63)	<0.001
Been found guilty in court since 16 years	1	1.53 (1.15, 2.04)	2.12 (1.59, 2.83)	<0.001

^a Reference category was never used illicit drugs. Odds ratios (ORs) are adjusted for sex, parental social class and the Rutter Malaise Inventory and at 16 years. When available, age 16-year (baseline) assessments of outcomes were added to multivariable models. This included adjusting for screening positive on the Rutter Malaise Inventory at 16 years for the outcome at 30-years of having seen a specialist for a psychiatric problem; units of alcohol consumed in the past week at 16 years for the outcome at 30-years of screening positive for alcohol dependency; and the number of cigarettes smoked per week at 16 years for the outcome at 30-years of smoking every day.