

**Joint Modelling Of Individual Trajectories, Within-Individual Variability And A Later Outcome: Systolic Blood Pressure Through Childhood And Left Ventricular Mass In Early Adulthood**

Richard M.A. Parker, George Leckie, Harvey Goldstein, Laura D. Howe, Jon Heron, Alun D. Hughes, David M. Phillippo, and Kate Tilling

Correspondence to Dr. Richard Parker, MRC Integrative Epidemiology Unit, University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, UK (email: [richard.parker@bristol.ac.uk](mailto:richard.parker@bristol.ac.uk))

Author affiliations: MRC Integrative Epidemiology Unit, University of Bristol, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, UK (Richard M.A Parker, Laura D. Howe, Jon Heron, and Kate Tilling); Population Health Sciences, Bristol Medical School, Oakfield House, Oakfield Grove, Bristol, BS8 2BN, UK (Richard M.A Parker, Laura D. Howe, Jon Heron, David M. Phillippo, and Kate Tilling); Centre for Multilevel Modelling, School of Education, University of Bristol, 35 Berkeley Square, Bristol, BS8 1JA, UK (George Leckie, and Harvey Goldstein); MRC Unit for Lifelong Health and Aging, Institute of Cardiovascular Science, University College London, London, WC1E 6HX, UK (Alun D. Hughes)

© The Author(s) 2020. Published by Oxford University Press on behalf of the Johns Hopkins Bloomberg School of Public Health. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

This work was supported by: the UK's Medical Research Council (MRC) grant MR/N027485/1, awarded to investigator K.T and funding R.M.A.P; the MRC and the University of Bristol support the MRC Integrative Epidemiology Unit (MC\_UU\_00011/3); A.D.H. received support from the Wellcome Trust (086676/7/08/Z) and the British Heart Foundation (PG/06/145 & CS/15/6/31468) and works in a unit that receives support from the MRC (Programme Code MC\_UU\_12019/1); D.M.P. was funded by MRC grant MR/P015298/1; the UK Medical Research Council and Wellcome (Grant ref: 102215/2/13/2) and the University of Bristol provide core support for ALSPAC. This publication is the work of the authors and they will serve as guarantors for the contents of this paper.

Conflict of interest: none declared.

Running head: Joint Modelling Of BP Variability And LVMI

ORIGINAL UNEDITED MANUSCRIPT

## ABSTRACT

Within-individual variability of repeatedly-measured exposures may predict later outcomes: e.g. blood pressure (BP) variability (BPV) is an independent cardiovascular risk factor above and beyond mean BP. Since two-stage methods, known to introduce bias, are typically used to investigate such associations, we introduce a joint modelling approach, examining associations of mean BP and BPV across childhood to left ventricular mass (indexed to height; LVMI) in early adulthood with data (collected 1990-2011) from the UK's Avon Longitudinal Study of Parents and Children cohort. Using multilevel models, we allow BPV to vary between individuals (a "random effect") as well as to depend on covariates (allowing for heteroscedasticity). We further distinguish within-clinic variability ("measurement error") from visit-to-visit BPV. BPV was predicted to be greater at older ages, at higher bodyweights, and in females, and was positively correlated with mean BP. BPV had a weak positive association with LVMI (10% increase in within-individual BP variance was predicted to increase LVMI by 0.21% (95% credible interval: -0.23%, 0.69%)), but this association became negative (-0.78%, 95% credible interval: -2.54%, 0.22%)) once the effect of mean BP on LVMI was adjusted for. This joint modelling approach offers a flexible method of relating repeatedly-measured exposures to later outcomes.

**KEYWORDS:**ALSPAC; Bayesian Analysis; Blood Pressure; Children; Joint model; Left Ventricular Hypertrophy; Longitudinal Studies; Young Adult

### ABBREVIATIONS

BMI body mass index

BPV (within-individual) blood pressure variability

BP blood pressure

CI credible interval

CVD cardiovascular disease

LVMI left ventricular mass indexed to height

SBP systolic blood pressure

SD standard deviation

When exposures vary over time – for example systolic blood pressure (BP) as measured over the life course – both the overall mean and the change in the exposure can affect later health outcomes (1). However, within-individual variability – i.e. the extent to which an individual’s measure fluctuates around this trend – may also be a risk factor (2).

Methodological difficulties have meant that within-individual variability in time-varying exposures have been seldom examined, and when they are it is often using two-stage procedures, known to introduce bias (3). An alternative, one-stage, approach is to use joint multilevel models to relate time-varying exposures (4) to a later outcome (5). These models have recently been extended to also examine within-individual variability (3). Such joint models lead to unbiased parameter estimates and correct standard errors: either when relating the mean and trajectory (6), or within-individual variability (3), of an exposure to a later outcome.

We illustrate this approach by examining the association of both mean BP and within-individual blood pressure variability (BPV) – or more specifically the estimated individual deviation away from the population mean BP and BPV (adjusted for covariates) – with an established biomarker of target organ damage: left ventricular mass indexed to height (LVMI). Whilst BPV can be measured across various time scales (7), here we focus on the longer-term, a.k.a. “visit-to-visit”. In adults, visit-to-visit BPV has been found to be an important predictor of subsequent cardiovascular disease (CVD) over and above mean level (8). Detecting those at risk of CVD early in life is important in order to design and administer preventative measures in a timely and targeted manner (9, 10). Mean systolic BP in childhood is positively associated with early signs of heart damage, such as left ventricular hypertrophy (11-13), and there is evidence from a US cohort of children that higher BPV – as estimated via the standard deviation (SD) of the measurements across all visits (conducted every 2-3 years) for each person – is associated with

adult hypertension, independent of mean BP levels (14). However, there has been little investigation of the role of BPV on early target organ damage, despite its potential utility as a predictive factor (15), and only limited analysis of the factors associated with childhood BPV (14, 16).

Our aim was to explore the factors associated with within-individual BPV in a UK cohort of children using multilevel analyses, and to extend these to joint models to investigate the association of both mean BP and BPV with LVMI in early adulthood. Note that, since the responses are both Gaussian, the model could also be referred to a multivariate Normal response model, with heteroscedastic error.

## METHODS

### Participants

The participants are from the UK's Avon Longitudinal Study of Parents And Children (ALSPAC), an ongoing prospective longitudinal birth cohort study (17-19). Pregnant women resident in Avon, with expected dates of delivery 4/1/1991 to 12/31/1992, were invited to take part in the study. 14,541 pregnancies, resulting in 13,988 children alive at 1-year of age, were initially enrolled, later bolstered by further eligible cases who had failed to join the study originally, increasing the number of enrolled pregnancies to 15,454, with 14,901 children alive at 1-year of age. The study website contains details of all available data through a fully searchable data dictionary and variable search tool (20).

5,217 of the participants attended an ALSPAC study clinic at mean age 17.7 years, of whom a random subsample of 2,047 (all singletons) had their LVMI (left ventricular mass in grammes, indexed to height in  $m^{2.7}$ , i.e.  $g/m^{2.7}$ ) measured via echocardiography (21, 22) (exclusion criteria

included pregnancy and congenital heart disease; see below for a comparison of the subsample modelled with the larger sample enrolled in ALSPAC, and Boyd *et al.* (17) for a discussion of attrition in ALSPAC). Echocardiography was performed using an HDI 5000 ultrasound machine (Phillips Healthcare, Amsterdam, The Netherlands) equipped with a P4-2 Phased Array ultrasound transducer by one of two experienced echocardiographers using a standard examination protocol (22).

1,988 of these participants had their systolic BP (SBP), height and weight and age recorded, on at least one prior occasion (research clinic). These clinics were at *c.* 7.5-, 9.5-, 10.5-, 11.5-, 13-, 15.5-years of age. At each clinic, the participant's SBP (in mmHg) was measured at least twice, using a validated electronic monitoring device and a cuff size appropriate for their upper arm circumference, with the participant sitting and at rest with the arm supported. Each of these measurements was available for our analysis from each clinic, apart from the clinic at *c.* 10.5 years of age, for which only the mean was available.

#### Statistical analysis

The covariates we included for SBP were age (for further details, see below), sex, an interaction between age and sex, weight, height, and a number of maternal characteristics including parity (the number of previous pregnancies resulting in either livebirth or stillbirth), age at delivery, and highest educational qualification. Sex, age and weight, as measured at the clinic at *c.* 17.7 years of age, were included as exposures for LVMI, but height was not, as LVMI is indexed to height. All continuous covariates were centred around their grand mean prior to analysis. Weight, as measured across childhood, and LVMI were log-transformed prior to analysis.

We used a joint model to relate BPV to subsequent log(LVMI). We describe this modelling approach below, building up model complexity using an example which assumes that SBP increases linearly with age across childhood (we later relax this assumption in the ALSPAC model). Equation 1 assumes that each individual has one measure at each occasion, but this is relaxed in equation 2 (and in the ALSPAC model).

$$\begin{aligned}
 y_{1ij} &= \beta_0 + \beta_1 t_{ij} + u_{0j} + u_{1j} t_{ij} + e_{ij} \\
 e_{ij} &\sim N(0, \sigma_{eij}^2), \quad \log(\sigma_{eij}^2) = \alpha_0 + \alpha_1 t_{ij} + u_{2j} \\
 y_{2j} &= \gamma_0 + \gamma_1 u_{0j} + \gamma_2 u_{1j} + \gamma_3 u_{2j} + u_{3j}
 \end{aligned} \tag{1}$$

$$\begin{pmatrix} u_{0j} \\ u_{1j} \\ u_{2j} \\ u_{3j} \end{pmatrix} \sim N \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma_{u0}^2 & & & \\ \sigma_{u01} & \sigma_{u1}^2 & & \\ \sigma_{u02} & \sigma_{u12} & \sigma_{u2}^2 & \\ 0 & 0 & 0 & \sigma_{u3}^2 \end{pmatrix} \right\}$$

In equation 1,  $y_{1ij}$  is the repeatedly-measured exposure (SBP) measured at clinic  $i$  ( $i = 1, \dots, n_j$ ) for individual  $j$  ( $j = 1, \dots, J$ ), with the covariate  $t_{ij}$  indicating the individual's age at that clinic. In the mean function for BP, there are individual-level random effects for the intercept ( $u_{0j}$ ; mean BP at  $t = 0$ ) and also for rate of change ( $u_{1j}$ ; BP slope), with within-individual (between-clinic) error  $e_{ij}$ . In standard multilevel models, the variance of the within-individual error is assumed to be constant (homoscedastic) across all observations (as  $\sigma_e^2$ ). Here it is allowed to depend on age ( $t_{ij}$ ), and also on an individual-level random effect  $u_{2j}$  (with a log-link ensuring that the within-individual variance remains positive). Within-individual variability in blood pressure, log(BPV), is therefore allowed to change with age (as estimated here via  $\alpha_1$ ). The random effect  $u_{2j}$  allows each individual to have their own estimate of log(BPV). It is also possible to allow for coefficients of covariates within the within-individual variability function to randomly-vary

across individual: for example allowing the coefficient for age ( $t_{ij}$ ) to randomly-vary to investigate whether its association with BPV differs between people (23). The submodel for the longitudinal process is a mixed-effects location scale model (24, 25). The inclusion of the individual-level outcome  $y_{2j}$ , denoting log(LVMI) measured at *c.* 17.7 years for individual  $j$  ( $j = 1, \dots, J$ ), results in a joint model, with shared random effects: here via the inclusion of mean BP, BP slope and log(BPV) as exposures in the linear model for the mean of log(LVMI). The model above has two levels, and assumes there is just one measure per person, per clinic. If more than one measurement of SBP is taken for each individual at each clinic session, however, then the model can be expanded as follows:

$$\begin{aligned}
 y_{1hij} &= \beta_0 + \beta_1 t_{ij} + u_{0j} + u_{1j} t_{ij} + e_{ij} + \varepsilon_{hij} \\
 e_{ij} &\sim N(0, \sigma_{eij}^2), \quad \log(\sigma_{eij}^2) = \alpha_0 + \alpha_1 t_{ij} + u_{2j} \\
 \varepsilon_{hij} &\sim N(0, \sigma_\varepsilon^2) \\
 y_{2j} &= \gamma_0 + \gamma_1 u_{0j} + \gamma_2 u_{1j} + \gamma_3 u_{2j} + u_{3j} \tag{2}
 \end{aligned}$$

$$\left( \begin{array}{c} u_{0j} \\ u_{1j} \\ u_{2j} \\ u_{3j} \end{array} \right) \sim N \left\{ \left( \begin{array}{c} 0 \\ 0 \\ 0 \\ 0 \end{array} \right), \left( \begin{array}{cccc} \sigma_{u0}^2 & & & \\ \sigma_{u01} & \sigma_{u1}^2 & & \\ \sigma_{u02} & \sigma_{u12} & \sigma_{u2}^2 & \\ 0 & 0 & 0 & \sigma_{u3}^2 \end{array} \right) \right\}$$

In equation 2,  $y_{1hij}$  denotes SBP as measured at (within-clinic) occasion  $h$  ( $h = 1, \dots, n_{ij}$ ) for clinic  $i$  ( $i = 1, \dots, I_j$ ), for individual  $j$  ( $j = 1, \dots, J$ ).  $\varepsilon_{hij}$  refers to the residual within-clinic error in the repeatedly-measured exposure, assumed to have constant variance  $\sigma_\varepsilon^2$ . This within-clinic measurement error, which we assume is not related to log(LVMI) ( $y_{2j}$ ), will incorporate both systematic (e.g. white coat effect / habituation) and random error.



Following previous work modelling SBP across these ALSPAC clinics, we allowed for a non-linear relationship between age and mean SBP by fitting a linear spline with a knot point at 12 years (26, 27). The spline variables were derived as follows (where  $t_{ij}$  = age in years) (28):

$$s_{1ij} = \begin{cases} t_{ij} & t_{ij} \leq 12 \\ 12 & t_{ij} > 12 \end{cases}$$

$$s_{2ij} = \begin{cases} 0 & t_{ij} \leq 12 \\ t_{ij} - 12 & t_{ij} > 12 \end{cases}$$

These spline terms were fitted as fixed effects in the mean function for SBP. Since allowing the coefficients of these terms to randomly-vary across individuals lead to convergence problems in some models, a linear term for age fitted across the whole age range was fitted as a random effect. We assumed that BPV had a linear association with age.

Web Appendix 1 includes estimates from sensitivity analyses designed to (a) investigate the influence of the submodel for log(LVMI) on the random effect estimates (Web Table 1); (b) compare a parameterisation modelling clinic mean BP with current models (Web Table 2); (c) check whether restricting the analysis sample to those with LVMI measures had any effect on model for change in SBP, examining possible selection bias (Web Table 3); (d) investigate sex as the only observed covariate (Web Table 4), for comparison with published findings elsewhere (14, 16).

We used Bayesian estimation via MCMC methods in Stan (2.19.1) (29, 30), called from R (31) using the rstan package (32). See Web Appendices 2-3 for examples and further details of estimation.

Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee and the Local Research Ethics Committees. Informed consent for the use of data collected via

questionnaires and clinics was obtained from participants following the recommendations of the ALSPAC Ethics and Law Committee at the time.

## RESULTS

### Cohort description

Two observations at the individual-level, eight at the clinic-level, and one at the measurement-level (36 measurement-level observations in total) were identified as outliers in quantile-quantile plots of residuals from preliminary models and were removed from the dataset prior to further analyses. The resulting dataset comprised 1,986 individuals attending 10,556 clinic sessions with 19,360 recorded BP measurements.

<Table 1 here>

Table 1 compares the participants included in the analysis with those 12,318 ALSPAC children who were not included (but were recruited in an eligible ALSPAC phase, singletons, and otherwise recorded as alive at 1 year of age). There was evidence that those included were more likely to be female, had mothers who had had fewer children and were more educated and healthier than those not included.

Figure 1 plots mean SBP against mean age for each clinic for those included in the model, indicating sample sizes for each clinic. The mean number of clinics attended was 5.3; the number of clinics attended by participants was 1: n = 46; 2: n = 47; 3: n = 68; 4: n = 142; 5: n = 454; 6: n = 1,229.

<Figure 1 here>

<Table 2 here>

Fixed effects: age and sex

Table 2 summarises 3-level joint analyses in which both SBP trajectory and log(LVMI) are simultaneously modelled, with age centred around the sample mean of 11.3 years (with some estimates presented as percentage change (33), as indicated; see Web Table 5 for original coefficient estimates). With just age and sex as observed covariates (in model A), mean SBP was predicted to increase with age. The rate of increase was similar for males and females up to the age of 12 years, with mean BP predicted to increase more steeply for males at older ages. With regard to within-individual variance, with each year of age, BPV was predicted to increase by 15.6% (95% credible interval (CI): 12.9%, 18.4%). Females' BPV was predicted to be 16.8% (95% CI: 4.5%, 30.4%) greater than that for males, across all time points. Age at time of echocardiography was estimated to have a modest negative association with LVMI, and females were predicted to have smaller values of LVMI.

Fixed effects: adding weight and height

In model B mean SBP was predicted to increase with greater weight and height. BPV was estimated to be larger with greater log(weight): specifically, for a 10% increase in weight, BPV is predicted to increase by 5.8% (95% CI: 2.1%, 9.6%). The estimated effect of height on log(BPV) was negative, with a credible interval including zero (BPV predicted to decrease by -0.4% (95% CI: -1.4%, 0.7%) per 1cm increase in height). Weight had a positive association with LVMI (LVMI predicted to increase by 0.7% (95% CI: 0.6%, 0.8%) per 1kg increase in weight).

Fixed effects: adding maternal characteristics

Finally, in model C, maternal characteristics were added as covariates in the mean function for SBP (with a slight drop in sample size due to fewer observations for these variables). This

predicted a modest positive association of mean SBP with mother's age at delivery, and an inverse association with higher parity, and highest educational qualifications.

#### Random effects

Focusing on Model C, this estimated a small negative correlation ( $r = -0.09$  (95% CI: -0.21, 0.03)) between mean BP and BP slope: i.e. those with higher mean BP at 11.3 years were predicted to have a smaller increase in their mean BP across age. A large positive correlation ( $r = 0.50$  (95% CI: 0.33, 0.71)) was estimated between mean BP and log(BPV): i.e. individuals with higher mean SBP at 11.3 years also tended to have more fluctuation in their SBP. The correlation between BP slope and log(BPV) was estimated as negative, and small ( $r = -0.05$  (95% CI: -0.34, 0.23)).

<Figure 2 here>

The SD of the BPV random effects on the log scale was estimated to be mean = 0.42mmHg (95% CI: 0.29mmHg, 0.53mmHg). Figure 2 plots examples from this model of three individuals randomly-drawn from the 25 with the lowest, and 25 with the highest, estimated random effect for log(BPV) (with an added constant on both the x- and y-axis to preserve anonymity), illustrating instances of individual-level patterns of SBP measurements at each end of the observed range of estimated BPV.

<Table 3 here>

Table 3 contrasts the estimates of the random effects from model C to those in which the effect of each set of random effects on log(LVMI) is examined without adjustment for the other random effects on log(LVMI) (but with the remaining structure of the model otherwise remaining the same; estimates presented as percentage change (33), as indicated; see Web Tables

6-7 for full estimates, including original coefficient estimates). When mean BP is the only random effect included as a predictor of LVMI, higher mean BP is associated with higher LVMI. The same is true when the random slope and the log(BPV) random effects are alternately added as predictors of LVMI. With regard to the log(BPV) predictor, a 10% increase in BPV is predicted to increase LVMI by 0.21% (95% CI: -0.23%, 0.69%). When all three random effects are included as predictors of LVMI, the estimated association between log(BPV) and LVMI becomes negative and larger in magnitude; specifically, a 10% increase in BPV is estimated to be associated with a decrease in LVMI by -0.78% (95% CI: -2.54%, 0.22%). This model also estimated a positive association of mean BP with LVMI (a 1mmHg increase in BP at the sample mean age was predicted to increase LVMI by 0.59% (95% CI: 0.12%, 1.37%)) and of change in mean BP with LVMI (a 1mmHg increase in the slope was predicted to increase LVMI by mean = 3.58% (95% CI: 0.01%, 7.92%)).

#### Sensitivity analyses

As Web Appendix 1 further discusses, a range of sensitivity analyses yielded substantively similar estimates to the models presented here.

#### DISCUSSION

Using data from a prospective longitudinal birth cohort study based in the UK, we used a joint modelling approach to examine the factors associated with visit-to-visit BPV across childhood, and the association of both mean BP and BPV with an established biomarker of target organ damage, LVMI, in early adulthood. BPV was estimated to be larger – i.e. individuals' SBP was estimated to fluctuate more greatly – at older ages, in females, and at greater bodyweights. BPV also had a large, positive correlation with mean BP (at the sample mean age of 11.3 years), but

only a very small correlation with the rate of change in BP (slope) across age. In a model which did not adjust for the effect of mean BP and its slope on LVMI, BPV was estimated to have a weak positive association with LVMI. When mean BP and its slope were also included as covariates, however, the direction of association between BPV and LVMI changed and the magnitude of the estimate increased. These models further estimated a positive association of mean BP with LVMI and of change in mean BP with LVMI.

Whilst we are not aware of other studies in this age range which have reported on the effect of weight and height on BPV, the effect of age, sex and race have been examined. Investigating BPV in 8-18 year-old children in Massachusetts, Rosner *et al.* fitted variance components models to BP measurements taken within- and across-visits (up to a maximum of 16 visits over four years), and subsequently examined the effect of age, sex and BP level on these two variance components in three-way ANOVAs (16). They found “no meaningful effects” of these covariates on variability of SBP (children with higher BP level had significantly more within-individual variability than those with lower BP, but this difference was judged small; correlation coefficient not reported). More recently, in an analysis of the Bogalusa Heart Study in the USA, Chen *et al.* measured visit-to-visit BPV in children aged 4-19 years by taking the SD of 4-8 measurements of clinic mean SBP from visits scheduled every 2-3 years (14). Males were estimated to have significantly higher SD(SBP) than females, and SD(SBP) was estimated to be significantly greater for blacks than for whites. Our supplementary analyses (Web Table 4) also indicated that females were estimated to have lower BPV than males, but not once the effect of time-varying covariates were adjusted for. In addition, and in keeping with our findings, Chen *et al.* found a positive correlation ( $r = 0.15$ ) between mean childhood BP level and SD(SBP) (14). Such a positive correlation has been characterised in studies of adults too (8, 34).

Whilst several studies have found a positive relationship between adult BPV, independent of mean BP, and later cardiovascular events such as stroke (8) and all-cause mortality (34), evidence for the relationship between BPV and target organ damage (as indicated by left ventricular hypertrophy, for example) is more equivocal (35, 36). A recent review described contrasting results with regard to the association of visit-to-visit BPV and cardiovascular outcomes, suggesting differences between (adult) study populations, such as underlying cardiovascular risk, may be an important determinant of such heterogeneous findings (7). With regard to the relationship between childhood BPV and adult biomarkers, Chen *et al.* found childhood visit-to-visit SD(SBP) to be significantly associated with adult hypertension (14). This remained the case after adjusting for mean childhood SBP, although elevated childhood SBP levels were found to be more predictive of adult hypertension than childhood BPV. We have previously demonstrated that in 17 year-olds, higher body mass index (BMI) is causally related to higher LVMI, suggesting that there is meaningful variation in cardiac structure measures in early adulthood that is likely to track across life and relate to later life cardiovascular health (37). Nevertheless, in the current study only a weak positive association between childhood BPV and LVMI in early adulthood was found, and this did not persist once childhood mean BP had been adjusted for. In fact, within strata of mean childhood SBP, higher BPV was associated with smaller values of LVMI. Whether it is appropriate to adjust for mean BP when assessing the association of BPV with LVMI somewhat depends on whether one has a causal model in mind which conceptualises mean BP as causing both BPV and LVMI (in which case adjustment for mean BP as a confounder might be appropriate), or whether no causal relationship is proposed between mean BP and BPV (i.e. other, unknown, factors cause mean BP and BPV, and each in turn has a causal relationship with later LVMI), in which case it may not be appropriate to adjust

for mean BP. If the purpose is prediction, on the other hand, then by simultaneously estimating the effect of mean BP, its slope, and BPV on the later outcome, we allow for a more complete appraisal of the association of repeatedly-measured BP with LVMI. Note that whether mean BP is adjusted for or not, the association of BPV with LVMI was estimated to be small, with appreciable uncertainty as to whether it was non-zero, or indeed of the opposite sign.

Our results also indicated that mean BP (at the sample mean age), and change in mean BP, were both positively associated with LVMI in early adulthood. Analyses of the Georgia Stress and Heart Study and the Bogalusa Heart Study have also found mean BP in childhood, and its slope from childhood to young adulthood and in adolescence, to be positively associated with left ventricular hypertrophy in adulthood (38-40).

If we assume that selection into our analysis sample only depends on measured variables such as sex of child, education and age of mother, then their inclusion as covariates will result in unbiased models. If, having conditioned on such covariates, inclusion in our analysis sample is related to the exposure (SBP) and outcome (LVMI), then there will be bias. *A priori*, this seems relatively unlikely, however: e.g. such biomarkers are not readily-observable in a manner which would typically influence one's participation in a study. Whilst there were fewer observations for the maternal (*cf.* child-based) characteristics, the proportion of missing data was relatively modest (8-9%), and estimates for parameters common to each model were substantively similar.

Selection bias could also arise from people missing specific measures. Again, this will only cause bias if the probability of the measure being missing depends on its (unobserved) value, having conditioned on the observed values of the exposure, outcome and covariates included in the models. This again seems relatively unlikely, however, given that participants are not typically privy to such biomarkers in a manner which might influence attendance.



The general class of model we have used could be employed to examine within-person variability in any repeatedly-measured exposure, relating it to later individual-level outcomes of interest within a joint model: e.g. the association between glycaemic variability and mortality (41), variability in gait and risk of falling (42), and variation in prostate specific antigen and prostate volume (43). We have focused on within-individual variability, but interest may lie in characterising within-clinic (inter-individual) variability instead. Also, individual-level outcomes (i.e.  $y_{2j}$ , in equations 1 and 2) which are binary, or time-to-event data (3), etc., can be modelled (with appropriate changes to distributional assumptions). Furthermore, the repeatedly-measured exposure might instead be intensively-sampled, with many observations per person (24) (adjusting for autocorrelation, if appropriate).

Whilst fitting joint models involves greater computational burden and complexity compared to a two-stage approach, the latter is known to introduce bias (3). Note, though, that a recent methodological development of the two-stage approach preserves uncertainty in estimates (44). A two-stage approach may appear more logical in cases where it is not feasible for the later outcome to influence the earlier longitudinal process, due to fears over allowing the future to cause the past (44, 45). The joint modelling framework, however, uses future measurements to improve the model for the past measurements, rather than to cause the past measurements. This is the case in a wide range of model types and applications (5, 46, 47): e.g. a growth curve model, where adding later waves revises estimates of true values for earlier occasions.

We have supplied example code to demonstrate how these models can be fitted using a Bayesian framework, with endeavours to fit such models being actively extended to other software packages (48). Joint models have been recently employed to examine within-individual variability (3). We further extend these by investigating the association of time-varying

covariates with within-individual variability, allowing us to examine residual BPV unexplained by known factors, and also by distinguishing between-clinic from within-clinic variability. As such, the joint modelling approach we have introduced offers a very flexible method of exploring the factors associated with within-individual variability.

#### ACKNOWLEDGMENTS

We are extremely grateful to all the families who took part in this study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses.

ORIGINAL UNEDITED MANUSCRIPT

## REFERENCES

1. Tielemans S, Geleijnse JM, Menotti A, et al. Ten-Year Blood Pressure Trajectories, Cardiovascular Mortality, and Life Years-Lost in 2 Extinction Cohorts: the Minnesota Business and Professional Men Study and the Zutphen Study. *J Am Heart Assoc* 2015;4(3).
2. Elliott MR, Sammel MD, Faul J. Associations between variability of risk factors and health outcomes in longitudinal studies. *Statistics in Medicine* 2012;31(23):2745-2756.
3. Barrett JK, Huille R, Parker RMA, et al. Estimating the association between blood pressure variability and cardiovascular disease: An application using the ARIC Study. *Statistics in Medicine* 2019;38(10):1855-1868.
4. Tu YK, Tilling K, Sterne JAC, et al. A critical evaluation of statistical approaches to examining the role of growth trajectories in the developmental origins of health and disease. *International Journal of Epidemiology* 2013;42(5):1327-1339.
5. Sayers A, Heron J, Smith A, et al. Joint modelling compared with two stage methods for analysing longitudinal data and prospective outcomes: A simulation study of childhood growth and BP. *Statistical Methods in Medical Research* 2017;26(1):437-452.
6. Sweeting MJ, Thompson SG. Joint modelling of longitudinal and time-to-event data with application to predicting abdominal aortic aneurysm growth and rupture. *Biometrical Journal* 2011;53(5):750-763.
7. Dolan E, O'Brien E. Is It Daily, Monthly, or Yearly Blood Pressure Variability that Enhances Cardiovascular Risk? *Current Cardiology Reports* 2015;17(11):93.
8. Rothwell PM, Howard SC, Dolan E, et al. Prognostic significance of visit-to-visit variability, maximum systolic blood pressure, and episodic hypertension. *Lancet* 2010;375(9718):895-905.

9. Berenson GS, Wattigney WA, Tracy RE, et al. Atherosclerosis of the aorta and coronary-arteries and cardiovascular risk-factors in persons aged 6 to 30 years and studied at necropsy (the Bogalusa Heart Study). *American Journal of Cardiology* 1992;70(9):851-858.
10. Berenson GS, Bogalusa Heart Study Res G. Childhood risk factors predict adult risk associated with subclinical cardiovascular disease: The Bogalusa Heart Study. *American Journal of Cardiology* 2002;90(10C):3L-7L.
11. Laird WP, Fixler DE. Left-ventricular hypertrophy in adolescents with elevated blood-pressure - assessment by chest roentgenography, electrocardiography, and echocardiography. *Pediatrics* 1981;67(2):255-259.
12. Malcolm DD, Burns TL, Mahoney LT, et al. Factors affecting left-ventricular mass in childhood - the Muscatine Study. *Pediatrics* 1993;92(5):703-709.
13. Daniels SR, Kimball TR, Morrison JA, et al. Effect of lean body-mass, fat mass, blood-pressure, and sexual-maturation on left-ventricular mass in children and adolescents - statistical, biological, and clinical-significance. *Circulation* 1995;92(11):3249-3254.
14. Chen W, Srinivasan SR, Ruan LT, et al. Adult Hypertension Is Associated With Blood Pressure Variability in Childhood in Blacks and Whites: The Bogalusa Heart Study. *American Journal of Hypertension* 2011;24(1):77-82.
15. Whincup PH, Nightingale CM, Owen CG, et al. Ethnic Differences in Carotid Intima-Media Thickness Between UK Children of Black African-Caribbean and White European Origin. *Stroke* 2012;43(7):1747-1754.
16. Rosner B, Cook NR, Evans DA, et al. Reproducibility and predictive values of routine blood-pressure measurements in children - comparison with adult values and implications

- for screening-children for elevated blood-pressure. *American Journal of Epidemiology* 1987;126(6):1115-1125.
17. Boyd A, Golding J, Macleod J, et al. Cohort Profile: The 'Children of the 90s'-the index offspring of the Avon Longitudinal Study of Parents and Children. *International Journal of Epidemiology* 2013;42(1):111-127.
  18. Fraser A, Macdonald-Wallis C, Tilling K, et al. Cohort Profile: The Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. *International Journal of Epidemiology* 2013;42(1):97-110.
  19. Northstone K, Lewcock M, Groom A, et al. The Avon Longitudinal Study of Parents and Children (ALSPAC): an updated on the enrolled sample of index children in 2019. *Wellcome Open Research* 2019 2019;4:51.
  20. Avon Longitudinal Study of Parents and Children (ALSPAC). ALSPAC: Explore data and samples. 2020. <http://www.bristol.ac.uk/alspac/researchers/our-data/>. Accessed February 6 2020.
  21. Timpka S, Hughes AD, Chaturvedi N, et al. Birth weight and cardiac function assessed by echocardiography in adolescence: Avon Longitudinal Study of Parents and Children. *Ultrasound in Obstetrics & Gynecology* 2019;54(2):225-231.
  22. Timpka S, Macdonald-Wallis C, Hughes AD, et al. Hypertensive Disorders of Pregnancy and Offspring Cardiac Structure and Function in Adolescence. *J Am Heart Assoc* 2016;5(11).
  23. Rast P, Hofer SM, Sparks C. Modeling Individual Differences in Within-Person Variation of Negative and Positive Affect in a Mixed Effects Location Scale Model Using BUGS/JAGS. *Multivariate Behavioral Research* 2012;47(2):177-200.

24. Hedeker D, Mermelstein RJ, Demirtas H. An application of a mixed-effects location scale model for analysis of ecological momentary assessment (EMA) data. *Biometrics* 2008;64(2):627-634.
25. Goldstein H, Leckie G, Charlton C, et al. Multilevel growth curve models that incorporate a random coefficient model for the level 1 variance function. *Stat Methods Med Res* 2018;27(11):3478-3491.
26. Staley JR, Bradley J, Silverwood RJ, et al. Associations of Blood Pressure in Pregnancy With Offspring Blood Pressure Trajectories During Childhood and Adolescence: Findings From a Prospective Study. *J Am Heart Assoc* 2015;4(5):12.
27. O'Keeffe LM, Simpkin AJ, Tilling K, et al. Sex-specific trajectories of measures of cardiovascular health during childhood and adolescence: A prospective cohort study. *Atherosclerosis* 2018;278:190-196.
28. Howe LD, Tilling K, Matijasevich A, et al. Linear spline multilevel models for summarising childhood growth trajectories: A guide to their application using examples from five birth cohorts. *Statistical Methods in Medical Research* 2016;25(5):1854-1874.
29. Stan Development Team. Stan statistical software, release 2.19.1. <http://mc-stan.org>: Stan Development Team, 2019.
30. Carpenter B, Gelman A, Hoffman MD, et al. Stan: A Probabilistic Programming Language. *Journal of Statistical Software* 2017;76(1):1-29.
31. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing, 2018.
32. Stan Development Team. RStan: the R interface to Stan, release 2.18.2. <http://mc-stan.org>: Stan Development Team, 2018.

33. UCLA: Statistical Consulting Group. FAQ How do I Interpret a Regression Model When Some Variables are Log Transformed? <https://stats.idre.ucla.edu/other/mult-pkg/faq/general/faqhow-do-i-interpret-a-regression-model-when-some-variables-are-log-transformed/>. Accessed 6th February 2020.
34. Muntner P, Shimbo D, Tonelli M, et al. The Relationship Between Visit-to-Visit Variability in Systolic Blood Pressure and All-Cause Mortality in the General Population Findings From NHANES III, 1988 to 1994. *Hypertension* 2011;57(2):160-166.
35. Vishram JKK, Dahlof B, Devereux RB, et al. Blood pressure variability predicts cardiovascular events independently of traditional cardiovascular risk factors and target organ damage: a LIFE substudy. *Journal of Hypertension* 2015;33(12):2422-2430.
36. Veloudi P, Blizzard CL, Head GA, et al. Blood Pressure Variability and Prediction of Target Organ Damage in Patients With Uncomplicated Hypertension. *American Journal of Hypertension* 2016;29(9):1046-1054.
37. Wade KH, Chiesa ST, Hughes AD, et al. Assessing the Causal Role of Body Mass Index on Cardiovascular Health in Young Adults Mendelian Randomization and Recall-by-Genotype Analyses. *Circulation* 2018;138(20):2187-2201.
38. Hao G, Wang XL, Treiber FA, et al. Blood Pressure Trajectories From Childhood to Young Adulthood Associated With Cardiovascular Risk Results From the 23-Year Longitudinal Georgia Stress and Heart Study. *Hypertension* 2017;69(3):435-442.
39. Toprak A, Wang HW, Chen W, et al. Relation of childhood risk factors to left ventricular hypertrophy (eccentric or concentric) in relatively young adulthood (from the Bogalusa Heart Study). *American Journal of Cardiology* 2008;101(11):1621-1625.

40. Zhang T, Li SX, Bazzano L, et al. Trajectories of Childhood Blood Pressure and Adult Left Ventricular Hypertrophy: The Bogalusa Heart Study. *Hypertension* 2018;72(1):93-101.
41. Hirakawa Y, Arima H, Zoungas S, et al. Impact of Visit-to-Visit Glycemic Variability on the Risks of Macrovascular and Microvascular Events and All-Cause Mortality in Type 2 Diabetes: The ADVANCE Trial. *Diabetes care* 2014.
42. Hausdorff JM, Rios DA, Edelberg HK. Gait variability and fall risk in community-living older adults: A 1-year prospective study. *Archives of Physical Medicine and Rehabilitation* 2001;82(8):1050-1056.
43. Nichols JH, Loeb S, Metter EJ, et al. The relationship between prostate volume and prostate-specific antigen variability: data from the Baltimore Longitudinal Study of Aging and the Johns Hopkins Active Surveillance Program. *BJU international* 2012;109(9):1304-1308.
44. Dzubur E, Ponnada A, Nordgren R, et al. MixWILD: A program for examining the effects of variance and slope of time-varying variables in intensive longitudinal data. *Behavior Research Methods* 2020;52:1403–1427.
45. Murawska M, Rizopoulos D, Lesaffre E. A Two-Stage Joint Model for Nonlinear Longitudinal Response and a Time-to-Event with Application in Transplantation Studies. *Journal of Probability and Statistics* 2012;Article ID 194194.
46. Steele F. Multilevel models for longitudinal data. *Journal of the Royal Statistical Society Series a-Statistics in Society* 2008;171:5-19.
47. Tsiatis AA, Davidian M. Joint modeling of longitudinal and time-to-event data: An overview. *Statistica Sinica* 2004;14(3):809-834.



48. Crowther MJ. merlin: mixed effects regression for linear and non-linear models (user-written Stata program), release 1.2.0. Boston College, MA: Statistical Software Components (SSC), 2019.

ORIGINAL UNEDITED MANUSCRIPT

Table 1. Comparing Characteristics of Children Later Undergoing Echocardiography, and Included in the Model, to those not Included in the Model, From the Avon Longitudinal Study of Parents And Children, SW England, 1990-2011.

Characteristic	Not included in model (N = 12,318)			Included in model (N = 1,986)		
	Valid N	%	Mean (SD)	Valid N	%	Mean (SD)
Sex of child						
Female	5,887	47.8		1,090	54.9	
Male	6,431	52.2		896	45.1	
Mother's education <sup>a,b</sup>						
CSE/none	2,226	21.7		212	11.5	
Vocational	1,048	10.2		148	8.0	
O level	3,564	34.8		618	33.4	
A level	2,227	21.7		498	27.0	
Degree	1,181	11.5		372	20.1	
Mother's BMI <sup>a,b</sup>	9,508		22.9 (3.9)	1,723		22.9 (3.6)
Cigarettes smoked per day by mother <sup>a,b</sup>	9,337		2.5 (5.5)	1,710		1.3 (4.3)
Mother's age at delivery (years) <sup>b</sup>	11,713		27.7 (5.0)	1,895		29.5 (4.6)
Parity <sup>a,b</sup>	10,745		0.9 (1.0)	1,848		0.7 (0.9)
Child's birthweight (kg) <sup>b</sup>	11,564		3.4 (0.5)	1,873		3.4 (0.5)

Abbreviations: BMI: body mass index; SD, standard deviation.

<sup>a</sup> Mother's education: highest educational attainment; Mother's BMI (weight (kg) / height (m)<sup>2</sup>): at 12 weeks gestation; Cigarettes smoked per day by mother: at 32 weeks gestation; Parity: mother's number of previous pregnancies resulting in either a livebirth or a stillbirth.

<sup>b</sup> Sample sizes for these variables do not sum to that for the total sample due to missing values.

Table 2. Estimates From Joint Models With Shared Random Effects, Analysing SBP and log(LVMI) as Outcomes, Presenting the Posterior Parameter Estimates of the Regression Coefficients (Except Where Indicated), Avon Longitudinal Study of Parents And Children, SW England, 1990-2011.

Parameter	A: Age & Sex <sup>a</sup>		B: Adding weight and height <sup>a</sup>		C: Adding maternal characteristics <sup>a</sup>	
	Mean	95% CI	Mean	95% CI	Mean	95% CI
Mean SBP: fixed effects						
Intercept	107.95	107.50, 108.40	107.93	107.52, 108.34	108.23	107.29, 109.16
≤12 years <sup>b</sup>	1.71	1.58, 1.85	-0.55	-0.81, -0.30	-0.56	-0.83, -0.30
>12 years <sup>b</sup>	6.00	5.80, 6.21	3.83	3.51, 4.16	3.83	3.49, 4.18
Female	-0.43	-1.02, 0.19	-0.40	-0.96, 0.16	-0.22	-0.81, 0.36
Female*≤12 years	0.07	-0.11, 0.25	-0.15	-0.33, 0.03	-0.17	-0.36, 0.01
Female*>12 years	-1.99	-2.28, -1.71	-1.26	-1.57, -0.95	-1.25	-1.58, -0.92
log(Weight, kg)			17.01	15.39, 18.64	16.56	14.84, 18.26
Height (cm)			0.04	0.00, 0.09	0.06	0.00, 0.11
Maternal characteristics						
Age at delivery (years)					0.01	-0.06, 0.09
Parity					-0.17	-0.50, 0.17
Highest ed.: vocational <sup>c</sup>					-0.04	-1.36, 1.30

Highest ed.: O-level <sup>c</sup>					0.05	-0.95, 1.03
Highest ed.: A-level <sup>c</sup>					-0.64	-1.70, 0.40
Highest ed.: Degree <sup>c</sup>					-1.39	-2.50, -0.27

Individual-level random effects for SBP

SD(mean BP) <sup>d</sup>	6.12	5.88, 6.37	5.37	5.14, 5.60	5.39	5.16, 5.63
SD(BP slope) <sup>d</sup>	0.66	0.55, 0.75	0.64	0.54, 0.73	0.65	0.54, 0.74
SD(log(BPV)) <sup>d</sup>	0.41	0.29, 0.51	0.39	0.26, 0.50	0.42	0.29, 0.53
Cor(mean BP, BP slope) <sup>d</sup>	-0.09	-0.20, 0.02	-0.11	-0.23, 0.00	-0.09	-0.21, 0.03
Cor(mean BP, log(BPV)) <sup>d</sup>	0.52	0.36, 0.71	0.49	0.31, 0.72	0.50	0.33, 0.71
Cor(BP slope, log(BPV)) <sup>d</sup>	-0.06	-0.34, 0.22	-0.04	-0.33, 0.26	-0.05	-0.34, 0.23

BPV: fixed effects

Intercept	3.22	3.12, 3.32	3.15	3.04, 3.25	3.13	3.02, 3.23
Age (years) <sup>e</sup>	15.61	12.87, 18.39	11.32	5.80, 16.98	10.80	5.05, 16.77
Female <sup>e</sup>	16.84	4.53, 30.40	17.64	4.66, 31.59	20.06	5.99, 35.46
log(Weight, kg) <sup>f</sup>			5.77	2.10, 9.55	5.24	1.42, 9.20
Height (cm) <sup>e</sup>			-0.37	-1.40, 0.70	-0.24	-1.36, 0.89
Residual within-clinic SD for SBP	5.62	5.54, 5.70	5.63	5.54, 5.71	5.61	5.53, 5.70

Mean for log(LVMI): fixed effects

Intercept	3.36	3.35, 3.37	3.33	3.31, 3.34	3.32	3.31, 3.33
-----------	------	------------	------	------------	------	------------

Age (years) <sup>e</sup>	-0.07	-2.85, 2.79	-1.65	-4.2, 0.94	-1.94	-4.61, 0.79
Female <sup>e</sup>	-11.00	-12.61, -9.36	-5.42	-7.08, -3.73	-4.95	-6.72, -3.17
Weight (kg) <sup>e</sup>			0.69	0.62, 0.76	0.70	0.63, 0.78
mean BP <sup>d,e</sup>	0.85	0.44, 1.44	0.57	0.09, 1.49	0.59	0.12, 1.37
BP slope <sup>d,e</sup>	4.38	0.88, 8.61	3.63	0.00, 8.35	3.58	0.01, 7.92
log(BPV) <sup>d,f</sup>	-0.61	-2.05, 0.40	-0.82	-3.08, 0.29	-0.78	-2.54, 0.22
Residual SD for log(LVMI)	0.20	0.19, 0.21	0.19	0.17, 0.19	0.19	0.18, 0.19

Abbreviations: BP: blood pressure; BPV: (within-individual) blood pressure variability; CI: credible interval; LVMI: left ventricular mass indexed to height.

<sup>a</sup> Sample sizes for Models A & B, N individuals: 1,986; N clinic visits: 10,556; N BP measurements: 19,360. Sample size for Model C, N individuals: 1,813; N clinic visits: 9,693; N BP measurements: 17,777.

<sup>b</sup> Linear spline terms, corresponding to change per year for age  $\leq 12$ , and for age  $>12$ , respectively.

<sup>c</sup> Reference category for mother's highest education qualification: CSE / None.

<sup>d</sup> mean BP denotes the individual-level, between-clinic, random effects for the intercept (at mean age 11.3 years) in the mean function for SBP; BP slope denotes the individual-level, between-clinic, random effects for age in the mean function for SBP; log(BPV) denotes the individual-level, between-clinic, random effects on the log-scale of the within-individual variance; i.e. these are the equivalent of  $u_0$ ,  $u_1$  and  $u_2$  in equation 2, respectively.

<sup>e</sup> estimates presented as percentage change in BPV / LVMI (on natural scale) per 1-unit increase in covariate, calculated (using posterior samples) as  $(\exp(\text{coefficient}) - 1) * 100$ ; see Web Table 5 for original coefficient estimates.

<sup>f</sup> estimates presented as percentage change in BPV / LVMI (on natural scale) per 10% increase in covariate (on natural scale), calculated (using posterior samples) as  $((1.10^{\text{coefficient}}) - 1) * 100$ ; see Web Table 5 for original coefficient estimates.

Table 3. Posterior Parameter Estimates From Joint Models in Which Mean BP, BP Slope, and log(BPV) are Included as Exposures for log(LVMI).<sup>a</sup> Avon Longitudinal Study of Parents And Children, SW England, 1990-2011.

Parameter	mean BP only		BP slope only		log(BPV) only		mean BP, BP slope, log(BPV) all included	
	Mean, %	95% CI	Mean, %	95% CI	Mean, %	95% CI	Mean, %	95% CI
mean BP <sup>b,c</sup>	0.19	0.00, 0.39					0.59	0.12, 1.37
BP slope <sup>b,c</sup>			2.40	-0.51, 5.56			3.58	0.01, 7.92
log(BPV) <sup>b,d</sup>					0.21	-0.23, 0.69	-0.78	-2.54, 0.22

Abbreviations: BP: blood pressure; BPV: (within-individual) blood pressure variability; CI: credible interval; LVMI: left ventricular mass indexed to height.

<sup>a</sup> These are selected parameter estimates from these models: see Web Tables 6-7 for full models.

<sup>b</sup> mean BP denotes the individual-level, between-clinic, random effects for the intercept (at mean age 11.3 years) in the mean function for SBP; BP slope denotes the individual-level, between-clinic, random effects for age in the mean function for SBP; log(BPV) denotes the individual-level, between-clinic, random effects for on the log-scale of the within-individual variance; i.e. these are the equivalent of  $u_0$ ,  $u_1$  and  $u_2$  in equation 2, respectively.

<sup>c</sup> estimates presented as percentage change in LVMI (on natural scale) per 1-unit change in covariate, calculated (using posterior samples) as  $(\exp(\text{coefficient}) - 1) * 100$ ; see Web Table 7 for original coefficient estimates.

<sup>d</sup> estimates presented as percentage change in LVMI (on natural scale) per 10% increase in covariate (on natural scale), calculated (using posterior samples) as  $((1.10^{\text{coefficient}}) - 1) * 100$ ; see Web Table 7 for original coefficient estimates.

## FIGURE LEGENDS

Figure 1. Mean clinic systolic blood pressure against mean age for clinics attended by those participants later undergoing echocardiography. Sample sizes (individuals per clinic), for the clinic at the youngest mean age through to the clinic at the oldest mean age, respectively, were  $n = 1,758$ ,  $n = 1,804$ ,  $n = 1,751$ ,  $n = 1,791$ ,  $n = 1,762$ , and  $n = 1,690$ . Avon Longitudinal Study of Parents And Children, SW England, 1990-2011. Bars: 1SD either side of mean.





Figure 2. Observed data, and predictions, for three randomly-selected individuals with amongst the lowest (A-C), and three individuals with amongst the highest (D-F), random effect estimates for BPV. Note that a constant has been added to each axis to preserve anonymity. Avon Longitudinal Study of Parents And Children, SW England, 1990-2011.



ORIGINAL UNEDITED MANUSCRIPT