

# ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: W. Riley, A. C. Jones, K. Singh, D. Browne and A. M. Stuart, *Chem. Commun.*, 2021, DOI: 10.1039/D1CC02587B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## COMMUNICATION

## Accessing novel fluorinated heterocycles with the hypervalent fluoroiodane reagent by solution and mechanochemical synthesis

William Riley,<sup>a</sup> Andrew C. Jones,<sup>b</sup> Kuldip Singh,<sup>a</sup> Duncan L. Browne,<sup>\*c</sup> and Alison M. Stuart<sup>\*a</sup>Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


**A new and efficient strategy for the rapid formation of novel fluorinated tetrahydropyridazines and dihydrooxazines has been developed by fluorocyclisation of  $\beta,\gamma$ -unsaturated hydrazones and oximes with the hypervalent fluoroiodane reagent. Mechanochemical synthesis delivered fluorinated tetrahydropyridazines in similar excellent yields to conventional solution synthesis, whereas fluorinated dihydrooxazines were prepared in much better yields by ball-milling.**

Over 85% of small molecule pharmaceuticals approved by the FDA in 2019 contained a nitrogen-based heterocycle and over a quarter of all FDA approved drug molecules on the market contain at least one fluorine atom.<sup>1a</sup> The prominence of heterocycles comes from their activity *in vivo*, where, specifically their structure provides rigidity to molecular shape and enables effective binding. The pharmaceutical industry is particularly interested in heterocycles with a high  $sp^3$  content<sup>1b</sup> and the C-F bond has a great influence on the pharmacokinetic properties of biologically active molecules.<sup>1c</sup>

Fluorinated *N*-heterocycles can be accessed in a single step by intramolecular fluorocyclisations of unsaturated substrates containing an internal nucleophile. The fluorocyclisations of  $\beta,\gamma$ -unsaturated hydrazones with Selectfluor produced the 5-membered, fluorinated dihydropyrazoles (Scheme 1A).<sup>2</sup> In one report, fluorinated tetrahydropyridazines were formed along with the dihydropyrazoles when R was an aryl group.<sup>2</sup> Similarly, the fluorocyclisations of  $\beta,\gamma$ -unsaturated oximes formed the 5-membered, fluorinated dihydroisoxazoles using either Selectfluor,<sup>3a,3b</sup> or hypervalent iodine reagent, PhI(OPiv)<sub>2</sub>, with HF-pyridine.<sup>3c</sup> The synthesis of the 6-membered, fluorinated dihydrooxazines has not been reported (Scheme 1B).

Since its inception, the hypervalent iodine(III) reagent, fluoroiodane **1** (Scheme 1C), has emerged as an excellent fluorinating reagent with a wide range of applications.<sup>4,5</sup> In many cases, fluoroiodane **1** exhibits differing reactivity to that

observed with fluoraza reagents such as Selectfluor. We recently employed fluoroiodane **1** in the synthesis of fluorinated lactones using either AgBF<sub>4</sub><sup>6a</sup> or hexafluoroisopropanol (HFIP)<sup>6b</sup> to activate the fluoroiodane reagent. Looking to further explore the utility of this reagent in fluorocyclisation reactions, we investigated the synthesis of fluorinated tetrahydropyridazines and dihydrooxazines. We also applied ball-milling to the fluoro-cyclisations since mechanochemical synthesis can minimise solvent requirement and in many cases, can provide enhanced and altered reactivity.<sup>7</sup> Ball-milling has shown promise in bromocyclisations<sup>8</sup> and in fluorination reactions, where Selectfluor can selectively mono- or di-fluorinate 1,3-dicarbonyl compounds,<sup>9</sup> fluorinate pyrazolone rings,<sup>10a</sup> and be used in asymmetric organocatalysis.<sup>10b</sup> Herein, we report the regioselective synthesis of novel fluorinated 6-membered



**Scheme 1** A) Synthesis of fluorinated dihydropyrazoles. B) Synthesis of fluorinated dihydroisoxazoles. C) Synthesis of fluorinated tetrahydropyridazines & dihydrooxazines.

<sup>a</sup> School of Chemistry, University of Leicester, Leicester, LE1 7RH, UK.

<sup>b</sup> Cardiff Catalysis Institute, School of Chemistry, Cardiff University, Cardiff, CF10 3AT, UK.

<sup>c</sup> School of Pharmacy, UCL, London, WC1X 1AX, UK.

E-mail: Alison.Stuart@le.ac.uk, Duncan.Browne@ucl.ac.uk

Electronic Supplementary Information (ESI) available: For experimental procedures, NMR spectra and CCDC 2083051 - 2083056, see DOI: 10.1039/x0xx00000x




heterocycles, tetrahydropyridazines and dihydrooxazines, alongside the use of mechanochemistry to minimise halogenated solvents and to improve the yields in the preparation of fluorinated dihydrooxazines.

We began our studies by investigating the fluorocyclisation of hydrazone **2a**. To our delight, **2a** was fluorocyclised at room temperature in 4 hours using fluoroiodane **1** (1.5 equiv.) with  $\text{AgBF}_4$  (1 equiv.) in  $\text{CH}_2\text{Cl}_2$  to produce the desired fluorinated tetrahydropyridazine **3a** in 91% yield (Table 1, entry 1). The reaction time could be reduced to just 15 minutes with minimal decrease in yield (entry 3). Alternatively, the amount of  $\text{AgBF}_4$  could be decreased to 0.2 equivalents in a 1 hour reaction to form the product in 86% yield (entry 4). On further investigation of solvent and temperature (entries 5-10), a set of metal-free conditions were identified using HFIP as the solvent and activator of fluoroiodane **1** giving **3a** in 93% yield after only 15 minutes (entry 10). Notably, whilst use of acetonitrile as solvent did furnish the desired fluorinated heterocyclic product, Ritter product **4** was also formed in 23% yield suggesting that a tertiary carbocation is a key intermediate in the fluorocyclisation. In entries 7 and 8 where product yield was lower at 40 °C, elimination byproduct **5** was also observed in 10-15% yield.


Using mechanochemistry the unsaturated hydrazones could efficiently undergo fluorocyclisations with fluoroiodane **1** (1.5 equiv.) and just 2 equivalents of HFIP (an 8 fold reduction in the amount of HFIP required, SI Table S3) to provide an alternative, solvent minimised method. For this protocol, a 10 mL stainless steel milling jar was used with a 2.5 g stainless steel milling ball, and the reaction mixture was milled at 30 Hz for 15 minutes.

With optimal conditions for both a solution and ball-milling



Entry	Solvent	$\text{AgBF}_4$ (equiv.)	Temp. (°C)	Time (h)	Yield of <b>3a</b> (%) <sup>b</sup>
1	$\text{CH}_2\text{Cl}_2$	1	RT	4	91
2	$\text{CH}_2\text{Cl}_2$	1	RT	1	91
3	$\text{CH}_2\text{Cl}_2$	1	RT	0.25	90
4	$\text{CH}_2\text{Cl}_2$	0.2	RT	1	86
5	$\text{CH}_2\text{Cl}_2$	-	40	1	30
6	$\text{CH}_3\text{CN}$	-	40	1	31*
7	HFIP	-	40	1	71**
8	HFIP <sup>c</sup>	-	40	1	76**
9	HFIP <sup>c</sup>	-	RT	1	93
10	HFIP <sup>c</sup>	-	RT	0.25	93
11 <sup>d</sup>	HFIP <sup>c</sup>	-	RT	1	0


\* also observed ( $\text{CH}_3\text{CN}$  solvent)    \*\* elimination byproduct    HFIP hexafluoroisopropanol



<sup>a</sup> Reaction conditions: unsaturated hydrazone **2a** (0.71 mmol), fluoroiodane **1** (1.07 mmol), 4 Å molecular sieves (180 mg), solvent (2.5 mL); <sup>b</sup> isolated yield; <sup>c</sup> HFIP (1.2 mL); <sup>d</sup> no fluoroiodane **1**

Table 1 Optimisation of  $\beta,\gamma$ -unsaturated hydrazone fluorocyclisation


method in hand, application to a range of unsaturated hydrazones was explored (Scheme 2). The reaction was tolerant for a series of aryl substituted hydrazones (Scheme 2, **3a-h**) with examples of both electron-donating (**3e**) and electron-withdrawing aryl groups (**3b**). Although synthesis of the thiophene substituted hydrazone proceeded efficiently (**3h**), the aromatic byproduct, 5-methyl-3-thiophen-2-yl pyridazine, was also formed in 16% yield. The fluorocyclisation proved successful on



Scheme 2 Scope of 5-fluorinated tetrahydropyridazine synthesis by solution and mechanochemical synthesis.



alkyl substituted hydrazones (**3i-k**) including benzyl and the more hindered isopropyl hydrazones. Bis-tetrahydropyridazine (**3l**) was prepared by a double fluorocyclisation. In all cases, metal-free protocols could be employed and delivered similar yields in comparison to reactions containing  $\text{AgBF}_4$ . The mechanochemical approach was applied to a small range of substrates (**3b-e**, **3k** and **3l**) and provided comparable excellent yields to those observed in solution demonstrating an efficient, solvent minimised method. Further studies into the activation of fluoroiodane **1** revealed that sub-stoichiometric HFIP could be used in  $\text{CH}_2\text{Cl}_2$  with minimal impact on yield (Scheme 3).



**Scheme 3** Substoichiometric HFIP studies.

Initial results with  $\beta,\gamma$ -unsaturated oximes demonstrated that they could undergo fluorocyclisation to form dihydrooxazines. This class of fluorinated heterocycle has not been reported previously. After optimisation for traditional solution conditions (SI Table S4), dihydrooxazine **7a** could be formed in 46% yield after 15 minutes with 1 equivalent of  $\text{AgBF}_4$  (Scheme 4A). Notably, dihydroisoxazole **8**, intercepted by the benzyl-alcohol backbone of the fluoroiodane reagent, was also isolated as a side product in 15% yield. In an attempt to improve yield and selectivity towards the desired product, attention was turned to mechanochemical techniques commencing with the conditions used for hydrazone fluorocyclisations (Scheme 2). Fluorocyclisation of unsaturated oxime **6b** using fluoroiodane **1** proceeded to give 36% of the desired product with 2 equivalents of HFIP (Table 2, entry 1). Increased loading of HFIP improved the yield of dihydrooxazine **7b**, but also increased the amount of unwanted dihydroisoxazole **9** (Table 2, entry 2). A small amount of oxazine **10**, resulting from elimination, was also observed. Switching to  $\text{AgBF}_4$  showed decreased activity initially (entry 3). However, using  $\text{AgBF}_4$  with 5 equivalents of  $\text{CH}_2\text{Cl}_2$  provided a dramatic improvement in the yield of dihydrooxazine **7b** (80% isolated yield), and reduced the formation of side products **9** and **10** to 1% (entry 5). Further studies on the two additives and reaction time revealed that 2 equivalents of  $\text{AgBF}_4$  and 5 equivalents of  $\text{CH}_2\text{Cl}_2$  in a 1 hour reaction was optimal (entries 6-10). In the absence of HFIP or  $\text{AgBF}_4$ , the reaction showed no activity and returned quantitative amounts of oxime **6b** (entry 11).

A range of  $\beta,\gamma$ -unsaturated oximes was examined and under traditional solution conditions, aryl unsaturated oximes could undergo fluorocyclisations in moderate yield (Scheme 4A, **7a-7d**, 42-53%). However, applying mechanochemical conditions to these substrates greatly improved the yields delivering products **7a-7d** in 71-80% yield. Alkyl oximes could also be tolerated (Scheme 4A, **7e-7f**), but oxime **6e** gave a lower yield as it contained a 1:1 mixture of (*Z*)- and (*E*)-isomers and only the (*Z*)-isomer fluorocyclised. Unsaturated oxime **6g** with a mono-substituted alkene selectively formed dihydroisoxazole **11**, as reported by Wang using  $\text{PhI}(\text{OPiv})_2$  and HF.pyridine.<sup>3c</sup> This

**A - Exploring solution conditions<sup>a</sup>**


**B - Exploring ball-mill conditions<sup>b</sup>**

Entry	Time (h)	Additive 1 (s)	Additive 2 (l)	Yield (%)		
				7b <sup>c</sup>	9 <sup>c</sup>	10 <sup>d</sup>
1	1	-	HFIP (2 equiv.)	36	8	4
2	1	-	HFIP (5 equiv.)	43	10	6
3	1	$\text{AgBF}_4$ (2 equiv.)	-	13	1	4
4	1	$\text{AgBF}_4$ (2 equiv.)	HFIP (5 equiv.)	26	5	2
5	1	$\text{AgBF}_4$ (2 equiv.)	$\text{CH}_2\text{Cl}_2$ (5 equiv.)	84 (80)	1	1
6	1	$\text{AgBF}_4$ (1 equiv.)	$\text{CH}_2\text{Cl}_2$ (5 equiv.)	53	1	1
7	1	$\text{AgBF}_4$ (3 equiv.)	$\text{CH}_2\text{Cl}_2$ (5 equiv.)	71	0	0
8	1	$\text{AgBF}_4$ (2 equiv.)	$\text{CH}_2\text{Cl}_2$ (2 equiv.)	47	1	1
9	0.5	$\text{AgBF}_4$ (2 equiv.)	$\text{CH}_2\text{Cl}_2$ (5 equiv.)	68	0	1
10	2	$\text{AgBF}_4$ (2 equiv.)	$\text{CH}_2\text{Cl}_2$ (5 equiv.)	82	0	0
11	1	-	$\text{CH}_2\text{Cl}_2$ (5 equiv.)	0	0	0

**a)** oxime (0.71 mmol), fluoroiodane **1** (1.07 mmol),  $\text{AgBF}_4$  (0.71 mmol), 4 Å molecular sieves (180 mg),  $\text{CH}_2\text{Cl}_2$  (0.4 mL); **b)** oxime (0.25 mmol), fluoroiodane **1** (0.375 mmol), mixer mill, 30 Hz; **c)** Yield determined by  $^{19}\text{F}$  NMR using benzotrifluoride as internal standard. **d)** Yield determined by  $^1\text{H}$  NMR using mesitylene as internal standard. Isolated yield in parentheses.


**Table 2** Mechanochemical optimisation of  $\beta,\gamma$ -unsaturated oxime fluorocyclisation

result supports our proposed mechanism (Scheme 4B) where there is initial formation of a 5-membered heterocycle **A**, followed by ring expansion via tertiary carbocation **B**, to afford the 6-membered fluorinated heterocycle. However, for mono-substituted unsaturated oximes, formation of a secondary carbocation is not favoured and instead attack of fluoride at the



**Scheme 4** A) Scope of 5-fluorodihydrooxazine synthesis. B) Proposed mechanism.





**Scheme 5** A) Scope of ball-milled intramolecular fluorolactonisation. B) Switching of cyclisation selectivity by choice of fluorinating reagent.

methylene site forms the 5-membered product. Good evidence that tertiary carbocation **B** is a key intermediate was provided by the formation of Ritter product **4** in the fluorocyclisation of **2a** in acetonitrile (Table 1, entry 6). Hara has also reported a similar ring expansion in the fluorocyclisation of 1-decen-5-ol with iodotoluene difluoride.<sup>11,12</sup>

It is intriguing that a rearrangement, driven by carbocation stability, still occurs under solvent minimised conditions in a ball-mill. On a similar thread, we also explored ball-milling techniques in the already established fluoroiodane mediated fluorolactonisation to furnish a range of  $\gamma$ - and  $\delta$ -lactones **13a-f** (Scheme 5A).<sup>6b</sup> These products can only be obtained from a reaction mechanism involving a fluorocyclisation and an aryl migration (Scheme 5B).<sup>6a</sup> It is remarkable that the aryl migration occurred in the ball-mill with a 6 fold reduction in the volume of HFIP. To demonstrate this further, treatment of unsaturated carboxylic acid **12f** with Selectfluor under ball-milling conditions for 1 hour at 30 Hz furnished fluorolactonised product **14** without an aryl migration, and so, the product could be switched by merely changing Selectfluor for fluoroiodane.

In summary, we have developed efficient protocols for accessing two new classes of fluorinated heterocycles using fluoroiodane reagent **1**. Alternative selectivity to that displayed by electrophilic fluorinating reagents (e.g. Selectfluor) is clearly demonstrated and thus, this study shows that fluoroiodane **1** is a complementary reagent for diversifying fluorination methodology. Mechanochemistry has provided a dramatic improvement in the yield and selectivity to fluorinated dihydrooxazines, and in so doing has highlighted that mechanistically complex reactions can be readily achieved in a ball-mill.

## Notes and references

View Article Online  
DOI: 10.1039/D1CC02587B

- a) B. G. de la Torre and F. Albericio, *Molecules*, 2020, **25**, 745.  
b) F. Lovering, J. Bikker and C. Humblet, *J. Med. Chem.*, 2009, **52**, 6752. c) S. Purser, P. R. Moore, S. Swallow and V. Gouverneur, *Chem. Soc. Rev.*, 2008, **37**, 320.
- J. Zhao, M. Jiang, J.-T. Liu, *Org. Chem. Front.*, 2018, **5**, 1155.
- a) Y.-Y. Liu, J. Yang, R.-J. Song and J.-H. Li, *Adv. Synth. Catal.*, 2014, **356**, 2913. b) J. Zhao, M. Jiang and J.-T. Liu, *Adv. Synth. Catal.*, 2017, **359**, 1626. c) W. Kong, Q. Guo, Z. Xu, G. Wang, X. Jiang and R. Wang, *Org. Lett.*, 2015, **17**, 3686.
- a) C. Y. Legault and J. Prévost, *Acta Cryst E*, 2012, **68**, o1238.  
b) V. Matoušek, E. Pietrasiak, R. Schwenk and A. Togni, *J. Org. Chem.*, 2013, **78**, 6763. c) G. C. Geary, E. G. Hope, K. Singh and A. M. Stuart, *Chem. Commun.*, 2013, **49**, 9263.
- a) N. O. Ilchenko, B. O. A. Tasch and K. J. Szabó, *Angew. Chem. Int. Ed.*, 2014, **53**, 12897. b) G. C. Geary, E. G. Hope, K. Singh and A. M. Stuart, *RSC Adv.*, 2015, **5**, 16501. c) W. Yuan and K. J. Szabó, *Angew. Chem. Int. Ed.*, 2015, **54**, 8533. d) A. Ulmer, C. Brunner, A. M. Arnold, A. Pöthig and T. Gulder, *Chem. Eur. J.*, 2016, **22**, 3660. e) W. Yuan, L. Eriksson and K. J. Szabó, *Angew. Chem. Int. Ed.*, 2016, **55**, 8410. f) N. O. Ilchenko, M. Hedberg and K. J. Szabó, *Chem. Sci.*, 2017, **8**, 1056. g) B. Yang, K. Chansaenpak, H. Wu, L. Zhu, M. Wang, Z. Li and H. Lu, *Chem. Commun.*, 2017, **53**, 3497. h) B. Xing, C. Ni and J. Hu, *Angew. Chem. Int. Ed.*, 2018, **57**, 9896. i) A. M. H. Abudken, E. G. Hope, K. Singh and A. M. Stuart, *Org. Biomol. Chem.*, 2020, **18**, 6140. j) S. Yang, S. Shi, Y. Chen, Z. Ding, *J. Org. Chem.*, 2021, **86**, doi: 10.1021/acs.joc.1c00159.
- a) G. C. Geary, E. G. Hope and A. M. Stuart, *Angew. Chem. Int. Ed.*, 2015, **54**, 14911. b) H. K. Minhas, W. Riley, A. M. Stuart, M. Urbonaitė, *Org. Biomol. Chem.*, 2018, **16**, 7170.
- a) J. L. Howard, Q. Cao and D. L. Browne, *Chem. Sci.*, 2018, **9**, 3080. b) G.-W. Wang, *Chem. Soc. Rev.*, 2013, **42**, 7668. c) A. Stolle, T. Szuppa, S. E. S. Leonhardt, B. Ondruschka, *Chem. Soc. Rev.*, 2011, **40**, 2317. d) T. Friščić, C. Mottillo, H. M. Titi, *Angew. Chem. Int. Ed.*, 2020, **59**, 1018. e) J. G. Hernández, C. Bolm, *J. Org. Chem.*, 2017, **82**, 4007. f) A. Porcheddu, E. Colacino, L. De Luca, F. Delogu, *ACS Catal.*, 2020, **10**, 8344. g) J. Andersen, J. Mack, *Green Chem.*, 2018, **20**, 1435. h) S. L. James, C. J. Adams, C. Bolm, D. Braga, P. Collier, T. Friščić, F. Grepioni, K. D. M. Harris, G. Hyett, W. Jones, A. Krebs, J. Mack, L. Maini, A. G. Orpen, I. P. Parkin, W. C. Shearouse, J. W. Steed and D. C. Waddell, *Chem. Soc. Rev.*, 2012, **41**, 413. i) M. Leonardi, M. Villacampa and J. C. Menéndez, *Chem. Sci.*, 2018, **9**, 2042. j) T.-X. Métro, J. Martinez and F. Lamaty, *ACS Sustainable Chem. Eng.*, 2017, **5**, 9599. k) A. C. Jones, W. I. Nicholson, H. R. Smallman and D. L. Browne, *Org. Lett.*, 2020, **22**, 7433. l) O. Eguagie, J. S. Vyle, P. F. Conlon, M. A. Gileay and Y. Liang, *Beilstein J. Org. Chem.*, 2018, **14**, 955. m) D. Tan and T. Friščić, *Eur. J. Org. Chem.*, 2018, **2018**, 18.
- J. Wong and Y.-Y. Yeung, *RSC Adv.*, 2021, **11**, 13564.
- a) J. L. Howard, Y. Sagatov, L. Repousseau, C. Schotten and D. L. Browne, *Green Chem.*, 2017, **19**, 2798. b) J. L. Howard, Y. Sagatov and D. L. Browne, *Tetrahedron*, 2018, **74**, 3118. c) Y. Wang, H. Wang, Y. Jiang, C. Zhang, J. Shao and D. Xu, *Green Chem.*, 2017, **19**, 1674.
- a) J. L. Howard, W. Nicholson, Y. Sagatov and D. L. Browne, *Beilstein J. Org. Chem.*, 2017, **13**, 1950. b) D. Křištofiková, M. Mečiarová, E. Rakovský, R. Šebesta, *ACS Sustainable Chem. Eng.*, 2020, **8**, 14417. c) B. R. Boswell, C. M. F. Mansson, J. M. Cox, Z. Jin, J. A. H. Romaniuk, K. P. Lindquist, L. Cegelski, Y. Xia, S. A. Lopez and N. Z. Burns, *Nat. Chem.*, 2021, **13**, 41.
- M. Sawaguchi, S. Hara, T. Fukuhara and N. Yoneda, *J. Fluorine Chem.*, 2000, **104**, 277.
- For ring expansion of cyclic ethers with *p*-iodotoluene difluoride see: T. Inagaki, Y. Nakamura, M. Sawaguchi, N. Yoneda, S. Ayuba, S. Hara, *Tetrahedron Lett.*, 2003, **44**, 4117.



Journal Name

COMMUNICATION

View Article Online  
DOI: 10.1039/D1CC02587B

Open Access Article. Published on 14 June 2021. Downloaded on 6/14/2021 2:43:31 PM.  
This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.



ChemComm Accepted Manuscript