

## Measurement of $B^0\text{-}\bar{B}^0$ Flavor Oscillations in Hadronic $B^0$ Decays

B. Aubert,<sup>1</sup> D. Boutigny,<sup>1</sup> J.-M. Gaillard,<sup>1</sup> A. Hicheur,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J.P. Lees,<sup>1</sup> P. Robbe,<sup>1</sup> V. Tisserand,<sup>1</sup>  
A. Palano,<sup>2</sup> A. Pompili,<sup>2</sup> G.P. Chen,<sup>3</sup> J.C. Chen,<sup>3</sup> N.D. Qi,<sup>3</sup> G. Rong,<sup>3</sup> P. Wang,<sup>3</sup> Y.S. Zhu,<sup>3</sup> G. Eigen,<sup>4</sup> B. Stugu,<sup>4</sup>  
G.S. Abrams,<sup>5</sup> A.W. Borgland,<sup>5</sup> A.B. Breon,<sup>5</sup> D.N. Brown,<sup>5</sup> J. Button-Shafer,<sup>5</sup> R.N. Cahn,<sup>5</sup> A.R. Clark,<sup>5</sup> M.S. Gill,<sup>5</sup>  
A.V. Gritsan,<sup>5</sup> Y. Groysman,<sup>5</sup> R.G. Jacobsen,<sup>5</sup> R.W. Kadel,<sup>5</sup> J. Kadyk,<sup>5</sup> L.T. Kerth,<sup>5</sup> Yu.G. Kolomensky,<sup>5</sup> J.F. Kral,<sup>5</sup>  
C. LeClerc,<sup>5</sup> M.E. Levi,<sup>5</sup> G. Lynch,<sup>5</sup> P.J. Oddone,<sup>5</sup> M. Pripstein,<sup>5</sup> N.A. Roe,<sup>5</sup> A. Romosan,<sup>5</sup> M.T. Ronan,<sup>5</sup>  
V.G. Shelkov,<sup>5</sup> A.V. Telnov,<sup>5</sup> W.A. Wenzel,<sup>5</sup> T.J. Harrison,<sup>6</sup> C.M. Hawkes,<sup>6</sup> D.J. Knowles,<sup>6</sup> S.W. O'Neale,<sup>6</sup>  
R.C. Penny,<sup>6</sup> A.T. Watson,<sup>6</sup> N.K. Watson,<sup>6</sup> T. Deppermann,<sup>7</sup> K. Goetzen,<sup>7</sup> H. Koch,<sup>7</sup> M. Kunze,<sup>7</sup> B. Lewandowski,<sup>7</sup>  
K. Peters,<sup>7</sup> H. Schmuecker,<sup>7</sup> M. Steinke,<sup>7</sup> N.R. Barlow,<sup>8</sup> W. Bhimji,<sup>8</sup> N. Chevalier,<sup>8</sup> P.J. Clark,<sup>8</sup> W.N. Cottingham,<sup>8</sup>  
B. Foster,<sup>8</sup> C. Mackay,<sup>8</sup> F.F. Wilson,<sup>8</sup> K. Abe,<sup>9</sup> C. Hearty,<sup>9</sup> T.S. Mattison,<sup>9</sup> J.A. McKenna,<sup>9</sup> D. Thiessen,<sup>9</sup> S. Jolly,<sup>10</sup>  
A.K. McKemey,<sup>10</sup> V.E. Blinov,<sup>11</sup> A.D. Bukin,<sup>11</sup> D.A. Bukin,<sup>11</sup> A.R. Buzykaev,<sup>11</sup> V.B. Golubev,<sup>11</sup> V.N. Ivanchenko,<sup>11</sup>  
A.A. Korol,<sup>11</sup> E.A. Kravchenko,<sup>11</sup> A.P. Onuchin,<sup>11</sup> S.I. Serednyakov,<sup>11</sup> Yu.I. Skovpen,<sup>11</sup> V.I. Telnov,<sup>11</sup>  
A.N. Yushkov,<sup>11</sup> D. Best,<sup>12</sup> M. Chao,<sup>12</sup> D. Kirkby,<sup>12</sup> A.J. Lankford,<sup>12</sup> M. Mandelkern,<sup>12</sup> S. McMahon,<sup>12</sup> D.P. Stoker,<sup>12</sup>  
K. Arisaka,<sup>13</sup> C. Buchanan,<sup>13</sup> S. Chun,<sup>13</sup> D.B. MacFarlane,<sup>14</sup> S. Prell,<sup>14</sup> Sh. Rahatlou,<sup>14</sup> G. Raven,<sup>14</sup> V. Sharma,<sup>14</sup>  
C. Campagnari,<sup>15</sup> B. Dahmes,<sup>15</sup> P.A. Hart,<sup>15</sup> N. Kuznetsova,<sup>15</sup> S.L. Levy,<sup>15</sup> O. Long,<sup>15</sup> A. Lu,<sup>15</sup> J.D. Richman,<sup>15</sup>  
W. Verkerke,<sup>15</sup> J. Beringer,<sup>16</sup> A.M. Eisner,<sup>16</sup> M. Grothe,<sup>16</sup> C.A. Heusch,<sup>16</sup> W.S. Lockman,<sup>16</sup> T. Pulliam,<sup>16</sup> T. Schalk,<sup>16</sup>  
R.E. Schmitz,<sup>16</sup> B.A. Schumm,<sup>16</sup> A. Seiden,<sup>16</sup> M. Turri,<sup>16</sup> W. Walkowiak,<sup>16</sup> D.C. Williams,<sup>16</sup> M.G. Wilson,<sup>16</sup>  
E. Chen,<sup>17</sup> G.P. Dubois-Felsmann,<sup>17</sup> A. Dvoretzki,<sup>17</sup> D.G. Hitlin,<sup>17</sup> S. Metzler,<sup>17</sup> J. Oyang,<sup>17</sup> F.C. Porter,<sup>17</sup> A. Ryd,<sup>17</sup>  
A. Samuel,<sup>17</sup> M. Weaver,<sup>17</sup> S. Yang,<sup>17</sup> R.Y. Zhu,<sup>17</sup> S. Devmal,<sup>18</sup> T.L. Geld,<sup>18</sup> S. Jayatilke,<sup>18</sup> G. Mancinelli,<sup>18</sup>  
B.T. Meadows,<sup>18</sup> M.D. Sokoloff,<sup>18</sup> T. Barillari,<sup>19</sup> P. Bloom,<sup>19</sup> M.O. Dima,<sup>19</sup> W.T. Ford,<sup>19</sup> U. Nauenberg,<sup>19</sup>  
A. Olivas,<sup>19</sup> P. Rankin,<sup>19</sup> J. Roy,<sup>19</sup> J.G. Smith,<sup>19</sup> W.C. van Hoek,<sup>19</sup> J. Blouw,<sup>20</sup> J.L. Harton,<sup>20</sup> M. Krishnamurthy,<sup>20</sup>  
A. Soffer,<sup>20</sup> W.H. Toki,<sup>20</sup> R.J. Wilson,<sup>20</sup> J. Zhang,<sup>20</sup> T. Brandt,<sup>21</sup> J. Brose,<sup>21</sup> T. Colberg,<sup>21</sup> M. Dickopp,<sup>21</sup>  
R.S. Dubitzky,<sup>21</sup> A. Hauke,<sup>21</sup> E. Maly,<sup>21</sup> R. Müller-Pfefferkorn,<sup>21</sup> S. Otto,<sup>21</sup> K.R. Schubert,<sup>21</sup> R. Schwierz,<sup>21</sup>  
B. Spaan,<sup>21</sup> L. Wilden,<sup>21</sup> D. Bernard,<sup>22</sup> G.R. Bonneaud,<sup>22</sup> F. Brochard,<sup>22</sup> J. Cohen-Tanugi,<sup>22</sup> S. Ferrag,<sup>22</sup>  
S. T'Jampens,<sup>22</sup> Ch. Thiebaut,<sup>22</sup> G. Vasileiadis,<sup>22</sup> M. Verderi,<sup>22</sup> A. Anjomshoa,<sup>23</sup> R. Bernet,<sup>23</sup> A. Khan,<sup>23</sup>  
D. Lavin,<sup>23</sup> F. Muheim,<sup>23</sup> S. Playfer,<sup>23</sup> J.E. Swain,<sup>23</sup> J. Tinslay,<sup>23</sup> M. Falbo,<sup>24</sup> C. Borean,<sup>25</sup> C. Bozzi,<sup>25</sup> S. Dittongo,<sup>25</sup>  
L. Piemontese,<sup>25</sup> E. Treadwell,<sup>26</sup> F. Anulli,<sup>27,\*</sup> R. Baldini-Ferrolì,<sup>27</sup> A. Calcaterra,<sup>27</sup> R. de Sangro,<sup>27</sup> D. Falciari,<sup>27</sup>  
G. Finocchiaro,<sup>27</sup> P. Patteri,<sup>27</sup> I.M. Peruzzi,<sup>27,\*</sup> M. Piccolo,<sup>27</sup> Y. Xie,<sup>27</sup> A. Zallo,<sup>27</sup> S. Bagnasco,<sup>28</sup> A. Buzzo,<sup>28</sup>  
R. Contri,<sup>28</sup> G. Crosetti,<sup>28</sup> M. Lo Vetere,<sup>28</sup> M. Macri,<sup>28</sup> M.R. Monge,<sup>28</sup> S. Passaggio,<sup>28</sup> F.C. Pastore,<sup>28</sup> C. Patrignani,<sup>28</sup>  
M.G. Pia,<sup>28</sup> E. Robutti,<sup>28</sup> A. Santroni,<sup>28</sup> S. Tosi,<sup>28</sup> M. Morii,<sup>29</sup> R. Bartoldus,<sup>30</sup> R. Hamilton,<sup>30</sup> U. Mallik,<sup>30</sup>  
J. Cochran,<sup>31</sup> H.B. Crawley,<sup>31</sup> P.-A. Fischer,<sup>31</sup> J. Lamsa,<sup>31</sup> W.T. Meyer,<sup>31</sup> E.I. Rosenberg,<sup>31</sup> G. Grosdidier,<sup>32</sup>  
C. Hast,<sup>32</sup> A. Höcker,<sup>32</sup> H.M. Lacker,<sup>32</sup> S. Laplace,<sup>32</sup> V. Lepeltier,<sup>32</sup> A.M. Lutz,<sup>32</sup> S. Plaszczynski,<sup>32</sup> M.H. Schune,<sup>32</sup>  
S. Trincaz-Duvoid,<sup>32</sup> G. Wormser,<sup>32</sup> R.M. Bionta,<sup>33</sup> V. Brigljević,<sup>33</sup> D.J. Lange,<sup>33</sup> M. Mugge,<sup>33</sup> K. van Bibber,<sup>33</sup>  
D.M. Wright,<sup>33</sup> A.J. Bevan,<sup>34</sup> J.R. Fry,<sup>34</sup> E. Gabathuler,<sup>34</sup> R. Gamet,<sup>34</sup> M. George,<sup>34</sup> M. Kay,<sup>34</sup> D.J. Payne,<sup>34</sup>  
R.J. Sloane,<sup>34</sup> C. Touramanis,<sup>34</sup> M.L. Aspinwall,<sup>35</sup> D.A. Bowerman,<sup>35</sup> P.D. Dauncey,<sup>35</sup> U. Egede,<sup>35</sup> I. Eschrich,<sup>35</sup>  
N.J.W. Gunawardane,<sup>35</sup> J.A. Nash,<sup>35</sup> P. Sanders,<sup>35</sup> D. Smith,<sup>35</sup> D.E. Azzopardi,<sup>36</sup> J.J. Back,<sup>36</sup> G. Bellodi,<sup>36</sup>  
P. Dixon,<sup>36</sup> P.F. Harrison,<sup>36</sup> R.J.L. Potter,<sup>36</sup> H.W. Shorthouse,<sup>36</sup> P. Strother,<sup>36</sup> P.B. Vidal,<sup>36</sup> G. Cowan,<sup>37</sup> S. George,<sup>37</sup>  
M.G. Green,<sup>37</sup> A. Kurup,<sup>37</sup> C.E. Marker,<sup>37</sup> P. McGrath,<sup>37</sup> T.R. McMahon,<sup>37</sup> S. Ricciardi,<sup>37</sup> F. Salvatore,<sup>37</sup> G. Vaitsas,<sup>37</sup>  
D. Brown,<sup>38</sup> C.L. Davis,<sup>38</sup> J. Allison,<sup>39</sup> R.J. Barlow,<sup>39</sup> J.T. Boyd,<sup>39</sup> A.C. Forti,<sup>39</sup> J. Fullwood,<sup>39</sup> F. Jackson,<sup>39</sup>  
G.D. Lafferty,<sup>39</sup> N. Savvas,<sup>39</sup> J.H. Weatherall,<sup>39</sup> J.C. Williams,<sup>39</sup> A. Farbin,<sup>40</sup> A. Jawahery,<sup>40</sup> V. Lillard,<sup>40</sup> J. Olsen,<sup>40</sup>  
D.A. Roberts,<sup>40</sup> J.R. Schieck,<sup>40</sup> G. Blaylock,<sup>41</sup> C. Dallapiccola,<sup>41</sup> K.T. Flood,<sup>41</sup> S.S. Hertzbach,<sup>41</sup> R. Kofler,<sup>41</sup>  
V.B. Koptchev,<sup>41</sup> T.B. Moore,<sup>41</sup> H. Staengle,<sup>41</sup> S. Willocq,<sup>41</sup> B. Brau,<sup>42</sup> R. Cowan,<sup>42</sup> G. Sciolla,<sup>42</sup> F. Taylor,<sup>42</sup>  
R.K. Yamamoto,<sup>42</sup> M. Milek,<sup>43</sup> P.M. Patel,<sup>43</sup> F. Palombo,<sup>44</sup> J.M. Bauer,<sup>45</sup> L. Cremaldi,<sup>45</sup> V. Eschenburg,<sup>45</sup>  
R. Kroeger,<sup>45</sup> J. Reidy,<sup>45</sup> D.A. Sanders,<sup>45</sup> D.J. Summers,<sup>45</sup> J.Y. Nief,<sup>46</sup> P. Taras,<sup>46</sup> H. Nicholson,<sup>4</sup> C. Cartaro,<sup>48</sup>  
N. Cavallo,<sup>48,†</sup> G. De Nardo,<sup>48</sup> F. Fabozzi,<sup>48</sup> C. Gatto,<sup>48</sup> L. Lista,<sup>48</sup> P. Paolucci,<sup>48</sup> D. Piccolo,<sup>48</sup> C. Sciacca,<sup>48</sup>  
J.M. LoSecco,<sup>49</sup> J.R.G. Alsmiller,<sup>50</sup> T.A. Gabriel,<sup>50</sup> J. Brau,<sup>51</sup> R. Frey,<sup>51</sup> E. Grauges,<sup>51</sup> M. Iwasaki,<sup>51</sup> N.B. Sinev,<sup>51</sup>  
D. Strom,<sup>51</sup> F. Colecchia,<sup>52</sup> F. Dal Corso,<sup>52</sup> A. Dorigo,<sup>52</sup> F. Galeazzi,<sup>52</sup> M. Margoni,<sup>52</sup> G. Michelon,<sup>52</sup> M. Morandin,<sup>52</sup>  
M. Posocco,<sup>52</sup> M. Rotondo,<sup>52</sup> F. Simonetto,<sup>52</sup> R. Stroili,<sup>52</sup> E. Torassa,<sup>52</sup> C. Voci,<sup>52</sup> M. Benayoun,<sup>53</sup> H. Briand,<sup>53</sup>  
J. Chauveau,<sup>53</sup> P. David,<sup>53</sup> Ch. de Vaissière,<sup>53</sup> L. Del Buono,<sup>53</sup> O. Hamon,<sup>53</sup> F. Le Diberder,<sup>53</sup> Ph. Leruste,<sup>53</sup>

J. Ocariz,<sup>53</sup> L. Roos,<sup>53</sup> J. Stark,<sup>53</sup> P. F. Manfredi,<sup>54</sup> V. Re,<sup>54</sup> V. Speziali,<sup>54</sup> E. D. Frank,<sup>55</sup> L. Gladney,<sup>55</sup> Q. H. Guo,<sup>55</sup> J. Panetta,<sup>55</sup> C. Angelini,<sup>56</sup> G. Batignani,<sup>56</sup> S. Bettarini,<sup>56</sup> M. Bondioli,<sup>56</sup> F. Bucci,<sup>56</sup> E. Campagna,<sup>56</sup> M. Carpinelli,<sup>56</sup> F. Forti,<sup>56</sup> M. A. Giorgi,<sup>56</sup> A. Lusiani,<sup>56</sup> G. Marchiori,<sup>56</sup> F. Martinez-Vidal,<sup>56</sup> M. Morganti,<sup>56</sup> N. Neri,<sup>56</sup> E. Paoloni,<sup>56</sup> M. Rama,<sup>56</sup> G. Rizzo,<sup>56</sup> F. Sandrelli,<sup>56</sup> G. Simi,<sup>56</sup> G. Triggiani,<sup>56</sup> J. Walsh,<sup>56</sup> M. Haire,<sup>57</sup> D. Judd,<sup>57</sup> K. Paick,<sup>57</sup> L. Turnbull,<sup>57</sup> D. E. Wagoner,<sup>57</sup> J. Albert,<sup>58</sup> P. Elmer,<sup>58</sup> C. Lu,<sup>58</sup> V. Miftakov,<sup>58</sup> S. F. Schaffner,<sup>58</sup> A. J. S. Smith,<sup>58</sup> A. Tumanov,<sup>58</sup> E. W. Varnes,<sup>58</sup> G. Cavoto,<sup>59</sup> D. del Re,<sup>59</sup> R. Faccini,<sup>59</sup> F. Ferrarotto,<sup>59</sup> F. Ferroni,<sup>59</sup> E. Lamanna,<sup>59</sup> M. A. Mazzoni,<sup>59</sup> S. Morganti,<sup>59</sup> G. Piredda,<sup>59</sup> F. Safai Tehrani,<sup>59</sup> M. Serra,<sup>59</sup> C. Voena,<sup>59</sup> S. Christ,<sup>60</sup> R. Waldi,<sup>60</sup> T. Adye,<sup>61</sup> N. De Groot,<sup>61</sup> B. Franek,<sup>61</sup> N. I. Geddes,<sup>61</sup> G. P. Gopal,<sup>61</sup> S. M. Xella,<sup>61</sup> R. Aleksan,<sup>62</sup> S. Emery,<sup>62</sup> A. Gaidot,<sup>62</sup> S. F. Ganzhur,<sup>62</sup> P.-F. Giraud,<sup>62</sup> G. Hamel Monchenault,<sup>62</sup> W. Kozanecki,<sup>62</sup> M. Langer,<sup>62</sup> G. W. London,<sup>62</sup> B. Mayer,<sup>62</sup> B. Serfass,<sup>62</sup> G. Vasseur,<sup>62</sup> Ch. Yèche,<sup>62</sup> M. Zito,<sup>62</sup> M. V. Purohit,<sup>63</sup> H. Singh,<sup>63</sup> A. W. Weidemann,<sup>63</sup> F. X. Yumiceva,<sup>63</sup> I. Adam,<sup>64</sup> D. Aston,<sup>64</sup> N. Berger,<sup>64</sup> A. M. Boyarski,<sup>64</sup> G. Calderini,<sup>64</sup> M. R. Convery,<sup>64</sup> D. P. Coupal,<sup>64</sup> D. Dong,<sup>64</sup> J. Dorfan,<sup>64</sup> W. Dunwoodie,<sup>64</sup> R. C. Field,<sup>64</sup> T. Glanzman,<sup>64</sup> S. J. Gowdy,<sup>64</sup> T. Haas,<sup>64</sup> T. Himel,<sup>64</sup> T. Hryn'ova,<sup>64</sup> M. E. Huffer,<sup>64</sup> W. R. Innes,<sup>64</sup> C. P. Jessop,<sup>64</sup> M. H. Kelsey,<sup>64</sup> P. Kim,<sup>64</sup> M. L. Kocian,<sup>64</sup> U. Langenegger,<sup>64</sup> D. W. G. S. Leith,<sup>64</sup> S. Luitz,<sup>64</sup> V. Luth,<sup>64</sup> H. L. Lynch,<sup>64</sup> H. Marsiske,<sup>64</sup> S. Menke,<sup>64</sup> R. Messner,<sup>64</sup> D. R. Muller,<sup>64</sup> C. P. O'Grady,<sup>64</sup> V. E. Ozcan,<sup>64</sup> A. Perazzo,<sup>64</sup> M. Perl,<sup>64</sup> S. Petrak,<sup>64</sup> H. Quinn,<sup>64</sup> B. N. Ratcliff,<sup>64</sup> S. H. Robertson,<sup>64</sup> A. Roodman,<sup>64</sup> A. A. Salnikov,<sup>64</sup> T. Schietinger,<sup>64</sup> R. H. Schindler,<sup>64</sup> J. Schwiening,<sup>64</sup> A. Snyder,<sup>64</sup> A. Soha,<sup>64</sup> S. M. Spanier,<sup>64</sup> J. Stelzer,<sup>64</sup> D. Su,<sup>64</sup> M. K. Sullivan,<sup>64</sup> H. A. Tanaka,<sup>64</sup> J. Va'vra,<sup>64</sup> S. R. Wagner,<sup>64</sup> A. J. R. Weinstein,<sup>64</sup> W. J. Wisniewski,<sup>64</sup> D. H. Wright,<sup>64</sup> C. C. Young,<sup>64</sup> P. R. Burchat,<sup>65</sup> C. H. Cheng,<sup>65</sup> T. I. Meyer,<sup>65</sup> C. Roat,<sup>65</sup> R. Henderson,<sup>66</sup> W. Bugg,<sup>67</sup> H. Cohn,<sup>67</sup> J. M. Izen,<sup>68</sup> I. Kitayama,<sup>68</sup> X. C. Lou,<sup>68</sup> F. Bianchi,<sup>69</sup> M. Bona,<sup>69</sup> D. Gamba,<sup>69</sup> L. Bosisio,<sup>70</sup> G. Della Ricca,<sup>70</sup> L. Lanceri,<sup>70</sup> P. Poropat,<sup>70</sup> G. Vuagnin,<sup>70</sup> R. S. Panvini,<sup>71</sup> C. M. Brown,<sup>72</sup> P. D. Jackson,<sup>72</sup> R. Kowalewski,<sup>72</sup> J. M. Roney,<sup>72</sup> H. R. Band,<sup>73</sup> E. Charles,<sup>73</sup> S. Dasu,<sup>73</sup> A. M. Eichenbaum,<sup>73</sup> H. Hu,<sup>73</sup> J. R. Johnson,<sup>73</sup> R. Liu,<sup>73</sup> F. Di Lodovico,<sup>73</sup> Y. Pan,<sup>73</sup> R. Prepost,<sup>73</sup> I. J. Scott,<sup>73</sup> S. J. Sekula,<sup>73</sup> J. H. von Wimmersperg-Toeller,<sup>73</sup> S. L. Wu,<sup>73</sup> Z. Yu,<sup>73</sup> T. M. B. Kordich,<sup>74</sup> and H. Neal<sup>74</sup>

(The BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

<sup>2</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

<sup>3</sup>Institute of High Energy Physics, Beijing 100039, China

<sup>4</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>5</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720

<sup>6</sup>University of Birmingham, Birmingham B15 2TT, United Kingdom

<sup>7</sup>Ruhr Universität Bochum, Institut für Experimentalphysik, D-44780 Bochum, Germany

<sup>8</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>9</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

<sup>10</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>11</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>12</sup>University of California at Irvine, Irvine, California 92697

<sup>13</sup>University of California at Los Angeles, Los Angeles, California 90024

<sup>14</sup>University of California at San Diego, La Jolla, California 92093

<sup>15</sup>University of California at Santa Barbara, Santa Barbara, California 93106

<sup>16</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064

<sup>17</sup>California Institute of Technology, Pasadena, California 91125

<sup>18</sup>University of Cincinnati, Cincinnati, Ohio 45221

<sup>19</sup>University of Colorado, Boulder, Colorado 80309

<sup>20</sup>Colorado State University, Fort Collins, Colorado 80523

<sup>21</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

<sup>22</sup>Ecole Polytechnique, F-91128 Palaiseau, France

<sup>23</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom

<sup>24</sup>Elon University, Elon University, North Carolina 27244-2010

<sup>25</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy

<sup>26</sup>Florida A&M University, Tallahassee, Florida 32307

<sup>27</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy

<sup>28</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy

<sup>29</sup>Harvard University, Cambridge, Massachusetts 02138

<sup>30</sup>University of Iowa, Iowa City, Iowa 52242

<sup>31</sup>Iowa State University, Ames, Iowa 50011-3160

<sup>32</sup>Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France

<sup>33</sup>Lawrence Livermore National Laboratory, Livermore, California 94550

- <sup>34</sup>University of Liverpool, Liverpool L69 3BX, United Kingdom  
<sup>35</sup>University of London, Imperial College, London SW7 2BW, United Kingdom  
<sup>36</sup>Queen Mary, University of London, E1 4NS, United Kingdom  
<sup>37</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom  
<sup>38</sup>University of Louisville, Louisville, Kentucky 40292  
<sup>39</sup>University of Manchester, Manchester M13 9PL, United Kingdom  
<sup>40</sup>University of Maryland, College Park, Maryland 20742  
<sup>41</sup>University of Massachusetts, Amherst, Massachusetts 01003  
<sup>42</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139  
<sup>43</sup>McGill University, Montréal, Quebec, Canada H3A 2T8  
<sup>44</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy  
<sup>45</sup>University of Mississippi, University, Mississippi 38677  
<sup>46</sup>Université de Montréal, Laboratoire René J.A. Lévêque, Montréal, Quebec, Canada H3C 3J7  
<sup>47</sup>Mount Holyoke College, South Hadley, Massachusetts 01075  
<sup>48</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy  
<sup>49</sup>University of Notre Dame, Notre Dame, Indiana 46556  
<sup>50</sup>Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831  
<sup>51</sup>University of Oregon, Eugene, Oregon 97403  
<sup>52</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy  
<sup>53</sup>Universités Paris VI et VII, Laboratoire de Physique Nucléaire H.E., F-75252 Paris, France  
<sup>54</sup>Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy  
<sup>55</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104  
<sup>56</sup>Università di Pisa, Scuola Normale Superiore and INFN, I-56010 Pisa, Italy  
<sup>57</sup>Prairie View A&M University, Prairie View, Texas 77446  
<sup>58</sup>Princeton University, Princeton, New Jersey 08544  
<sup>59</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy  
<sup>60</sup>Universität Rostock, D-18051 Rostock, Germany  
<sup>61</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom  
<sup>62</sup>DAPNIA, Commissariat à l'Energie Atomique/Saclay, F-91191 Gif-sur-Yvette, France  
<sup>63</sup>University of South Carolina, Columbia, South Carolina 29208  
<sup>64</sup>Stanford Linear Accelerator Center, Stanford, California 94309  
<sup>65</sup>Stanford University, Stanford, California 94305-4060  
<sup>66</sup>TRIUMF, Vancouver, British Columbia, Canada V6T 2A3  
<sup>67</sup>University of Tennessee, Knoxville, Tennessee 37996  
<sup>68</sup>University of Texas at Dallas, Richardson, Texas 75083  
<sup>69</sup>Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy  
<sup>70</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy  
<sup>71</sup>Vanderbilt University, Nashville, Tennessee 37235  
<sup>72</sup>University of Victoria, Victoria, British Columbia, Canada V8W 3P6  
<sup>73</sup>University of Wisconsin, Madison, Wisconsin 53706  
<sup>74</sup>Yale University, New Haven, Connecticut 06511

(Received 31 December 2001; published 20 May 2002)

Flavor oscillations of neutral  $B$  mesons have been studied in  $e^+e^-$  annihilation data collected with the  $BABAR$  detector at center-of-mass energies near the  $\Upsilon(4S)$  resonance. The data sample used for this purpose consists of events in which one  $B^0$  meson is reconstructed in a hadronic decay mode, while the flavor of the recoiling  $B^0$  is determined with a tagging algorithm that exploits the correlation between the flavor of the heavy quark and the charges of its decay products. From the time development of the observed mixed and unmixed final states, we determine the  $B^0-\bar{B}^0$  oscillation frequency  $\Delta m_d$  to be  $0.516 \pm 0.016(\text{stat}) \pm 0.010(\text{syst}) \text{ ps}^{-1}$ .

DOI: 10.1103/PhysRevLett.88.221802

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh, 14.40.Nd

In the standard model,  $B^0-\bar{B}^0$  mixing [1] occurs through second-order weak diagrams involving the exchange of up-type quarks, with the top quark contributing the dominant amplitude. A measurement of the mass difference  $\Delta m_d$  between the mass eigenstates is therefore sensitive to the value of the Cabibbo-Kobayashi-Maskawa matrix element  $V_{td}$  [2,3]. Mixing in the neutral  $B$  meson system was first seen almost fifteen years ago [4], and  $\Delta m_d$

has been measured with both time-integrated and time-dependent techniques [5].

In this Letter, we present a measurement of time-dependent mixing based on a sample of  $32 \times 10^6 B\bar{B}$  pairs recorded at the  $\Upsilon(4S)$  resonance with the  $BABAR$  detector at the Stanford Linear Accelerator Center. This study and a related  $CP$  asymmetry measurement [6] are described in more detail in Ref. [7]. At the PEP-II asymmetric-energy

$e^+e^-$  collider, the  $Y(4S)$  provides a source of  $B^0\bar{B}^0$  pairs moving along the  $e^-$  beam direction ( $z$  axis) with a known Lorentz boost of  $\beta\gamma = 0.55$ , which allows a new technique for determining  $\Delta m_d$  with a high purity sample of fully reconstructed  $B^0$  mesons.

The  $B^0\bar{B}^0$  mixing probability, for a given  $B^0$  lifetime  $\tau$ , is a function of  $\Delta m_d$  and the proper decay-time difference  $\Delta t$  between the two neutral  $B$  mesons produced in a coherent  $P$ -wave state in the  $Y(4S)$  event. The result is a time-dependent probability to observe *unmixed* (+),  $B^0\bar{B}^0$ , or *mixed* (-),  $B^0B^0$  and  $\bar{B}^0\bar{B}^0$ , events:

$$\text{Prob}(B^0\bar{B}^0 \rightarrow B^0\bar{B}^0, B^0B^0 \text{ or } \bar{B}^0\bar{B}^0) \propto e^{-|\Delta t|/\tau} \times (1 \pm \cos\Delta m_d\Delta t). \quad (1)$$

The effect can be measured by reconstructing one  $B$  in a flavor eigenstate, referred to as  $B_{\text{rec}}$ , while the remaining charged particles originating from the decay of the other  $B$ , referred to as  $B_{\text{tag}}$ , are used to identify, or “tag,” its flavor as a  $B^0$  or  $\bar{B}^0$ . The charges of identified leptons and kaons are the primary indicators, although other information in the event can also be used to identify the flavor of  $B_{\text{tag}}$ . The time difference  $\Delta t = t_{\text{rec}} - t_{\text{tag}} \approx \Delta z/\beta\gamma c$  is determined from the separation  $\Delta z$  of the decay vertices for the flavor-eigenstate and tagging  $B$  along the boost direction.

The value of  $\Delta m_d$  is extracted from a tagged flavor-eigenstate  $B^0$  sample with a simultaneous unbinned maximum likelihood fit to the  $\Delta t$  distributions of mixed and unmixed events. There are two principal experimental complications to the probability distribution [Eq. (1)]. First, the tagging algorithm, which classifies events into categories  $i$  depending on the source of the available tagging information, incorrectly identifies the flavor of  $B_{\text{tag}}$  with a probability  $w_i$  with reduction of the observed amplitude for the oscillation by a factor  $(1 - 2w_i)$ . Second, the resolution for  $\Delta t$  is comparable to the oscillation period and must be well understood. The probability density functions for the unmixed and mixed signal events,  $\mathcal{H}_{\pm, \text{sig}}$ , can be expressed as the convolution of the underlying  $\Delta t$  distribution for the  $i$ th tagging category,

$$h_{\pm}(\Delta t; \Delta m_d, w_i) = \frac{e^{-|\Delta t|/\tau}}{4\tau} [1 \pm (1 - 2w_i) \cos\Delta m_d\Delta t],$$

with a  $\Delta t$  resolution function  $\mathcal{R}$  containing parameters  $\hat{a}_j$ . A log-likelihood function is then constructed by summing  $\ln\mathcal{H}_{\pm, \text{sig}}$  over all events within each of the tagging categories. The likelihood is maximized to extract simultaneously the mistag rates  $w_i$ , the resolution function parameters  $\hat{a}_j$ , and the mixing parameter  $\Delta m_d$ .

The *BABAR* detector is described in detail elsewhere [8]. Charged particles are detected and their momenta measured by a combination of a 40-layer drift chamber (DCH) and a five-layer silicon vertex tracker (SVT) embedded in a 1.5-T solenoidal magnetic field. A detector of internally reflected Cherenkov radiation (DIRC) is used

for charged hadron identification. Kaons are identified with a neural network based on the likelihood ratios in the SVT and DCH, derived from  $dE/dx$  measurements, and in the DIRC, calculated by comparing the observed and expected pattern of Cherenkov light for either kaons or pions. A finely segmented CsI(Tl) electromagnetic calorimeter (EMC) is used to detect photons and neutral hadrons, and to identify electrons. Electron candidates are required to have a ratio of EMC energy to track momentum, an EMC cluster shape, DCH  $dE/dx$ , and DIRC Cherenkov angle consistent with expectation. The instrumented flux return (IFR) contains resistive plate chambers for muon and neutral hadron identification. Muon candidates are required to have IFR hits located along the extrapolated DCH track, an IFR penetration length, and an energy deposit in the EMC consistent with the muon hypothesis.

Neutral  $B$  mesons are reconstructed in a sample of multihadron events in the flavor eigenstate decay modes  $D^{(*)-}\pi^+$ ,  $D^{(*)-}\rho^+$ ,  $D^{(*)-}a_1^+$ , and  $J/\psi K^{*0}$ . The decay channels  $K^+\pi^-$ ,  $K^+\pi^-\pi^0$ ,  $K^+\pi^+\pi^-\pi^-$ , and  $K_S^0\pi^+\pi^-$  are used to reconstruct  $\bar{D}^0$  candidates, while the modes  $K^+\pi^-\pi^-$  and  $K_S^0\pi^-$  are used for  $D^-$  candidates. Charged  $D^{*-}$  candidates are formed by combining a  $\bar{D}^0$  with a soft  $\pi^-$ . Finally, the  $B^0$  candidates are formed by combining a  $D^{*-}$  or  $D^-$  candidate with a  $\pi^+$ ,  $\rho^+$  ( $\rho^+ \rightarrow \pi^+\pi^0$ ), or  $a_1^+$  ( $a_1^+ \rightarrow \pi^+\pi^-\pi^+$ ); likewise,  $B^0 \rightarrow J/\psi K^{*0}$  candidates are reconstructed from combinations of  $J/\psi$  candidates, in the decay modes  $e^+e^-$  and  $\mu^+\mu^-$ , with a  $K^{*0}$  ( $K^{*0} \rightarrow K^+\pi^-$ ). The selection and reconstruction of these decays is described in detail in Ref. [9].

Neutral  $B$  candidates are identified by the difference  $\Delta E$  between the energy of the candidate and the beam energy  $\sqrt{s}/2$  in the center-of-mass frame, and the beam-energy substituted mass  $m_{\text{ES}}$ , calculated from  $\sqrt{s}/2$  and the reconstructed momentum of the  $B$  candidate. Candidates are selected by requiring  $m_{\text{ES}} > 5.2 \text{ GeV}/c^2$  and  $\Delta E$  within  $\pm 2.5$  standard deviations of 0 (typically  $|\Delta E| < 40 \text{ MeV}$ ). When multiple candidates in a given event are selected (with probability of about 0.25%), only the one with the smallest  $|\Delta E|$  is retained.

After the daughter tracks of the  $B_{\text{rec}}$  are removed, the remaining tracks are analyzed to determine the flavor of the  $B_{\text{tag}}$ . Events are assigned a lepton tag if they contain an identified lepton with a center-of-mass momentum greater than 1.0 or 1.1  $\text{GeV}/c$  for electrons and muons, respectively, thereby selecting mostly primary leptons. If the sum of charges of all identified kaons is nonzero, the event is assigned a kaon tag. The final two tags involve a multivariable analysis based on a neural network, which is trained to identify primary leptons, kaons, and soft pions, and the momentum and charge of the track with the maximum center-of-mass momentum. Depending on the output of the neural net, events are assigned either an NT1 (more certain) tag, an NT2 (less certain) tag, or are not tagged at all (about 30% of all events) and excluded from the analysis.

Tagging assignments are made mutually exclusive by the hierarchical use of the tags. Events with a lepton tag and no conflicting kaon tag are assigned to the lepton category. If no lepton tag exists, but the event has a kaon tag, it is assigned to the kaon category. Otherwise, events with neural network tags are assigned to corresponding neural network categories.

The decay time difference  $\Delta t$  between  $B$  decays is determined from the measured separation  $\Delta z = z_{\text{rec}} - z_{\text{tag}}$  along the  $z$  axis between the reconstructed  $B_{\text{rec}}(z_{\text{rec}})$  and flavor-tagging decay  $B_{\text{tag}}(z_{\text{tag}})$  vertex. This measured  $\Delta z$  is converted into  $\Delta t$  with the use of the known  $Y(4S)$  boost, including a correction on an event-by-event basis for the direction of the  $B$  mesons with respect to the  $z$  direction in the  $Y(4S)$  frame. The  $\Delta t$  resolution is dominated by the  $z$  resolution of the tag vertex position. After removal of the  $B_{\text{rec}}$  daughters, the  $B_{\text{tag}}$  vertex is formed from all remaining tracks in the event except kaons, which are mostly  $D$  meson decay products. An additional constraint is provided by the calculated  $B_{\text{tag}}$  production point and three momentum, determined from the momentum of the  $B_{\text{rec}}$  candidate, its decay vertex, the average position of the interaction point, and the  $Y(4S)$  boost. Tracks with a large contribution to the  $\chi^2$  are iteratively removed until those remaining ( $\geq 1$ ) have a reasonable fit probability or all tracks are removed. Only events with a reconstructed  $B_{\text{tag}}$  vertex,  $|\Delta t| < 20$  ps and  $\sigma_{\Delta t} < 1.4$  ps are retained (about 84%), where  $\sigma_{\Delta t}$  is the measurement error derived from the vertex fits.

The distribution of  $m_{\text{ES}}$  for the selected candidates is shown in Fig. 1, where the result of a fit with a Gaussian distribution for the signal and an ARGUS function [10] for the background is also displayed. The fitted number of signal events and their purity (for  $m_{\text{ES}} > 5.27$   $\text{GeV}/c^2$ ) are  $6347 \pm 89$  and  $(85.8 \pm 0.5)\%$ , respectively. The sample composition by tagging category is given in Table I.

In the likelihood fit, the  $\Delta t$  resolution function is approximated by a sum of three Gaussian distributions (core, tail, and outlier) with different means and widths,




FIG. 1. Distribution of  $m_{\text{ES}}$  for all  $B^0$  candidates with a flavor tag and a reconstructed tag vertex.

$$\mathcal{R}(\delta_t; \hat{a}) = \sum_{k=1}^2 \frac{f_k}{S_k \sigma_{\Delta t} \sqrt{2\pi}} \exp\left(-\frac{(\delta_t - b_k \sigma_{\Delta t})^2}{2(S_k \sigma_{\Delta t})^2}\right) + \frac{f_3}{\sigma_3 \sqrt{2\pi}} \exp\left(-\frac{\delta_t^2}{2\sigma_3^2}\right),$$

where  $\delta_t = \Delta t - \Delta t_{\text{true}}$ . The sum of the fractions  $f_k$  is constrained to unity. For the core and tail Gaussians, the widths  $\sigma_k = S_k \times \sigma_{\Delta t}$  are the event-by-event measurement errors multiplied by overall scale factors  $S_k$ . The scale factor of the tail Gaussian is fixed to the Monte Carlo value since it is strongly correlated with the other resolution function parameters. The third Gaussian, with a fixed width of  $\sigma_3 = 8$  ps, accounts for outlier events with incorrectly reconstructed vertices (less than 1% of events). A separate core bias coefficient  $b_{1,i}$  is allowed for each tagging category  $i$  to account for small shifts due to inclusion of charm decay products in the tag vertex, while a common bias coefficient  $b_2$  is used for the tail component. These offsets are proportional to  $\sigma_{\Delta t}$  since both the size of the bias and the resolution for  $z_{\text{tag}}$  depend kinematically on the polar angle of the flight direction of the charm daughter. The tail and outlier fractions and the scale factors are assumed to be the same for all decay modes, since the  $z_{\text{tag}}$  measurement dominates the resolution for  $\Delta t$ . This assumption is confirmed by Monte Carlo studies. Separate resolution parameters are used for two different data-reconstruction periods, referred to as Run1 and Run2, which mainly differ in vertex performance and tracking efficiency.

In the presence of backgrounds, which are dominated by continuum  $e^+e^-$  and  $B\bar{B}$  combinatorial sources, additional terms are added to the signal PDF  $\mathcal{H}_{\pm, \text{sig}}$  for various background types,

$$\mathcal{H}_{\pm, i} = f_{i, \text{sig}} \mathcal{H}_{\pm, \text{sig}} + \sum_{j=\text{bkgd}} f_{i, j} \mathcal{B}_{\pm, i, j}(\Delta t; \hat{b}_{\pm, i, j}),$$

where the background PDFs  $\mathcal{B}_{\pm, i, j}$  provide an empirical description for the possible  $\Delta t$  behavior of background events in each tagging category  $i$ . The background  $\Delta t$  types considered are a zero lifetime component and a nonoscillatory component with an empirical nonzero lifetime. We fit for separate resolution function parameters for signal and background to minimize correlations. The fraction of background events for each tagging category and background source is given by  $f_{i, j}$ , while  $\hat{b}_{\pm, i, j}$  are parameters used to characterize each source of background

TABLE I. Signal yields per tagging category, obtained from the  $m_{\text{ES}}$  distributions after all selection requirements. The purity is quoted for  $m_{\text{ES}} > 5.27$   $\text{GeV}/c^2$ .

Category	Tagged	Purity (%)
lepton	$1097 \pm 34$	$96.0 \pm 0.7$
kaon	$3156 \pm 63$	$84.6 \pm 0.7$
NT1	$798 \pm 31$	$88.9 \pm 1.2$
NT2	$1293 \pm 43$	$79.4 \pm 1.3$

by tagging category for mixed and unmixed events. The signal probability  $f_{i,\text{sig}}$  is determined from the measured event  $m_{\text{ES}}$  on the basis of a separate fit to the observed  $m_{\text{ES}}$  distribution in tagging category  $i$ . The sum of signal and background fractions is forced to unity.

Altogether, the likelihood fit involves a total of 44 parameters, including  $\Delta m_d$ , the average mistag fraction and the difference between  $B^0$  and  $\bar{B}^0$  for each tagging category (8), parameters for the signal  $\Delta t$  resolution (16), and parameters for background time dependence (5),  $\Delta t$  resolution (6), and effective dilutions (8). The value of  $\Delta m_d$  was kept hidden throughout the analysis until all analysis details and the systematic errors were finalized, to eliminate possible experimenter's bias.

The results from the likelihood fit to the tagged  $B^0$  sample are summarized in Table II. The probability to obtain a likelihood smaller than that observed is 44%, evaluated with a parametrized Monte Carlo technique. The value of  $\Delta m_d$  given by the fit, prior to final corrections, is  $\Delta m_{d,\text{fit}} = 0.525 \pm 0.016 \text{ ps}^{-1}$ . One method for displaying the result of the full likelihood fit is to use the observed mixing asymmetry,

$$\mathcal{A}_{\text{mix}}(\Delta t) = \frac{N_{\text{unmixed}}(\Delta t) - N_{\text{mixed}}(\Delta t)}{N_{\text{unmixed}}(\Delta t) + N_{\text{mixed}}(\Delta t)}.$$

The unit amplitude for the otherwise pure cosine dependence of  $\mathcal{A}_{\text{mix}}$  is diluted by the mistag probability and the experimental resolution for  $\Delta t$ . The observed  $\Delta t$  distributions of both the mixed and unmixed events, and their asymmetry  $\mathcal{A}_{\text{mix}}$ , are shown along with projections of the likelihood fit result in Fig. 2.

Since the parameters of the  $\Delta t$  resolution for both signal and backgrounds are free parameters in the fit, their contribution to the uncertainty on  $\Delta m_d$  is included as part of the statistical error. Remaining systematic errors arise from the choice of the signal  $\Delta t$  resolution description, its capability to handle outliers and various worst-case SVT misalignment scenarios ( $\pm 0.005 \text{ ps}^{-1}$ ), and by approximations and uncertainties in the  $\Delta z$  to  $\Delta t$  conversion from the absolute  $z$  scale of the detector and PEP-II boost (less than  $\pm 0.002 \text{ ps}^{-1}$ ). Systematic errors due to background include the choice of its  $\Delta t$  distribution and resolution description ( $\pm 0.002 \text{ ps}^{-1}$ ), variation of the sum of background fractions from the separate  $m_{\text{ES}}$  fits, and the uncertainty on the magnitude of the small  $B^+$  component of

TABLE II. Results for  $\Delta m_d$  and a subset of the parameters obtained from the likelihood fit to the  $\Delta t$  distributions.  $\Delta m_d$  includes small corrections described in the text.

Parameter	Fit value	Parameter	Fit value
$\Delta m_d$	$0.516 \pm 0.016$		
$w$ (Lepton)	$0.079 \pm 0.014$	$w$ (NT1)	$0.219 \pm 0.022$
$w$ (Kaon)	$0.166 \pm 0.012$	$w$ (NT2)	$0.344 \pm 0.020$
$S_1$ (Run1)	$1.37 \pm 0.09$	$S_1$ (Run2)	$1.18 \pm 0.11$
$f_2$ (Run1)	$0.014 \pm 0.020$	$f_2$ (Run2)	$0.015 \pm 0.010$
$f_3$ (Run1)	$0.008 \pm 0.004$	$f_3$ (Run2)	$0.000 \pm 0.014$

the signal ( $\pm 0.002 \text{ ps}^{-1}$ ). A correction of  $-0.002 \text{ ps}^{-1}$ , derived from data, is made to account for the small variation of the background composition as a function of  $m_{\text{ES}}$ , which affects the background  $\Delta t$  distribution. The statistical error ( $\pm 0.002 \text{ ps}^{-1}$ ) on this correction is included as a systematic uncertainty. An additional correction of  $-0.007 \text{ ps}^{-1}$  is applied for a bias observed in fully simulated Monte Carlo events. The bias is mainly due to correlations between the mistag rate and the  $\Delta t$  resolution that are not explicitly incorporated into the likelihood function. The systematic error assigned to this correction includes contributions from the statistical precision of the Monte Carlo study ( $\pm 0.003 \text{ ps}^{-1}$ ), model variations due to uncertain branching fractions and lifetimes of the tag-side  $D$  mesons and the assumed fraction of wrong-sign kaons produced in  $B$  decays ( $\pm 0.001 \text{ ps}^{-1}$ ), and variation of the requirement on the maximum allowed value of  $\sigma_{\Delta t}$  ( $\pm 0.003 \text{ ps}^{-1}$ ). Finally, the variation of the fixed  $B^0$  lifetime within the known errors [5] leads to a systematic uncertainty of  $\pm 0.006 \text{ ps}^{-1}$ .




FIG. 2. Distributions of  $\Delta t$  data for the selected (a) unmixed and (b) mixed events [ $m_{\text{ES}}(B_{\text{rec}}) > 5.27 \text{ GeV}/c^2$ ], with projections of the likelihood fit (solid) and the contribution of the background (dashed) overlaid. The time-dependent mixing asymmetry  $\mathcal{A}_{\text{mix}}(|\Delta t|)$  is shown in (c).

In conclusion, a new technique involving the time-difference distribution of a tagged sample of fully reconstructed neutral  $B$  decays has been used to determine the  $B^0\text{-}\bar{B}^0$  mixing frequency  $\Delta m_d$  to be

$$\Delta m_d = 0.516 \pm 0.016(\text{stat}) \pm 0.010(\text{syst}) \text{ ps}^{-1}.$$

This is one of the single most precise measurements available, with an error still dominated by the sample size. The sample consists almost entirely of neutral  $B$  mesons, with excellent control of both flavor tagging for the recoil  $B$  and measurement of the vertex separation between reconstructed and tagged  $B$  meson. The result is consistent with the current world average [5] and a recent  $BABAR$  measurement with a dilepton sample [11]. The analysis shares the same flavor-eigenstate sample, and tagging and vertexing algorithms as used for the determination of  $\sin 2\beta$ , thereby providing an essential validation for the reported  $\sin 2\beta$  result [6].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support  $BABAR$ . The collaborating institutions thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF (Germany), INFN (Italy), NFR (Norway), MIST (Russia),

and PPARC (United Kingdom). Individuals have received support from the A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

---

\*Also with Università di Perugia, Perugia, Italy.

†Also with Università della Basilicata, Potenza, Italy.

- [1] The symbol  $B^0$  refers to the  $B_d$  meson; charge conjugate modes are implied throughout this paper.
- [2] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] For a recent discussion of the relation, see C. Gay, Annu. Rev. Nucl. Part. Sci. **50**, 577 (2000).
- [4] UA1 Collaboration, C. Albajar *et al.*, Phys. Lett. B **186**, 247 (1987); ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **192**, 245 (1987).
- [5] D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- [6]  $BABAR$  Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 151802 (2001).
- [7]  $BABAR$  Collaboration, B. Aubert *et al.*,  $BABAR\text{-PUB-01/03}$ , hep/ex-0201020, Phys. Rev. D (to be published).
- [8]  $BABAR$  Collaboration, B. Aubert *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [9]  $BABAR$  Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 201803 (2001).
- [10] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **241**, 278 (1990).
- [11]  $BABAR$  Collaboration, B. Aubert *et al.*, following Letter, Phys. Rev. Lett. **88**, 221803 (2002).