

Analysis of the Quantum Numbers J^{PC} of the $X(3872)$ Particle

- A. Abulencia,²⁴ J. Adelman,¹³ T. Affolder,¹⁰ T. Akimoto,⁵⁶ M. G. Albrow,¹⁷ D. Ambrose,¹⁷ S. Amerio,⁴⁴ D. Amidei,³⁵
 A. Anastassov,⁵³ K. Anikeev,¹⁷ A. Annovi,¹⁹ J. Antos,¹⁴ M. Aoki,⁵⁶ G. Apollinari,¹⁷ J.-F. Arguin,³⁴ T. Arisawa,⁵⁸
 A. Artikov,¹⁵ W. Ashmanskas,¹⁷ A. Attal,⁸ F. Azfar,⁴³ P. Azzi-Bacchetta,⁴⁴ P. Azzurri,⁴⁷ N. Bacchetta,⁴⁴ W. Badgett,¹⁷
 A. Barbaro-Galtieri,²⁹ V. E. Barnes,⁴⁹ B. A. Barnett,²⁵ S. Baroiant,⁷ V. Bartsch,³¹ G. Bauer,³³ F. Bedeschi,⁴⁷ S. Behari,²⁵
 S. Belforte,⁵⁵ G. Bellettini,⁴⁷ J. Bellinger,⁶⁰ A. Belloni,³³ D. Benjamin,¹⁶ A. Beretvas,¹⁷ J. Beringer,²⁹ T. Berry,³⁰
 A. Bhatti,⁵¹ M. Binkley,¹⁷ D. Bisello,⁴⁴ R. E. Blair,² C. Blocker,⁶ B. Blumenfeld,²⁵ A. Bocci,¹⁶ A. Bodek,⁵⁰ V. Boisvert,⁵⁰
 G. Bolla,⁴⁹ A. Bolshov,³³ D. Bortoletto,⁴⁹ J. Boudreau,⁴⁸ A. Boveia,¹⁰ B. Brau,¹⁰ L. Brigliadori,⁵ C. Bromberg,³⁶
 E. Brubaker,¹³ J. Budagov,¹⁵ H. S. Budd,⁵⁰ S. Budd,²⁴ S. Budroni,⁴⁷ K. Burkett,¹⁷ G. Busetto,⁴⁴ P. Bussey,²¹ K. L. Byrum,²
 S. Cabrera,^{16, o} M. Campanelli,²⁰ M. Campbell,³⁵ F. Canelli,¹⁷ A. Canepa,⁴⁹ S. Carillo,^{18, i} D. Carlsmith,⁶⁰ R. Carosi,⁴⁷
 M. Casarsa,⁵⁵ A. Castro,⁵ P. Catastini,⁴⁷ D. Cauz,⁵⁵ M. Cavalli-Sforza,³ A. Cerri,²⁹ L. Cerrito,^{43, m} S. H. Chang,²⁸
 Y. C. Chen,¹ M. Chertok,⁷ G. Chiarelli,⁴⁷ G. Chlachidze,¹⁵ F. Chlebana,¹⁷ I. Cho,²⁸ K. Cho,²⁸ D. Chokheli,¹⁵ J. P. Chou,²²
 G. Choudalakis,³³ S. H. Chuang,⁶⁰ K. Chung,¹² W. H. Chung,⁶⁰ Y. S. Chung,⁵⁰ M. Ciljak,⁴⁷ C. I. Ciobanu,²⁴ M. A. Ciocci,⁴⁷
 A. Clark,²⁰ D. Clark,⁶ M. Coca,¹⁶ G. Compostella,⁴⁴ M. E. Convery,⁵¹ J. Conway,⁷ B. Cooper,³⁶ K. Copic,³⁵ M. Cordelli,¹⁹
 G. Cortiana,⁴⁴ F. Crescioli,⁴⁷ C. Cuenna Almenar,⁷ J. Cuevas,^{11, l} R. Culbertson,¹⁷ J. C. Cully,³⁵ D. Cyr,⁶⁰ S. DaRonco,⁴⁴
 M. Datta,¹⁷ S. D'Auria,²¹ T. Davies,²¹ M. D'Onofrio,³ D. Dagenhart,⁶ P. de Barbaro,⁵⁰ S. De Cecco,⁵² A. Deisher,²⁹
 G. De Lentdecker,^{50, c} M. Dell'Orso,⁴⁷ F. Delli Paoli,⁴⁴ L. Demortier,⁵¹ J. Deng,¹⁶ M. Deninno,⁵ D. De Pedis,⁵²
 P. F. Derwent,¹⁷ G. P. Di Giovanni,⁴⁵ C. Dionisi,⁵² B. Di Ruza,⁵⁵ J. R. Dittmann,⁴ P. DiTuro,⁵³ C. Dörr,²⁶ S. Donati,⁴⁷
 M. Donega,²⁰ P. Dong,⁸ J. Donini,⁴⁴ T. Dorigo,⁴⁴ S. Dube,⁵³ J. Efron,⁴⁰ R. Erbacher,⁷ D. Errede,²⁴ S. Errede,²⁴ R. Eusebi,¹⁷
 H. C. Fang,²⁹ S. Farrington,³⁰ I. Fedorko,⁴⁷ W. T. Fedorko,¹³ R. G. Feild,⁶¹ M. Feindt,²⁶ J. P. Fernandez,³² R. Field,¹⁸
 G. Flanagan,⁴⁹ A. Foland,²² S. Forrester,⁷ G. W. Foster,¹⁷ M. Franklin,²² J. C. Freeman,²⁹ I. Furic,¹³ M. Gallinaro,⁵¹
 J. Galyardt,¹² J. E. Garcia,⁴⁷ F. Garberson,¹⁰ A. F. Garfinkel,⁴⁹ C. Gay,⁶¹ H. Gerberich,²⁴ D. Gerdes,³⁵ S. Giagu,⁵²
 P. Giannetti,⁴⁷ A. Gibson,²⁹ K. Gibson,⁴⁸ J. L. Gimmell,⁵⁰ C. Ginsburg,¹⁷ N. Giokaris,^{15, a} M. Giordani,⁵⁵ P. Giromini,¹⁹
 M. Giunta,⁴⁷ G. Giurgiu,¹² V. Glagolev,¹⁵ D. Glenzinski,¹⁷ M. Gold,³⁸ N. Goldschmidt,¹⁸ J. Goldstein,^{43, b} A. Golossanov,¹⁷
 G. Gomez,¹¹ G. Gomez-Ceballos,¹¹ M. Goncharov,⁵⁴ O. González,³² I. Gorelov,³⁸ A. T. Goshaw,¹⁶ K. Goulianatos,⁵¹
 A. Gresele,⁴⁴ M. Griffiths,³⁰ S. Grinstein,²² C. Grossi-Pilcher,¹³ R. C. Group,¹⁸ U. Grundler,²⁴ J. Guimaraes da Costa,²²
 Z. Gunay-Unalan,³⁶ C. Haber,²⁹ K. Hahn,³³ S. R. Hahn,¹⁷ E. Halkiadakis,⁵³ A. Hamilton,³⁴ B.-Y. Han,⁵⁰ J. Y. Han,⁵⁰
 R. Handler,⁶⁰ F. Happacher,¹⁹ K. Hara,⁵⁶ M. Hare,⁵⁷ S. Harper,⁴³ R. F. Harr,⁵⁹ R. M. Harris,¹⁷ M. Hartz,⁴⁸
 K. Hatakeyama,⁵¹ J. Hauser,⁸ A. Heijboer,⁴⁶ B. Heinemann,³⁰ J. Heinrich,⁴⁶ C. Henderson,³³ M. Herndon,⁶⁰ J. Heuser,²⁶
 D. Hidas,¹⁶ C. S. Hill,^{10, b} D. Hirschbuehl,²⁶ A. Hocker,¹⁷ A. Holloway,²² S. Hou,¹ M. Houlden,³⁰ S.-C. Hsu,⁹
 B. T. Huffman,⁴³ R. E. Hughes,⁴⁰ U. Husemann,⁶¹ J. Huston,³⁶ J. Incandela,¹⁰ G. Introzzi,⁴⁷ M. Iori,⁵² Y. Ishizawa,⁵⁶
 A. Ivanov,⁷ B. Iyutin,³³ E. James,¹⁷ D. Jang,⁵³ B. Jayatilaka,³⁵ D. Jeans,⁵² H. Jensen,¹⁷ E. J. Jeon,²⁸ S. Jindariani,¹⁸
 M. Jones,⁴⁹ K. K. Joo,²⁸ S. Y. Jun,¹² J. E. Jung,²⁸ T. R. Junk,²⁴ T. Kamon,⁵⁴ P. E. Karchin,⁵⁹ Y. Kato,⁴² Y. Kemp,²⁶
 R. Kephart,¹⁷ U. Kerzel,²⁶ V. Khotilovich,⁵⁴ B. Kilminster,⁴⁰ D. H. Kim,²⁸ H. S. Kim,²⁸ J. E. Kim,²⁸ M. J. Kim,¹²
 S. B. Kim,²⁸ S. H. Kim,⁵⁶ Y. K. Kim,¹³ N. Kimura,⁵⁶ L. Kirsch,⁶ S. Klimentko,¹⁸ M. Klute,³³ B. Knuteson,³³ B. R. Ko,¹⁶
 K. Kondo,⁵⁸ D. J. Kong,²⁸ J. Konigsberg,¹⁸ A. Korytov,¹⁸ A. V. Kotwal,¹⁶ A. Kovalev,⁴⁶ A. C. Kraan,⁴⁶ J. Kraus,²⁴
 I. Kravchenko,³³ M. Kreps,²⁶ J. Kroll,⁴⁶ N. Krumnack,⁴ M. Kruse,¹⁶ V. Krutelyov,¹⁰ T. Kubo,⁵⁶ S. E. Kuhlmann,²
 T. Kuhr,²⁶ Y. Kusakabe,⁵⁸ S. Kwang,¹³ A. T. Laasanen,⁴⁹ S. Lai,³⁴ S. Lami,⁴⁷ S. Lammel,¹⁷ M. Lancaster,³¹ R. L. Lander,⁷
 K. Lannon,⁴⁰ A. Lath,⁵³ G. Latino,⁴⁷ I. Lazzizzera,⁴⁴ T. LeCompte,² J. Lee,⁵⁰ J. Lee,²⁸ Y. J. Lee,²⁸ S. W. Lee,^{54, n}
 R. Lefèvre,³ N. Leonardo,³³ S. Leone,⁴⁷ S. Levy,¹³ J. D. Lewis,¹⁷ C. Lin,⁶¹ C. S. Lin,¹⁷ M. Lindgren,¹⁷ E. Lipeles,⁹
 A. Lister,⁷ D. O. Litvintsev,¹⁷ T. Liu,¹⁷ N. S. Lockyer,⁴⁶ A. Loginov,⁶¹ M. Loreti,⁴⁴ P. Loverre,⁵² R.-S. Lu,¹ D. Lucchesi,⁴⁴
 P. Lujan,²⁹ P. Lukens,¹⁷ G. Lungu,¹⁸ L. Lyons,⁴³ J. Lys,²⁹ R. Lysak,¹⁴ E. Lytken,⁴⁹ P. Mack,²⁶ D. MacQueen,³⁴
 R. Madrak,¹⁷ K. Maeshima,¹⁷ K. Makhoul,³³ T. Maki,²³ P. Maksimovic,²⁵ S. Malde,⁴³ G. Manca,³⁰ F. Margaroli,⁵
 R. Marginean,¹⁷ C. Marino,²⁶ C. P. Marino,²⁴ A. Martin,⁶¹ M. Martin,²¹ V. Martin,^{21, g} M. Martinez,³ T. Maruyama,⁵⁶
 P. Mastrandrea,⁵² T. Masubuchi,⁵⁶ H. Matsunaga,⁵⁶ M. E. Mattson,⁵⁹ R. Mazini,³⁴ P. Mazzanti,⁵ K. S. McFarland,⁵⁰
 P. McIntyre,⁵⁴ R. McNulty,^{30, f} A. Mehta,³⁰ P. Mehtala,²³ S. Menzemer,^{11, h} A. Menzione,⁴⁷ P. Merkel,⁴⁹ C. Mesropian,⁵¹
 A. Messina,³⁶ T. Miao,¹⁷ N. Miladinovic,⁶ J. Miles,³³ R. Miller,³⁶ C. Mills,¹⁰ M. Milnik,²⁶ A. Mitra,¹ G. Mitselmakher,¹⁸
 A. Miyamoto,²⁷ S. Moed,²⁰ N. Moggi,⁵ B. Mohr,⁸ R. Moore,¹⁷ M. Morello,⁴⁷ P. Movilla Fernandez,²⁹ J. Mühlstädt,²⁹
 A. Mukherjee,¹⁷ Th. Muller,²⁶ R. Mumford,²⁵ P. Murat,¹⁷ J. Nachtman,¹⁷ A. Nagano,⁵⁶ J. Naganoma,⁵⁸ I. Nakano,⁴¹

- A. Napier,⁵⁷ V. Necula,¹⁸ C. Neu,⁴⁶ M. S. Neubauer,⁹ J. Nielsen,²⁹ T. Nigmanov,⁴⁸ L. Nodulman,² O. Norniella,³ E. Nurse,³¹ S. H. Oh,¹⁶ Y. D. Oh,²⁸ I. Oksuzian,¹⁸ T. Okusawa,⁴² R. Oldeman,³⁰ R. Orava,²³ K. Osterberg,²³ C. Pagliarone,⁴⁷ E. Palencia,¹¹ V. Papadimitriou,¹⁷ A. A. Paramonov,¹³ B. Parks,⁴⁰ S. Pashapour,³⁴ J. Patrick,¹⁷ G. Paulette,⁵⁵ M. Paulini,¹² C. Paus,³³ D. E. Pellett,⁷ A. Penzo,⁵⁵ T. J. Phillips,¹⁶ G. Piacentino,⁴⁷ J. Piedra,⁴⁵ L. Pinera,¹⁸ K. Pitts,²⁴ C. Plager,⁸ L. Pondrom,⁶⁰ X. Portell,³ O. Poukhov,¹⁵ N. Pounder,⁴³ F. Prakoshyn,¹⁵ A. Pronko,¹⁷ J. Proudfoot,² F. Ptchos,^{19,e} G. Punzi,⁴⁷ J. Pursley,²⁵ J. Rademacker,^{43,b} A. Rahaman,⁴⁸ N. Ranjan,⁴⁹ S. Rappoccio,²² B. Reisert,¹⁷ V. Rekovic,³⁸ P. Renton,⁴³ M. Rescigno,⁵² S. Richter,²⁶ F. Rimondi,⁵ L. Ristori,⁴⁷ A. Robson,²¹ T. Rodrigo,¹¹ E. Rogers,²⁴ S. Rolli,⁵⁷ R. Roser,¹⁷ M. Rossi,⁵⁵ R. Rossin,¹⁸ A. Ruiz,¹¹ J. Russ,¹² V. Rusu,¹³ H. Saarikko,²³ S. Sabik,³⁴ A. Safonov,⁵⁴ W. K. Sakamoto,⁵⁰ G. Salamanna,⁵² O. Saltó,³ D. Saltzberg,⁸ C. Sánchez,³ L. Santi,⁵⁵ S. Sarkar,⁵² L. Sartori,⁴⁷ K. Sato,¹⁷ P. Savard,³⁴ A. Savoy-Navarro,⁴⁵ T. Scheidle,²⁶ P. Schlabach,¹⁷ E. E. Schmidt,¹⁷ M. P. Schmidt,⁶¹ M. Schmitt,³⁹ T. Schwarz,⁷ L. Scodellaro,¹¹ A. L. Scott,¹⁰ A. Scribano,⁴⁷ F. Scuri,⁴⁷ A. Sedov,⁴⁹ S. Seidel,³⁸ Y. Seiya,⁴² A. Semenov,¹⁵ L. Sexton-Kennedy,¹⁷ A. Sfyrla,²⁰ M. D. Shapiro,²⁹ T. Shears,³⁰ P. F. Shepard,⁴⁸ D. Sherman,²² M. Shimojima,^{56,k} M. Shochet,¹³ Y. Shon,⁶⁰ I. Shreyber,³⁷ A. Sidoti,⁴⁷ P. Sinervo,³⁴ A. Sisakyan,¹⁵ J. Sjolin,⁴³ A. J. Slaughter,¹⁷ J. Slaunwhite,⁴⁰ K. Sliwa,⁵⁷ J. R. Smith,⁷ F. D. Snider,¹⁷ R. Snihur,³⁴ M. Soderberg,³⁵ A. Soha,⁷ S. Somalwar,⁵³ V. Sorin,³⁶ J. Spalding,¹⁷ F. Spinella,⁴⁷ T. Spreitzer,³⁴ P. Squillacioti,⁴⁷ M. Stanitzki,⁶¹ A. Staveris-Polykalas,⁴⁷ R. St. Denis,²¹ B. Stelzer,⁸ O. Stelzer-Chilton,⁴³ D. Stentz,³⁹ J. Strologas,³⁸ D. Stuart,¹⁰ J. S. Suh,²⁸ A. Sukhanov,¹⁸ H. Sun,⁵⁷ T. Suzuki,⁵⁶ A. Taffard,²⁴ R. Takashima,⁴¹ Y. Takeuchi,⁵⁶ K. Takikawa,⁵⁶ M. Tanaka,² R. Tanaka,⁴¹ M. Tecchio,³⁵ P. K. Teng,¹ K. Terashi,⁵¹ J. Thom,^{17,d} A. S. Thompson,²¹ E. Thomson,⁴⁶ P. Tipton,⁶¹ V. Tiwari,¹² S. Tkaczyk,¹⁷ D. Toback,⁵⁴ S. Tokar,¹⁴ K. Tollefson,³⁶ T. Tomura,⁵⁶ D. Tonelli,⁴⁷ S. Torre,¹⁹ D. Torretta,¹⁷ S. Tourneur,⁴⁵ W. Trischuk,³⁴ R. Tsuchiya,⁵⁸ S. Tsuno,⁴¹ N. Turini,⁴⁷ F. Ukegawa,⁵⁶ T. Unverhau,²¹ S. Uozumi,⁵⁶ D. Usynin,⁴⁶ S. Vallecorsa,²⁰ N. van Remortel,²³ A. Varganov,³⁵ E. Vataga,³⁸ F. Vázquez,^{18,i} G. Velev,¹⁷ G. Veramendi,²⁴ V. Veszpremi,⁴⁹ R. Vidal,¹⁷ I. Vila,¹¹ R. Vilar,¹¹ T. Vine,³¹ I. Vollrath,³⁴ I. Volobouev,^{29,n} G. Volpi,⁴⁷ F. Würthwein,⁹ P. Wagner,⁵⁴ R. G. Wagner,² R. L. Wagner,¹⁷ J. Wagner,²⁶ W. Wagner,²⁶ R. Wallny,⁸ S. M. Wang,¹ A. Warburton,³⁴ S. Waschke,²¹ D. Waters,³¹ W. C. Wester III,¹⁷ B. Whitehouse,⁵⁷ D. Whiteson,⁴⁶ A. B. Wicklund,² E. Wicklund,¹⁷ G. Williams,³⁴ H. H. Williams,⁴⁶ P. Wilson,¹⁷ B. L. Winer,⁴⁰ P. Wittich,^{17,d} S. Wolbers,¹⁷ C. Wolfe,¹³ T. Wright,³⁵ X. Wu,²⁰ S. M. Wynne,³⁰ A. Yagil,¹⁷ K. Yamamoto,⁴² J. Yamaoka,⁵³ T. Yamashita,⁴¹ C. Yang,⁶¹ U. K. Yang,^{13,j} Y. C. Yang,²⁸ W. M. Yao,²⁹ G. P. Yeh,¹⁷ J. Yoh,¹⁷ K. Yorita,¹³ T. Yoshida,⁴² G. B. Yu,⁵⁰ I. Yu,²⁸ S. S. Yu,¹⁷ J. C. Yun,¹⁷ L. Zanello,⁵² A. Zanetti,⁵⁵ I. Zaw,²² X. Zhang,²⁴ J. Zhou,⁵³ and S. Zucchelli⁵

(CDF Collaboration)

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*²*Argonne National Laboratory, Argonne, Illinois 60439, USA*³*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*⁴*Baylor University, Waco, Texas 76798, USA*⁵*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*⁶*Brandeis University, Waltham, Massachusetts 02254, USA*⁷*University of California, Davis, Davis, California 95616, USA*⁸*University of California, Los Angeles, Los Angeles, California 90024, USA*⁹*University of California, San Diego, La Jolla, California 92093, USA*¹⁰*University of California, Santa Barbara, Santa Barbara, California 93106, USA*¹¹*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*¹²*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*¹³*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*¹⁴*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*¹⁵*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*¹⁶*Duke University, Durham, North Carolina 27708, USA*¹⁷*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*¹⁸*University of Florida, Gainesville, Florida 32611, USA*¹⁹*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*²⁰*University of Geneva, CH-1211 Geneva 4, Switzerland*²¹*Glasgow University, Glasgow G12 8QQ, United Kingdom*²²*Harvard University, Cambridge, Massachusetts 02138, USA*²³*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*

- ²⁴University of Illinois, Urbana, Illinois 61801, USA
²⁵The Johns Hopkins University, Baltimore, Maryland 21218, USA
²⁶Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany
²⁷High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan
²⁸Center for High Energy Physics: Kyungpook National University, Taegu 702-701, Korea; Seoul National University, Seoul 151-742, Korea and SungKyunKwan University, Suwon 440-746, Korea
²⁹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA
³⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom
³¹University College London, London WC1E 6BT, United Kingdom
³²Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
³³Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
³⁴Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8 and University of Toronto, Toronto, Canada M5S 1A7
³⁵University of Michigan, Ann Arbor, Michigan 48109, USA
³⁶Michigan State University, East Lansing, Michigan 48824, USA
³⁷Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
³⁸University of New Mexico, Albuquerque, New Mexico 87131, USA
³⁹Northwestern University, Evanston, Illinois 60208, USA
⁴⁰The Ohio State University, Columbus, Ohio 43210, USA
⁴¹Okayama University, Okayama 700-8530, Japan
⁴²Osaka City University, Osaka 588, Japan
⁴³University of Oxford, Oxford OX1 3RH, United Kingdom
⁴⁴University of Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy
⁴⁵LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
⁴⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁴⁷Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena and Scuola Normale Superiore, I-56127 Pisa, Italy
⁴⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
⁴⁹Purdue University, West Lafayette, Indiana 47907, USA
⁵⁰University of Rochester, Rochester, New York 14627, USA
⁵¹The Rockefeller University, New York, New York 10021, USA
⁵²Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, University of Rome "La Sapienza," I-00185 Roma, Italy
⁵³Rutgers University, Piscataway, New Jersey 08855, USA
⁵⁴Texas A&M University, College Station, Texas 77843, USA
⁵⁵Istituto Nazionale di Fisica Nucleare, University of Trieste/Udine, Italy
⁵⁶University of Tsukuba, Tsukuba, Ibaraki 305, Japan
⁵⁷Tufts University, Medford, Massachusetts 02155, USA
⁵⁸Waseda University, Tokyo 169, Japan
⁵⁹Wayne State University, Detroit, Michigan 48201, USA
⁶⁰University of Wisconsin, Madison, Wisconsin 53706, USA
⁶¹Yale University, New Haven, Connecticut 06520, USA

(Received 21 December 2006; published 28 March 2007)

We present an analysis of angular distributions and correlations of the $X(3872)$ particle in the exclusive decay mode $X(3872) \rightarrow J/\psi\pi^+\pi^-$ with $J/\psi \rightarrow \mu^+\mu^-$. We use 780 pb^{-1} of data from $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ collected with the CDF II detector at the Fermilab Tevatron. We derive constraints on spin, parity, and charge conjugation parity of the $X(3872)$ particle by comparing measured angular distributions of the decay products with predictions for different J^{PC} hypotheses. The assignments $J^{PC} = 1^{++}$ and 2^{-+} are the only ones consistent with the data.

DOI: 10.1103/PhysRevLett.98.132002

PACS numbers: 14.40.Gx, 12.39.Mk, 13.25.Gv

The recent discovery of the $X(3872)$ particle [1,2] has revived general interest in charmonium spectroscopy. The exact nature of this particle is still unknown. Attempts to explain the $X(3872)$ particle as a conventional bound quark-antiquark state have shortcomings, such as deviations from mass predictions or violation of isospin conservation [3]. The close proximity of the $X(3872)$ particle

mass to the $D^0\bar{D}^{*0}$ mass threshold has raised the question whether the $X(3872)$ particle is an exotic form of matter [3]. The determination of the quantum numbers spin J , parity P , and charge conjugation parity C is of vital importance for establishing the nature of the $X(3872)$ particle. The evidence for the decay mode $X(3872) \rightarrow J/\psi\gamma$ [4] and the measurement of the dipion mass distribution [5], which

is in agreement with the decay mode $X(3872) \rightarrow J/\psi \rho^0$, are consistent with a C -even assignment. Reference [6] observes an enhancement in the $D^0 \bar{D}^0 \pi^0$ mass spectrum and concludes that, if assigned to the $X(3872)$ particle, low values for the spin quantum number are favored. Neglecting effects from model uncertainties in the dipion mass spectrum (see [5]), preliminary results from [7] favor $J^{PC} = 1^{++}$. In this Letter we report the angular distributions in the decay $X(3872) \rightarrow J/\psi \pi^+ \pi^-$, $J/\psi \rightarrow \mu^+ \mu^-$, and compare them with predictions for different J^{PC} states. The analysis is independent of any specific model of the internal structure of the $X(3872)$ particle. We consider all allowed states up to spin two and C -odd spin three states.

We use a sample of $p\bar{p}$ collisions at a center-of-mass energy of $\sqrt{s} = 1.96$ TeV with an integrated luminosity of 780 pb^{-1} collected with the CDF II detector at the Fermilab Tevatron. The CDF II detector [8] consists of a magnetic spectrometer surrounded by electromagnetic and hadronic calorimeters and muon detectors. The tracking system is composed of a silicon microstrip detector [9] surrounded by an open-cell drift chamber called the central outer tracker (COT) [10]. We detect muons in planes of multiwire drift chambers [11] in the pseudorapidity range $|\eta| \leq 1.0$. The $J/\psi \rightarrow \mu^+ \mu^-$ decays used in this analysis are recorded using a dimuon trigger, which requires two oppositely charged COT tracks matched to muon chamber track segments with an invariant mass from 2.7 to 4.0 GeV/c^2 .

The basic event selection is described in [2,5], although we do not cut on the dipion mass. Additional criteria are imposed on the number of candidates per event, the transverse momentum p_T of the $X(3872)$ particle candidate ($>6 \text{ GeV}/c$), the p_T of the J/ψ ($>4 \text{ GeV}/c$), and the kinetic energy released in the $X(3872)$ particle decay, $Q = m(J/\psi \pi\pi) - m(J/\psi) - m(\pi\pi)$ ($<100 \text{ MeV}/c^2$), where $m(J/\psi)$ is from [12]. The cuts are chosen to optimize the significance $S/\sqrt{S+B}$ of the observed signal, where S and B are the fitted number of signal and combinatorial background events in a $\pm 1.5\sigma$ window centered on the $X(3872)$ particle mass. The resulting distribution of the invariant $J/\psi \pi^+ \pi^-$ mass is shown in Fig. 1.

To simulate the decays of $X(3872)$ particle states with specific J^{PC} assumptions, we first generate phase-space decays of $X(3872) \rightarrow J/\psi \pi^+ \pi^-$, $J/\psi \rightarrow \mu^+ \mu^-$. Detector effects are included using parameterized efficiencies and acceptances. This sample is weighted according to each specific J^{PC} hypothesis using the corresponding matrix element \mathcal{M}_{tot} described below.

The decay of the narrow $X(3872)$ particle is modeled as the sequential two-body decay chain $X(3872) \rightarrow J/\psi(\pi^+ \pi^-)$, $J/\psi \rightarrow \mu^+ \mu^-$ and the decay of the intermediate $(\pi^+ \pi^-)$ state to $\pi^+ \pi^-$. Assuming low relative angular momentum between the pions and conservation of C parity, the intermediate pion state can be in either a relative S -wave ($(\pi\pi)_S$) or a P -wave (ρ^0) state. \mathcal{M}_{tot} is




FIG. 1 (color online). The $J/\psi \pi^+ \pi^-$ mass spectrum after optimizing the selection cuts, fitted by a double Gaussian function for the $\psi(2S)$ (left), a Gaussian function for the $X(3872)$ particle (right), and a second order polynomial for the combinatorial background.

formed by the product of a matrix element \mathcal{M}_i for each decay and a term $T(m(\pi\pi))$, which describes the mass dependence of the intermediate $(\pi^+ \pi^-)$ system. Because of the very narrow width of the intermediate J/ψ , we can neglect the J/ψ mass dependence.

With fixed helicities the angular dependence of a two-body decay amplitude is given by the Wigner function $D_{\lambda_i, \lambda_{i,1}-\lambda_{i,2}}^{J_i}$ [13,14], where J_i and λ_i are the spin and helicity of the decaying particle, and $\lambda_{i,1}$ and $\lambda_{i,2}$ are the helicities of the child particles in the parent rest frame. The function is multiplied by two Clebsch-Gordan coefficients, coupling the spins of the child particles to their summed spin S_i , and S_i with their relative angular momentum L_i to J_i .

In general, in the $X(3872) \rightarrow J/\psi(\pi^+ \pi^-)$ decay there is more than one combination to form J from L and S in a parity-conserving way. Of the independent amplitudes corresponding to these combinations, only the ones with lowest L , assumed to be dominant, are taken into account. If more than one amplitude remains, mixing parameters are introduced to describe the physical state. Since the virtual photon in the $J/\psi \rightarrow \mu^+ \mu^-$ decay can be treated as transverse, helicity combinations with $\lambda_{\mu^+} - \lambda_{\mu^-} = 0$ are neglected.

The dependence of \mathcal{M}_{tot} on the dipion mass has model ambiguities. Therefore, we do not use the information from $m(\pi\pi)$ to distinguish between different J^{PC} hypotheses. The influence of the $m(\pi\pi)$ model on the angular distributions via acceptance effects is very small. Nevertheless we choose for all J^{PC} hypotheses the same model for the $m(\pi\pi)$ -dependent terms, which agrees with the $m(\pi\pi)$ spectrum measurement. In this way, no hypothesis is rejected due to a wrong $m(\pi\pi)$ model. In detail, we fix $T(m(\pi\pi))$ to a relativistic Breit-Wigner formula with mass and width of a ρ^0 [12]. Following [5], we also fix the momentum dependence of the matrix element of the




FIG. 2 (color online). Definition of the decay angles. The polar angles (θ) are calculated from the parent momenta and the child momenta in the corresponding parent rest frame.

$(\pi^+\pi^-) \rightarrow \pi^+\pi^-$ decay to $k^* \cdot f_1(k^*)$, where k^* is the magnitude of the three-momentum of one of the pions in the $(\pi^+\pi^-)$ rest frame and $f_1(k^*)$ is a Blatt-Weisskopf form factor [15] to counter the divergence for rising k^* . This form factor has the effective size r of the particle as a free parameter which we set to a common choice of $r = 1$ fm.

A weight is formed from the square of the total matrix element \mathcal{M}_{tot} by averaging over all initial state helicities assuming unpolarized $X(3872)$ particle production, incoherently summing over all final state helicities, and coherently summing over all intermediate state helicities.

The decay is described by the decay angles θ_X , $\theta_{J/\psi}$, $\phi_{J/\psi}$, $\theta_{\pi\pi}$, $\phi_{\pi\pi}$, and $\Delta\Phi$, as defined in Fig. 2. For unpolarized $X(3872)$ particle production and because of rotational symmetry, the J^{PC} of the $X(3872)$ particle and the $(\pi^+\pi^-)$ system affect the distribution of only four variables: $m(\pi\pi)$, $\cos(\theta_{J/\psi})$, $\cos(\theta_{\pi\pi})$, and $\Delta\Phi$.

The angular distributions are analyzed with a three-dimensional fit to take into account their correlations. From simulation studies, the optimal binning is determined to be three bins in $|\Delta\Phi - \pi| - \frac{\pi}{2}|$, and two bins in each of $|\cos(\theta_{J/\psi})|$ and $|\cos(\theta_{\pi\pi})|$, where absolute values are used to exploit final state charge symmetry. The invariant $J/\psi\pi^+\pi^-$ mass spectrum is fitted in each of the resulting 12 bins in a mass window of ± 110 MeV/ c^2 around the $X(3872)$ particle position using a binned maximum likelihood fit, where the bin width is 2.5 MeV/ c^2 . The distribution is described by a Gaussian function for the $X(3872)$ particle and a second order polynomial for the background. The position and width of the Gaussian function describing the $X(3872)$ particle are first determined from a fit to the full invariant mass spectrum and are then fixed in the subsequent fits. We compare the fitted yield as a function of the angular variables with the predictions for different J^{PC} assignments by forming a χ^2 based on statistical uncertainties of the measurement. We determine the normalization of the simulated distributions from the measurement so that 11 degrees of freedom remain.

The decay amplitude for the state with $J^{PC} = 1^{-+}$ consists of three LS -terms with the same L value; the $J^{PC} = 2^{-+}$ state has an amplitude with two LS -terms (see Table I). None of the 1^{-+} terms describes the data alone, so we fit for a mixed state by minimizing the χ^2 . For the 2^{-+} state, the amplitude for $S = 1$ is sufficient to describe the data.

TABLE I. Result of the $X(3872)$ particle angular analysis. Listed are the state, the decay mode, the L and S quantum numbers of the $J/\psi-(\pi^+\pi^-)$ system, the χ^2 with 11 degrees of freedom and the χ^2 probability.

J^{PC}	decay	LS	χ^2 (11 d.o.f.)	χ^2 prob.
1^{++}	$J/\psi\rho^0$	01	13.2	0.28
2^{-+}	$J/\psi\rho^0$	11,12	13.6	0.26
1^{--}	$J/\psi(\pi\pi)_S$	01	35.1	2.4×10^{-4}
2^{+-}	$J/\psi(\pi\pi)_S$	11	38.9	5.5×10^{-5}
1^{+-}	$J/\psi(\pi\pi)_S$	11	39.8	3.8×10^{-5}
2^{--}	$J/\psi(\pi\pi)_S$	21	39.8	3.8×10^{-5}
3^{+-}	$J/\psi(\pi\pi)_S$	31	39.8	3.8×10^{-5}
3^{--}	$J/\psi(\pi\pi)_S$	21	41.0	2.4×10^{-5}
2^{++}	$J/\psi\rho^0$	02	43.0	1.1×10^{-5}
1^{-+}	$J/\psi\rho^0$	10,11,12	45.4	4.1×10^{-6}
0^{+-}	$J/\psi\rho^0$	11	104	3.5×10^{-17}
0^{+-}	$J/\psi(\pi\pi)_S$	11	129	$\leq 1 \times 10^{-20}$
0^{++}	$J/\psi\rho^0$	00	163	$\leq 1 \times 10^{-20}$

Table I shows the χ^2 for each J^{PC} assignment. We find that only the assignments $J^{PC} = 1^{++}$ and 2^{-+} are able to describe the data. All other states are rejected by more than 3 standard deviations (χ^2 prob. $\leq 2.7 \times 10^{-3}$). Figure 3 shows the measurement and the expected distribution for four of the assignments.

An important cross-check of the analysis is to verify whether the correct result is obtained for the $\psi(2S)$, with




FIG. 3 (color online). Measured 3D angular distribution with acceptance corrected predictions for $J^{PC} = 0^{++}$ (solid line), 1^{++} (dotted line), 2^{-+} (dashed line), and 1^{--} (dash-dotted line). The plot is divided into 2×2 regions, corresponding to intervals of $|\cos(\theta_{J/\psi})|$ and $|\cos(\theta_{\pi\pi})|$. Each region shows the distribution of $||\Delta\Phi - \pi| - \frac{\pi}{2}|$ in 3 bins. The bin contents have been scaled to the same bin volume.




FIG. 4. Total χ^2 for different analysis variations on the y-axis, explained in the text. Vertical bars are added for visual guidance. The χ^2 values of the spin 0 states are all above 100. The 2^{--} and 3^{+-} states have the same angular distribution as the 1^{+-} state.

known quantum numbers $J^{PC} = 1^{--}$, which decays into the same exclusive final state as the $X(3872)$ particle. For the 1^{--} assignment, the fit probability is 1.5%. Using the $\psi(2S)$ model of Novikov and Shifman [16], which includes a small D -wave admixture in the description of the $(\pi^+ \pi^-)$ system, the fit probability is 17.9%. The sensitivity to such a small admixture is only present in the high statistics $\psi(2S)$ sample. The next best model $J^{PC} = 2^{++}$ has a fit probability of 0.58%, and all other hypotheses that were tested yielded fit probabilities smaller than 2×10^{-6} .

We vary several inputs to the fitting procedure and the model of the $X(3872)$ particle to investigate the stability of the χ^2 . Figure 4 shows the resulting χ^2 values for the different J^{PC} hypotheses for the variations investigated. The default analysis is shown as variation (1). The following effects are considered: (2)/(3) decrease/increase the fit window by $20 \text{ MeV}/c^2$, (4)/(5) decrease/increase the bin width to $2.0/2.86 \text{ MeV}/c^2$, (6)/(7) vary fixed $X(3872)$ particle mass by $\pm 1\sigma$, (8)/(9) vary fixed $X(3872)$ particle width by $\pm 1\sigma$.

To evaluate the contribution to the systematic uncertainty from our choice of the $m(\pi\pi)$ spectrum, the following variations are considered: (10) fix form-factor r to 0.001 fm , (11) fix form-factor r to 100.0 fm , (12) use simple phase space for $m(\pi\pi)$.

Finally, systematic uncertainty due to details concerning the simulation has been considered by varying distributions for (13) p_T and (14) η of the $X(3872)$ particle, switching off (15) a p_T dependent efficiency correction for the pions, (16) a ϕ dependent correction of the COT, and (17) an

effective η correction used to model the position of the generated primary vertex. All variations are consistent with 1^{++} and 2^{+-} being the only likely assignments.

A conventional explanation for the $X(3872)$ resonance is a charmonium ($c\bar{c}$) state. In this picture, the state with $J^{PC} = 1^{++}$ could be identified with the χ'_{c1} and the assignment $J^{PC} = 2^{-+}$ with the η_{c2} . An exotic interpretation is that the $X(3872)$ particle is a molecular state or that a significant four-quark interaction contributes to the wavefunction [17]. The result of this analysis is compatible with the models of a molecular state developed by Tornqvist [18] and Swanson [19], who predict the quantum numbers $J^{PC} = 1^{++}$ for a bound $D\bar{D}^*$ state.

In summary, a spin-parity analysis of the $X(3872)$ particle in the final state $\mu^+ \mu^- \pi^+ \pi^-$ has been performed. The method of helicity amplitudes has been used to analyze $X(3872) \rightarrow J/\psi(\pi\pi)_S$ and $X(3872) \rightarrow J/\psi\rho^0$ transitions. Using a χ^2 approach to compare expected angular distributions with measured distributions, it is found that only the C -even assignments $J^{PC} = 1^{++}$ and 2^{-+} , both decaying via $J/\psi\rho^0$, describe the data. All other states are excluded at 99.7% confidence level.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, UK; the Institut National de Physique Nucléaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Comisión Interministerial de Ciencia y Tecnología, Spain; the European Community's Human Potential Programme under Contract No. HPRN-CT-2002-00292; and the Academy of Finland.

^aVisiting scientist from University of Athens.

^bVisiting scientists from University of Bristol.

^cVisiting scientist from University Libre de Bruxelles.

^dVisiting scientists from Cornell University.

^eVisiting scientist from University of Cyprus.

^fVisiting scientist from University of Dublin.

^gVisiting scientist from University of Edinburgh.

^hVisiting scientist from University of Heidelberg.

ⁱVisiting scientists from Universidad Iberoamericana.

^jVisiting scientist from University of Manchester.

^kVisiting scientist from Nagasaki Institute of Applied Science.

^lVisiting scientist from University de Oviedo.

^mVisiting scientist from University of London, Queen Mary and Westfield College.

ⁿVisiting scientists from Texas Tech University.

^oVisiting scientist from IFIC (CSIC-Universitat de Valencia).

[1] S. K. Choi *et al.* (Belle Collaboration), Phys. Rev. Lett. **91**, 262001 (2003).

[2] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **93**, 072001 (2004).

[3] T. Barnes and S. Godfrey, Phys. Rev. D **69**, 054008 (2004).

[4] K. Abe *et al.* (Belle Collaboration), hep-ex/0505037; B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **74**, 071101 (2006).

[5] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **96**, 102002 (2006).

[6] G. Gokhroo *et al.* (Belle Collaboration), Phys. Rev. Lett. **97**, 162002 (2006).

[7] K. Abe *et al.* (Belle Collaboration), hep-ex/0505038.

[8] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).

- [9] C. S. Hill, Nucl. Instrum. Methods Phys. Res., Sect. A **530**, 1 (2004); A. Affolder *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **453**, 84 (2000); A. Sill, Nucl. Instrum. Methods Phys. Res., Sect. A **447**, 1 (2000).
- [10] T. Affolder *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **526**, 249 (2004).
- [11] G. Ascoli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **268**, 33 (1988).
- [12] S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [13] S. U. Chung, Phys. Rev. D **57**, 431 (1998).
- [14] D. A. Varshalovich, A. N. Moskalev, and V. Khersonsky, *Quantum Theory of Angular Momentum* (World Scientific, Singapore, 1988).
- [15] J. M. Blatt and V. Weisskopf, *Theoretical Nuclear Physics* (John Wiley, New York, 1952).
- [16] V. A. Novikov and M. A. Shifman, Z. Phys. C **8**, 43 (1981).
- [17] A. De Rujula, H. Georgi, and S. L. Glashow, Phys. Rev. Lett. **38**, 317 (1977); M. B. Voloshin and L. B. Okun, JETP Lett. **23**, 333 (1976).
- [18] N. A. Tornqvist, Phys. Lett. B **590**, 209 (2004).
- [19] E. S. Swanson, Phys. Lett. B **588**, 189 (2004).