

Correction to “Relative importance of meridional and zonal sea surface temperature gradients for the onset of the ice ages and Pliocene-Pleistocene climate evolution”

Christopher M. Brierley and Alexey V. Fedorov

Received 2 September 2010; published 19 November 2010.

Citation: Brierley, C. M., and A. V. Fedorov (2010), Correction to “Relative importance of meridional and zonal sea surface temperature gradients for the onset of the ice ages and Pliocene-Pleistocene climate evolution,” *Paleoceanography*, 25, PA4214, doi:10.1029/2010PA002052.

[1] In the paper “Relative importance of meridional and zonal sea surface temperature gradients for the onset of the ice ages and Pliocene-Pleistocene climate evolution” by Christopher M. Brierley and Alexey V. Fedorov (*Paleoceanography*, 25, PA2214, doi:10.1029/2009PA001809, 2010), there was a slight inaccuracy in Figure 9. The corrected image is included here.




Figure 9. The impact of increasing meridional and zonal SST gradients on the $P - E$ field over Africa and surrounding regions. (top) Impact of increasing the meridional SST gradient on (right) boreal summer and (left) boreal winter and (bottom) consequences of establishing modern zonal SST gradients on the same seasons. The colors show changes in $P - E$ over land (in mm d^{-1}). The lines show changes in $P - E$ with a contour interval of 2 mm d^{-1} over the ocean; the bold line corresponds to 0, and negative contours are dashed.